

**TABLE 2-3. Estimated time (T) to rebuild relative to alternative 2009-2010 OYs for depleted West Coast groundfish species.**

Species	Ttarget in the FMP	OY Alternative	Median Time to Rebuild	2009-10 OY (mt)	T @ F=0	Current Tmax	Re-estimated Tmax
Bocaccio (S of 40°10')	2026	1	2020	0	2020	2032	2033
		2	2022	223			
		3	2023	295			
		4	2026	475			
Canary	2063	1	2019	0	2019	2071	2035
		2	2020	55			
		3	2020	95			
		4	2021	155			
		5	2023	327			
Cowcod	2039	1	2061	0	2061	2074	2098
		2	2065	2			
		3	2072	4			
		4	2080	7			
Darkblotched	2011	1	2018	0	2018	2033	2040
		2	2022	162			
		3	2030	303			
		4	2031	321			
		5	2040	388			
POP	2017	1	2010	0	2010	2043	2042
		2	2010	134			
		3	2011	169			
		4	2011	195			
		5	2017	982			
Widow	2015	1	2009	0	2009	2027	2023
		2	2009	367			
		3	2009	516			
		4	2009	4,195			
Yelloweye	2084	1	2049	0	2049	2096	2090
		2	2082	13			
		3	2082	Ramp-down			
		4	2090	15			










Figure 2-2. Average 2009-2010 optimum yields (mt) vs. predicted rebuilding times for overfished species.