

GROUND FISH MANAGEMENT TEAM REPORT ON MANAGEMENT MEASURES –  
 PART 2

**Table 1.** Tentatively adopted 2011-2012 Estimates of research, tribal, and incidental open access. Set asides for EFPs.

<b>Category</b>	<b>Bocaccio South 40'10</b>	<b>Canary</b>	<b>Cowcod South 40'10</b>	<b>Dark- blotched</b>	<b>POP</b>	<b>Widow</b>	<b>Yelloweye</b>	<b>Petrals</b>
Tribal Whiting Trawl		4.3		0.1	7.2	5	0	
Tribal Mid-water Trawl		3.6			0	40	0	
Tribal Bottom Trawl		0.8			3.7	0	0	45.4
Tribal Troll		0.5			0		0	
Tribal Fixed Gear		0.3			0	0	2.3	
Open Access Incidental	0.7	2	0	15	0.1	3.3	0.2	1
Research	1.7	7.2	0.1	2.1	1.8	1.6	3.3	17
EFP	11	1.3	0.2	1.5	0.1	11	0.4	2
<b>Subtotal</b>	<b>13.4</b>	<b>20</b>	<b>0.3</b>	<b>18.7</b>	<b>12.9</b>	<b>60.9</b>	<b>6.2</b>	<b>65.4</b>

**Table 2.** June 2010 Scorecard after inseason action taken at this meeting.

**Projected mortality impacts (mt) of overfished groundfish species for 2010 updated based on June inseason action.**

Fishery	Bocaccio b/	Canary	Cowcod	Dkbl	POP	Widow	Yelloweye
<b>Limited Entry Trawl - Non-whiting</b>	<b>8.0</b>	<b>12.7</b>	<b>0.3</b>	<b>191.4</b>	<b>93.8</b>	<b>15.5</b>	<b>0.3</b>
<b>Limited Entry Trawl - Whiting</b>							
At-sea whiting motherships a/		3.3		6.0	0.5	67.0	0.0
At-sea whiting cat-proc a/		4.8		8.5	0.5	95.0	0.0
Shoreside whiting a/		5.9		10.5	4.7	117.0	0.0
Tribal whiting		4.3		0.0	7.2	5.0	0.0
<b>Tribal</b>							
Midwater Trawl		3.6		0.0	0.0	40.0	0.0
Bottom Trawl		0.8		0.0	3.7	0.0	0.0
Troll		0.5		0.0	0.0		0.0
Fixed gear		0.3		0.0	0.0	0.0	2.3
<b>Fixed Gear Sablefish</b>	0.0	2.5	0.0	4.5	0.4	0.0	0.9
<b>Fixed Gear Nearshore</b>	0.3	3.6	0.0	0.0	0.0	0.3	1.1
<b>Fixed Gear Other</b>	5.0	0.0	0.0	9.0	0.0	0.7	0.0
<b>Open Access: Incidental Groundfish</b>	0.8	1.7	0.0	15.0	0.0	<b>3.3</b>	0.3
<b>Recreational Groundfish e/</b>							
WA		20.9					<b>4.9</b>
OR						1.0	
CA	67.3	22.9	0.3			6.2	<b>2.7</b>
<b>EFPs</b>	11.0	1.3	0.2	1.5	0.1	11.0	<b>0.2</b>
<b>Research: Includes NMFS trawl shelf-slope surveys, the IPHC halibut survey, and expected impacts from SRPs and LOAs.</b>							
	2.0	4.5	0.2	2.0	2.0	5.7	<b>1.3</b>
<b>TOTAL</b>	94.4	93.5	1.0	248.4	112.9	367.7	14.0
<b>2010 OY f/</b>	288	105	4.0	<b>330</b>	200	509	<b>14</b>
<b>Difference</b>	193.6	11.5	3.0	81.6	87.1	141.3	0.0
<b>Percent of OY</b>	32.8%	89.1%	25.0%	75.3%	56.5%	72.2%	100.0%
Key		= either not applicable; trace amount (<0.01 mt); or not reported in available data sources.					
a/ Non-tribal whiting values for canary, darkblotched, and widow reflect bycatch limits for the non-tribal whiting sectors. All other species' impacts are projected from the GMT's whiting impact projection model. The Council may elect to change these bycatch limits when setting final whiting management measures in March 2010 or under any inseason action at any of their future meetings.							
b/ South of 40°10' N. lat.							
e/ Values in scorecard represent projected impacts for all species except canary and yelloweye rockfish, which are the prescribed harvest guidelines.							
f/ 2009 and 2010 OYs are the same except for darkblotched (291 mt in 2010), POP (200 mt in 2010), and widow (509 mt in 2010).							