

Projected mortality (mt) of overfished groundfish species updated with Council action on non-tribal whiting bycatch limits for canary rockfish and widow rockfish in March 2010.

Fishery	Bocaccio a/	Canary	Cow cod	Dkbl	POP	Widow	Yelloweye
Limited Entry Trawl - Non-whiting	16.1	21.3	1.5	230.6	100.8	21.6	0.6
Limited Entry Trawl - Whiting							
At-sea w hiting motherships b/		3.3		6.0	0.5	67	0.0
At-sea w hiting cat-proc b/		4.8		8.5	0.5	95	0.0
Shoreside w hiting b/		5.9		10.5	4.7	117	0.0
Tribal w hiting		4.3		0.0	7.2	5.0	0.0
Tribal							
Midwater Trawl		3.6		0.0	0.0	40.0	0.0
Bottom Trawl		0.8		0.0	3.7	0.0	0.0
Troll		0.5		0.0	0.0		0.0
Fixed gear		0.3		0.0	0.0	0.0	2.3
Fixed Gear Sablefish	0.0	2.5	0.0	4.5	0.4	0.0	0.9
Fixed Gear Nearshore	0.3	3.6	0.0	0.0	0.0	0.3	1.3
Fixed Gear Other	5.0	0.0	0.0	9.0	0.0	0.7	0.0
Open Access: Incidental Groundfish	2.0	0.9	0.0	0.0	0.0	4.0	0.3
Recreational Groundfish c/							
WA		20.9					5.1
OR						1.0	
CA	67.3	22.9	0.3			6.2	2.8
EFPs	11.0	1.3	0.2	1.5	0.1	11.0	0.4
Research: Includes NMFS trawl shelf-slope surveys, the IPHC halibut survey, and expected impacts from SRPs and LOAs.							
	2.0	4.5	0.2	2.0	2.0	5.7	3.3
TOTAL	103.7	101.3	2.2	272.6	119.9	374.5	17.0
2010 OY	288	105	4.0	291	200	509	17
Difference	184.3	3.7	1.8	18.4	80.1	134.5	0.0
Percent of OY	36.0%	96.5%	55.0%	93.7%	60.0%	73.6%	100.0%
Key	= either not applicable; trace amount (<0.01 mt); or not reported in available						

a/ South of 40°10' N. lat.

b/ Non-tribal whiting values for canary, darkblotched, and widow reflect Council recommended bycatch limits for the non-tribal whiting sectors. All other species' impacts are projected from the GMT's whiting impact projection model. The Council may elect to change these bycatch limits under any inseason action at any of their future meetings.

c/ Values in scorecard represent projected impacts for all species except canary and yellow eye rockfish, which are the prescribed harvest guidelines.