

REVISED TABLE 2.1-1. Acceptable biological catches (ABCs) and total catch optimum yield (OY) alternatives (mt) for 2003 for the West Coast under the Council-proposed alternatives. (Overfished stocks in CAPS).

Stock	2002 ABCs/OYs		2003 ABCs and OY Alternatives					
	ABC	No Action OY	ABC	Low OY	Medium OY	High OY	Alloc. Cm. OY	Council OY
LINGCOD	745	577	841	555	651	725	651	651
Pacific Cod	3,200	3,200	3,200			3,200		3,200
PACIFIC WHITING	166,000	129,600	188,000	129,60	148,200	173,600	148,200	148,200
Sablefish								
North of Conception	4,644	4,367	8,209	4,477	7,455	8,187	5,000	6,500
Conception INPFC area	333	229	441	233	323	346	249	294
PACIFIC OCEAN PERCH	640	350	689	311	377	496	377	377
Shortbelly Rockfish	13,900	13,900	13,900			13,900		13,900
WIDOW ROCKFISH	3,727	856	3,871	656	832	916	832	832
CANARY ROCKFISH								
(50% Comm.-50% Rec.)	228	93	256	30	41	45	41	41 ¹
(80% Comm.-20% Rec.)			309	38	52	57		
(20% Comm.-80% Rec.)			218	20	34	37		
Chilipepper Rockfish	2,700	2,000	2,700			2,000		2,000
BOCACCI ²	122	100	198	0	5.8	≤20	≤20	≤20
Splitnose Rockfish	615	461	615			461		461
Yellowtail Rockfish	3,146	3,146	3,146			3,146		3,146
Shortspine Thornyhead	1,004	955	1,004			955		955
Longspine Thornyhead	2,461	2,461	2,461			2,461		2,461
S. of Pt. Conception	390	195	390			195		195
COWCOD (S. Concep)	5	2.4	5			2.4		2.4
N. Concep & Monterey	19	2.4	19			2.4		2.4
DARKBLOTCHED	187	168	205	100	184	205	172	172
YELLOWEYE ³	27	13.5	52	2.1	13.5	27	22	22
Minor Rockfish North	4,795	3,115	4,795			3,115		3,115 ⁴
Minor Rockfish South								
Alternative 1								
Near-Shore:								452
Shallow Near-Shore								105
CA Scorpionfish								85
Deeper Near-Shore								262
Shelf (Remaining+Other)								714
Slope (Remaining+Other)								639
Total	3,506	2,015	3,506			2,015		1,805
Alternative 2								
Near-Shore:								541
Shallow Near-Shore								105
CA Scorpionfish								85
Deeper Near-Shore								351
Shelf (Remaining+Other)								714
Slope (Remaining+Other)								639
Total	3,506	2,015	3,506			2,015		1,894

(CONTINUED)

¹The canary rockfish OY is subject to change dependent upon final council adoption of commercial:recreational catch sharing.

²The medium and high OYs are not supported by a revised rebuilding analysis (MacCall and He 2002) that is scheduled for SSC review at the September Council meeting. The *Medium OY* alternative is based on the June 2002 version of the rebuilding analysis. The *High OY* and *Alloc. Cm.-Preferred OY* alternatives are based on a recent decision by NOAA Fisheries that bocaccio do not conform to National Standard Guidelines; the harvest limit specified is estimated to achieve rebuilding beyond T_{MAX} and is supported by Magnuson-Stevens Act objectives.

³The *High OY* and *Alloc. Cm.-Preferred OY* alternatives are based on a new rebuilding analysis (Methot et al. 2002) that is scheduled for SSC review at the September Council meeting.

⁴Subject to change.

REVISED TABLE 2.1-1. Acceptable biological catches (ABCs) and total catch optimum yield (OY) alternatives (mt) for 2003 for the West Coast under the Council-proposed alternatives. (Overfished stocks in CAPS).

Stock	2002 ABCs/OYs		2003 ABCs and OY Alternatives					
	ABC	No Action OY	ABC	Low OY	Medium OY	High OY	Alloc. Cm. OY	Council OY
Remaining Rockfish North	2,727		2,727					
Black	1,115		1,115					
Bocaccio	318		318					
Chilipepper - Eureka	32		32					
Redstripe	576		576					
Sharpchin	307		307					
Silvergrey	38		38					
Splitnose	242		242					
Yellowmouth	99		99					
Remaining Rockfish South	854		854					
Bank	350		350					
Blackgill	343		343					
Sharpchin	45		45					
Yellowtail	116		116					
Other Rockfish North	2,068		2,068					
South	2,652		2,652					
Dover Sole	8,510	7,440	8,510			7,440		7,440
English Sole	3,100		3,100					
Petrale Sole	2,762		2,762					
Arrowtooth Flounder	5,800		5,800					
Other Flatfish	7,700		7,700					
Other Fish	14,700		14,700					

PFMC
09/13/02

ABCs and for 2003, Council Meeting Draft #2, Friday the 13th, 2002.

All amounts in metric tons	2003 Total ABC		Optimum Yield (OY)			Open-Access			Limited-entry			Limited-entry				
	Total	Landed	Tribal	Rec.	2003 Comp.	Res.	Non-GF ¹	Non-tribal Comm.	%	Total catch	Landed catch	Total catch	At-sea Bycatch	Landed	Total catch	
																Catch
Lingcod	841	651	582	5.2	355	3	4.3	284	19%	54	43	230	0.27 at-sea 70,296	184	shore 50,904	
Whiting	188,000	148,200	25,000			200	1,800	121,200								
Sablefish NoC Conception Area	8,209 441	6,500 294	650 271		11.1	53.0	18.5	5,767 294	9.4%	542	499	5,225	15	4,383	2,364	2,019
Dover sole	8,510	7,440	7,006		62.4	58.0	2.0	7,318						7,006		
English sole	3,100															
Petrale sole	2,762															
Arrowtooth flounder	5,800															
Other flatfish	7,700															
Thornyheads																
Shortspine N. of Pt.C	1,004	955	755	3.0		9.0		941	0.27%	3	3	939		751		
Longspine Conception	2,461 390	2,461 195	2,029 195		8.9	18.0		2,434 0				2,434 0		2,020 195		
Widow 60%	3,871	832	513	45	5	1.5	0.0	781	3.0%	23	20	757	182	488		
POP 60%	689	377	314			3.0		374				374		314		
Yellowtail	3,146	3,146	1,978	400	15	8.0	5.8	2,717	8.3%	226	189	2,492	300	1,773		
Chilipepper	2,700	2,000	1,682		15			1,985	44.3%	879	739	1,106		929		
Splitnose (Rosefish)	615	461	387									461		387		
Canary 60%	256	41	30	2.5	19	1	2.2	17	12.3%	2.0	1.7	15	3	10		
Canary 60%	272	44	34	2.5	16	1	2.2	23	12.3%	2.8	2.3	20	3	14		
Canary 61% C-39%R																
Bocaccio	198	20	17		5	0.2	0.8	14	44.3%	6	5	8		7		
Cowcod (Conception)	5	2.4	0					0				0		0		
Cowcod (Monterey)	19	2.4	0					0				0		0		
Cowcod (total)	24	4.8					0.1	0				0		0		
Darkblotched 80%; Tmid:	205	172	132			1.6	0	170		0	0	170	5	132		
Yelloweye Tmid:50%	52	22		3.0	7.7	0.6	3.3	7		1.1	0.0	2.8		2.2		

1/ This column reflects estimate revisions made during the Council meeting for groundfish bycatch in non-groundfish fisheries.

ABCs and OYs for 2003, Council Meeting Draft #2, Friday the 13th, 2002.

All amounts in metric tons	2003 Total/ABC		Optimum Yield (OY)										Open-Access		Limited-entry		Limited-entry	
	Catch	Landed	Tribal	Rec.	2003 Comp.	Res.	Non-GF ¹	Non-tribal Comm.	%	Total catch	Landed catch	Total catch	At-sea Bycatch	Landed	Total catch	Trawl	FG	
																		Landed
Minor Sebastes 2002 OY	987	971																
Near-shore (OR:2000 cap)	530	396		396			135											
OR : Black/Blue HG	497			384			113											
OR : Other NS HG	33			11			22											
Near-shore (N.CA.:2000 cap)	108			54			54											
N.CA : Black/Blue HG	88.3			44			44											
N.CA : Other NS HG	19.5			10			10											
Near-shore (OR+N.CA.:2000 cap)	638			319			319											
N.CA : Black/Blue HG	585			293			293											
N.CA : Other NS HG	53			26			26											
Near-shore (WA. combined species)	290			290														
Shelf (Remaining+Other)	968	10	10	10			948											
Slope (Remaining+Other)	1,160	0	4				1,156	0.1										
South							0											
Near-shore (Rev 8-23) Alt. #1B	541	433		433			108											
Shallow NS	105	66		66			39	0.4										
CA Scorpionfish	85	64		64			21	0.2										
Deeper NS	351	303		303			48	0.4										
Shelf (Remaining+Other)	714	0		60			639											
Slope (Remaining+Other)	639																	

1/ This column reflects estimate revisions made during the Council meeting for groundfish bycatch in non-groundfish fisheries.