

**SOUTHERN NEARSHORE ROCKFISH (south of 40⁰10' N. lat.)
 OPTIMUM YIELD
 AND HARVEST GUIDELINES FOR 2003**

*Maria
 Vojkanich*

Scenario #1B

**Comm.: Rec. Ratio for shallow Sps. based upon 1983-89 & 93-99 landings
 Deeper NS RF % adjusted for overall
 allotment = 20/80**

	MT	Comm %	Comm MT	Rec %	Rec MT
Shallow NS Rockfishes	104.8	37.0	38.8	63.0	66.1
CA Scorpionfish	84.9	24.8	21.0	75.2	63.8
Deeper NS Rockfishes	262.0	11.6	30.4	88.4	231.6
Total	451.7	20.0	90.2	80.0	361.5

Scenario #1B (Revised)

**Comm.: Rec. Ratio for shallow Sps. based upon 1983-89 & 93-99
 landings
 Deeper NS RF % adjusted
 for overall allotment = 20/80**

	MT	Comm %	Comm MT	Rec %	Rec MT
Shallow NS Rockfishes	104.8	37.0	38.8	63.0	66.1
CA Scorpionfish	84.9	24.8	21.0	75.2	63.8
Deeper NS Rockfishes	351.0	13.7	48.0	86.3	303.0
Total	540.7	20.0	107.8	80.0	432.9

NORTHERN CALIFORNIA NEARSHORE ROCKFISH (north of 40⁰10' N. lat.)

HARVEST GUIDELINES FOR 2003

California Catch of Selected Marine species Groups (metric tons)

From 40⁰10' N. Lat. To Oregon Border

Year	Black & Blue Rockfish			Other Nearshore Rockfish		
	Com	Rec	Total	Com	Rec	Total
1994	141.5	102.8	244.4	6.5	12.3	18.8
1995	165.6	93.3	258.9	23.6	6.0	29.6
1996	115.9	71.2	187.1	18.5	6.9	25.4
1997	138.5	47.7	186.2	28.3	6.1	34.4
1998	81.3	59.8	141.1	15.0	4.0	19.0
1999	59.7	67.1	126.8	28.7	9.6	38.3
Mean 1994-99	117.1	73.7	190.8	20.1	7.5	27.6
Mean 94-99 x 0.5	58.5	36.8	95.4	10.1	3.7	13.8
2000	38.6	49.7	88.3	12.4	7.1	19.5
2001 ¹	104.7	141.4	246.1	22.2	5.6	27.8

Source: PacFIN and RecFIN. Note: Commercial "Other Nearshore Rockfish
 in 2000 and 2001 likely contain some black rockfish

1 Note: The 2001 commercial estimates are very preliminary as of this summary date.
 It is estimated that these data represent 70 - 80% of the total.