

2007 Projected mortality impacts (mt) under current regulations. March 2007 update prior to inseason adjustments. a/

1/31/2007

Fishery	Bocaccio b/	Canary	Cowcod	Dkbl	POP	Widow	Yelloweye
Limited Entry Trawl- Non-whiting	47.9	20.0	2.1	194.3	71.6	0.7	0.1
Limited Entry Trawl- Whiting							
At-sea whiting motherships					1.0		0.0
At-sea whiting cat-proc		4.7		25.0	2.9	200.0	0.0
Shoreside whiting					1.8		0.0
Tribal whiting		0.7		0.0	0.6	6.1	0.0
Tribal							
Midwater Trawl		1.8		0.0	0.0	40.0	0.0
Bottom Trawl		0.8		0.0	3.7	0.0	0.0
Troll		0.5		0.0	0.0		0.0
Fixed gear		0.3		0.0	0.0	0.0	2.3
Limited Entry Fixed Gear		1.1		1.3	0.4		2.9
Sablefish	13.4		0.0			0.0	
Non-Sablefish			0.1			0.5	
Open Access: Directed Groundfish		1.0					
Sablefish DTL	0.0			0.2	0.1	0.0	0.5
Nearshore (North of 40°10' N. lat.)	0.0	1.7	0.1	0.0	0.0	0.1	2.0
Nearshore (South of 40°10' N. lat.)	0.0			0.0	0.0		
Other	10.6			0.0	0.0	0.0	0.1
Open Access: Incidental Groundfish							
CA Halibut	0.1	0.0		0.0	0.0		
CA Gillnet c/	0.5			0.0	0.0	0.0	
CA Sheephead c/				0.0	0.0	0.0	0.0
CPS- wetfish c/	0.3						
CPS- squid d/							
Dungeness crab c/	0.0		0.0	0.0	0.0		
HMS b/		0.0	0.0	0.0			
Pacific Halibut c/	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pink shrimp	0.1	0.1	0.0	0.0	0.0	0.1	0.1
Ridgeback prawn	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Salmon troll	0.2	0.8	0.0	0.0	0.0	0.3	0.2
Sea Cucumber	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Spot Prawn (trap)							
Recreational Groundfish e/							
WA		5.7					6.2
OR						1.4	
CA	98.0	8.3	0.4			8.0	1.7
Research: Includes NMFS trawl shelf-slope surveys, the IPHC halibut survey, and expected impacts from SRPs and LOAs. f/							
	2.0	7.5	0.1	3.8	3.6	0.9	2.0
TOTAL	173.2	55.0	2.8	224.7	85.7	258.1	18.2
2007 OY	218	44.0	4.0	290	150	368	23
Difference	44.8	-11.0	1.2	65.4	64.3	110.0	4.8
Percent of OY	79.4%	125.0%	70.0%	77.5%	57.1%	70.1%	79.0%
Key	= either not applicable; trace amount (<0.01 mt); or not reported in available						

a/ All numbers reflect projected annual total catches except that the non-tribal "Limited Entry Trawl- Whiting" numbers are the total bycatch caps for canary, darkblotched, and widow rockfish. Only cells in bold font borders have been updated.

b/ South of 40°10' N. lat.

c/ Mortality estimates are not hard numbers; based on the GMT's best professional judgment.

d/ Bycatch amounts by species unavailable, but bocaccio occurred in 0.1% of all port samples and other rockfish in another 0.1% of all port samples (and squid fisheries usually land their whole catch).

e/ Values in scorecard represent projected impacts. However, harvest guidelines for 2007 are as follows: canary in WA and OR combined = 8.2 mt and in CA = 9.0 mt; yelloweye in WA and OR combined = 6.8 mt and in CA = 2.1 mt.

f/ Research projections only updated for canary rockfish in November 2006. The other species' updates will be updated in March 2007.