


PRESEASON REPORT I

STOCK ABUNDANCE ANALYSIS FOR 2008 OCEAN SALMON FISHERIES



Pacific Fishery Management Council
7700 NE Ambassador Place, Suite 101
Portland, OR 97220-1384
(503) 820-2280
www.pcouncil.org

FEBRUARY 2008

ACKNOWLEDGMENTS

SALMON TECHNICAL TEAM

MR. DELL SIMMONS, CHAIR

National Marine Fisheries Service, Lacey, Washington

MR. ALLEN GROVER, VICE CHAIR

California Department of Fish and Game, Santa Rosa, California

MR. CRAIG FOSTER

Oregon Department of Fish and Wildlife, Clackamas, Oregon

DR. ROBERT KOPE

National Marine Fisheries Service, Seattle, Washington

MR. KEITH LUTZ

Northwest Indian Fisheries Commission, Olympia, Washington

MR. DOUG MILWARD

Washington Department of Fish and Wildlife, Olympia, Washington

MR. MICHAEL MOHR

National Marine Fisheries Service, Santa Cruz, California

MR. HENRY YUEN

U.S. Fish and Wildlife Service, Vancouver, Washington

PACIFIC FISHERY MANAGEMENT COUNCIL STAFF

MR. CHUCK TRACY

MS. RENEE DORVAL

MS. CARRIE MONTGOMERY

MS. KIM MERYDITH

The Salmon Technical Team and the Council staff express their thanks for the expert assistance provided by Ms. Wendy Beeghley and Ms. Cindy LeFleur, Washington Department of Fish and Wildlife; Mr. Eric Schindler, Oregon Department of Fish and Wildlife; Ms. Melodie Palmer-Zwahlen and Ms. Jennifer Simon, California Department of Fish and Game; Ms. Sandy Zeiner, Northwest Indian Fisheries Commission; Dr. Michael O'Farrell, National Marine Fisheries Service, Southwest Science Center, and numerous other agency and tribal personnel in completing this report.

This document may be cited in the following manner:

Pacific Fishery Management Council. 2008. *Preseason Report I: Stock Abundance Analysis for 2008 Ocean Salmon Fisheries*. (Document prepared for the Council and its advisory entities.) Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, Oregon 97220-1384.



A report of the Pacific Fishery Management Council pursuant to National Oceanic and Atmospheric Administration Award Number NA05NMF4410008.

TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES	iii
LIST OF FIGURES	iv
LIST OF ACRONYMS AND ABBREVIATIONS.....	v
INTRODUCTION	1
CHAPTER I - Abundance Projections.....	3
CHAPTER II - CHINOOK SALMON ASSESSMENT.....	15
CHINOOK STOCKS SOUTH OF CAPE FALCON	15
SACRAMENTO RIVER FALL CHINOOK SALMON.....	15
Predictor Description	15
KLAMATH RIVER FALL CHINOOK	16
Predictor Description	16
Predictor Performance	16
2008 Stock Status.....	16
Evaluation of 2007 Regulations on 2008 Stock Abundance.....	16
OTHER CALIFORNIA COASTAL CHINOOK STOCKS	16
OREGON COASTAL CHINOOK STOCKS.....	17
North Migrating Chinook	17
South/Local Migrating Chinook	18
Evaluation of 2007 Regulations on 2008 Stock Abundance.....	19
CHINOOK STOCKS NORTH OF CAPE FALCON.....	19
Columbia River Fall Chinook.....	19
Predictor Description and Past Performance	19
2008 Stock Status.....	20
Evaluation of 2007 Regulations on 2008 Stock Abundance.....	21
Washington Coastal Chinook.....	21
Predictor Description and Past Performance	21
2008 Stock Status.....	21
Puget Sound Chinook	21
2008 Stock Status.....	21
Evaluation of 2007 Regulations on 2008 Stock Abundance.....	22
CHAPTER III - COHO SALMON ASSESMENT.....	39
COLUMBIA RIVER AND OREGON/CALIFORNIA COASTAL COHO	39
(OREGON PRODUCTION INDEX AREA)	39
Public Hatchery Coho	39
Predictor Description	40
Predictor Performance	40
2008 Stock Status.....	40
Oregon Coastal Natural Coho.....	40
Predictor Description	41
Predictor Performance	42
2008 Stock Status.....	42
Private Hatchery Coho	42

TABLE OF CONTENTS (continued)

	<u>Page</u>
Salmon Trout Enhancement Hatchery Coho Smolt Program	42
Predictor Description	42
Predictor Performance	42
2008 Stock Status.....	42
Lower Columbia River Natural.....	42
Predictor Description	42
Predictor Performance	43
2008 Stock Status.....	43
Oregon Production Index Area Summary of 2008 Stock Status.....	43
WASHINGTON COASTAL AND PUGET SOUND COHO STOCKS	43
Predictor Description and Past Performance	43
2008 Stock Status.....	43
Washington Coastal Coho.....	43
Puget Sound	46
SELECTIVE FISHERY CONSIDERATIONS	47
EVALUATION OF 2007 REGULATIONS ON 2008 STOCK ABUNDANCE.....	48
Oregon Production Index Area	48
North of the Oregon Production Index Area.....	48
CHAPTER IV - FRASER RIVER AND PUGET SOUND PINK SALMON ASSESSMENTS.....	61
APPENDIX A SUMMARY OF COUNCIL STOCK MANAGEMENT GOALS	63
APPENDIX B OREGON PRODUCTION INDEX DATA	79
APPENDIX C SALMON HARVEST ALLOCATION SCHEDULES	85
APPENDIX D CVI PREDICTOR: EXCLUSION OF 2005 DATA POINT	97

LIST OF TABLES

		<u>Page</u>
TABLE I-1.	Preseason adult Chinook salmon stock forecasts in thousands of fish	4
TABLE I-2.	Preseason adult coho salmon stock forecasts in thousands of fish	7
TABLE I-3.	Achievement of conservation objectives for natural stocks listed in Table 3-1 of the Pacific Coast Salmon Plan	9
TABLE II-1.	Indices of annual abundance and ocean fishery impacts on California Central Valley Chinook in thousands of fish.....	23
TABLE II-2.	Comparisons of preseason forecast and postseason estimates for the CVI in thousands of fish	24
TABLE II-3.	Klamath River fall Chinook ocean abundance (thousands), harvest rate, and river run size estimates (thousands) by age.....	25
TABLE II-4.	Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook	26
TABLE II-5.	Summary of management objectives and predictor performance for Klamath River fall Chinook	28
TABLE II-6.	Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook	29
TABLE II-7.	Rogue River fall Chinook inriver run and ocean population indices.....	31
TABLE II-8.	Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish	32
TABLE II-9.	Comparison of preseason and postseason forecasts of Puget Sound run size for summer/fall Chinook	35
TABLE III-1.	Preliminary 1996-2008 preseason and postseason coho stock Stratified Random Sampling abundance estimates for Oregon production index area stocks in thousands of fish	50
TABLE III-2.	Oregon production index (OPI) area coho harvest impacts, spawning, abundance, and exploitation rate estimates by SRS accounting in thousands of fish.....	52
TABLE III-3.	Preseason and postseason estimates of ocean escapements for selected Washington coastal adult natural coho stocks in thousands of fish	53
TABLE III-4.	Preseason and postseason estimates of ocean escapements for selected Puget Sound adult natural coho stocks in thousands of fish	54
TABLE III-5.	Mass marking of 2005 brood coho available to 2008 Council fisheries. The mark used is an adipose fin clip	55
TABLE III-6.	Projected coho mark rates for 2008 fisheries under base period fishing patterns (% marked)	56
TABLE III-7.	Estimated ocean escapements for critical natural and Columbia River hatchery coho stocks (thousands of fish) based on preliminary 2008 preseason abundance forecasts and 2007	57
TABLE III-8.	Comparison of Lower Columbia natural (LCN), Oregon coastal natural (OCN), and Rogue/Klamath (RK) coho projected harvest mortality and exploitation rates by fishery under Council-adopted 2007 regulations and preliminary 2008 preseason abundance estimates	58
TABLE III-9.	Maximum allowable fishery impact rate for OCN coho under Amendment 13 matrix (Appendix A, Table A-2) and the OCN work group matrix (Appendix A, Table A-3) based on parent escapement levels by stock component and marine survival category.....	59
TABLE IV-1.	Estimated annual run sizes (odd-numbered years 1977-2007) for Fraser River and Puget Sound pink salmon in millions of fish	61

LIST OF FIGURES

	<u>Page</u>
FIGURE I-1. Selected preseason vs. postseason forecasts for Chinook stocks with significant contribution to Council area fisheries.	11
FIGURE I-2a. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.	12
FIGURE I-2b. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.	13
FIGURE II-1. Regression estimator for CVI based on previous year's river return of age-two Central Valley Chinook, 1990-2007 with 2005 data point excluded.....	37
FIGURE II-2. Spawning escapements of adult Sacramento River fall Chinook, 1970-2007, and the goal range for the stock of 122,000 to 180,000 adult fish.....	37
FIGURE II-3. Regression estimators for Klamath River fall Chinook ocean abundance (September 1) based on that year's river return of same cohort	38

LIST OF ACRONYMS AND ABBREVIATIONS

BY	brood year
CDFG	California Department of Fish and Game
CoTC	Coho Technical Committee (of the PSC)
Council	Pacific Fishery Management Council
CRFMP	Columbia River Fishery Management Plan
CVI	Central Valley Index
CWT	coded-wire tag
EEZ	exclusive economic zone (from 3-200 miles from shore)
ESA	Endangered Species Act
ESU	evolutionarily significant unit
FMP	fishery management plan
FRAM	Fishery Regulatory Assessment Model
ISBM	individual stock-based management
Jack CR	Columbia River jacks
Jack OC	Oregon coastal and Klamath River Basin jacks
KMZ	Klamath management zone (ocean zone between Humbug Mountain and Horse Mountain where management emphasis is on Klamath River fall Chinook)
KOHM	Klamath Ocean Harvest Model
KRFC	Klamath River fall Chinook
LCN	lower Columbia River natural (coho)
LRB	lower Columbia River bright (Chinook)
LRH	lower Columbia River hatchery (tule fall Chinook returning to hatcheries below Bonneville Dam)
LRW	lower Columbia River wild (bright fall Chinook spawning naturally in tributaries below Bonneville Dam)
MCB	mid-Columbia River brights (bright hatchery fall Chinook released below McNary Dam)
MOC	mid-Oregon coast
MSY	maximum sustainable yield
NA	not available
NEPA	National Environmental Policy Act
NMFS	National Marine Fisheries Service
NOC	north Oregon coast
OCN	Oregon coastal natural (coho)
OCNL	Oregon coastal natural lake
OCNR	Oregon coastal natural river
ODFW	Oregon Department of Fish and Wildlife
OPI	Oregon Production Index (coho salmon stock index south of Leadbetter Point)
OPIH	Oregon Production Index public hatchery
PRIH	Private hatchery
PSC	Pacific Salmon Commission
PST	Pacific Salmon Treaty
RER	rebuilding exploitation rate
RK	Rogue/Klamath (coho)
RMP	Resource Management Plan (for exemption from ESA section 9 take prohibitions under limit 6 of the 4(d) rule)
SAB	Select Area brights
SCH	Spring Creek Hatchery (tule fall Chinook returning to Spring Creek Hatchery)
SRS	Stratified Random Sampling
STEP	Salmon Trout Enhancement Program

LIST OF ACRONYMS AND ABBREVIATIONS (continued)

STT	Salmon Technical Team (formerly the Salmon Plan Development Team)
URB	upper river brights (naturally spawning bright fall Chinook normally migrating past McNary Dam)
VSI	visual stock identification
WCVI	West Coast Vancouver Island
WDFW	Washington Department of Fish and Wildlife

INTRODUCTION

This is the second report in an annual series of four reports prepared by the Salmon Technical Team (STT) of the Pacific Fishery Management Council (Council) to document and help guide salmon fishery management off the coasts of Washington, Oregon, and California. This report will be formally reviewed at the Council's March meeting. The third and fourth reports in this series will be developed at the close of the March and April Council meetings, respectively, to analyze the impacts of the Council's proposed and final ocean salmon fishery management recommendations for 2008.

This report provides 2008 salmon stock abundance projections, and an analysis of the impacts of 2007 regulations, or regulatory procedures, on the projected 2008 abundance. This analysis is analogous to that of a no-action alternative in a National Environmental Policy Act (NEPA) analysis, and is intended to give perspective in developing 2008 management measures. The report focuses on Chinook, coho, and pink salmon stocks that have been important in determining Council fisheries in recent years, and on stocks listed under the Endangered Species Act (ESA) with established National Marine Fisheries Service (NMFS) ESA consultation standards.

Chapter I provides a summary of stock abundance projections. Chapters II and III provide detailed stock-by-stock analyses of abundance, a description of prediction methodologies, and accuracy of past abundance predictions for Chinook and coho salmon, respectively. Chapter IV summarizes abundance information for pink salmon. Four appendices provide supplementary information as follows: Appendix A provides a summary of Council stock management goals; Appendix B contains pertinent data for Oregon production index (OPI) area coho; Appendix C contains the Council's current harvest allocation schedules, and; Appendix D details updated forecasting methodologies.

In 2002, the Pacific Salmon Commission (PSC) reached agreement on a management regime that constrains total fishery exploitation rates on key management units of naturally spawning coho salmon originating in Southern British Columbia, Puget Sound, and the Washington Coast. The agreement calls for the PSC Coho Technical Committee (CoTC) to develop a regional coho fishery planning model for application beginning in 2005. The CoTC has agreed to use Coho Fishery Regulation Assessment Model (FRAM) as the core for an initial version of the regional coho fishery planning model to provide a consistent basis for fishery planning processes in the United States and Canada.

STT Concerns

A Conservation Alert is triggered when a stock is forecast to fall short of its conservation objective. While no Conservation Alert has been triggered this year, the STT is concerned that 2007 observed escapements for several stocks are outside the bounds of the data. In 2007, Central Valley Chinook had a record low number of jacks in the escapement last year. Sacramento River fall Chinook are the only component of the Central Valley stock for which the Council has a conservation objective. Sacramento River fall Chinook normally make up the majority of the Central Valley stock, but this year, they constituted only 32% of the jack return used to forecast the CVI. Klamath River fall Chinook also had record low jack returns in 2007, and are thus outside the bounds of the data used to forecast ocean age-3 abundance.

The STT does not make a quantitative forecast of the Oregon coast fall Chinook. In the past, the STT has relied on the recent increasing trend in escapement, and the fact that the stock consistently met or exceeded its goal for many years, to justify an expectation that the stock would continue meet its conservation objective. The escapement index for north migrating Oregon coast fall Chinook has declined sharply for the past four years and the stocks failed to meet their post-season escapement goal in 2007 for the first time since 1983.

The fact that so many of the stocks south of Cape Falcon are experiencing declining trends suggests that recent ocean conditions have been very unfavorable for survival. The STT is concerned that the 2008 forecasts for stocks south of Cape Falcon may be overly optimistic.

CHAPTER I - ABUNDANCE PROJECTIONS

Abundance expectations in 2008 are summarized for key Chinook and coho salmon stocks in Tables I-1 and I-2, respectively. A cursory comparison of preseason forecast and postseason abundance estimates for selected stocks is presented in Figures I-1 and I-2. More detailed analyses of this subject are covered in Chapter II (Chinook) and III (coho). Information on pink salmon abundance, which is only significant in odd-numbered years, is contained in Chapter IV. Council Salmon Fishery Management Plan (FMP) management goals are presented in Table I-3 and Appendix A, Table A-1.

In addition to the key stocks with abundance projections listed in Tables I-1 and I-2, Council management decisions for the 2008 ocean salmon fishing seasons may be constrained by other stocks, such as those listed under the ESA or subject to the PSC agreement, which may not have abundance projections made, or do not have abundance projections available in time for inclusion in this report. These include the following ESU's: Sacramento River Winter, Central Valley Spring, California Coastal, Lower Columbia River, and Snake River Fall Chinook; and Central California and Southern Oregon/Northern California coho, as well as Interior Fraser (including Thompson River) coho.

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 1 of 3)

Production Source and Stock or Stock Group	2000	2001	2002	2003	2004	2005	2006	2007	2008	Methodology for 2008 Prediction and Source
California Central Valley (Index)										
Sacramento and San Joaquin Basins, Fall, Late Fall, Spring, and Winter Run	790.4	649.4	825.4	1,108.1	831.8	1,678.3	632.5	499.9	157.1	Linear regression analysis of river age-2 jacks on CVI of the following year. Data point 2005 excluded for 2008 CVI forecast. CDFG staff.
Klamath River (Ocean Abundance)										
Fall Run	389.9	435.5	362.5	310.2	216.3	239.8	110.0	546.2	190.7	Linear regression analysis of age-specific ocean abundance estimates on river runs of same cohort. KRTAT.
Oregon Coast										
North and South/Local Migrating	Estimates Not Made									None.
Columbia River (Ocean Escapement)										
Upriver Spring	134.0	364.6	333.7	145.4	360.7	254.1 ^{aj}	88.4	78.5	269.3	Age-specific linear regressions of cohort returns in previous run years. WDFW staff.
Willamette Spring	59.9	61.0	73.8	109.8	109.4	116.9	46.5	52.0	34.0	Age-specific linear regressions of cohort returns in previous run years. ODFW staff.
Sandy Spring	3.8	4.0	4.3	4.8	5.2	7.4	8.2	7.9	6.8	Recent year average. ODFW staff.
Cowlitz Spring	2.0	1.0	3.1	4.9	15.9	12.7	3.0	6.4	5.2	Age-specific linear regressions of cohort returns in previous run years. WDFW staff.
Kalama Spring	1.4	1.0	1.6	3.6	6.0	4.5	1.5	4.0	3.7	Age-specific linear regressions of cohort returns in previous run years. WDFW staff.
Lewis Spring	2.6	2.8	2.0	3.1	5.4	7.6	1.8	5.9	3.5	Age-specific linear regressions of cohort returns in previous run years. WDFW staff.
Upriver Summer	33.3	24.5	77.7	87.6	102.8	62.4 ^{aj}	49.0	45.6	52.0	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.
URB Fall	171.1	127.2	281.0	280.4	292.2	352.2	253.9	182.4	162.5	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.
SCH Fall	21.9	56.6	144.4	96.9	138.0	114.1	50.0	21.8	87.2	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.
LRW Fall	3.5	16.7	18.7	24.6	24.1	20.2	16.6	10.1	3.8	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.
LRH Fall	23.7	32.2	137.6	115.9	77.1	74.1	55.8	54.9	59.0	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.
MCB Fall	50.6	43.5	96.2	104.8	90.4	89.4	88.3	68.0	54.0	Age-specific average cohort ratios/cohort regressions. Columbia River TAC.

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 2 of 3)

Production Source and Stock or Stock Group		2000	2001	2002	2003	2004	2005	2006	2007	2008	Methodology for 2008 Prediction and Source	
Washington Coast (Ocean Escapement)												
Willapa Bay	Natural	4.2	4.3	3.7	2.4	4.1	3.2	2.0	2.0	2.5	Mean return per release by age class adjusted for brood performance through 2007 return year. WDFW staff.	
	Hatchery	18.9	17.8	18.8	14.2	14.7	17.4	29.8	29.8	27.0		
Quinault Spring/Summer	Natural	NA	NA	NA	NA	NA	NA	NA	NA	NA	Age specific mean cohort ratios and linear regression analysis using recent 5 year mean.	
Quinault Fall	Hatchery	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Queets Spring/Summer	Natural	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Queets Fall	Natural	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Hoh Spring/Summer	Hatchery	NA	NA	NA	NA	NA	NA	NA	NA	NA		
	Natural	NA	NA	NA	NA	NA	NA	NA	NA	0.9		
Hoh Fall	Natural	NA	NA	NA	NA	NA	NA	NA	NA	2.9		Age specific mean cohort ratios and linear regression analysis, means of all years subtracting out the high and low years.
Quillayute Spring	Hatchery	NA	NA	NA	NA	NA	NA	NA	NA	1.7		
Quillayute Summer/Fall	Natural	NA	NA	NA	NA	NA	NA	NA	NA	6.3	Mean return per release, adjusted means for 5-6 year olds. Summer: Recent 5 year mean return per spawner. Fall: Recent year mean return rates from cohort analysis.	
<i>North Coast Totals</i>												
Spring/Summer	Natural	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Fall	Natural	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Spring/Summer	Hatchery	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Fall	Hatchery	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Puget Sound^{b/}												
Nooksack/Samish	Hatchery	19.0	34.9	52.8	45.8	34.2	19.5	16.9	18.8	35.3	Brood release times average return/release rate (2004-07 return years) adjusted for forecast performance.	
East Sound Bay	Hatchery	5.0	1.6	1.6	1.6	0.8	0.4	0.4	0.4	0.8	Brood release times average return/release rate (2005-07 return years).	
Skagit	Natural	7.3	9.1	13.8	13.7 ^{ci}	20.4 ^{ci}	23.4 ^{ci}	24.1 ^{ci}	15.0 ^{ci}	23.8 ^{ci}	Age-specific average cohort return rate method, averaged with environmental predictor model-based forecast.	
	Hatchery	0.0	0.0	0.0	0.0 ^{ci}	0.5 ^{ci}	0.7 ^{ci}	0.6 ^{ci}	1.1 ^{ci}	0.7 ^{ci}		
Stillaguamish	Natural	2.0 ^{di}	1.7 ^{di}	2.0 ^{di}	2.0 ^{di}	3.3 ^{di}	2.0 ^{di}	1.6 ^{di}	1.9 ^{di}	1.1 ^{di}	Supplemental fish forecast based on observed survival rates for tagged fish (1986-1993 brood years). Natural-origin based on recruits per spawner observed for 1974-99 brood years. Forecast is combination of supplemental plus natural origin.	
	Hatchery	6.2	4.1	6.8 ^{di}	9.4 ^{di}	10.1 ^{di}	9.9 ^{di}	9.6 ^{di}	8.7 ^{di}	8.8 ^{di}		
Snohomish	Natural	6.0	5.8 ^{di}	6.7 ^{di}	5.5 ^{di}	15.7 ^{di}	14.2 ^{di}	8.7 ^{di}	12.3 ^{di}	6.5 ^{di}	Average total recruitment based on TRT A and P tables. For Skykomish used BYs 1994-1998 applied to 2002-2005 BY age returns, adjusted by the ratio actual/expected 2007 escapement.	
	Hatchery	6.2	4.1	6.8 ^{di}	9.4 ^{di}	10.1 ^{di}	9.9 ^{di}	9.6 ^{di}	8.7 ^{di}	8.8 ^{di}		
Tulalip	Hatchery	5.0	5.5	5.8 ^{di}	6.0 ^{di}	7.6 ^{di}	9.2 ^{di}	10.0 ^{di}	8.1 ^{di}	4.1 ^{di}	CWT survival rates (1986-1991) multiplied by release numbers for brood years 2002-2005, adjusted by the ratio actual/expected 2007 escapement.	

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 3 of 3)

Production Source and Stock or Stock Group		2000	2001	2002	2003	2004	2005	2006	2007	2008	Methodology for 2008 Prediction and Source
South Puget Sound	Natural	17.5	16.2	16.9	19.6	17.5	17.7	21.3	17.0	21.1	Puyallup-based predicted return at age calculated for return years 1993-2005, multiplied by average difference between forecasts and run sizes from 1999 to 2006. For Nisqually, recent 5-year average (2002-2006).
	Hatchery	77.5	73.7	90.8	86.6	86.5	83.1	85.8	92.1	101.3	Average return at age multiplied by cohort release for Green and 10E. Average of two different methods for Carr Inlet, (1) 1980-2005 mean return/smolt released multiplied by 2002 brood smolts released, and (2) 1980-2006 mean return/pound released multiplied by 2004 brood pounds released.
Hood Canal	Natural	19.2	2.7	2.9 ^{c/}	3.6 ^{c/}	2.4 ^{c/}	3.1 ^{c/}	2.5 ^{c/}	3.8 ^{c/}	2.6 ^{c/}	Natural fish based on the Hood Canal terminal run reconstruction-based relative contribution of the individual Hood Canal management units in the 2004-2007 return years.
	Hatchery		22.6	21.1 ^{c/}	30.2 ^{c/}	27.2 ^{c/}	27.5 ^{c/}	27.7 ^{c/}	43.6 ^{c/}	34.2 ^{c/}	Brood 2004 fingerling lbs released from WDFW facilities in 2005, multiplied by the average of postseason estimated terminal area return rates (total terminal run / hatchery fingerling lbs released three years previous) for the last eight return years (2000-2007), excluding return year 2005 in which the return rate was a statistical outlier.
Hoko	Natural	----- Included in Juan de Fuca Forecast -----								1.1 ^{a/}	Sibling regressions;
Strait of Juan de Fuca	Natural	1.1	3.5	3.6 ^{c/}	3.4 ^{c/}	3.6 ^{c/}	4.2 ^{c/}	4.2 ^{c/}	4.4 ^{c/}	3.2	Four-year average 2003-2006 of terminal run size. Elwha estimate is a combination of hatchery and wild fish.
	Hatchery	2.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	

a/ Beginning in 2005, the upriver spring/summer designation was changed, with stream type Snake Basin summer fish being combined with the spring stock.

b/ Forecast is Puget Sound run size available to U.S. net fisheries. Does not include fish caught in troll and recreational fisheries.

c/ Terminal run forecast.

d/ Expected spawning escapement without fishing.

TABLE I-2. Preseason adult coho salmon stock forecasts in thousands of fish. (Page 1 of 2)

Production Source and Stock or Stock Group		2000	2001	2002	2003	2004	2005	2006	2007	2008	Methodology for 2008 Prediction and Source
OPI Area (Total Abundance)		727.9	1,758.7	434.1	984.6	777.9	542.9	460.2	870.7	289.5	Sum of stock component estimates.
(California and Oregon Coasts and Columbia River)											
OPI Public	Hatchery	671.4	1,707.6	361.7	863.1	623.9	389.9	398.8	593.6	216.1	A new method was developed to estimate coho abundances for the hatchery components of the Columbia River and the Oregon Coast. The new method is based on the 86-92 base period and "backwards" FRAM runs for recent years. See text in Chapter III for details.
	Columbia River Early	326.3	1,036.5	161.6	440.0	313.6	284.6	245.8	424.9	110.3	
	Columbia River Late	278.0	491.8	143.5	377.9	274.7	78.0	113.8	139.5	86.4	
	Coastal N. of Cape Blanco	48.5	127.3	36.6	29.3	16.6	11.5	8.6	7.0	1.7	
	Coastal S. of Cape Blanco	18.6	52.0	20.0	15.9	19.0	15.8	30.6	22.2	17.7	
Lower Columbia River	Natural	NA	NA	NA	NA	NA	NA	NA	21.5	13.4	A new method was developed to estimate LCR wild coho. The method relies on the 86-92 base period and "backward" FRAM runs for recent years. See text in
Oregon Coast (OCN)	Natural	55.9	50.1	71.8	117.9	150.9	152.0	60.8	255.4	60.0	Prediction for 2008 is equal to 2007 observed return.
STEP	Hatchery	0.6	1.0	0.6	3.6	3.1	1.0	0.6	0.2	0.0	No forecast for 2008; releases discontinued.
Washington Coast											
Willapa	Natural	9.9	21.6	21.6	31.8	36.7	35.9	30.3	24.4	35.1	A variety of methods were used for 2008, primarily based on smolt production and survival. See text in Chapter III for details.
	Hatchery	19.6	36.1	40.4	57.5	55.0	56.4	37.7	37.2	25.5	
Grays Harbor	Natural	47.8	51.3	55.4	58.0	117.9	91.1	67.3	59.4	42.7	
	Hatchery	75.8	67.1	56.8	64.0	67.8	54.4	52.4	74.0	53.1	
Quinalt	Natural	4.4	8.7	29.4	47.7	50.5	44.9	28.8	18.6	17.4	
	Hatchery	7.4	10.8	12.3	20.6	18.2	33.6	34.5	22.7	24.5	
Queets	Natural	2.7	12.0	12.5	24.0	18.5	17.1	8.3	13.6	10.2	
	Hatchery	11.8	10.0	16.0	24.9	17.1	17.4	11.9	19.1	10.3	
	Supplemental ^{b/}	0.8	NA	2.0	1.3	2.5	2.4	-	-	-	
	(Flood)										
Hoh	Natural	3.5	8.5	8.5	12.5	8.1	7.6	6.4	5.4	4.3	
Quillayute Fall	Natural	8.7	23.0	22.3	24.9	21.2	18.6	14.6	10.8	10.5	
	Hatchery	13.9	15.3	15.0	15.2	20.9	22.1	10.4	18.1	13.0	

TABLE I-2. Preseason adult coho salmon stock forecasts in thousands of fish. (Page 2 of 2)

Production Source and Stock or Stock Group		2000	2001	2002	2003	2004	2005	2006	2007	2008	Methodology for 2008 Prediction and Source	
Quillayute Summer	Natural	1.6	0.6	1.2	1.8	1.1	0.8	1.1	1.0	1.1	A variety of methods were used for 2008, primarily based on smolt production and survival. See text in Chapter III and Joint WDFW and tribal annual reports on Puget Sound Coho Salmon Forecast Methodology for details.	
	Hatchery	5.4	5.3	4.9	5.4	6.1	6.1	4.0	6.4	4.2		
North Coast Independent Tributaries	Natural	5.1	8.1	6.4	14.8	12.7	8.5	8.1	3.2	3.2		
	Hatchery	11.7	8.1	8.1	11.0	4.3	5.6	3.2	4.1	5.0		
WA Coast Total	Natural	83.7	133.8	157.3	215.5	266.7	224.5	164.9	136.4	124.5		
	Hatchery	146.4	152.7	155.5	199.9	191.9	198.0	154.1	181.6	135.7		
Puget Sound												
Strait of Juan de Fuca	Natural	13.5	21.4	21.2	20.1	35.7	20.7	26.1	29.9	24.1		
	Hatchery	13.6	14.4	14.0 ^{a/}	24.0 ^{a/}	28.7 ^{a/}	26.5 ^{a/}	20.5	18.4	9.5		
Nooksack-Samish	Natural	14.9	12.4	22.0	16.4	27.5	17.0	18.3	5.2	14.8		
	Hatchery	65.5	44.4	105.4	66.2	75.5	89.5	81.1	53.1	47.1		
Skagit	Natural	30.2	87.2	98.5	116.6	155.8	61.8	106.6	26.8	61.4		
	Hatchery	10.3	10.1	14.1	10.4	22.8	9.1	22.5	8.9	18.3		
Stillaguamish	Natural	17.7	24.4	19.7	37.8	38.0	56.7	45.0	69.2	31.0		
	Hatchery	-	-	-	1.3	0.5	0.2	1.2	0.0	0.1		
Snohomish	Natural	53.0	129.6	123.1	203.0	192.1	241.6	139.5	98.9	92.0		
	Hatchery	62.1	60.9	60.3	35.4	48.3	59.1	96.4	25.7	53.5		
South Sound	Natural	11.7	29.5	40.4	103.6	61.3	45.7	45.3	18.2	27.3		
	Hatchery	121.8	172.6	222.5	315.6	288.4	222.2	256.1	181.7	170.0		
Hood Canal	Natural	61.0	62.0	34.9	32.4	98.7	98.4	59.4	42.4	30.4		
	Hatchery	38.5	33.5	31.3 ^{a/}	48.0 ^{a/}	43.1 ^{a/}	60.6 ^{a/}	57.9	54.8	35.0		
Puget Sound Total	Natural	202.0	366.5	359.8	529.9	609.2	541.9	440.2	290.6	281.0		
	Hatchery	311.8	335.9	447.6	501.0	507.3	465.2	535.7	342.6	333.5		

a/ Strait of Juan de Fuca and Hood Canal Hatchery numbers in 2002-2005 include natural coho from secondary (hatchery) management zones.

b/ Program ended in 2005.

TABLE I-3. Achievement of conservation objectives for natural stocks listed in Table 3-1 of the Pacific Coast Salmon Plan. Bolded numbers indicate a failure to meet the conservation objective. Stocks listed under the Endangered Species Act are not included. (Page 1 of 2)

Stock and Conservation Objective (thousands of spawners; spawners per mile; impact or replacement rate)	Observed or Projected Conservation Achievement (postseason estimates of thousands of spawners or spawners per mile; preseason or postseason impact or replacement rate)										Overfishing Criteria		
	CHINOOK	2000	2001	2002	2003	2004	2005	2006	2007 ^{ai}	2008 ^{bi}	Alert ^{ci}	Concern ^{di}	Exception ^{ei}
Sacramento River Fall 122.0 - 180.0 adult spawners	416.8	546.1	775.5	521.6	283.6	394.0	267.9	88.0	68.4	No	No	No	
Klamath River Fall - < 66%-67% avg. spawner reduction rate but no less than 35.0 adult natural spawners annually	82.7	77.8	65.6	87.6	24.1	26.8	30.2	59.7	26.9	No	Yes	No	
Southern, Central and Northern Oregon Coast Spring and Fall No less than 60 adult spawners/mile ^{fi}	85.0	203.0	268.0	297.0	211.0	118.0	106.0	42.0	NA	No	No	No	
Upper Columbia River Bright Fall 43.5 adults over McNary Dam Council area base period impacts <4%	66.4	110.5	141.7	180.0	170.6	134.8	91.0	58.7	>43.5	No	No	Exp. Rate	
Columbia River Summer Chinook 80.0 to 90.0 adults over Bonneville Dam Council area base period impacts <2%	30.6	76.2	127.4	114.8	NA	NA	NA	NA	NA	NA	NA	NA	
In 2004 state and tribal co-managers changed the stock definition from Chinook passing Bonneville Dam after May 31 to Chinook passing Bonneville Dam after June 14, and the goal changed to 29,000 at the river mouth	23.2	54.9	92.8	83.1	65.4	60.1	76.2	37.2	>29.0	No	No	Exp. Rate	
Grays Harbor Fall - 14.6 adult spawners (MSP)	9.3	9.5	11.3	19.4	31.8	19.5	17.1	NA	NA ^{gi}	No	No	Exp. Rate	
Grays Harbor Spring - 1.4 adult spawners	3.1	2.9	2.6	1.9	5.0	2.1	2.5	NA	NA ^{gi}	No	No	Exp. Rate	
Queets Fall - no less than 2.5 adult spawners (MSY)	3.6	2.3	2.1	4.1	3.6	3.1	2.3	1.9	NA ^{gi}	No	No	Exp. Rate	
Queets Spring/Summer - no less than 0.7 adult spawners	0.2	0.5	0.7	0.2	0.6	0.3	0.3	0.4	NA ^{gi}	Limited ^{ei}	No	Exp. Rate	
Hoh Fall - no less than 1.2 adult spawners (MSY)	1.7	2.6	4.4	1.6	3.2	4.2	1.5	1.7	2.9	No	No	Exp. Rate	
Hoh Spring/Summer - no less than 0.9 adult spawners	0.5	1.2	2.5	1.2	1.8	1.2	0.9	0.8	0.9	No	No	Exp. Rate	
Quillayute Fall - no less than 3.0 adult spawners (MSY)	3.7	5.1	6.1	7.4	3.8	6.4	5.6	2.9	5.5	No	No	Exp. Rate	
Quillayute Spring/Summer - 1.2 adult spawners (MSY)	1.0	1.2	1.0	1.2	1.1	0.9	0.6	NA	2.5	Limited ^{ei}	No	Exp. Rate	

TABLE I-3. Achievement of conservation objectives for natural stocks listed in Table 3-1 of the Pacific Coast Salmon Plan. Bolded numbers indicate a failure to meet the conservation objective. Stocks listed under the Endangered Species Act are not included. (Page 2 of 2)

Stock and Conservation Objective (thousands of spawners; spawners per mile; impact or replacement rate)	Observed or Projected Conservation Achievement (postseason estimates of thousands of spawners or spawners per mile; preseason or postseason impact or replacement rate)									Overfishing Criteria		
	COHO	2000	2001	2002	2003	2004	2005	2006	2007 ^{a/}	2008 ^{b/}	Alert ^{c/}	Concern ^{d/}
Grays Harbor - 35.4 adult spawners (MSP)	38.1	79.1	108.7	83.9	60.7	44.1	14.4	23.7	>35.4	No	No	No
Queets - 5.8 to 14.5 adult spawners (MSY range) Includes supplemental adults prior to 2006.	8.6	24.9	13.8	10.6	8.7	6.5	5.4	5.3	>5.8	No	No	No
Hoh - 2.0 to 5.0 adult spawners (MSY range)	6.8	10.8	9.0	6.3	4.7	4.7	1.3	3.1	>2.0	No	No	No
Quillayute Fall - 6.3 to 15.8 adult spawners (MSY range)	13.3	18.9	23.0	14.8	13.4	11.5	5.6	5.6	>6.3	No	No	No
Western Strait of Juan de Fuca - 11.9 adult spawners	16.9	34.3	20.6	12.4	12.0	6.8	>11.9	>11.9	>11.9	No	No	No
Eastern Strait of Juan de Fuca - 0.95 adult spawners	2.1	2.6	2.5	2.9	8.5	3.4	>0.95	>0.95	>0.95	No	No	No
Hood Canal - 21.5 adult spawners (MSP)	27.2	94.8	69.3	170.3	146.9	38.1	13.8	>21.5	15.0	No	No	No
Skagit - 30.0 adult spawners (MSP)	62.9	87.0	56.0	69.2	138.8	34.7	14.5	>30.0	41.5	No	No	No
Stillaguamish - 17.0 adult spawners (MSP)	28.3	73.6	27.3	45.7	59.2	25.8	8.5	38.7	20.4	No	No	No
Snohomish - 70.0 adult spawners (MSP)	94.2	261.8	161.6	182.7	252.8	109.0	75.8	117.9	61.9	No	No	No

a/ Preliminary data.

b/ Preliminary approximations based on preseason abundance projections and last year's regulations or season structures.

c/ Conservation Alert - triggered during the annual preseason process if a natural stock or stock complex, listed in Table 3-1 of the salmon FMP, is projected to fall short of its conservation objective (MSY, MSY proxy, MSP, or floor in the case of some harvest rate objectives [e.g., 35,000 natural Klamath River fall Chinook spawners]).

Actions for Stocks that are not Exceptions - The Council will close salmon fisheries within its jurisdiction which impact the stocks, except in the case of Washington coastal and Puget Sound salmon stocks and fisheries managed under U.S. District Court orders. In these cases, the Council may allow fisheries which meet annual spawner targets developed through relevant U.S. v. Washington, Hoh v. Baldrige, and subsequent U.S. District Court ordered processes and plans, that may vary from the MSY or MSP conservation objectives. For all natural stocks that meet the conservation alert criteria, the Council will notify pertinent fishery and habitat managers, advising that the stock may be temporarily depressed or approaching an overfishing concern (depending on its recent conservation status), and request state and tribal fishery managers identify the probable causes, if known. If the stock has not met its conservation objective in the previous two years, the Council will request state and tribal managers to do a formal assessment of the primary factors leading to the shortfalls and report to the Council no later than the March meeting prior to the next salmon season.

d/ Overfishing concern - triggered if, in three consecutive years, the postseason estimates indicate a natural stock, listed in Table 3-1 of the salmon FMP, has fallen short of its conservation objective (MSY, MSP, or spawner floor as noted for some harvest rate objectives).

Actions required for Stocks that are not Exceptions - Within one year, the STT to recommend and the Council to adopt management measures to end the overfishing concern and recover the stock in as short a time as possible, preferably within ten years or less. The HC to provide recommendations for habitat restoration and enhancement measures within a suitable time frame.

e/ Exception -application of the conservation alert and overfishing criteria and subsequent Council actions do not apply for (1) hatchery stocks, (2) natural stocks with a cumulative adult equivalent exploitation rate of less than 5% in ocean fisheries under Council jurisdiction during the FRAM base periods, and (3) stocks listed under the ESA.

Conservation Alert and Overfishing Concern Actions for Natural Stocks that are Exceptions (those with exploitation rates limited to less than 5% in base period Council-area ocean fisheries) - Use the expertise of STT and HC to confirm negligible impacts of proposed Council fisheries, identify factors which have led to the decline or low abundance (e.g., fishery impacts outside Council jurisdiction, or degradation or loss of essential fish habitat) and monitor abundance trends and total harvest impact levels. Council action will focus on advocating measures to improve stock productivity, such as reduced interceptions in non-Council managed fisheries, and improvements in spawning and rearing habitat, fish passage, flows, and other factors affecting overall stock survival.

f/ Based on the sum of south/local and north migrating spawners per mile weighted by the total number of miles surveyed for each of the two components (2.2 miles for south/local and 7.5 miles for northern stocks).

g/ Preseason forecasts are not available for Washington coastal Chinook stocks.







FIGURE I-1. Selected preseason vs. postseason forecasts for Chinook stocks with significant contribution to Council area fisheries.







FIGURE I-2a. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.







FIGURE I-2b. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.

CHAPTER II - CHINOOK SALMON ASSESSMENT

CHINOOK STOCKS SOUTH OF CAPE FALCON

SACRAMENTO RIVER FALL CHINOOK SALMON

Predictor Description

The Council's Salmon FMP sets the escapement goal for Sacramento River fall Chinook as a range from 122,000 to 180,000 adults. This stock comprises approximately 80-90 percent of the escapement of all Chinook stocks that return to Central Valley streams and hatcheries. The Central Valley Index (CVI), which provides an annual index of abundance for the combined Central Valley Chinook stocks, is the sum of ocean fishery Chinook harvests in the area south of Point Arena plus the Central Valley adult Chinook spawning escapement (Table II-1). The CVI harvest index is the ocean harvest landed south of Point Arena divided by the CVI, and has varied significantly since it was first calculated in 1970 (Table II-1). From 1970–1986 it tracked ocean harvest and ranged from 0.50–0.73. From 1987–1995 it held steady at 0.70–0.79, while ocean harvest ebbed to a low in 1992. From 1996–2006 it again tracked ocean harvest in a generally declining pattern, reaching a low of 0.26 in 2001. The CVI harvest index was 0.48 in 2007.

Prior to 1989, the STT based its projection of the CVI on recent CVI levels (with general consideration given for brood year natural escapements), hatchery releases, and the previous year jack returns. Between 1989 and 1991, several predictors of the CVI were evaluated, including weight and number of juveniles in hatchery releases and previous year jack returns. Since 1991, the STT has used a linear regression of the CVI on the previous year's Central Valley age-2 return to forecast the CVI (Figure II-1). The 2005 data point was excluded from the CVI predictor in 2008 because it has excessive leverage on the predictor and is not informative to prediction of the CVI at lower jack abundances (see Appendix D for details).

Predictor Performance

For the 1985–2007 period, the CVI preseason forecast ranged from 0.49 to 2.16 times its postseason value (Table II-2). The 2007 CVI preseason forecast of 499,900 fish was about 2.16 times greater than its postseason estimate of 232,000 fish (Table II-2).

2008 Stock Status

A total of 5,939 age-2 Chinook are estimated to have returned to the Central Valley in 2007, the lowest return on record. Sacramento River fall Chinook normally make up the majority of the Central Valley stock, but this year they constituted only 32% of the age-2 return used to forecast the CVI. The resulting 2008 CVI forecast is 157,100 adult Chinook (Figure II-1), and is the lowest CVI forecast on record.

Evaluation of 2007 Regulations on 2008 Stock Abundance

A repeat of 2007 regulations is expected to result in a CVI harvest index equal to last year (0.48). Applying the complement of this fraction (1-0.48) to the 2008 CVI forecast of 157,100 fish and multiplying that quantity by the typical proportion of Central Valley adult Chinook spawners that are Sacramento River fall run fish (0.84, five-year average), yields a 2008 adult escapement forecast of 68,400 Sacramento River fall Chinook, which is well below the lower end of the escapement goal range (Figure II-2).

KLAMATH RIVER FALL CHINOOK

Predictor Description

For Klamath River fall Chinook, linear regressions are used to relate September 1 ocean abundance estimates of age-3, age-4, and age-5 fish to that year's river run size estimates of age-2, age-3, and age-4 fish, respectively (Table II-3). Historical abundance estimates were derived from a cohort analysis of CWT information (brood years 1979-2003). The y-intercept of the regressions is constrained to zero, which gives the biologically reasonable expectation that a river run size of zero predicts an ocean abundance remainder of zero for the same cohort. The abundance of age-2 fish is not forecasted because no precursor to age-2 fish of that brood is available. Ocean fisheries harvest small numbers of age-2 Klamath River fall Chinook.

Predictor Performance

Since 1985, the preseason ocean abundance forecasts for age-3 fish have ranged from 0.33 to 2.72 times the postseason estimates; for age-4 fish from 0.47 to 2.60 times the postseason estimates; and for the adult stock as a whole from 0.34 to 2.03 times the postseason estimates (Table II-4). The September 1, 2006 age-3 forecast (515,400) was 0.99 times its postseason estimate (521,400). The age-4 forecast (26,100) was the lowest on record and was 0.80 times its postseason estimate (32,500); and the age-5 forecast (4,700) was 1.63 times its postseason estimate (2,900) (Table II-4).

Management of Klamath River fall Chinook harvest since 1986 has attempted to achieve specific harvest rates on fully-vulnerable age-4 and age-5 fish in ocean and river fisheries (Table II-5). The Council has used a combination of quotas and time/area restrictions in ocean fisheries in an attempt to meet the harvest rate objective set each year. Since 1992, fisheries have been managed to achieve 50/50 allocation between tribal and non-tribal fisheries. Tribal and recreational river fisheries have been managed on the basis of adult Chinook quotas.

The Council's FMP conservation objective for Klamath River fall Chinook (Amendment 9) permits a natural spawner reduction rate via fisheries of no more than 0.67, with a minimum escapement of 35,000 natural spawning adults. The plan allows for any ocean and river harvest allocation that meets the spawner reduction rate constraint, provided it also meets the minimum escapement goal. The regulations adopted in 2007 were expected to result in 35,000 natural spawning adults and an age-4 ocean harvest rate of 16.0 percent. Postseason estimates of these quantities were 59,700 natural spawning adults and, an age-4 ocean harvest rate of 21.0 percent (Table II-6).

2008 Stock Status

The forecast September 1, 2007 (preseason) ocean abundance of Klamath River fall Chinook salmon is 31,600 age-3 fish, the age-4 forecast is 157,200 and the age-5 forecast is 1,900 fish.

Late-season ocean fisheries in 2007 (September-November) were estimated to have harvested zero age-3, 3,700 age-4, and 800 age-5 Klamath River fall Chinook. This harvest will be deducted from the ocean fishery's allocation in determining the 2008 allowable ocean harvest.

Evaluation of 2007 Regulations on 2008 Stock Abundance

A repeat of 2007 fishery regulations, including a river recreational harvest allocation of 26 percent (of the nontribal adult harvest) and a tribal allocation of 50 percent (of the overall adult harvest), would be expected to result in 26,900 natural area adult spawners, which fails to meet the spawner floor objective. The forecasted age-4 ocean harvest rate of 16.8 percent also fails to meet the NMFS ESA consultation standard for California coastal Chinook. If the ocean fisheries (recreational and commercial) were closed

from January through August 2008 between Cape Falcon and Point Sur, and the Klamath River fisheries (tribal and recreational) were closed in 2008, the expected number of natural area adult spawners would be 74,300, with an expected age-4 ocean harvest rate of 2.4 percent (due to ocean harvest that already occurred in the September through November 2007 period).

OTHER CALIFORNIA COASTAL CHINOOK STOCKS

Other California coastal streams that support fall Chinook stocks, which contribute to ocean fisheries off Oregon and California, include the Smith, Little, Mad, Eel, and Mattole rivers, and Redwood Creek. Except for the Smith River, these stocks are included in the California coastal Chinook ESU, which is listed as threatened under the ESA. Current information is insufficient to forecast the ocean abundance of these stocks, however, the NMFS ESA consultation standard restricts the Klamath River fall Chinook age-4 ocean harvest rate to no more than 16.0 percent to limit impacts on these stocks. As indicated in the previous section, the postseason estimate of this rate for 2007 is 21.0 percent, with a preseason forecast of 16.0 percent. If the ocean fishery was closed from January through August 2008 between Cape Falcon and Point Sur, the expected age-4 ocean harvest rate for 2008 would be 2.4 percent (due to ocean harvest that already occurred in the September through November 2007 period).

OREGON COASTAL CHINOOK STOCKS

Oregon coastal Chinook stocks are categorized into two major subgroups based on ocean migration patterns. Although their ocean harvest distributions overlap somewhat, they have been labeled as either north or south/local migrating.

North Migrating Chinook

North migrating Chinook stocks include stocks north of and including the Elk River, with the exception of Umpqua River spring Chinook. Based on CWT analysis, the populations from ten major North Oregon Coast (NOC) river systems from the Nehalem through the Siuslaw Rivers are harvested primarily in ocean fisheries off British Columbia, Canada and Southeast Alaska, and to a much lesser degree in Council area and terminal area (state waters) fisheries off Washington and Oregon. CWT analysis indicates populations from five major mid-Oregon Coast (MOC) systems, from the Coos through the Elk Rivers, are harvested primarily in ocean fisheries off British Columbia, Canada, Washington, and Oregon with minor contributions to California fisheries.

Predictor Description and 2008 Stock Status

Quantitative abundance predictions are not made for these stocks for use in annual development of Council area fishery regulations. Qualitative expectations of abundance are based on parental year spawner escapements and hatchery indicator stock data used in the PSC management process.

Natural spawner escapement is assessed yearly from the Nehalem through Sixes rivers. Peak spawning counts of adults are obtained from standard index areas on these rivers and monitored to assess stock trends (*Review of 2007 Ocean Salmon Fisheries*, Chapter II, Table II-4 and Figure II-3). Natural fall Chinook stocks from both the NOC and MOC dominate production from this subgroup. Also present in lesser numbers are naturally-produced spring Chinook stocks from several rivers, and hatchery fall and/or spring Chinook released in the Trask, Nestucca, Salmon, Alsea, and Elk Rivers.

North Oregon Coast

Since 1986, the Salmon River Hatchery production has been CWT'd for use primarily as a PSC indicator stock for the NOC stock component. Because these fish are harvested in fisheries north of the Council management area, the STT has not reviewed the procedure by which this indicator stock is used in estimating annual stock status. The annual spawner counts have been decreasing since 2002 despite

excellent parental escapements indices in 2001 to 2004 (*Review of 2007 Ocean Salmon Fisheries*, Appendix B, Table B-11). If this trend continues, the 2008 NOC stock abundance is expected to be less than the 2007 abundance.

Mid-Oregon Coast

Since 1992, the Elk River Hatchery production has been CWT'd for use as a PSC indicator stock for the MOC stock component. Age specific ocean abundance forecasts for 2008 are not currently available. The STT has not undertaken a review of the methods used by Oregon Department of Fish and Wildlife (ODFW) staff in preparing these abundance forecasts.

The annual spawner counts have been decreasing since 2004 despite excellent parental escapements indices in 2001 to 2004 (*Review of 2007 Ocean Salmon Fisheries*, Appendix B, Table B-11). If this trend continues, the 2008 MOC stock abundance is expected to be less than the 2007 abundance.

Based on the density index of total spawners, the generalized expectation for NOC and MOC stocks in 2008 is below recent years average abundance. Specifically, the 2007 spawner density in standard survey areas for the NOC averaged 23 spawners per mile; well below the lower bound of the FMP aggregate goal of 60 to 90 spawners per mile. Moreover, escapements in the NOC escapement indicator basins of the Nehalem, Siletz, and Siuslaw have failed to achieve PSC agreed-to escapement goals in 2007. The escapement of fall Chinook to the Nehalem basin has failed to reach its PSC agreed-to escapement goal (6,989) for the past 2 years. The MOC average spawner per mile from standard survey areas was 20 adult spawners per mile, again, well below the goal of 60 to 90 spawners per mile. Fall Chinook escapement goals are currently under development for the South Umpqua and Coquille basins of the MOC. (*Review of 2007 Ocean Salmon Fisheries*, Appendix B, Table B-11).

South/Local Migrating Chinook

South/local migrating Chinook stocks include Rogue River spring and fall Chinook, fall Chinook from smaller rivers south of the Elk River, and Umpqua River spring Chinook. These stocks are important contributors to ocean fisheries off Oregon and northern California. Umpqua River spring Chinook contributes to a lesser degree to fisheries off Washington, British Columbia, Canada, and southeast Alaska.

Rogue River fall Chinook contribute to ocean fisheries principally as age-3 through age-5 fish. Mature fish enter the river each year from mid-July through October, with the peak of the run occurring during August and September.

Umpqua and Rogue rivers spring Chinook contribute to ocean fisheries primarily as age-3 fish. Mature Chinook enter the rivers primarily during April and May and generally prior to annual ocean fisheries. Quantitative abundance predictions are not made for these stocks.

Natural fall Chinook stocks from river systems south of the Elk River and spring Chinook stocks from the Rogue and Umpqua Rivers dominate production from this subgroup. Also present in lesser numbers are hatchery fall Chinook, primarily from the Chetco River. Substantial releases of hatchery spring Chinook occur in both the Rogue and Umpqua Rivers.

Predictor Description and 2008 Stock Status

Quantitative abundance predictions are not made for these stocks, although an abundance index for Rogue River fall Chinook has been developed. General trends in stock abundance for southern Oregon coastal Chinook stocks are assessed through escapement indices (*Review of 2007 Ocean Salmon Fisheries*, Chapter II, Table II-4 and Figure II-3).

Carcass recovery numbers in Rogue River index surveys that covered a large proportion of the total spawning area were available for 1977-2004. Using Klamath Ocean Harvest Model (KOHM) methodology, these carcass numbers, allocated into age-classes from scale data, were used to estimate the Rogue Ocean Population Index (ROPI) for age-3 to age-5 fish. A linear regression using the escapement estimates (all ages) in year i based on seining at Huntley Park (1976-2003) to predict the ROPI in year $i + 1$ (1977-2004) was developed. The 2007 Huntley Park escapement estimate and the resulting 2008 ROPI forecast was then scaled to the historical carcass survey-based ROPI. The 2008 ROPI forecast (11,600) consisting of age-3 (6,600), age-4 (4,300) and age-5 (700) are based on the average annual age-class strengths of the carcass-based ROPIs from 1991-2004. This data-set was truncated at 1991 because significant harvest restrictions that could affect age structure began that year. The 2008 ROPI is lower than the recent year average of 17,400, Table II-7.

Other Stocks

Information is insufficient to forecast the abundance of fall Chinook from other small systems south of the Elk River. These stocks are minor contributors to general season mixed stock ocean fisheries.

Evaluation of 2007 Regulations on 2008 Stock Abundance

The FMP conservation objective for Oregon coast Chinook is 150,000 to 200,000 natural adult spawners, and attainment of this goal is assessed using peak spawner counts of 60 to 90 fish per mile in nine standard index reaches. The aggregate stock has been meeting or exceeding this goal since 1984 and has been generally increasing. However, since reaching a peak in 2003, the escapement has been declining. In 2007, the stock failed to meet its goal for the first time in 23 years. No forecast is available for this stock, but given recent trends, it seems likely that it would fail to meet its goal again in 2008 under 2007 fishing seasons.

CHINOOK STOCKS NORTH OF CAPE FALCON

Columbia River Fall Chinook

Predictor Description and Past Performance

Columbia River fall Chinook stocks typically form the largest contributing stock group to Council Chinook fisheries north of Cape Falcon. Abundance of these stocks is a major factor in determining impacts of fisheries on weak natural stocks critical to Council area management. Abundance predictions are made for five major fall stock units characterized as being hatchery or natural production, and originating above or below Bonneville Dam. The upriver brights (URB) and lower river wild (LRW) are primarily naturally-produced stocks. The lower river hatchery (LRH) tule, Spring Creek Hatchery (SCH) tule, and mid-Columbia brights (MCB) are primarily hatchery-produced stocks. The MCB include the lower river bright (LRB) stock as a small naturally-produced component. LRB spawn in the mainstem Columbia River near Beacon Rock and are believed to have originated from MCB hatchery strays. The tule stocks generally mature at an earlier age than the bright fall stocks and do not migrate as far north. Minor stocks include the Select Area brights (SAB), a Big Creek Hatchery stock originally from Rogue River stock.

Preseason estimates of Columbia River fall Chinook stock abundance, used by the STT to assess the Council's adopted fishery regulations, are based on age-specific and stock-specific forecasts of annual ocean escapement (return to the Columbia River). These forecasts are developed by the Columbia River Technical Advisory Committee (TAC). Columbia River return forecast methodologies used for Council management are identical to those used for planning Columbia River fall season fisheries, although minor updates to Council estimates of inriver run size may occur prior to finalization of the inriver fishery plans.

The 2008 return of each fall Chinook stock group is estimated using relationships between successive age groups within a cohort. The database for these relationships was constructed by combining age-specific estimates of escapement and inriver fishery catches for years since 1964 (except for MCB, which started in 1980). Typically, only the more recent broods are used in the current predictions. Fall Chinook stock identification in the Columbia River mixed stock fisheries is determined by sampling catch and escapement for such factors as CWT recovery and visual stock identification (VSI). Age composition estimates are based on CWT data and scale reading of fishery and escapement samples, where available. These stock and age data for Columbia River fall Chinook are the basis for the return data presented in the *Review of 2007 Ocean Salmon Fisheries* (Appendix B, Tables B-15 through B-20). The 2007 returns for the five fall Chinook stocks listed in this report may differ somewhat from those provided in the *Review of 2007 Ocean Salmon Fisheries*, since ocean escapement estimates may have been updated after that report was printed.

Performance of the preliminary inriver run size estimation methodology can be assessed, in part, by examining the differences between preseason and postseason estimates (Table II-8). The recent 10-year average March preliminary preseason estimates as a percentage of the postseason estimates for the URB, LRW, LRH, SCH, and MCB stock estimates are 1.02, 1.06, 0.84, 1.03, and 0.98 respectively. The only March preliminary preseason estimate to show a bias was LRH, which has been under predicted between 1994 and 2006. The other four stocks have been both over and under predicted.

Ocean escapement estimates developed for the March Council meeting do not take into account variations in marine harvest. The STT combines the initial inriver run size (ocean escapement; Table II-8) with expected Council area fishery harvest levels and stock distribution patterns to produce adjusted ocean escapement estimates based on the proposed ocean fishing regulations. These revised estimates are available at the end of the Council preseason planning process in April and should provide a more accurate prediction of ocean escapement.

2008 Stock Status

The preliminary forecast for 2008 URB fall Chinook ocean escapement is 162,500 adults. If the forecast is realized, it would be about 144 percent of last year's return and about 70 percent of the recent 10-year average of 232,640.

No preseason forecast for 2008 ocean escapement of ESA-listed Snake River wild fall Chinook is currently available. However, the Columbia River TAC is expected to develop a run size estimate for this stock prior to the April Council meeting.

Ocean escapement of LRW fall Chinook in 2008 is forecast at 3,800 adults. If the forecast is realized, it would be about 88 percent of last year's return, and about 26 percent of the recent 10-year average return of 14,890. The forecast is the third lowest since at least 1984, and less than the spawning escapement goal of 5,700 in the North Fork Lewis River.

The preliminary forecast for 2008 ocean escapement of LRH fall Chinook is for a return of 59,000 adults, which would be 180 percent of last year's return and 74 percent of the recent 10-year average of 79,620.

Ocean escapement of SCH fall Chinook in 2008 is forecast at 87,200 adults. If the forecast is realized, it would be about six times last year's return and near the recent 10-year average of 86,820.

The preliminary forecast for the 2008 ocean escapement of MCB fall Chinook is 54,000 adults. If the forecast is realized, it would be about 115 percent of last year's return and about 67 percent of the recent 10-year average of 80,340.

Evaluation of 2007 Regulations on 2008 Stock Abundance

Applying 2007 regulations to the projected 2008 abundance of Columbia River fall Chinook would result in ocean escapements meeting spawning escapement goals for all major stocks except LRW. Compared to actual 2007 returns, the 2008 ocean escapement forecasts are higher for all stocks except LRW. Compared to 2007 forecast ocean escapement, the 2008 forecasts are higher for LRH and SCH, but lower for LRW, URB, and MCB stocks.

Washington Coastal Chinook

Predictor Description and Past Performance

Council fisheries have only minor impacts on Washington coastal Chinook stocks, and except for Willapa Bay Chinook, Hoh River Chinook and Quillayute River Chinook, forecast data is unavailable at the time this report is published; therefore, preseason abundance estimates are not presented. However, abundance estimates are provided for Washington Coastal stocks in subsequent preseason fishery impact assessment reports prepared by the STT.

2008 Stock Status

The 2008 Willapa Bay hatchery fall Chinook ocean escapement abundance forecast is 27,047, which is slightly less than the 2007 prediction of 29,846. The 2008 natural fall Chinook ocean escapement forecast is 2,516, up from last year's 2,012 prediction.

For the Hoh River, the 2008 natural spring/summer Chinook ocean escapement abundance forecast is 892. The natural fall Chinook forecast is predicted to be 2,873.

The 2008 Quillayute hatchery spring Chinook forecast for ocean escapement abundance is 1,745 and the natural summer/fall Chinook abundance forecast is for a return of 6,264.

Puget Sound Chinook

Run-size expectations for various Puget Sound stock management units are listed in Table I-1. A comparison of preseason and postseason forecasts for recent years is detailed in Table II-9. The STT has not undertaken a review of the methods employed by state and tribal staffs in preparing these abundance forecasts. Methodologies for estimates are described in the annual Puget Sound management reports (starting in 1993, reports are available by Puget Sound management unit, not by individual species). Forecasts for Puget Sound stocks generally assume production is dominated by age-4 adults. Puget Sound Chinook were listed as threatened under the ESA in March 1999. Southern U.S. fisheries that impact Puget Sound Chinook are constrained by terms of a Resource Management Plan (RMP), and are exempted from ESA Section 9 take prohibitions under Limit 6 of the 4(d) rule.

2008 Stock Status

Spring Chinook

Spring Chinook originating in Puget Sound are expected to remain depressed. Runs in the Nooksack, Skagit, White, and Dungeness rivers are of particular concern.

Summer/Fall Chinook

The 2008 preliminary forecast for Puget Sound summer/fall stocks is for a return of 245,268 Chinook, slightly higher than the 2007 preseason forecast of 227,300. The 2008 natural Chinook return forecast of 59,154 is higher than the 2007 forecast of 54,000. Changes in the abundance of individual stocks from various production areas are detailed in Table I-1.

Natural stocks from Puget Sound had experienced improved survival in recent years. However, natural returns to several major populations, including Snohomish and Skagit were significantly lower in 2007 than has been observed for recent years. Fishery management for Puget Sound Chinook has changed from an escapement goal basis to the use of stock specific exploitation rates and “critical abundance thresholds.” This new approach is evaluated on an annual basis through the RMP.

Evaluation of 2007 Regulations on 2008 Stock Abundance

Council fisheries north of Cape Falcon have only a minor impact on most stocks that originate in Washington coastal and Puget Sound rivers. These stocks have northerly marine distribution patterns and are therefore impacted primarily by Canadian and Alaskan fisheries. An evaluation of 2007 Council area regulations on projected 2008 abundance would not provide a useful comparison of ocean escapement.

TABLE II-1. Indices of annual abundance and ocean fishery impacts on California Central Valley Chinook in thousands of fish. (Page 1 of 1)

Year	Ocean Chinook Landings South of Pt. Arena			Hatchery and Natural Escapements of Central Valley Adults			CVI Abundance (Ocean Landings + Escapement)		CVI Harvest Index (%) ^{b/}
	Troll	Sport	Total	Fall	Other ^{a/}	Total			
1970	226.8	111.1	337.9	186.3	55.6	241.9	579.8	58	
1971	150.7	166.3	317.0	196.2	65.4	261.6	578.6	55	
1972	229.8	187.6	417.4	104.6	47.6	152.3	569.7	73	
1973	422.5	180.9	603.4	225.4	34.0	259.4	862.8	70	
1974	282.7	141.6	424.3	207.3	42.3	249.6	673.9	63	
1975	234.4	92.7	327.1	162.3	56.5	218.9	546.0	60	
1976	237.9	68.6	306.4	172.0	45.6	217.7	524.1	58	
1977	263.8	76.6	340.4	165.6	43.0	208.6	549.1	62	
1978	291.0	65.9	356.9	129.8	19.9	149.7	506.6	70	
1979	234.1	108.5	342.6	171.9	10.9	182.9	525.5	65	
1980	294.3	77.1	371.4	148.4	34.0	182.4	553.8	67	
1981	289.9	73.8	363.7	196.9	21.8	218.7	582.4	62	
1982	426.1	122.5	548.6	182.4	38.9	221.3	769.9	71	
1983	178.2	53.0	231.2	129.9	14.4	144.3	375.4	62	
1984	221.7	78.7	300.3	205.8	16.9	222.7	523.0	57	
1985	212.3	121.8	334.1	312.7	20.7	333.4	667.4	50	
1986	502.5	114.8	617.3	262.9	41.3	304.1	921.4	67	
1987	446.8	152.8	599.7	202.8	21.6	224.4	824.1	73	
1988	830.5	130.4	960.9	244.9	26.6	271.5	1,232.4	78	
1989	363.8	130.9	494.7	155.0	18.0	173.0	667.7	74	
1990	336.2	112.6	448.8	105.7	14.0	119.7	568.6	79	
1991	254.6	62.1	316.7	118.3	16.4	134.6	451.3	70	
1992	160.3	66.7	227.0	82.6	4.2	86.8	313.8	72	
1993	259.7	99.3	359.0	139.6	6.0	145.7	504.6	71	
1994	290.4	165.8	456.2	169.5	6.6	176.0	632.2	72	
1995	670.6	354.6	1,025.2	302.2	16.5	318.6	1,343.8	76	
1996	348.8	129.3	478.1	307.6	12.9	320.5	798.6	60	
1997	482.2	208.4	690.6	368.0	46.6	414.6	1,105.2	62	
1998	221.6	114.4	336.0	254.2	55.8	310.0	646.0	52	
1999	259.7	76.4	336.1	408.9	21.4	430.3	766.4	44	
2000	447.6	146.4	594.0	459.9	34.9	494.8	1,088.8	55	
2001	172.6	59.9	232.5	575.6	74.0	649.5	882.0	26	
2002	312.9	134.7	447.6	804.4	40.1	844.5	1,292.0	35	
2003	239.0	69.7	308.7	541.7	46.3	588.0	896.7	34	
2004	362.9	175.1	538.0	296.7	34.9	331.6	869.6	62	
2005	287.9	103.5	391.5	415.3	42.9	458.2	849.7	46	
2006	58.9	65.9	124.8	276.5	33.6	310.1	434.9	29	
2007 ^{c/}	88.6	23.0	111.6	90.4	30.0	120.4	232.0	48	

a/ Spring run of the current calendar year and late fall and winter runs of the following calendar year.

b/ Ocean harvest landed south of Pt. Arena as a percent of the CVI.

c/ Preliminary.

TABLE II-2. Comparisons of preseason forecast and postseason estimates for the CVI in thousands of fish. (Page 1 of 1)

Year	Preseason Forecast	Postseason Estimate	Pre/Postseason
1985	524.8	667.4	0.79
1986	546.5	921.4	0.59
1987	592.9	824.1	0.72
1988	707.1	1,232.4	0.57
1989	625-885	667.7	0.94-1.33
1990	500-900	568.6	0.88-1.58
1991	466.0	451.3	1.03
1992	452.0	313.8	1.44
1993	501.0	504.6	0.99
1994	503.0	632.2	0.80
1995	654.0	1,343.8	0.49
1996	533.0	798.6	0.67
1997	849.0	1,105.2	0.77
1998	1,051.0	646.0	1.63
1999	847.7	766.4	1.11
2000	790.4	1,088.8	0.73
2001	649.4	882.0	0.74
2002	825.4	1,292.0	0.64
2003	1,108.1	896.7	1.24
2004	831.8	869.6	0.96
2005	1,678.3	849.7	1.98
2006	632.5	434.9	1.45
2007	499.9	232.0	2.16
2008	157.1 ^{a/}	-	-

a/ CVI predictor excludes 2005 data point.

TABLE II-3. Klamath River fall Chinook ocean abundance (thousands), harvest rate, and river run size estimates (thousands) by age. (Page 1 of 1)

Year (t)	Ocean Abundance Sept. 1 (t-1)			Annual Ocean Harvest Rate Sept. 1 (t-1) - Aug. 31 (t)		Klamath Basin River Run (t)				
	Age-3	Age-4	Total	Age-3	Age-4	Age-2	Age-3	Age-4	Age-5	Total Adults
1981	493.2	57.0	550.2	0.21	0.53	28.2	64.1	14.4	1.8	80.3
1982	566.2	133.4	699.6	0.30	0.52	39.4	30.1	33.9	2.6	66.6
1983	316.5	116.3	432.9	0.19	0.60	3.8	35.9	20.7	0.9	57.5
1984	156.6	83.4	240.0	0.08	0.38	8.3	21.7	24.4	1.1	47.2
1985	376.5	56.6	433.1	0.11	0.24	69.4	32.9	25.7	5.8	64.4
1986	1,305.8	141.8	1,447.6	0.18	0.46	44.6	162.9	29.8	2.3	195.0
1987	782.0	342.6	1,124.6	0.16	0.43	19.1	89.7	112.6	6.8	209.1
1988	756.9	235.5	992.4	0.20	0.39	24.1	101.2	86.5	3.9	191.6
1989	370.3	177.7	548.0	0.15	0.36	9.1	50.4	69.6	4.3	124.3
1990	176.1	104.1	280.3	0.30	0.55	4.4	11.6	22.9	1.3	35.9
1991	69.4	37.2	106.6	0.03	0.18	1.8	10.0	21.6	1.1	32.7
1992	39.5	28.2	67.7	0.02	0.07	13.7	6.9	18.8	1.0	26.7
1993	168.5	15.0	183.5	0.05	0.16	7.6	48.3	8.2	0.7	57.2
1994	119.9	41.7	161.6	0.03	0.09	14.4	37.0	26.0	1.0	64.0
1995	784.3	28.7	813.0	0.04	0.14	22.8	201.9	18.3	2.6	222.8
1996	192.3	225.5	417.8	0.05	0.16	9.5	38.8	136.7	0.3	175.8
1997	140.4	62.8	203.3	0.01	0.06	8.0	35.0	44.2	4.6	83.7
1998	154.8	44.9	199.7	0.00	0.09	4.6	59.2	29.7	1.7	90.6
1999	129.4	30.5	159.8	0.01	0.09	19.2	29.2	20.5	1.3	51.0
2000	617.6	44.3	661.9	0.06	0.10	10.2	187.1	30.5	0.5	218.1
2001	357.1	133.9	491.0	0.03	0.09	11.3	99.1	88.2	0.2	187.4
2002	514.5	99.5	614.0	0.02	0.15	9.2	94.6	62.5	3.7	160.8
2003	401.1	192.6	593.7	0.08	0.21	3.8	94.3	96.8	0.9	191.9
2004	160.2	105.3	265.6	0.12	0.34	9.7	33.2	40.7	5.3	79.2
2005	190.6	38.2	228.8	0.02	0.20	2.3	43.8	17.5	3.9	65.2
2006	88.7 ^{a/}	63.4	152.1	0.01 ^{a/}	0.10	26.9	18.5	41.6	1.3	61.4
2007	521.4 ^{b/}	32.5 ^{a/}	553.9	NA ^{c/}	0.21 ^{a/}	1.7	112.2	16.7	1.6	130.5

a/ Preliminary: incomplete cohort data (age-5 unavailable).

b/ Preliminary: incomplete cohort data (age-4 and age-5 unavailable).

c/ Not estimated: incomplete cohort data (age-4 and age-5 unavailable).

TABLE II-4. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 1 of 2)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
Age-3			
1985	113,000	276,000	0.41
1986	426,000 ^{b/}	1,305,782	0.33
1987	511,800	782,032	0.65
1988	370,800	756,908	0.49
1989	450,600	370,328	1.22
1990	479,000	176,133	2.72
1991	176,200	69,442	2.54
1992	50,000	39,485	1.27
1993	294,400	168,473	1.75
1994	138,000	119,913	1.15
1995	269,000	784,279	0.34
1996	479,800	192,290	2.50
1997	224,600	140,421	1.60
1998	176,000	154,819	1.14
1999	84,800	129,355	0.66
2000	349,600	617,573	0.57
2001	187,200	357,085	0.52
2002	209,000	514,524	0.41
2003	171,300	401,092	0.43
2004	72,100	160,243	0.45
2005	185,700	190,568	0.97
2006	44,100	88,652	0.50
2007 ^{c/}	515,400	521,412	0.99
2008	31,600	-	-
Age-4			
1985	56,875	57,500	0.99
1986	66,250	141,772	0.47
1987	206,125	342,555	0.60
1988	186,375	235,535	0.79
1989	215,500	177,655	1.21
1990	50,125	104,131	0.48
1991	44,625	37,172	1.20
1992	44,750	28,181	1.59
1993	39,125	15,028	2.60
1994	86,125	41,736	2.06
1995	47,000	28,725	1.64
1996	268,500	225,526	1.19
1997	53,875	62,830	0.86
1998	46,000	44,889	1.02
1999	78,750	30,468	2.58
2000	38,875	44,346	0.88
2001	247,000	133,869	1.85
2002	143,800	99,464	1.45
2003	132,400	192,598	0.69
2004	134,500	105,346	1.28
2005	48,900	38,239	1.28
2006	63,700	63,446	1.00
2007 ^{c/}	26,100	32,494	0.80
2008	157,200	-	-

TABLE II-4. Comparisons of preseason forecasts and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 2 of 2)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
Age-5			
1985	NA	11,187	NA
1986	NA	5,855	NA
1987	5,250	19,443	0.27
1988	13,250	14,669	0.90
1989	10,125	9,627	1.05
1990	7,625	7,776	0.98
1991	1,500	2,774	0.54
1992	1,250	1,444	0.87
1993	1,125	1,759	0.64
1994	500	1,462	0.34
1995	2,000	3,805	0.53
1996	1,125	787	1.43
1997	7,875	8,859	0.89
1998	3,250	2,389	1.36
1999	2,000	2,106	0.95
2000	1,375	1,051	1.31
2001	1,250	258	4.84
2002	9,700	6,970	1.39
2003	6,500	1,917	3.39
2004	9,700	17,196	0.56
2005	5,200	6,893	0.75
2006	2,200	5,242	0.42
2007 ^{c/}	4,700	2,886	1.63
2008	1,900	-	-
Total Adults			
1985	169,875	344,687	0.49
1986	492,250	1,453,409	0.34
1987	723,175	1,144,030	0.63
1988	570,425	1,007,112	0.57
1989	676,225	557,610	1.21
1990	536,750	288,040	1.86
1991	222,325	109,388	2.03
1992	96,000	69,110	1.39
1993	334,650	185,260	1.81
1994	224,625	163,111	1.38
1995	318,000	816,809	0.39
1996	749,425	418,603	1.79
1997	286,350	212,110	1.35
1998	225,250	202,097	1.11
1999	165,550	161,929	1.02
2000	389,850	662,970	0.59
2001	435,450	491,212	0.89
2002	362,500	620,958	0.58
2003	310,200	595,607	0.52
2004	216,300	282,785	0.76
2005	239,800	235,700	1.02
2006	110,000	157,340	0.70
2007 ^{c/}	546,200	556,792	0.98
2008	190,700	-	-

a/ Original preseason forecasts for years 1985-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the assumed May 1 (t) number by the Sept. 1 (t-1) through May 1 (t) survival rate in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ A scalar of 0.75 was applied to the jack count because, (1) most jacks returned to the Trinity River, and (2) the jack count was outside the database range.

c/ Preliminary.

TABLE II-5. Summary of management objectives and predictor performance for Klamath River fall Chinook. (Page 1 of 1)

Year(t)	Preseason Ocean Abundance Forecast ^{a/}		Postseason Ocean Abundance Estimate		Preseason Age-4 Harvest Rate Forecast ^{b/}		Postseason Age-4 Harvest Rate Estimate ^{c/}		Preseason Adult Harvest Forecast		Postseason Adult Harvest Estimate	
	Sept. 1 (t-1)		Sept. 1 (t-1)		Harvest Rate Forecast ^{b/}		Harvest Rate Estimate ^{c/}		Harvest Forecast		Harvest Estimate	
	Age-3	Age-4	Age-3	Age-4	Ocean	River	Ocean	River	Ocean	River	Ocean	River
1986	426,000	66,250	1,305,782	141,772	0.28	0.50	0.46	0.67	72,000	37,700	304,512	46,154
1987	511,800	206,125	782,032	342,555	0.28	0.53	0.43	0.44	121,200	78,200	277,104	73,265
1988	370,800	186,375	756,908	235,535	0.31	0.53	0.39	0.52	114,100	65,400	254,444	73,854
1989	450,600	215,500	370,328	177,655	0.30	0.49	0.36	0.70	128,100	67,600	125,523	54,340
1990	479,000	50,125	176,133	104,131	0.30	0.49	0.55	0.36	85,100	31,200	114,911	11,459
1991	176,200	44,625	69,442	37,172	0.13	0.28	0.18	0.45	16,700	12,800	9,871	13,581
1992	50,000	44,750	39,485	28,181	0.06	0.15	0.07	0.27	4,200	4,200	3,140	6,787
1993	294,400	39,125	168,473	15,028	0.12	0.43	0.16	0.49	20,100	22,500	11,354	12,808
1994	138,000	86,125	119,913	41,736	0.07	0.20	0.09	0.29	10,400	14,300	8,889	13,524
1995	269,000	47,000	784,279	28,725	0.07	0.32	0.14	0.19	13,500	18,500	32,230	21,637
1996	479,800	268,500	192,290	225,526	0.17	0.66	0.16	0.39	88,400	129,100	45,147	69,241
1997	224,600	53,875	140,421	62,830	0.10	0.43	0.06	0.26	17,600	26,500	8,657	17,764
1998	176,000	46,000	154,819	44,889	0.07	0.29	0.09	0.30	10,200	14,800	5,012	17,897
1999	84,800	78,750	129,355	30,468	0.10	0.28	0.09	0.45	12,300	18,100	5,126	16,942
2000	349,600	38,875	617,573	44,346	0.11	0.53	0.10	0.25	24,000	32,400	42,336	35,066
2001	187,200	247,000	357,085	133,869	0.14	0.61	0.09	0.29	45,600	105,300	21,783	50,780
2002	209,000	143,800	514,524	99,464	0.13	0.57	0.15	0.26	30,000	70,900	29,436	35,069
2003	171,300	132,400	401,092	192,598	0.16	0.50	0.21	0.28	30,600	52,200	71,124	39,715
2004	72,100	134,500	160,243	105,346	0.15	0.38	0.34	0.48	26,500	35,800	64,264	29,807
2005	185,700	48,900	190,568	38,239	0.08	0.16	0.20	0.19	7,100	9,600	13,228	10,001
2006	44,100	63,700	88,652	63,446	0.11	0.23	0.10	0.18	10,000	10,000	10,457	10,345
2007 ^{d/}	515,400	26,100	521,412	32,494	0.16	0.63	0.21	0.56	30,200	51,400	28,551	33,282
2008	31,600	157,200	-	-	-	-	-	-	-	-	-	-

a/ Original preseason forecasts for years 1986-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the May 1 (t) number by the assumed Sept. 1 (t-1) through May 1 (t) survival rate assumed in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ Ocean harvest rate forecast is the fraction of the predicted ocean abundance expected to be harvested Sept. 1 (t-1) through August 31(t). River harvest rate forecast is the fraction of the predicted river run expected to be harvested in river fisheries. Original ocean harvest rate forecasts for year (t), 1986-2001, were based on a May 1 (t) ocean abundance denominator; converted to Sept. 1 (t-1) abundance denominator by multiplying former values by 0.8 (the assumed age-4 survival rate between Sept. 1 (t-1) and May 1 (t) in those years).

c/ Ocean harvest rate is the fraction of the postseason ocean abundance harvested Sept. 1 (t-1) through August 31 (t). River harvest rate is the fraction of the river run harvested by river fisheries.

d/ Postseason estimates are preliminary.

TABLE II-6. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 1 of 2)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))							River Fisheries (t)		
	KMZ			North of	South of	Ocean		Net	Sport	Total
	Troll	Sport	Subtotal	KMZ	KMZ	Subtotal	Total			
HARVEST (numbers of fish)										
Age-3										
1986	35,630	4,876	40,506	73,913	122,913	196,826	237,332	8,100	18,100	26,200
1987	17,231	5,083	22,314	42,875	56,362	99,237	121,551	11,400	11,400	22,800
1988	15,996	5,164	21,160	24,312	107,949	132,261	153,421	12,500	15,600	28,100
1989	6,462	11,793	18,255	15,368	23,750	39,118	57,373	2,700	900	3,600
1990	81	4,357	4,438	36,578	11,006	47,584	52,022	1,300	1,400	2,700
1991	0	1,022	1,022	343	810	1,153	2,175	2,123	1,277	3,400
1992	0	0	0	971	0	971	971	970	251	1,221
1993	0	822	822	833	6,424	7,257	8,079	5,426	2,917	8,343
1994	42	604	646	0	3,387	3,387	4,033	4,543	965	5,508
1995	0	999	999	12,211	14,808	27,019	28,018	11,840	5,536	17,376
1996	0	0	0	0	9,312	9,312	9,312	12,363	3,661	16,024
1997	0	232	232	620	1,215	1,835	2,067	2,166	2,736	4,902
1998	0	6	6	298	466	764	770	2,231	5,781	8,012
1999	63	180	243	1,262	433	1,695	1,938	4,981	1,748	6,729
2000	404	3,282	3,686	8,730	25,206	33,936	37,622	22,458	4,893	27,351
2001	113	105	218	2,765	6,088	8,853	9,071	17,885	7,294	25,179
2002	220	783	1,003	1,623	9,912	11,535	12,538	11,734	6,258	17,992
2003	173	679	852	2,026	27,312	29,338	30,190	6,996	5,061	12,057
2004	403	971	1,374	9,902	7,337	17,239	18,613	4,679	2,051	6,730
2005	0	568	568	889	2,381	3,270	3,838	4,394	1,641	6,035
2006 ^{al}	0	465	465	31	332	363	828	2,388	13	2,401
2007 ^{al}	719	7,518	8,237	4,111	8,649	12,760	20,997	17,422	5,356	22,778
Age-4										
1986	7,797	1,120	8,917	23,560	32,131	55,691	64,608	17,000	2,900	19,900
1987	21,727	4,427	26,154	71,123	48,812	119,935	146,089	41,000	8,500	49,500
1988	11,867	3,598	15,465	26,950	50,278	77,228	92,693	38,600	6,200	44,800
1989	6,062	9,735	15,797	32,428	16,608	49,036	64,833	41,000	7,700	48,700
1990	4,000	2,916	6,916	39,760	10,608	50,368	57,284	6,000	2,200	8,200
1991	0	1,001	1,001	1,513	4,135	5,648	6,649	7,593	2,016	9,609
1992	171	55	226	1,781	12	1,793	2,019	4,360	723	5,083
1993	0	0	0	849	1,615	2,464	2,464	3,786	243	4,029
1994	0	1,124	1,124	1,168	1,499	2,667	3,791	6,666	818	7,484
1995	0	242	242	1,879	1,771	3,650	3,892	2,957	480	3,437
1996	773	3,464	4,237	10,336	20,738	31,074	35,311	43,959	9,080	53,039
1997	3	172	175	463	2,995	3,458	3,633	8,734	2,586	11,320
1998	0	105	105	4,062	0	4,062	4,167	7,164	1,822	8,986
1999	15	381	396	1,667	696	2,363	2,759	8,789	494	9,283
2000	117	895	1,012	2,484	1,076	3,560	4,572	6,733	756	7,489
2001	1,312	1,604	2,916	5,830	3,927	9,757	12,673	20,759	4,819	25,578
2002	1,938	827	2,765	3,226	9,416	12,642	15,407	11,929	4,063	15,992
2003	834	918	1,752	8,154	30,002	38,156	39,908	22,754	4,592	27,346
2004	1,422	1,215	2,637	11,667	21,960	33,627	36,264	17,623	1,751	19,374
2005	247	317	564	5,355	1,910	7,265	7,829	3,048	304	3,352
2006	196	725	921	4,267	984	5,251	6,172	7,569	42	7,611
2007 ^{al}	259	2,248	2,507	1,945	2,361	4,306	6,813	8,923	471	9,394

TABLE II-6. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 2 of 2)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))						River Fisheries (t)			
	KMZ			North of	South of	Ocean	River Fisheries (t)			
	Troll	Sport	Subtotal	KMZ	KMZ	Subtotal	Total	Net	Sport	Total
HARVEST RATE^{b/}										
Age-3										
1986	0.03	0.00	0.03	0.06	0.09	0.15	0.18	0.05	0.11	0.16
1987	0.02	0.01	0.03	0.05	0.07	0.13	0.16	0.13	0.13	0.25
1988	0.02	0.01	0.03	0.03	0.14	0.17	0.20	0.12	0.15	0.28
1989	0.02	0.03	0.05	0.04	0.06	0.11	0.15	0.05	0.02	0.07
1990	0.00	0.02	0.03	0.21	0.06	0.27	0.30	0.11	0.12	0.23
1991	0.00	0.01	0.01	0.00	0.01	0.02	0.03	0.21	0.13	0.34
1992	0.00	0.00	0.00	0.02	0.00	0.02	0.02	0.14	0.04	0.18
1993	0.00	0.00	0.00	0.00	0.04	0.04	0.05	0.11	0.06	0.17
1994	0.00	0.01	0.01	0.00	0.03	0.03	0.03	0.12	0.03	0.15
1995	0.00	0.00	0.00	0.02	0.02	0.03	0.04	0.06	0.03	0.09
1996	0.00	0.00	0.00	0.00	0.05	0.05	0.05	0.32	0.09	0.41
1997	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.06	0.08	0.14
1998	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.14
1999	0.00	0.00	0.00	0.01	0.00	0.01	0.01	0.17	0.06	0.23
2000	0.00	0.01	0.01	0.01	0.04	0.05	0.06	0.12	0.03	0.15
2001	0.00	0.00	0.00	0.01	0.02	0.02	0.03	0.18	0.07	0.25
2002	0.00	0.00	0.00	0.00	0.02	0.02	0.02	0.12	0.07	0.19
2003	0.00	0.00	0.00	0.01	0.07	0.07	0.08	0.07	0.05	0.13
2004	0.00	0.01	0.01	0.06	0.05	0.11	0.12	0.14	0.06	0.20
2005	0.00	0.00	0.00	0.00	0.01	0.02	0.02	0.10	0.04	0.14
2006 ^{a/}	0.00	0.01	0.01	0.00	0.00	0.00	0.01	0.13	0.00	0.13
2007 ^{a/}	0.00	0.01	0.02	0.01	0.02	0.02	0.04	0.16	0.05	0.20
Age-4										
1986	0.05	0.01	0.06	0.17	0.23	0.39	0.46	0.57	0.10	0.67
1987	0.06	0.01	0.08	0.21	0.14	0.35	0.43	0.36	0.08	0.44
1988	0.05	0.02	0.07	0.11	0.21	0.33	0.39	0.45	0.07	0.52
1989	0.03	0.05	0.09	0.18	0.09	0.28	0.36	0.59	0.11	0.70
1990	0.04	0.03	0.07	0.38	0.10	0.48	0.55	0.26	0.10	0.36
1991	0.00	0.03	0.03	0.04	0.11	0.15	0.18	0.35	0.09	0.45
1992	0.01	0.00	0.01	0.06	0.00	0.06	0.07	0.23	0.04	0.27
1993	0.00	0.00	0.00	0.06	0.11	0.16	0.16	0.46	0.03	0.49
1994	0.00	0.03	0.03	0.03	0.04	0.06	0.09	0.26	0.03	0.29
1995	0.00	0.01	0.01	0.07	0.06	0.13	0.14	0.16	0.03	0.19
1996	0.00	0.02	0.02	0.05	0.09	0.14	0.16	0.32	0.07	0.39
1997	0.00	0.00	0.00	0.01	0.05	0.06	0.06	0.20	0.06	0.26
1998	0.00	0.00	0.00	0.09	0.00	0.09	0.09	0.24	0.06	0.30
1999	0.00	0.01	0.01	0.05	0.02	0.08	0.09	0.43	0.02	0.45
2000	0.00	0.02	0.02	0.06	0.02	0.08	0.10	0.22	0.02	0.25
2001	0.01	0.01	0.02	0.04	0.03	0.07	0.09	0.24	0.05	0.29
2002	0.02	0.01	0.03	0.03	0.09	0.13	0.15	0.19	0.06	0.26
2003	0.00	0.00	0.01	0.04	0.16	0.20	0.21	0.24	0.05	0.28
2004	0.01	0.01	0.03	0.11	0.21	0.32	0.34	0.43	0.04	0.48
2005	0.01	0.01	0.01	0.14	0.05	0.19	0.20	0.17	0.02	0.19
2006	0.00	0.01	0.01	0.07	0.02	0.08	0.10	0.18	0.00	0.18
2007 ^{a/}	0.01	0.07	0.08	0.06	0.07	0.13	0.21	0.53	0.03	0.56

a/ Preliminary.

b/ Ocean harvest rates are the fraction of Sept 1(t-1) ocean abundance harvested in these fisheries. River harvest rates are the fraction of the river run (t) harvested in these fisheries.

TABLE II-7. Rogue River fall Chinook inriver run and ocean population indices. (Page 1 of 1)

Return Year	Inriver Run Index in Thousands of Fish ^{a/}					Ocean Impact Rate by Age ^{b/}		Ocean Population Index in Thousands of Fish ^{c/}			
	Age-2	Age-3	Age-4	Age-5	Total ^{d/}	Age-3	Age-4-5	Age-3	Age-4	Age-5	Total
1977	2.4	1.0	0.3	0.0	3.7	0.23	0.55	9.7	1.4	0.1	11.2
1978	1.0	6.1	2.3	0.1	9.5	0.23	0.55	38.7	5.2	0.2	44.1
1979	0.2	1.0	6.5	0.0	7.7	0.23	0.55	7.8	18.8	0.1	26.7
1980	0.4	0.2	0.9	0.6	2.1	0.23	0.55	5.2	4.0	1.4	10.6
1981	1.1	3.3	1.0	0.3	5.7	0.21	0.53	9.2	3.0	0.7	12.9
1982	0.7	1.3	1.3	0.1	3.4	0.30	0.52	9.8	2.9	0.3	13.0
1983	0.3	1.1	1.5	0.0	2.9	0.19	0.60	8.6	4.4	0.1	13.1
1984	0.4	1.2	1.8	0.1	3.5	0.08	0.38	9.9	4.7	0.2	14.8
1985	2.5	1.3	3.5	0.6	7.9	0.11	0.25	9.7	6.3	0.9	16.9
1986	3.1	12.5	2.3	0.5	18.4	0.18	0.46	71.3	5.9	1.0	78.2
1987	2.6	7.8	18.1	0.4	28.9	0.16	0.43	80.3	36.3	0.6	117.2
1988	1.4	4.8	25.2	1.5	32.9	0.20	0.39	17.3	47.9	2.5	67.7
1989	0.5	1.3	4.0	2.0	7.8	0.15	0.36	8.4	7.2	3.2	18.8
1990	0.0	0.3	1.4	0.2	1.9	0.30	0.55	6.0	4.7	0.5	11.2
1991	0.2	0.4	1.9	0.5	3.0	0.03	0.18	3.5	3.2	0.6	7.3
1992	0.5	0.3	1.5	0.5	2.8	0.02	0.07	4.4	2.4	0.6	7.4
1993	0.3	3.5	1.5	0.5	5.8	0.05	0.16	16.1	3.2	0.6	19.9
1994	0.5	0.8	5.8	0.9	8.0	0.03	0.09	3.0	9.5	0.9	13.4
1995	0.2	0.6	1.4	2.0	4.2	0.04	0.13	4.3	1.7	2.3	8.3
1996	0.1	0.4	1.8	0.1	2.4	0.05	0.16	2.4	2.8	0.1	5.3
1997	0.1	0.3	1.0	0.3	1.7	0.01	0.06	5.2	1.5	0.3	7.0
1998	0.0	0.5	2.8	0.3	3.6	0.00	0.09	3.8	3.9	0.3	8.0
1999	0.2	0.3	1.6	0.5	2.6	0.01	0.09	1.5	2.7	0.6	4.8
2000	0.2	2.0	0.8	0.6	3.6	0.06	0.10	9.9	0.9	0.6	11.4
2001	0.8	2.3	4.2	0.0	7.3	0.03	0.09	14.1	5.9	0.0	20.0
2002	0.9	4.0	7.1	0.8	12.7	0.02	0.15	32.2	9.1	0.9	42.2
2003	0.9	2.3	12.0	0.4	15.6	0.08	0.21	14.4	22.1	0.5	37.0
2004	0.4	0.6	4.9	2.9	8.8	0.12	0.34	3.9	9.7	4.4	18.0
2005 ^{f/}	NA	NA	NA	NA	NA	0.02	0.20	7.6	5.0	0.8	13.4
2006 ^{f/}	NA	NA	NA	NA	NA	0.01	0.11	4.9 ^{e/}	3.2	0.5	8.6 ^{e/}
2007 ^{f/}	NA	NA	NA	NA	NA	0.04	0.21	5.8 ^{e/}	3.8 ^{e/}	0.6	10.2 ^{e/}
2008 ^{f/}	NA	NA	NA	NA	NA	-	-	6.6 ^{g/}	4.3 ^{g/}	0.7 ^{g/}	11.6 ^{g/}

a/ Index based on carcass counts in spawning survey index areas. Carcass counts in 1978, 1979, and 1980 adjusted for prespawning mortality. Age composition developed from carcass scale sampling.

b/ Exploitation rates since 1981 are based on Klamath River fall Chinook cohort analysis, 1977-1980 based on 1981-1983 average.

c/ Based on cohort reconstruction methods. Index values for 2008 predicted from regression equations; postseason estimates are not available.

d/ Excludes age-6 fish.

e/ Preliminary, complete cohort not available, mean maturity rate used to derive estimate.

f/ Spawning surveys were discontinued 2005.

g/ Preseason forecast.

TABLE II-8. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 1 of 3)

Year	March Preseason Forecast ^{a/}	April STT Modeled Forecast ^{b/}	Postseason Return	March Pre/Postseason	April Pre/Postseason
URB					
1984	90.10	93.00	131.40	0.69	0.71
1985	159.10	159.10	196.40	0.81	0.81
1986	285.90	286.10	281.60	1.02	1.02
1987	436.40	436.40	420.70	1.04	1.04
1988	450.70	446.50	339.90	1.33	1.31
1989	234.00	231.80	261.30	0.90	0.89
1990	127.20	126.90	153.60	0.83	0.83
1991	88.80	88.90	103.30	0.86	0.86
1992	68.40	66.30	81.00	0.84	0.82
1993	84.50	82.70	102.90	0.82	0.80
1994	85.40	94.70	132.80	0.64	0.71
1995	103.70	125.00	106.50	0.97	1.17
1996	88.90	94.20	143.20	0.62	0.66
1997	166.40	158.00	161.70	1.03	0.98
1998	150.80	141.80	142.30	1.06	1.00
1999	147.50	102.10	166.10	0.89	0.61
2000	171.10	208.20	155.70	1.10	1.34
2001	127.20	132.70	232.60	0.55	0.57
2002	281.00	273.80	276.90	1.01	0.99
2003	280.40	253.20	373.20	0.75	0.68
2004	292.20	287.00	367.90	0.79	0.78
2005	352.20	354.60	268.70	1.31	1.32
2006	253.90	249.10	230.40	1.10	1.08
2007	182.40	185.20	112.60	1.62	1.64
2008	162.50	-	-	-	-
LRW					
1984	16.70	NA	13.30	1.26	NA
1985	12.90	NA	13.30	0.97	NA
1986	15.70	NA	24.50	0.64	NA
1987	29.20	NA	37.90	0.77	NA
1988	43.30	42.10	41.70	1.04	1.01
1989	27.30	26.90	38.60	0.71	0.70
1990	23.70	23.40	20.30	1.17	1.15
1991	12.70	12.70	19.80	0.64	0.64
1992	17.40	16.70	12.50	1.39	1.34
1993	12.50	11.90	13.30	0.94	0.89
1994	14.70	13.20	12.20	1.20	1.08
1995	12.40	11.50	16.00	0.78	0.72
1996	8.80	8.10	14.60	0.60	0.55
1997	7.50	7.20	12.30	0.61	0.59
1998	8.10	7.00	7.30	1.11	0.96
1999	2.60	2.50	3.30	0.79	0.76
2000	3.50	2.70	10.20	0.34	0.26
2001	16.70	18.50	15.70	1.06	1.18
2002	18.70	18.30	24.90	0.75	0.73
2003	24.60	23.40	26.00	0.95	0.90
2004	24.10	24.20	22.30	1.08	1.09
2005	20.20	21.40	16.80	1.20	1.27
2006	16.60	16.60	18.10	0.92	0.92
2007	10.10	10.00	4.30	2.35	2.33
2008	3.80	-	-	-	-

TABLE II-8. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 2 of 3)

Year	March Preseason	April STT Modeled	Postseason Return	March	April
	Forecast ^{a/}	Forecast ^{b/}		Pre/Postseason	Pre/Postseason
LRH					
1984	70.40	89.00	102.40	0.69	0.87
1985	81.50	86.70	111.00	0.73	0.78
1986	171.60	173.90	154.80	1.11	1.12
1987	294.90	298.70	344.10	0.86	0.87
1988	267.70	246.50	309.90	0.86	0.80
1989	104.90	97.50	130.90	0.80	0.74
1990	68.50	65.50	60.00	1.14	1.09
1991	71.40	73.10	62.70	1.14	1.17
1992	113.20	121.50	62.60	1.81	1.94
1993	79.30	77.70	52.30	1.52	1.49
1994	36.10	46.50	53.60	0.67	0.87
1995	35.80	42.40	46.40	0.77	0.91
1996	37.70	48.30	75.50	0.50	0.64
1997	54.20	68.70	57.40	0.94	1.20
1998	19.20	22.50	45.30	0.42	0.50
1999	34.80	38.20	40.00	0.87	0.96
2000	23.70	26.40	27.00	0.88	0.98
2001	32.20	30.50	94.30	0.34	0.32
2002	137.60	133.00	156.40	0.88	0.85
2003	115.90	116.90	155.00	0.75	0.75
2004	77.10	79.00	108.90	0.71	0.73
2005	74.10	78.44	78.30	0.95	1.00
2006	55.80	57.50	58.30	0.96	0.99
2007	54.90	54.40	32.70	1.68	1.66
2008	59.00	-	-	-	-
SCH					
1984	21.30	27.00	47.50	0.45	0.57
1985	34.90	37.10	33.20	1.05	1.12
1986	16.00	16.20	16.60	0.96	0.98
1987	9.10	9.20	9.10	1.00	1.01
1988	6.50	5.90	12.00	0.54	0.49
1989	29.50	23.00	26.80	1.10	0.86
1990	27.30	23.70	18.90	1.44	1.25
1991	56.30	61.40	52.40	1.07	1.17
1992	40.90	41.30	29.50	1.39	1.40
1993	19.90	18.20	16.80	1.18	1.08
1994	20.20	28.90	18.50	1.09	1.56
1995	17.50	22.50	33.80	0.52	0.67
1996	27.60	35.40	33.10	0.83	1.07
1997	21.90	25.70	27.40	0.80	0.94
1998	14.20	14.20	20.20	0.70	0.70
1999	65.80	61.00	50.20	1.31	1.22
2000	21.90	26.90	20.50	1.07	1.31
2001	56.60	61.90	125.00	0.45	0.50
2002	144.40	136.00	160.80	0.90	0.85
2003	96.90	101.90	180.60	0.54	0.56
2004	138.00	150.00	175.30	0.79	0.86
2005	114.10	115.79	93.10	1.23	1.24
2006	50.00	51.80	27.90	1.79	1.86
2007	21.80	21.30	14.60	1.49	1.46
2008	87.20	-	-	-	-

TABLE II-8. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 3 of 3)

Year	March Preseason	April STT Modeled	Postseason Return	March	April
	Forecast ^{a/}	Forecast ^{b/}		Pre/Postseason	Pre/Postseason
	MCB				
1990	69.50	69.30	58.90	1.18	1.18
1991	48.40	48.50	35.40	1.37	1.37
1992	42.50	40.70	31.10	1.37	1.31
1993	33.00	32.30	27.50	1.20	1.17
1994	23.90	26.70	33.70	0.71	0.79
1995	25.00	30.00	34.20	0.73	0.88
1996	40.80	43.20	59.70	0.68	0.72
1997	72.10	61.90	59.00	1.22	1.05
1998	47.80	44.90	36.80	1.30	1.22
1999	38.30	27.70	50.70	0.76	0.55
2000	50.60	61.60	36.80	1.38	1.67
2001	43.50	45.30	76.40	0.57	0.59
2002	96.20	91.80	108.40	0.89	0.85
2003	104.80	94.60	150.20	0.70	0.63
2004	90.40	88.80	117.60	0.77	0.76
2005	89.40	89.73	98.00	0.91	0.92
2006	88.30	86.60	80.40	1.10	1.08
2007	68.00	69.10	46.90	1.45	1.47
2008	54.00	-	-	-	-

a/ March preseason forecasts are ocean escapements based on terminal run size and stock-specific cohort relationships affected by the historical "normal" ocean fisheries, generally between 1979 and the most recent adequately complete broods.

b/ STT modeled forecasts adjust March preseason forecasts for Council-adopted ocean regulations each year and should provide a more accurate estimate of expected ocean escapement.

TABLE II-9. Comparison of preseason and postseason forecasts of Puget Sound run size for summer/fall Chinook.^{af} (Page 1 of 2)

Year	Preseason			Postseason			Preseason			Postseason			Preseason			Postseason		
	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason
Nooksack-Samish Hatchery and Natural																		
1993	50.4	32.3	1.53	3.2	3.8	0.84	1.0	1.4	0.71	14.0	6.9	2.00						
1994	46.6	28.1	1.66	3.2	0.7	4.00	1.3	5.5	0.30	8.4	5.9	1.27						
1995	38.5	22.3	1.73	3.5	0.2	17.50	1.6	3.4	0.48	5.0	9.2	0.52						
1996	27.0	29.2	0.92	1.7	0.5	2.43	1.0	1.2	0.83	7.1	10.9	0.58						
1997	34.0	41.7	0.99	1.2	1.2	1.00	0.1	0.0	-	6.4	6.1	1.03						
1998	28.0	31.5	0.95	0.5	0.3	1.67	0.0	0.0	-	6.6	15.0	0.44						
1999	27.0	42.1	0.66	2.3	0.3	7.67	0.0	0.0	-	7.6	5.3	1.46						
2000	19.0	32.6	0.57	5.0	0.1	50.00	0.0	0.0	-	7.3	17.3	0.42						
2001	34.9	64.7	0.55	1.6	0.9	16.00	0.0	0.0	-	9.1	14.1	0.65						
2002	52.8	54.3	0.99	1.6	0.9	2.29	0.0	0.1	-	13.8	20.0	0.69						
2003	45.8	30.0	1.51	1.6	0.2	8.00	0.0	0.3	-	13.7	10.3	1.38						
2004	34.2	17.9	1.83	0.8	0.0	-	0.5	0.0	-	20.3	24.3	0.83						
2005	14.5	15.9	1.07	0.4	0.0	13.30	0.7	0.4	3.50	23.4	23.4	0.99						
2006 ^{bf}	16.9	30.7	0.55	0.4	0.0	-	0.6	0.4	1.51	24.1	22.5	1.07						
2007	18.8	NA	NA	0.4	NA	NA	1.1	NA	NA	15.0	NA	NA						
2008	35.3	-	-	0.8	-	-	0.7	-	-	23.8	-	-						
East Sound Bay Hatchery																		
Skagit Hatchery																		
Skagit Natural																		
Stillaguamish Natural																		
1993	NA	1.3	-	1.6	2.7	0.59	4.9	5.5	0.86	2.8	1.4	2.00						
1994	NA	1.3	-	1.8	5.4	0.33	4.5	5.0	0.90	2.8	1.8	1.47						
1995	1.8	0.9	1.29	2.2	4.0	0.37	4.3	4.0	0.73	2.3	8.5	0.56						
1996	1.3	1.2	0.57	6.7	4.6	0.73	4.2	5.9	0.53	2.7	11.5	0.68						
1997	1.6	1.2	1.33	7.7	12.0	2.85	5.2	4.4	1.18	4.0	8.7	0.47						
1998	1.6	1.6	1.07	6.5	4.7	5.91	5.6	6.4	0.88	2.5	7.2	0.35						
1999	1.5	1.1	1.36	7.8	4.7	4.88	5.6	4.8	1.17	4.5	15.2	0.30						
2000	2.0	1.7	1.18	6.2	1.9	4.13	6.0	6.1	0.98	5.0	8.3	0.60						
2001	1.7	1.4	1.21	4.1	0.9	5.86	5.8	8.4	0.69	5.5	5.1	1.08						
2002	2.0	1.6	1.25	6.8	2.6	2.62	6.7	7.3	0.92	5.8	5.2	1.32						
2003	2.0	1.0	2.00	9.4	5.8	47.00	5.5	5.6	0.98	6.0	8.7	0.80						
2004	2.2	1.5	1.47	10.1	6.2	1.63	15.7	10.7	0.92	7.6	5.7	1.31						
2005	2.0	1.0	2.00	9.9	3.7	2.68	14.2	4.6	3.16	9.2	7.4	1.24						
2006 ^{bf}	1.6	1.3	1.26	9.6	4.9	1.97	8.7	8.4	1.03	10.0	4.0	2.51						
Snohomish Hatchery																		
Snohomish Natural																		
Tulalip Hatchery																		

TABLE II-9. Comparison of preseason and postseason forecasts of Puget Sound run size for summer/fall Chinook.^{a/} (Page 2 of 2)

Year	South Puget Sound Hatchery			South Puget Sound Natural			Strait of Juan de Fuca Hatchery			Strait of Juan de Fuca Natural		
	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason
1993	61.8	43.1	1.68	26.5	9.6	1.34	0.7	1.0	3.50	3.1	1.6	1.29
1994	52.7	49.9	1.08	18.0	10.5	0.60	3.9	1.2	2.44	1.0	1.0	2.00
1995	49.6	75.4	0.67	21.7	24.9	0.63	3.0	0.7	30.00	0.9	2.3	0.33
1996	51.9	53.2	0.89	19.0	16.5	0.53	2.8	1.4	14.00	0.9	2.0	0.29
1997	65.1	38.3	1.40	18.2	15.9	0.88	2.2	1.0	7.33	0.8	2.9	0.23
1998	67.8	49.6	1.24	21.8	14.6	0.79	1.7	1.7	1.00	0.9	2.1	0.47
1999	59.4	67.3	0.71	19.6	33.5	1.15	1.9	0.7	2.71	0.9	2.7	0.33
2000	77.5	47.4	1.39	17.5	39.5	1.26	2.0	1.2	1.67	1.1	1.7	0.65
2001	73.7	76.6	0.76	16.2	44.6	0.80	0.0	1.7	NA	3.5	2.0	1.75
2002	90.8	69.2	1.07	16.9	58.5	0.79	0.0	1.6	NA	3.6	2.2	0.97
2003	86.6	56.6	1.14	19.6	31.0	1.28	0.0	1.3	NA	3.4	2.8	0.72
2004	86.5	66.4	1.16	17.5	24.5	0.61	0.0	1.4	NA	3.5	4.1	0.85
2005	83.1	73.7	0.95	17.7	19.1	0.46	0.0	1.4	NA	4.2	2.0	2.00
2006 ^{b/}	85.8	105.1	0.82	21.3	29.3	0.73	0.0	1.2	NA	4.2	3.0	1.39
2007	83.0	NA	NA	17.0	NA	NA	0.0	NA	NA	4.4	NA	NA
2008	101.6	-	-	21.1	-	-	0.0	-	-	4.5	-	-
	Hood Canal Hatchery and Natural											
1993												
1994	11.7	4.7	2.44									
1995	11.5	3.7	3.03									
1996	3.9	9.9	0.41									
1997	9.0	8.1	1.10									
1998	2.7	7.8	0.34									
1999	6.7	16.3	0.41									
2000	14.0	29.0	0.47									
2001	19.2	20.1	0.90									
2002	25.3	26.6	1.31									
2003	24.0	39.6	0.76									
2004	29.6	36.5	0.86									
2005	30.5	41.1	1.36									
2006 ^{b/}	30.2	68.1	0.44									
2007	47.5	47.7	1.00									
2008	36.8	-	-									

a/ Puget Sound run size is defined as the run available to Puget Sound net fisheries. Does not include fish caught by troll and recreational fisheries inside Puget Sound.

b/ Postseason returns are preliminary.




FIGURE II-1. Regression estimator for CVI based on previous year's river return of age-two Central Valley Chinook, 1990-2007 with 2005 data point omitted. Years shown are CVI year. Numbers in plot denote calendar year t.




FIGURE II-2. Spawning escapements of adult Sacramento River fall Chinook, 1970-2007, and the goal range for the stock of 122,000 to 180,000 adult fish.






FIGURE II-3. Regression estimators for Klamath River fall Chinook ocean abundance (September 1) based on that year's river return of same cohort. Numbers in plots denote brood years.

CHAPTER III - COHO SALMON ASSESMENT

COLUMBIA RIVER AND OREGON/CALIFORNIA COASTAL COHO

(OREGON PRODUCTION INDEX AREA)

The majority of coho harvested in the OPI area originate from stocks produced in rivers located within the OPI area (Leadbetter Point, Washington, to the U.S./Mexico border). These stocks include hatchery and natural production from the Columbia River, Oregon Coast, and northern California, and are divided into the following components: (1) public hatchery (OPIH), (2) Oregon coastal natural (OCN), including river and lake components, (3) hatchery smolt production from the Oregon coastal Salmon Trout Enhancement Program (STEP), Lower Columbia natural (LCN), and (5) natural and hatchery stocks south of Cape Blanco, Oregon, which include the Rogue, Klamath, and Northern California coastal stocks.

A stratified random sampling (SRS) study implemented in 1990 indicated an overestimation of annual OCN spawner escapement, which had previously been based on index surveys. Because the stock composition of the OPI area ocean impacts is based on the proportions of the OPI ocean escapements, a reduction in OCN spawner escapement meant that traditional OCN ocean impacts and abundances were overestimated, while traditional ocean impact and abundance estimates for other OPI area stocks had been underestimated. Starting in 1992, the Council adopted an abundance adjustment procedure for use in assessing fishery impacts. This procedural change, based on improved estimates of OCN spawner escapements, adjusted traditional index abundances of the other OPI area stocks. To achieve targeted exploitation rates and spawner escapement goals, the various OPI area stock abundance index predictions were scaled in the Coho FRAM to reflect the results of the ongoing OCN spawner study and are referred to as SRS abundances. In 1998, after eight years of SRS abundance estimates, the historic OPI data set was rescaled to reflect the revised OCN abundance estimates.

Beginning in 1999, with the availability of a long-term data set in SRS values, all five OPI area stock abundances were projected in SRS accounting. Direct comparisons of 2008 abundance forecasts with recent year SRS abundance projections, both preseason and postseason, are reported in Table III-1. All fishery impacts and escapements from the coho FRAM are reported in SRS values.

Beginning in 2008, a new method was developed to estimate coho abundances for both the natural and hatchery components of the Columbia River and the Oregon coast. The traditional method of stock abundance estimation used only catch data from Leadbetter Point, Washington, to the U.S./Mexico border. This estimation technique was not consistent with the methods used in the FRAM. The Mixed Stock Model (MSM) used for constructing the FRAM base period data was used to estimate the contribution of various coho stocks, including the OPI area stocks, to ocean fisheries and was based on coded-wire tags (CWTs) and associated tag rates. The MSM includes all fisheries that impact a particular stock and therefore should provide a better overall accounting of total harvest and mortality of both Columbia River and Oregon coast coho stocks. The new run size estimates are based on the 1986 – 1992 base period and “backwards” FRAM runs for more recent years. The Oregon Production Index Technical Team (OPITT) has decided to use the MSM run reconstruction database for future accounting and predictions.

Public Hatchery Coho

OPI area public hatchery coho smolt production occurs primarily in Columbia River facilities and net pens. Several facilities located in Oregon coastal rivers and in the Klamath River Basin, California, collectively produce fewer coho. OPI area smolt releases since 1960 are reported by geographic area in Appendix B, Table B-1.

Predictor Description

Since 1988, the OPIH stock predictor was a multiple linear regression with the following variables: Columbia River jacks (Jack CR), Oregon coastal and Klamath River Basin jacks (Jack OC), and a correction term for delayed smolts released from Columbia River hatcheries (Jack CR * [SmD/SmCR]).

The 2008 stock prediction differs slightly from that used in previous years. Because of the shorter data set (1986-2007 vs. 1970-2007) and the near-total phase-out of coastal coho salmon hatcheries the factor for Oregon and California jacks (JackOC) was not significant in the regression. A simplified model with all OPI Jacks combined in one term (Jack OPI) was used, and all parameters were significant.

The OPIH stock predictor is partitioned into Columbia River early and late stocks based on the proportion of the 2007 jack returns to each area adjusted for stock specific maturation rates. The coastal hatchery stock is partitioned into northern and southern coastal stock components. The northern OPIH coastal stock is comprised of hatchery production from the central Oregon Coast. The southern OPIH coastal stock is comprised of hatchery production from the Rogue River basin in southern Oregon and the Klamath and Trinity basins in northern California. The 2008 partition was based on the “backwards” FRAM 2007 ocean abundance proportion.

For the 2008 abundance prediction, the data base includes 1986-2007 recruits. It also includes 1985-2006 jack returns. The model is:

$$\text{OPIH}(t) = a + b * \text{Jack OPI}(t-1) + c * (\text{Jack CR}(t-1) * [\text{SmD}(t-1) / \text{SmCR}(t-1)])$$

Where:

$$a = -42.645724$$

$$b = 15.885113$$

$$c = 37.059908$$

$$\text{adjusted } r^2 = 0.87$$

The OPIH stock data set and a definition of the above terms are presented in Appendix B, Table B-2.

Predictor Performance

Recent year OPIH stock preseason abundance predictions, partitioned by production area and as a total, are compared with postseason estimates in Table III-1. The 2007 preseason abundance prediction of 593,600 OPIH coho was 125 percent of the preliminary postseason estimate of 476,500 coho.

Since 1983, the OPIH predictor has performed well. The years with the highest variations were due principally to high interannual variability in the jack to adult ratios.

2008 Stock Status

Using the appropriate values from Appendix B, Table B-2, the OPIH abundance prediction for 2008 is 216,100 coho, 36 percent of the 2007 prediction and 45 percent of the preliminary 2007 postseason estimate.

Oregon Coastal Natural Coho

The OCN stock is composed of natural production north of Cape Blanco, Oregon from OCNR and OCNL systems, which are predicted independently.

Predictor Description

Oregon Coastal Natural Rivers

From 1988-1993, the abundance of OCNR index coho was predicted using a modified Ricker spawner-recruit model. The predictor related OCNR recruits to the parent brood stock size incorporating an adjustment for ocean survival based on OPI hatchery smolt to jack survival the previous year. Due to a tendency to over-predict abundances, the database in the predictor was shortened from 1970-1991 to 1980-1991 starting with 1992 predictions.

Because of concern that the adopted OCNR model did not adequately incorporate environmental variability, an alternative model was used to predict the 1994 and 1995 index abundances. The model used ocean upwelling, sea surface temperatures, and year to predict OCNR index coho abundance. The year term was included in the model to reflect an observed decline in stock productivity.

For 1996-1998, the environmental based model without the year component was used in predicting OCNR stock abundances. In addition, the predictions were in SRS rather than traditional index accounting. The OCNR environmental variables are annual deviation from the mean April-June Bakun upwelling index at 42° N latitude (UpAnom), and annual deviation from the mean January sea surface temperature at Charleston, Oregon (JanAnom).

For 1999-2002, the environmental-based model with the year component included was used to predict OCNR stock abundances.

For 2003-2007, the same environmental-based model without the year component that was used for 1996-1998 was used in predicting OCNR abundance.

The OCNR stock data set and a definition of the terms are presented in Appendix B, Table B-4.

In 2008, the OPITT adopted a new abundance time series based on MSM run reconstructions and “backwards” FRAM modeling. This time series starts in 1986, in contrast to the SRS time series that starts in 1970. There is much less contrast in the environmental variables in the shorter time period than there was in the longer period. In addition, there appears to be a weaker relationship between abundance and the environmental variables in recent years. A third consideration is that the MSM estimates of OCN abundance are considerably higher than the OPITT estimates and additional work is needed to determine the superior estimate.

For 2008, several models using the MSM time series were considered. These all tended to predict higher abundances than what would reasonably be expected and none were statistically significant. In the absence of a satisfactory model, the OPITT examined patterns in ocean conditions and hatchery jack returns and determined that the 2007 postseason abundance estimate was the most appropriate forecast for 2008.

Oregon Coastal Natural Lakes

Since 1988, the abundance of OCNL index coho has been predicted using the most recent three-year average adult stock abundance. OCNL coho production occurs from three lake systems (Tenmile, Siltcoos, and Tahkenitch). Production from these systems has declined substantially from the levels observed during 1950-1973, but has steadily increased in recent years. The 2007 abundance estimate was 10,400. Following the same reasoning used for the OCN Rivers predictor, the OPITT chose to use the 2007 postseason abundance for the 2008 preseason prediction instead of using a three year average.

Predictor Performance

Recent-year OCN preseason SRS abundance predictions are compared to postseason estimates in Table III-1. Since 2000 the OCN predictor has under estimated abundance except for 2005 and 2007. The 2007 preseason abundance prediction of 255,400 OCN coho was 426 percent of the preliminary postseason estimate of 60,000 coho.

2008 Stock Status

The 2008 preseason prediction for OCN (river and lake systems combined) is 60,000 coho, 23 percent of the 2007 preseason prediction and the same as the 2007 postseason estimate (Table III-1). The 2008 preseason SRS prediction for OCNR and OCNL components are 50,000 and 10,000 coho, respectively.

Private Hatchery Coho

There have been no Oregon coastal PRIH coho smolt releases since 1990. Thus, there is no PRIH recruitment in 2007.

Salmon Trout Enhancement Hatchery Coho Smolt Program

Predictor Description

From 1988 to 2007, preseason abundance predictions for Oregon coastal STEP index coho smolt production facilities have been based on the Council-approved procedure. This procedure involved multiplying the average smolt to adult survival rate by the ratio of the current OPI jack survival to the previous year's OPI jack survival.

The 2007 prediction used the observed 2002-2003 brood smolt-to-adult survival rate applied to the 2004 brood smolt production.

Predictor Performance

Recent-year STEP preseason abundance predictions are compared to postseason estimates in Table III-1. For 2007, there were no reported returns for the preseason abundance prediction of 2008 coho.

2008 Stock Status

Due to changes with the STEP program, releases for this forecast were discontinued after the 2004 brood and no forecast was made for 2008 (Table III-1).

Lower Columbia River Natural

Predictor Description

The 2008 prediction for the Clackamas and Sandy Rivers is based on the recent 3-year cohort averages. The forecast for other Oregon lower Columbia River populations are recent cohort averages and average of recent year abundances. The total Oregon lower Columbia natural coho forecast to terminal areas of 3,200 was expanded by the recent 2-year OPI harvest rate to produce an ocean abundance estimate of 3,900.

The 2008 prediction for the Washington lower Columbia natural coho populations are derived by combining estimates of natural smolt production based on watershed area and a predicted 2005 brood year marine survival rate. The 2008 adult ocean abundance forecast is 9,500 coho.

Predictor Performance

The LCN stock predictor methodology was developed in 2007. The preseason abundance compared to the postseason estimate is presented in Table III-1. The 2007 preseason abundance prediction of 21,500 LCN coho was 111 percent of the preliminary postseason estimate of 19,400 coho

2008 Stock Status

The 2008 prediction for LCN coho is 13,400 coho (Table III-1). This ocean abundance estimate includes both Oregon and Washington LCN components.

Oregon Production Index Area Summary of 2008 Stock Status

The 2008 combined OPI area stock abundance is predicted to be 276,100 coho, which is 33 percent of the 2007 preseason prediction of 849,200 coho and 51 percent of the 2007 preliminary postseason estimate of 536,500 coho. The 2008 OPI area predictions are compared to historical abundances in Table III-2.

WASHINGTON COASTAL AND PUGET SOUND COHO STOCKS

Predictor Description and Past Performance

A variety of preseason abundance estimators currently are employed for Washington coastal and Puget Sound coho stocks (Table I-2). These estimators are used to forecast preseason abundance of adult ocean recruits.

The performance of preseason abundance forecasts (adult ocean recruits) cannot be evaluated at this time because postseason run reconstructions for U.S. and Canadian coho production units have not been completed. A comparison of expected preseason and postseason ocean escapements for Washington coastal and Puget Sound stocks in recent years is presented in Tables III-3 and III-4. Postseason estimates of 2007 ocean escapements for some of these stocks are not available at this time. The comparison of preseason and postseason estimates of ocean escapement reflects annual errors in abundance estimates, deviations in ocean fisheries from preseason expectations, and variations in ocean distributions of stocks as described in the introduction. Fishery impact levels anticipated preseason may be substantially different than those that actually occur.

2008 Stock Status

Washington Coastal Coho

Willapa Bay

The 2008 Willapa Bay hatchery coho abundance forecast is 25,511 ocean recruits compared to a 2007 preseason forecast of 37,228. The hatchery forecast is based on the regression of 1998-2007 hatchery terminal returns on the 1997-2006 jack returns. The natural coho forecast is 35,063 ocean recruits, based on the regression of wild terminal returns in 1997, 1999, 2003, 2005, and 2006 on the previous year's ($n-1$) hatchery jack returns.

Grays Harbor

Preseason abundance forecasts are made for natural fish throughout the system and for hatchery fish returning to three freshwater rearing complexes and three saltwater net-pen sites. The forecasts include fish originating from numerous volunteer production projects. The abundance forecast for Grays Harbor natural stock coho for 2008 is 42,651 ocean age-3 recruits. The forecast for hatchery stock ocean abundance is 53,051 ocean age-3 recruits.

The natural coho forecast consists of an estimate of smolt production in the Humptulips and Chehalis basins multiplied by a smolt to adult survival rate. The smolt production estimate is calculated using 66 smolts per female multiplied by the number of female spawners, which is ½ of the observed 2005 total Grays Harbor natural escapement. The smolt production estimate is then divided by the total available square miles of habitat in the entire Grays Harbor Basin. This results in 587 smolts per square mile. The 587 smolts per square mile is then multiplied by the number of square miles in each component basin (Humptulips and Chehalis). That result is then multiplied by the survival estimate of 0.0285, which is based on a regression of postseason survival estimates on trawl survey catch off Oregon and Washington for 1999-2006.

The hatchery coho forecast consists of an estimate of smolt releases from on- and off-station sites, multiplied by a smolt to adult survival rate. For 2008, the on-station releases totaled 2,208,100 from Bingham Creek, Satsop Springs, Lake Aberdeen and Humptulips Hatcheries. Off station releases totaled 805,100 from Grays Harbor and Lower Chehalis net-pens, and other lower and upper Chehalis River rearing and release sites. The survival estimate of 0.0187 was the mean 2004-2007 survival rate for on-station releases. The survival rate was scaled by a factor of 0.5 (0.0093) for off-station releases.

Quinault River

The 2008 forecast for Quinault natural coho is 17,441 ocean recruits, a 6 percent decrease from the 2007 forecast of 18,600. This forecast is based on the mean estimate of recent ocean recruits for 2001, 2003, 2004, 2005, 2006, and 2007 resulting from the recent Quinault Department of Fisheries work to re-develop the Quinault coho run reconstruction estimates.

The Quinault hatchery coho forecast is 24,540 ocean recruits, a 7 percent increase from the 2007 forecast of 22,735. This return is from a smolt release of 649,573, and is based on a survival rate of 3.8 percent, which lies between the recent five year mean rate for Queets River hatchery rates and somewhat higher rates indicated from recent Quinault Department of Fisheries work to re-develop the Quinault River coho run reconstruction estimates.

Queets River

The 2008 Queets natural coho forecast is 10,182 ocean recruits, a decrease of 25 percent compared to the 2007 forecast level of 13,551. This forecast represents the estimated smolt production (301,250) multiplied by an expected survival rate of 3.38 percent. The estimate of survival rate is based on a regression of Queets wild coho ocean survival rates against September trawl survey data of juvenile coho collected off the coast of Oregon and Washington.

The 2008 Queets hatchery (Salmon River) coho forecast is 10,334 ocean recruits, a decrease of 46 percent compared to the 2007 forecast of 19,138. This forecast is based on a smolt release of 467,680 multiplied by the 2000-2003 brood year average observed marine survival rate (2.2 percent). Approximately 79 percent of the fish released from the Salmon River facility were marked with an adipose fin clip.

Hoh River

The Hoh River natural coho forecast is 4,349 ocean recruits, a decrease of 20 percent compared to the 2007 forecast of 5,400. This forecast is based on estimated smolt production per square mile of watershed from the Clearwater tributary to the Queets River (430 smolts/square mile), multiplied by the size of the Hoh watershed (299 square miles), for a total of 128,678 smolts. The total wild smolt production prediction was then multiplied by an expected survival rate of 3.38 percent. The marine survival rate was based on the regression of eight years of marine survival data for Queets wild coho against trawl survey catch of yearling coho off the WA/OR coast in September of 2007. This estimate of

3.8 percent is consistent with an upward trend in marine survival observed for coho stocks originating from more northerly systems.

No hatchery production is projected for the Hoh system for 2008.

Quillayute River

The Quillayute River summer natural and hatchery coho forecasts for 2008 are 1,115 and 4,228 ocean recruits, respectively. The natural component run size is based on the estimated total summer coho smolt production (27,877) and a projected ocean survival rate of 4.0 percent, which was derived from a regression of survival rates for Queets and Bingham Creek wild coho against catches of juvenile coho in September trawl surveys off the Washington and Oregon coast. The estimate of 4.0 percent is consistent with an upward trend in marine survival observed for coho stocks originating from more northerly streams. The hatchery component run forecast was based on a projected marine survival rate of 2.0 percent, which was derived from review of the relative performance of hatchery vs. wild coho in recent years, multiplied by a release of 211,400 smolts. Approximately 100 percent of the fish were marked with an adipose fin clip. The 2008 forecast abundance of natural summer coho is 1 percent higher than the 2007 forecast, while the hatchery forecast is 34 percent lower than the 2007 forecast level.

The Quillayute River fall natural and hatchery coho forecasts are 10,529 and 12,988 ocean recruits, respectively. The 2008 forecast abundance of natural Quillayute fall coho is 3 percent lower, and the hatchery forecast 28 percent lower, than their respective 2007 forecast levels. The forecast for the natural component is based on the estimated total fall coho smolt production (263,227) multiplied by an expected marine survival rate of 4.0 percent, which was derived from a regression of survival rates for Queets and Bingham Creek wild coho against catches of juvenile coho in September trawls off of the Washington and Oregon coast. The estimate of 4.0 percent is consistent with an upward trend in marine survival observed for coho stocks originating from more northerly streams. The fall hatchery production forecast was based on the same prediction of marine survival (2.0 percent) used for the summer hatchery coho forecast, multiplied by a release of 649,400 smolts. Approximately 88.7 percent of the hatchery fish were marked with an adipose fin clip.

The basin total coho smolt production estimate (summer and fall stocks) was derived by multiplying the 1987, 1988, and 1990 out-migration year average smolt production for the Quillayute system (306,000) by a multiplier (0.95) which represents the proportion of production from the Clearwater in those years. Smolt production was apportioned according to brood year natural spawning escapements of summer and fall coho, to yield the smolt estimates for each natural population.

North Washington Coast Independent Tributaries

Production from several smaller rivers and streams along the North Washington Coast (Waatch River, Sooes River, Ozette River, Goodman Creek, Mosquito Creek, Cedar Creek, Kalaloch Creek, Raft River, Camp Creek, Duck Creek, Moclips River, Joe Creek, Copalis River, Conner Creek), which flow directly into the Pacific Ocean, is forecast as an aggregate. Generally, stock assessment programs on these systems are minimal. The 2008 forecast of natural coho production for these independent streams is 3,180, based on a prediction of 375 smolts per square mile of watershed drainage, 424 square miles of watershed, and an expected marine survival rate of 2.0 percent. The marine survival projection was derived from jack-to-adult return information collected at the WDFW Bingham Creek research station.

The hatchery forecast of 5,007 ocean recruits is developed from linear regression model estimates of marine survival, predicted by the jack return rate for coho from the Makah National Fish Hatchery. The predicted marine survival for the brood year 2005 was multiplied by the 2005 brood year smolt release

(224,579) from the Makah National Fish Hatchery. The entire 2005 brood year release was marked with an adipose fin clip.

Puget Sound

The 2008 total hatchery and natural coho ocean recruit forecast for the Puget Sound region of 614,547 is 3 percent below the 2007 forecast of 633,153. The hatchery coho forecast of 333,543 is 3 percent below the 2007 forecast of 342,529, and the natural coho forecast of 281,004 is 3 percent below the 2007 forecast of 290,624.

Puget Sound hatchery forecasts for 2008 were generally the product of 2005 brood year (BY) smolt releases from each facility, and a predicted marine survival rate for each program. Marine survival rates were typically based on recent year average survival rates derived from CWT recovery information and/or run reconstructions. Forecasts for natural Puget Sound coho stocks were generally derived by measured or predicted smolt production from each major watershed or region, multiplied by stock-specific marine survival rate predictions based on jack return models, recruits/smolt or adult models, or other information.

Strait of Juan de Fuca

The 2008 forecasts for Strait of Juan de Fuca natural and hatchery coho ocean recruits are 24,111 and 9,483, respectively. The natural coho forecast was derived by multiplying the estimated 2005 brood natural smolt production for the region by a predicted ocean age-3 marine survival rate of 6.2 percent. The hatchery forecasts were based on applying hatchery-specific ocean age-3 recruitment rate predictions (0.9 percent for Dungeness, 0.7 percent for Elwha) to the 2005 BY smolt releases for each hatchery. The recruitment rate predictions were based on recent year averages of cohort reconstruction-based recruits/smolt for the aggregate natural stock, and each hatchery production unit.

Nooksack-Samish

The 2008 forecasts for Nooksack-Samish natural and hatchery coho ocean recruits are 14,800 and 47,118, respectively. The natural coho forecast is the product of projected natural smolt production from each stream basin in the region, multiplied by a marine survival rate expectation of 8 percent. The natural coho marine survival rate prediction is based on the average Baker River (Skagit basin) indicator stock CWT-based recruits/smolt rate, adjusted against the Big Beef Creek jack-based marine survival prediction. The hatchery forecasts are based on the 2001-2003 BY average recruits/smolt rate for Kendall Creek Hatchery (1.8 percent), applied to the 2005 BY smolt releases for each facility in the region.

Skagit

The 2008 forecasts for Skagit River natural and hatchery coho ocean recruits are 61,444 and 18,340 (16,744 from in-river hatchery production, 1,596 from Oak Harbor net-pens), respectively. The natural coho forecast is the product of measured smolt production from the Skagit basin multiplied by a marine survival rate expectation of 8.2 percent. The natural coho marine survival rate is based on the average of the 1989-2003 BY (odd years only) Skagit wild recruits/smolt rate adjusted by marine environmental and NMFS ocean juvenile salmonid sampling information related to the parent brood. New environmental and juvenile salmonid indicator data were incorporated into the marine survival rate forecast this year to account for generally poor juvenile coho abundance and erratic ocean conditions observed by NMFS researchers during 2007 off the coasts of Washington and Oregon. The hatchery forecasts are based on an average marine survival rate of the 1989-2003 BY (odd years only) Cascade Hatchery CWT-based recruits/smolt rate adjusted by marine environmental and NMFS ocean juvenile salmonid sampling information related to the parent brood, as was done for the wild forecast.

Stillaguamish

The 2008 forecast for Stillaguamish River natural coho ocean recruits is 31,000. The natural coho forecast is based upon an adult/recruit spawner production model, which contains a recruitment rate adjustment variable based on the deviation pattern in Wallace River Hatchery and South Fork Skykomish River natural coho recruits/smolt rates.

Snohomish

The 2008 forecast for Snohomish River natural coho ocean recruits is 92,000. The Snohomish regional hatchery coho forecast is 53,457; 6,198 for Skykomish River/Wallace River Hatchery facility releases, 44,519 for the Tulalip Bay facility, and 2,740 for the Edmonds and Possession net-pen projects. The natural coho forecast used the measured smolt production for the river basin multiplied by a survival rate expectation of 4.1 percent (see below), averaged with the output from a recruits-per-spawner model. The hatchery forecasts are based on a marine survival rate of 4.1 percent applied to the 2005 BY smolt releases. This value was taken from the observed survival rate for 2003 BY Wallace Hatchery releases. This was a low return rate relative to most return years for this facility, and was expected to represent continued low survival expectations for the coming return year.

South Sound

The 2008 forecasts for South Sound region natural and hatchery coho ocean recruits are 27,286 and 170,022, respectively. The natural coho forecast is the product of projected smolt production from each of the stream basins in the region multiplied by marine survival rate expectation of 5-7 percent for natural coho in the region, with the lowest survival rates expected for deep south sound populations, as per the trend observed for recent years. The marine survival prediction was based upon review of the Big Beef Creek and Deschutes River indicator stock survival data trends, and review of hatchery and natural fish survival rate information from around the region. The hatchery coho forecasts are typically based on the 2001-2003 or 2002-2003 BY average CWT based recruits/smolt rate for each facility, applied to the 2005 BY smolt releases. The expected survival rates range from 4.2-6.1 percent for central Puget Sound hatchery programs, to 0.6 percent -3.1 percent for the deep South Sound region.

Hood Canal

The 2008 forecasts for Hood Canal region natural and hatchery coho ocean recruits are 30,363 and 35,042, respectively. The natural coho forecast is based on a regression of Big Beef Creek jacks versus Hood Canal natural coho run sizes. The hatchery coho forecasts are based on the 1995-2003 BY average cohort reconstruction-based recruits/smolt rates for each facility, applied to the 2005 BY smolt releases (5.2 percent for George Adams Hatchery, 1.7 percent for Port Gamble Net Pens, 5.2 percent for the Quilcene National Fish Hatchery, and 2.7 percent for the Quilcene Bay Net Pens). A moving average of the most recent 3-year marine survival rate is typically used for forecasting hatchery coho production in this region, but concerns regarding the most recent three years (2001-2003 BY) being higher than what likely occurred in 2007, and generally higher than expected to occur in 2008, resulted in a decision by the co-managers to use a longer term marine survival average for forecasting.

SELECTIVE FISHERY CONSIDERATIONS

As the region has moved forward with mass marking of hatchery coho salmon stocks, selective fishing options have become an important consideration for fishery managers. Table III-5 summarizes estimates of mass mark rates for coho stocks from Southern British Columbia, Canada to the Oregon Coast, based on preseason abundance forecasts. Agencies have released coho mass marked with adipose clips from the 2005 brood, making these fish available to 2008 fisheries (Table III-6).

EVALUATION OF 2007 REGULATIONS ON 2008 STOCK ABUNDANCE

Escapements and fishery impacts were estimated using coho FRAM. Abundance forecasts for 2008 were updated for Washington and Oregon stocks, but forecasts for Canadian stocks are unchanged from those employed for 2007 planning. Updated forecasts for Canadian stocks are expected to become available in March 2008. To provide information on the effect of changes in abundance forecasts, the final 2007 pre-season regulatory package for ocean and inside fisheries was applied to 2008 projections of abundance.

Oregon Production Index Area

Ocean fisheries were modeled with 2007 Council regulations and 2007 expectations for non-Council area fisheries. Under this scenario, expected exploitation rates are 42.7 percent on OCN coho and 18.2 percent on Rogue/Klamath hatchery coho. Expected spawner escapement is 35,044 for OCN coho (Tables III-7 and III-8). For Columbia River hatchery coho stocks, the predicted ocean exploitation rate (excluding Buoy 10) is 33.7 percent on the Columbia River early stock and 36.4 percent on the Columbia River late stock. Predicted ocean escapements (after Buoy 10) into the Columbia River in 2008 under this exercise show that under 2007 ocean regulations, Columbia River early would not meet hatchery egg take goals; Columbia River late coho are expected to meet hatchery egg take goals (without inside fishing).

Based on parent escapement levels and observed OPI smolt-to-jack survival for 2005 brood OPI smolts, the total allowable OCN coho exploitation rate for 2008 fisheries is no greater than 15 percent under FMP Amendment 13 and no greater than 8 percent under the matrix developed by the OCN work group. (Table III-9; Appendix A, Tables A-2 and A-3). The total allowable Rogue/Klamath hatchery coho marine exploitation rate is 13.0 percent (NMFS ESA consultation standard).

Lower Columbia River natural (LCN) coho were listed as Endangered under the Oregon state ESA in 1999 and have been managed under a state Recovery Plan harvest rate matrix since 2001. LCN coho were listed as threatened under the Federal ESA in 2005. From 2001 through 2005, Oregon coast hatchery stocks were used as a surrogate in FRAM; in 2006 and 2007, unmarked Columbia River hatchery stocks were used as a surrogate in FRAM. In 2007, NMFS allowed a 20.0 percent exploitation rate in marine area and mainstem Columbia River fisheries combined. The 20.0 percent exploitation rate was split by managers to allow one-third for inriver fisheries and two-thirds for all marine fisheries. Under 2007 fishery regulations and 2008 abundances the exploitation rate is predicted to be 34.7 percent for marine fisheries (excluding the Buoy 10 fishery) using combined unmarked Columbia River hatchery stocks as the proxy. Based on guidance from NMFS the allowable exploitation rate on LCN coho in 2008 is no more than 8 percent in marine and mainstem Columbia River fisheries.

North of the Oregon Production Index Area

Ocean escapement expectations in relation to management goals for selected naturally-spawning coho stocks, given 2008 pre-season abundance forecasts and 2007 pre-season projections for fishing patterns, are presented in Table III-7. The 2008 forecasts for Canadian coho stocks are not available, but are assumed to be at 2007 levels for this analysis. More detailed fishery management goals for Council area coho stocks are listed in Appendix A, Table A-1.

Under 2007 regulations, 2008 ocean escapements for natural coho stocks north of the OPI index area are expected to be at levels that would permit attainment of FMP escapement goals for all U.S. stocks except Hood Canal and Snohomish coho. In addition, all annual management objectives for stocks subject to the PSC agreement would be met except Hood Canal coho. The Hood Canal coho exploitation rate is predicted to be 51 percent (the Council area portion of this is 7.2 percent) under this exercise and the allowable rate for 2008 is 45 percent. The exploitation rate by U.S. fisheries south of the Canadian border

on Interior Fraser coho is projected to be 12.3 percent, exceeding the anticipated 10.0 percent allowable exploitation rate under the 2002 PST Coho Agreement. The Council area fisheries portion is 7.3 percent.

Coho bycatch during Puget Sound fisheries directed at chum and sockeye salmon will also be a consideration for preseason planning.

TABLE III-1. Preliminary 1996-2008 preseason and postseason coho stock Stratified Random Sampling abundance estimates for Oregon production index area stocks in thousands of fish. (Page 1 of 2)

Stock	Year	Preseason	Postseason	Preseason/Postseason ^{a/}
Oregon Production Index Area Hatchery Total	1996	309.2	182.6	1.69
	1997	376.1	215.3	1.75
	1998	118.4	203.6	0.58
	1999	559.2	319.6	1.75
	2000	671.4	677.1	0.99
	2001	1,707.6	1,395.5	1.22
	2002	361.7	660.1	0.55
	2003	863.1	952.5	0.91
	2004	623.9	634.6	0.98
	2005	389.9	443.1	0.88
	2006	398.8	440.6	0.91
	2007	593.6	476.5	1.25
	2008	216.1	-	-
Columbia River Early	1996	142.2	98.0	1.45
	1997	206.9	129.8	1.59
	1998	63.8	126.4	0.50
	1999	325.5	174.9	1.86
	2000	326.3	378.0	0.86
	2001	1,036.5	815.9	1.27
	2002	161.6	324.7	0.50
	2003	440.0	645.7	0.68
	2004	313.6	389.0	0.81
	2005	284.6	282.7	1.01
	2006	245.8	251.4	0.98
	2007	424.9	291.0	1.46
	2008	110.3	-	-
Columbia River Late	1996	114.4	30.8	3.71
	1997	86.5	53.7	1.61
	1998	24.9	47.3	0.53
	1999	140.9	120.7	1.17
	2000	278.0	260.1	1.07
	2001	491.8	488.3	1.01
	2002	143.5	271.8	0.53
	2003	377.9	248.0	1.52
	2004	274.7	203.0	1.35
	2005	78.0	111.6	0.70
	2006	113.8	156.3	0.73
	2007	139.5	171.0	0.82
	2008	86.4	-	-
Oregon Coastal North of Cape Blanco	1996	38.5	28.0	1.38
	1997	60.4	19.0	3.18
	1998	21.6	19.7	1.10
	1999	59.4	14.4	4.13
	2000	48.5	23.4	2.07
	2001	127.3	46.9	2.71
	2002	36.6	41.6	0.88
	2003	29.3	34.5	0.85
	2004	16.6	21.7	0.77
	2005	11.5	10.7	1.07
	2006	8.6	7.9	1.09
	2007	7.0	1.3	5.38
	2008	1.7	-	-

TABLE III-1. Preliminary 1996-2008 preseason and postseason coho stock Stratified Random Sampling abundance estimates for Oregon production index area stocks in thousands of fish. (Page 2 of 2)

Stock	Year	Preseason	Postseason	Preseason/Postseason
Oregon and California Coastal South of Cape Blanco				
	1996	14.2	25.8	0.55
	1997	22.3	12.8	1.74
	1998	8.1	10.2	0.79
	1999	33.4	9.6	3.48
	2000	18.6	15.6	1.19
	2001	52.0	46.0	1.13
	2002	20.0	22.0	0.91
	2003	15.9	24.3	0.65
	2004	19.0	29.9	0.64
	2005	15.8	38.1	0.41
	2006	30.6	25.0	1.22
	2007	22.2	13.2	1.68
	2008	17.7	-	-
Lower Columbia River Natural				
	2007	21.5	19.4	1.11
	2008	13.4	-	-
Oregon Coastal Natural				
	1996	63.2	86.1	0.73
	1997	86.4	27.8	3.11
	1998	47.2	29.2	1.62
	1999	60.7	51.9	1.17
	2000	55.9	69.0	0.81
	2001	50.1	163.2	0.31
	2002	71.8	304.5	0.24
	2003	117.9	278.8	0.42
	2004	150.9	197.0	0.77
	2005	152.0	150.1	1.01
	2006	60.8	116.4	0.52
	2007	255.4	60.0	4.26
	2008	60.0	-	-
Salmon Trout Enhancement Program				
	1996	0.4	1.2	0.33
	1997	1.3	0.3	4.33
	1998	0.2	0.3	0.67
	1999	0.7	0.4	1.75
	2000	0.6	0.5	1.20
	2001	1.0	1.4	0.71
	2002	0.6	3.0	0.20
	2003	3.6	3.6	1.00
	2004	3.1	1.0	3.10
	2005	1.0	0.4	2.50
	2006	0.6	0.1	6.00
	2007	0.2	0.0	-
	2008	0.2	-	-

a/ Postseason estimates are based on preliminary data, and not all stocks have been updated with final estimates.

TABLE III-2. Oregon production index (OPI) area coho harvest impacts, spawning, abundance, and exploitation rate estimates by SRS accounting in thousands of fish.^{a/}
(Page 1 of 1)

Year or Avg.	Oregon and California Coastal Returns							Ocean	OCN Exploitation
	Ocean Fisheries ^{b/}		Hatcheries and Freshwater		Private	Columbia River	Abundance	Exploitation Rate	Rate Based on
	Troll	Sport	Harvest ^{c/}	OCN Spaw ners	Hatcheries	Returns		Based on OPI	Postseason
							Abundance ^{d/}	FRAM	
1970-1975	1,629.6	558.4	45.8	55.2	-	460.4	2,749.3	0.80	-
1976-1980	1,253.6	555.0	31.2	31.1	26.1	263.3	2,155.1	0.83	-
1981	830.9	339.9	34.1	32.6	117.8	170.3	1,555.0	0.81	-
1982	740.9	300.4	37.1	76.2	184.7	453.1	1,763.4	0.62	-
1983	429.6	275.0	18.2	22.7	133.9	109.7	1,070.0	0.79	-
1984	95.8	174.2	51.2	74.4	115.4	424.7	881.5	0.32	-
1985	166.4	280.4	45.4	73.9	332.0	366.2	1,373.4	0.43	-
1986	643.5	320.6	81.8	70.0	453.7	1,548.2	3,026.7	0.34	-
1987	469.1	296.2	45.3	30.1	119.3	316.3	1,377.9	0.60	-
1988	844.7	297.2	62.3	56.8	116.1	670.7	1,989.2	0.56	-
1989	646.9	425.5	62.3	46.4	46.9	711.8	1,871.2	0.55	-
1990	277.6	357.1	30.6	24.3	35.6	196.1	1,128.5	0.69	-
1991	450.6	469.9	84.0	38.6	35.1	934.3	1,823.2	0.45	-
1992	67.5	256.5	52.8	44.4	-	215.9	610.0	0.51	-
1993	13.2	140.8	40.6	55.7	-	113.9	342.1	0.42	-
1994	2.7	3.0	30.0	49.6	-	168.9	250.5	0.02	0.07
1995	5.4	43.5	38.6	57.7	-	74.1	215.9	0.23	0.12
1996	7.0	31.8	47.9	78.6	-	113.0	297.3	0.15	0.08
1997	5.5	22.4	27.2	31.7	-	148.3	204.6	0.12	0.12
1998	3.5	12.8	29.7	34.1	-	168.7	265.2	0.06	0.08
1999	3.6	36.5	20.9	50.4	-	274.1	414.0	0.12	0.09
2000	25.9	74.6	32.9	79.6	-	547.6	901.0	0.13	0.07
2001	38.1	216.8	82.5	182.9	-	1,108.3	1,438.6	0.16	0.07
2002	14.9	118.7	56.3	268.4	-	499.7	990.5	0.14	0.12
2003	28.8	252.4	47.8	235.0	-	677.2	1,183.6	0.23	0.14
2004	26.2	159.4	38.7	194.4	-	442.6	826.8	0.25	0.15
2005	10.5	58.2	42.8	164.1	-	341.8	592.1	0.12	0.11
2006	4.5	47.5	31.7	132.8	-	384.1	557.1	0.06	0.10
2007 ^{e/}	26.7	128.5	11.7	57.1	-	318.6	536.6	0.31	0.11

a/ The OPI area includes ocean and inside harvest impacts and escapement to streams and lakes south of Leadbetter Point, Washington.

b/ Includes estimated nonretention mortality: troll fishery-hook-and-release mortality for 1982-2005 and drop-off mortality for all years; sport fishery-hook-and-release mortality for 1994-2005 and drop-off mortality for all years.

c/ Includes returns from Salmon-Trout Enhancement Program (STEP) smolt releases.

d/ Ocean fishery impacts on private hatchery stock and returns to private hatcheries are excluded in calculating the OPI area stock aggregate ocean exploitation rate index.

e/ Preliminary.

TABLE III-3. Preseason and postseason estimates of ocean escapements for selected Washington coastal adult natural coho stocks in thousands of fish. (Page 1 of 1)

Year	Preseason			Postseason			Preseason			Postseason			Preseason			Postseason								
	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason						
	Quillayute River Fall						Hoh River						Queets River						Grays Harbor^{a/}					
1984	7.0	11.0	0.64	2.7	7.7	0.35	5.2	9.7	0.54	28.7	103.8	0.28												
1985	19.2	15.8	1.22	6.6	5.2	1.27	11.3	6.0	1.88	56.4	25.1	2.25												
1986	6.1	17.1	0.36	3.9	6.4	0.61	5.2	5.8	0.90	51.6	33.3	1.55												
1987	11.7	23.8	0.49	5.5	7.2	0.76	9.0	8.9	1.01	103.3	55.7	1.85												
1988	10.4	9.1	1.14	2.0	2.6	0.77	4.7	4.5	1.04	26.4	58.0	0.46												
1989	14.5	11.1	1.31	5.7	5.4	1.06	6.2	5.4	1.15	43.0	60.9	0.71												
1990	15.2	9.5	1.60	5.1	4.5	1.13	5.9	7.1	0.83	48.3	57.3	0.84												
1991	8.8	10.6	0.83	3.4	5.4	0.63	7.9	8.6	0.92	138.0	108.7	1.27												
1992	12.5	13.6	0.92	4.9	5.0	0.98	5.6	7.0	0.80	48.4	40.9	1.18												
1993	7.6	4.7	1.62	4.8	1.9	2.53	6.5	5.4	1.20	84.7	37.3	2.27												
1994	7.0	6.4	1.09	3.0	1.4	2.14	3.6	1.2	3.00	31.3	11.8	2.65												
1995	8.5	14.3	0.59	4.4	5.4	0.81	7.2	7.3	0.99	64.4	58.9	1.09												
1996	9.2	14.6	0.63	3.0	5.8	0.52	5.4	10.7	0.50	82.7	82.4	1.00												
1997	5.1	5.0	1.02	1.6	1.4	1.14	2.4	2.0	1.20	14.8	18.9	0.78												
1998	7.4	17.0	0.44	3.2	5.2	0.62	4.5	4.6	0.98	27.1	41.2	0.66												
1999	12.8	19.5	0.66	2.8	6.3	0.44	3.7	5.0	0.74	50.3	38.9	1.29												
2000	8.2	17.7	0.46	3.3	8.8	0.38	2.5	8.3	0.30	44.2	40.8	1.08												
2001	20.6	36.7	0.56	7.6	14.8	0.51	10.6	27.8	0.38	46.6	73.5	0.63												
2002	18.5	34.7	0.53	6.9	11.2	0.62	10.2	16.1	0.63	50.3	117.2	0.43												
2003	21.2	25.2	0.84	10.4	8.1	1.28	19.6	11.2	1.75	52.3	107.9	0.48												
2004	17.7	25.1	0.71	6.6	6.3	1.05	14.7	11.1	1.32	101.1	93.1	1.09												
2005	16.1	22.1	0.73	6.4	8.2	0.78	14.1	9.8	1.44	78.5	49.8	1.58												
2006	13.0	11.5	1.13	5.6	3.1	1.81	7.1	5.4	1.31	60.3	19.8	3.05												
2007 ^{b/}	10.8	9.8	1.10	5.4	5.2	1.04	13.6	NA	NA	59.4	NA	NA												
2008 ^{b/}	10.5	-	-	4.3	-	-	10.2	-	-	41.6	-	-												

a/ The source for postseason return estimates is Washington Department of Fish and Wildlife.

b/ Postseason returns are preliminary.

TABLE III-4. Preseason and postseason estimates of ocean escapements for selected Puget Sound adult natural coho stocks in thousands of fish. (Page 1 of 1)

Year	Preseason			Postseason			Preseason			Postseason		
	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason	Forecast	Return	Pre/Postseason
	Skagit River^{a/}			Stilliguamish River^{a/}			Hood Canal^{b/}					
1984	29.6	37.2	0.80	NA	26.9	NA	NA	57.5	NA	NA	NA	NA
1985	26.1	31.3	0.83	NA	34.4	NA	NA	38.5	NA	NA	NA	NA
1986	43.5	73.4	0.59	37.0	49.9	0.74	NA	82.2	NA	NA	NA	NA
1987	33.0	41.2	0.80	29.7	46.3	0.64	NA	71.7	NA	NA	NA	NA
1988	29.6	29.9	0.99	24.5	35.4	0.69	18.2	15.5	1.2	NA	NA	NA
1989	31.2	27.6	1.13	24.5	13.5	1.81	36.8	25.5	1.4	NA	NA	NA
1990	37.6	25.9	1.45	30.8	34.1	0.90	43.9	14.2	3.1	NA	NA	NA
1991	40.8	11.8	3.46	32.9	11.3	2.91	17.6	15.3	1.2	NA	NA	NA
1992	35.7	9.5	3.76	18.7	18.0	1.04	10.1	19.9	0.5	NA	NA	NA
1993	28.1	14.5	1.94	24.5	10.6	2.31	39.5	16.7	2.4	NA	NA	NA
1994	17.9	30.5	0.59	10.2	30.3	0.34	13.5	57.0	0.2	NA	NA	NA
1995	30.0	16.2	1.85	32.7	20.4	1.60	19.3	41.1	0.5	NA	NA	NA
1996	26.7	8.6	3.07	29.8	10.1	2.44	15.4	53.6	0.4	NA	NA	NA
1997	34.2	40.4	0.85	15.7	14.1	1.14	38.1	109.2	0.4	NA	NA	NA
1998	41.1	83.2	0.48	37.7	31.2	1.23	87.3	132.1	0.7	NA	NA	NA
1999	53.4	34.1	1.44	27.3	7.5	3.64	45.2	17.6	2.4	NA	NA	NA
2000	24.7	74.7	0.35	15.0	31.2	0.46	50.4	41.2	1.2	NA	NA	NA
2001	46.9	105.0	0.41	18.1	80.6	0.22	40.6	123.8	0.4	NA	NA	NA
2002	79.9	67.7	1.31	14.5	30.5	0.48	25.6	79.6	0.3	NA	NA	NA
2003	97.9	87.9	1.12	27.7	49.8	0.56	25.8	201.6	0.1	NA	NA	NA
2004	130.9	166.7	0.76	26.6	66.0	0.40	79.7	223.8	0.4	NA	NA	NA
2005	48.4	50.7	1.39	41.8	29.9	1.62	79.6	57.6	2.1	NA	NA	NA
2006	106.6	18.9	5.65	45.0	23.6	1.91	59.4	37.8	1.6	NA	NA	NA
2007 ^{c/}	26.8	NA	NA	69.2	38.7	1.79	42.4	NA	NA	NA	NA	NA
2008 ^{c/}	61.4	-	-	31.0	-	-	30.4	-	-	NA	NA	NA

a/ Post-season numbers for 1996-to-present represent terminal run sizes. Pre-season values for 2001 forward are for April age-3 ocean runsize before fishing.

b/ Post-season numbers for 1996-to-present represent Ocean age-3 runsizes. Pre-season values for 2001 forward are for April age-3 ocean runsize before fishing.

c/ Preliminary.

TABLE III-5. Mass marking of 2005 brood coho available to 2008 Council fisheries. The mark used is an adipose fin clip. (Page 1 of 1)

Region	Ocean Recruits		Percent Mass Marked
	Natural	Hatchery	
PUGET SOUND STOCKS:			
Nooksack-Samish and 7/7A Independent	14,800	47,118	69.7%
Skagit	61,444	18,340	20.0%
Stillaguamish	31,000	80	0.3%
Snohomish	92,000	53,457	26.3%
South Puget Sound Normal	27,286	161,978	82.4%
South Puget Sound Delayed	0	8,044	97.7%
Hood Canal	30,363	34,425	45.8%
Strait of Juan de Fuca and Area 9	24,111	9,484	19.3%
Puget Sound Total	281,004	332,926	48.4%
WASHINGTON COASTAL STOCKS:			
North Coast Independent Tributaries	3,180	5,007	61.2%
Quillayute Summer	1,115	4,228	79.1%
Quillayute Fall	10,529	12,988	49.0%
Hoh	4,349	0	0.0%
Queets	10,182	10,333	39.8%
Quinault	17,441	24,540	49.6%
Grays Harbor	42,651	53,051	54.0%
Willapa Bay	35,063	25,511	39.9%
Washington Coastal Total	124,510	135,658	48.3%
COLUMBIA RIVER STOCKS:			
Columbia River Early	7,243	103,057	69.3% ^{a/}
Columbia River Late	6,189	80,211	75.8% ^{a/}
Columbia River Total	13,432	183,268	72.2% ^{a/}
OREGON COASTAL	60,000	19,400	24.4%
SOUTHERN BRITISH COLUMBIA STOCKS^{b/}:			
Georgia Strait Mainland	81,408	20,044	17.5%
Georgia Strait Vancouver Island	122,304	1,701	0.7%
Johnstone Strait	57,098	7,269	5.6%
Southwest Vancouver Island	32,650	30,377	26.0%
Northwest Vancouver Island	176,545	8,024	0.1%
Lower Fraser River	5,601	89,665	65.9%
Interior Fraser River	14,177	854	0.7%
Southern British Columbia Total	489,782	157,934	15.2%

a/ Columbia River estimate of percent mass marked do not include natural production.

b/ For this assessment, the percent mass marked was assumed to be the same as in 2007.

TABLE III-6. Projected coho mark rates for 2008 fisheries under base period fishing patterns (% marked). (Page 1 of 1)

Area	Fishery	June	July	August	Sept
Canada					
Johnstone Strait	Recreational	-	12%	9%	-
West Coast Vancouver Island	Recreational	27%	14%	10%	8%
North Georgia Strait	Recreational	24%	23%	23%	18%
South Georgia Strait	Recreational	28%	27%	20%	21%
Juan de Fuca Strait	Recreational	37%	40%	42%	39%
Johnstone Strait	Troll	32%	21%	14%	18%
NW Vancouver Island	Troll	19%	19%	21%	25%
SW Vancouver Island	Troll	38%	36%	41%	45%
Georgia Strait	Troll	34%	34%	35%	28%
Puget Sound					
Strait of Juan de Fuca (Area 5)	Recreational	48%	49%	47%	49%
Strait of Juan de Fuca (Area 6)	Recreational	49%	46%	47%	46%
San Juan Island (Area 7)	Recreational	44%	35%	38%	34%
North Puget Sound (Areas 6 & 7A)	Net	-	31%	31%	37%
Council Area					
Neah Bay (Area 4/4B)	Recreational	39%	48%	49%	54%
LaPush (Area 3)	Recreational	50%	50%	56%	43%
Westport (Area 2)	Recreational	56%	57%	56%	56%
Columbia River (Area 1)	Recreational	67%	65%	62%	65%
Tillamook	Recreational	56%	53%	49%	43%
New port	Recreational	53%	49%	48%	32%
Coos Bay	Recreational	43%	40%	31%	16%
Brookings	Recreational	34%	25%	22%	13%
Neah Bay (Area 4/4B)	Troll	50%	48%	50%	54%
LaPush (Area 3)	Troll	48%	54%	51%	60%
Westport (Area 2)	Troll	47%	50%	55%	59%
Columbia River (Area 1)	Troll	59%	57%	56%	61%
Tillamook	Troll	55%	51%	55%	50%
New port	Troll	51%	51%	51%	47%
Coos Bay	Troll	45%	41%	38%	23%
Brookings	Troll	29%	26%	28%	46%
Columbia River					
Buoy 10	Recreational	-	-	-	68%

TABLE III-7. Estimated ocean escapements for critical natural and Columbia River hatchery coho stocks (thousands of fish) based on preliminary 2008 preseason abundance forecasts and 2007 Council regulations.^{a/} (Page 1 of 1)

Stock	Ocean Escapement Estimates Under 2007 Regulations ^{b/}		2008 Spawning Escapement Goal ^{c/}
	2008 Preseason Abundance	2007 Preseason Abundance	
Natural Coho Stocks			
Skagit	49.1	18.8	30.0 ^{d/}
Stillaguamish	24.0	50.2	17.0 ^{d/}
Snohomish	69.2	66.4	70.0 ^{d/}
Hood Canal	18.6	29.8	21.5 ^{d/}
Strait of Juan de Fuca	21.3	26.9	12.8 ^{d/}
Quillayute Fall	9.5	9.7	6.3 - 15.8
Hoh	3.4	4.7	2.0 - 5.0
Queets	7.4	11.5	5.8 - 14.5
Grays Harbor	36.8	53.4	35.4
LCN	13.4 (34.7%)	20.0 (7.0%)	Exploitation Rate ≤8.0%
OCN	35.0 (42.7%)	240.0 (6.2%)	Exploitation Rate ≤8.0%
R/K	NA (18.2%)	NA (2.9%)	Exploitation Rate ≤13.0%
Hatchery Coho Stocks			
Columbia Early	23.7	343.6	18.6
Columbia Late	18.2	93.4	11.9

a/ Quota levels include harvest and hooking mortality estimates used in planning the Council's 2007 ocean fisheries and a coho catch for the Canadian troll fishery off the West Coast of Vancouver Island (WCVI).

b/ 2007 preseason regulations include the following coho quota fisheries: Treaty Indian troll - 38,000 non-selective; non-Indian troll - 22,400 selective; recreational north of Cape Falcon - 117,600 selective; recreational Cape Falcon to OR/CA border - 50,000 selective; troll Cape Falcon to OR/CA border - 10,000 non-selective. Ocean escapement is generally the estimated number of coho escaping ocean fisheries and entering freshwater. For Puget Sound stocks, ocean escapement is the estimated number of coho entering Area 4B which are available for U.S. net fisheries in Puget Sound and spawning escapement after impacts associated with the Canadian and Puget Sound troll and recreational fisheries have been deducted. For the OCN coho stock, this value represents the estimated spawner escapement in SRS accounting. For Columbia River hatchery and LCN stocks, ocean escapement represents the number of coho before the Buoy 10 fishery; the LCN exploitation rate shown is the total ocean fisheries exploitation rate, which had an ER forecast of 13.3% and an ESA limit of 20% including in mainstem Columbia River fisheries.

c/ Goals represent Salmon FMP conservation objectives, ESA consultation standards, or hatchery escapement needs. Spawning escapement goals are not directly comparable to ocean escapement because the latter occur before inside fisheries.

d/ Annual management goals may be determined by the state and tribal co-managers during the preseason planning process, and expressed in terms of total mortality exploitation rate constraints.

TABLE III-8. Comparison of Lower Columbia natural (LCN), Oregon coastal natural (OCN), and Rogue/Klamath (RK) coho projected harvest mortality and exploitation rates by fishery under Council-adopted 2007 regulations and preliminary 2008 preseason abundance estimates. (Page 1 of 1)

Fishery	Projected Harvest Mortality and Exploitation Rate					
	LCN		OCN		RK	
	Number	Percentage	Number	Percentage	Number	Percentage
SOUTHEAST ALASKA	0	0.0%	0	0.0%	0	0.0%
BRITISH COLUMBIA	0	0.0%	112	0.2%	21	0.2%
PUGET SOUND/STRAITS	27	0.2%	93	0.2%	0	0.0%
NORTH OF CAPE FALCON						
Recreational	2,144	16.0%	3,846	6.3%	15	0.1%
Treaty Indian Troll	295	2.2%	902	1.5%	0	0.0%
Non-Indian Troll	469	3.5%	1,214	2.0%	2	0.0%
SOUTH OF CAPE FALCON						
Recreational:	978	7.3%				
Cape Falcon to Humbug Mt.			8,176	13.4%	131	0.9%
Humbug Mt. to Horse Mt. (KMZ)			1,828	3.0%	1,216	8.7%
Fort Bragg			851	1.4%	341	2.4%
South of Pt. Arena			798	1.3%	218	1.6%
Troll:	697	5.2%				
Cape Falcon to Humbug Mt.			6,448	10.5%	311	2.2%
Humbug Mt. to Horse Mt. (KMZ)			120	0.2%	62	0.4%
Fort Bragg			184	0.3%	66	0.5%
South of Pt. Arena			794	1.3%	125	0.9%
BUOY 10	764	5.7%	349	0.6%	0	0.0%
ESTUARY/FRESHWATER	NA	NA	453	0.7%	31	0.2%
TOTAL	5,373	42.0%	26,168	42.7%	2,539	18.2%

TABLE III-9. Maximum allowable fishery impact rate for OCN coho under Amendment 13 matrix (Appendix A, Table A-2) and the OCN work group matrix (Appendix A, Table A-3) based on parent escapement levels by stock component and marine survival category.^{a/} (Page 1 of 1)

Fishery Year (t)	Estimated OCN Coho Spawners by Stock Component					Hatchery Jack Survival Rate (t-1)	Amendment 13 Matrix			OCN Work Group Matrix ^{b/}		
	Parent Spawner Year (t-3)	Northern	North-Central	South-Central	Southern		Marine Survival Category	Parental Spawner Category	Maximum Allowable Impacts	Marine Survival Category	Parental Spawner Category	Maximum Allowable Impacts
1998	1995	3,900	13,600	36,500	3,800	0.04%	Low	Very Low	≤10-13%	Extremely Low	Very Low	≤8%
1999	1996	3,300	18,100	52,600	4,600	0.10%	Med	Very Low	≤15%	Low	Critical	0-8%
2000	1997	2,100	2,800	18,400	8,300	0.12%	Med	Very Low	≤15%	Low	Critical	0-8%
2001	1998	2,600	3,300	25,900	2,300	0.27%	Med	Very Low	≤15%	Medium	Critical	0-8%
2002	1999	8,900	11,800	28,300	1,400	0.09%	Med	Low	≤15%	Low	Low	≤15%
2003	2000	17,900	14,300	36,500	11,000	0.20%	Med	Low	≤15%	Med	Low	≤15%
2004	2001	33,500	25,200	112,000	12,200	0.14%	Med	Low	≤15%	Med	Low	≤15%
2005	2002	52,500	104,000	104,100	7,800	0.11%	Med	High	≤20%	Low	High	≤15%
2006	2003	59,600	68,900	99,800	6,800	0.12%	Med	High	≤20%	Low	High	≤15%
2007	2004	33,100	40,400	96,400	24,500	0.17%	Med	Med	≤20%	Med	Med	≤20%
2008	2005	16,500	51,400	86,300	10,000	0.07%	Low	High	≤15%	Extremely Low	High	≤8%
2009	2006	24,100	21,200	83,500	3,900	-	-	Med	-	-	Low	-
2010	2007	15,100	10,000	26,800	5,200	-	-	Low	-	-	Very Low	-

a/ Under the NMFS ESA consultation standards, the southern stock component is managed for a total allowable Marine Exploitation rate of 13%, as represented by Rogue/Klamath hatchery stocks, which is separate from these OCN coho impact rates.

b/ Developed by the OCN work group as a result of the 2000 Review of Amendment 13.

CHAPTER IV - FRASER RIVER AND PUGET SOUND PINK SALMON ASSESSMENTS

Pink salmon do not contribute significantly to Council ocean fisheries in even numbered years. Two major runs comprise the pink salmon population available to Council fisheries during odd-numbered years: the Fraser River (British Columbia) run, which is more abundant, and the Puget Sound run. The 2007 run size forecast for Fraser pinks was 19.6 million fish, above the forecast of 16.3 million in 2005. Timing of the 2005 and 2003 Fraser pink runs was earlier than normal. The 2007 Puget Sound pink salmon run size forecast was 3.34 million; with 3.3 million natural and 3,800 hatchery fish.

The only self sustaining even-year run known to occur in Washington is from the Snohomish River. The 2008 forecast for the 4B run size is 3,600.

Table IV-1 provides a summary of recent run sizes.

TABLE IV-1. Estimated annual run sizes (odd-numbered years 1977-2007) for Fraser River and Puget Sound pink salmon in millions of fish. (Page 1 of 1)

Year	Puget Sound	Fraser River ^{a/}
1977	0.88	8.21
1979	1.32	14.40
1981	0.50	18.69
1983	1.01	15.35
1985	1.76	19.10
1987	1.57	7.17
1989	1.93	16.63
1991	1.09	22.33
1993	1.06	17.01
1995	2.11	12.88
1997	0.44	8.20
1999	0.95	3.59
2001	3.50	21.17
2003	2.30	26.00
2005	1.23	10.00
2007 ^{b/}	3.34	11.00

a/ Total run size.

b/ Preliminary.

**APPENDIX A
SUMMARY OF COUNCIL STOCK MANAGEMENT GOALS**

LIST OF TABLES

	<u>Page</u>
TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries	63
TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13.....	75
TABLE A-3. Fishery impact rate criteria for OCN coho stock components based on the harvest matrix resulting from the OCN work group 2000 review of Amendment 13.....	76

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 1 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p>CALIFORNIA CENTRAL VALLEY - All fall, late-fall, winter, and spring stocks of the Sacramento and San Joaquin Rivers and their tributaries. Management of this stock complex is based primarily on Sacramento River fall Chinook, which includes a large hatchery component and natural Sacramento River winter Chinook, which are listed as endangered. The San Joaquin system has been severely degraded by water development projects and pollution. Natural populations of spring Chinook there have been extirpated, and remaining spawning areas are utilized primarily by fall Chinook, which have comprised <10% of the total Central Valley fall run.</p>			
<p>Sacramento River Fall</p>	<p>122,000-180,000 natural and hatchery adult spawners (MSY proxy adopted 1984). This objective is intended to provide adequate escapement of natural and hatchery production for Sacramento and San Joaquin fall and late-fall stocks based on habitat conditions and average run-sizes as follows: Sacramento River 1953-1960; San Joaquin River 1972-1977 (ASETF 1979; PFMC 1984; SRFCRT 1994). The objective is less than the estimated basin capacity of 240,000 spawners (Hallock 1977), but greater than the 118,000 spawners for maximum production estimated on a basin by basin basis before Oroville and Nimbus Dams (Reisenbichler 1986).</p>	<p>Yes. A conservation alert or overfishing concern will be based on a failure to meet 122,000 adult spawners.</p>	<p>Contributes to ocean fisheries off California, southern and central Oregon, Washington, and British Columbia. Council management actions on this stock are directed at fisheries south of Pt. Arena; impacts on this stock between Pt. Arena and Horse Mt. are incidental to management measures directed at Klamath River fall Chinook.</p>
<p>Sacramento River Spring Threatened (1999)</p>	<p>Listed as threatened under ESA. NMFS ESA consultation standard/recovery plan. Present level of ocean fishery impacts limited by measures constraining harvest on Sacramento River winter and Klamath River fall Chinook.</p>	<p>No. NMFS ESA consultation standard provides interim rebuilding program MSY criteria undefined.</p>	<p>Contributes to ocean fisheries off California, but also known to occur off Oregon. Ocean fishery impacts primarily incidental to harvest of Sacramento River fall Chinook and may be lower due to differences in run timing. Stock has been affected by man-caused loss and deterioration of freshwater habitat.</p>
<p>Sacramento River Winter Endangered (1994)</p>	<p>Listed as endangered under ESA. NMFS ESA consultation standard specifies duration and timing of commercial and recreational fisheries south of Pt. Arena.</p>	<p>No. NMFS ESA consultation standard provides interim rebuilding program.</p>	<p>Believed to contribute predominantly to ocean fisheries south of Pt. Arena. Ocean fishery impacts incidental to harvest of Sacramento River fall Chinook.</p>
<p>NORTHERN CALIFORNIA COAST - All fall and spring stocks of California streams north of the entrance to San Francisco Bay. Management of this stock complex is based primarily on meeting spawning escapements for natural fall Chinook. Limited data is available except for the Klamath River. An assessment and monitoring program is under consideration by CDFG for stocks originating from the Smith, Eel, Mattole, and Mad Rivers, which might provide a more thorough management basis for the future. There are significant water diversion problems in several drainages. In the Klamath River Basin, there is significant hatchery production of fall Chinook, and less so of spring Chinook, resulting primarily from mitigation programs for dams constructed in both Upper Klamath and Trinity Rivers.</p>			
<p>Eel, Mattole, Mad, and Smith Rivers (Fall and Spring) Eel, Mattole, and Mad River stocks - Threatened (1999)</p>	<p>Eel, Mattole, and Mad River stocks listed as threatened under ESA. Data insufficient to define MSY criteria. Indices of spawning abundance limited to one tributary of the Mad River and two tributaries of the Eel River. NMFS ESA consultation standard/recovery plan for Eel, Mattole, and Mad River stocks requires that the projected ocean harvest rates on age-4 Klamath River fall Chinook not exceed 16.0%.</p>	<p>Eel, Mattole, and Mad - No. NMFS ESA consultation standard provides interim rebuilding program MSY criteria undefined. Smith - Indirectly. Data insufficient to define MSY criteria. CDFG developing an assessment and monitoring program.</p>	<p>Very limited management data available. Believed to occur in ocean fisheries off northern California and southern Oregon. Ocean fishery impacts incidental to fisheries for Sacramento and Klamath Rivers fall Chinook. No preseason or postseason abundance estimates available.</p>

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 2 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
Klamath River Fall (Klamath and Trinity Rivers)	33% to 34% of potential adult natural spawners, but no fewer than 35,000 naturally spawning adults in any one year. Brood escapement rate must average 33% to 34% over the long-term, but an individual brood may vary from this range to achieve the required tribal/nontribal annual allocation. Objective designed to allow a wide range of spawner escapements from which to develop an MSY objective or proxy while protecting the stock during prolonged periods of reduced productivity. Adopted 1988 based on Hubbell and Boydstun (1985); KRTT (1986); PFMC (1988); minor technical modifications in 1989 and 1996 (Table I-1). Natural spawners to maximize recruitment are estimated at 41,000 to 106,000 adults (Hubbell and Boydstun 1985).	Yes. A conservation alert or overfishing concern will be based on a failure to meet the 35,000 floor.	Contributes primarily to ocean fisheries from Humbug Mt., Oregon to Horse Mt., California (the KMZ) and to Klamath River tribal and recreational fisheries. Coastwide impacts are considered in meeting allocation requirements for Indian tribes with federally recognized fishing rights and the inland fishery. Specific management measures for this stock generally are implemented from Pt. Sur, California to Cape Falcon, Oregon.
Klamath River Spring (Klamath and Trinity Rivers)	Undefined. Productive potential believed to be protected by fishery management objective for Klamath River fall Chinook, which includes an inside allocation to tribal and sport fisheries.	Indirectly. MSY criteria undefined.	Little information available on ocean distribution. Believed to occur in ocean fisheries off northern California and southern Oregon (based on Trinity River Hatchery fish).
OREGON COAST - All fall and spring stocks from Oregon streams south of the Columbia River. No preseason abundance estimates available. Management based primarily on an aggregate objective of 150,000 to 200,000 natural adult spawners (attainment of objective based on a postseason estimate of 60 to 90 natural adult spawners per mile in nine standard index streams). This objective is based on optimal escapement estimates for individual coastal rivers at habitat capacity (Thompson 1977). Lower end of the objective range is nearly twice the estimated MSY spawning escapement of 79,000 fall Chinook adults based on stock recruit analysis (McGie 1982). Significant hatchery production also exists within the coastal streams. Far-north migrating, naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.			
Southern Oregon (Aggregate of fall and spring stocks in all streams south of Elk River; Rogue River fall stock is used to indicate relative abundance and ocean contribution rates)	Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982). ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council.	Yes, based on postseason estimates of <60 natural adult spawners per mile. Conservation promoted by the objective for Klamath River fall Chinook, which includes a large inside allocation component that reduces ocean fishery exploitation rate in areas inhabited by these fish, and by ESA consultation standard for California coastal Chinook, which limits projected ocean harvest rates on age-4 Klamath River fall Chinook to $\leq 16.0\%$.	Medium abundance. Data limited except for Rogue River fall stock. Stocks migrate southerly or remain local, and fall Chinook contribute to ocean fisheries off northern California and Oregon, less so for spring stocks.
Central and Northern Oregon (Aggregate of fall and spring stocks in all streams from the Elk River to just south of the Columbia River)	Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982). ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council.	Yes, based on postseason estimates of <60 natural adult spawners per mile.	Variable between high and medium abundance. Stocks migrate northward and contribute to ocean fisheries off British Columbia and southeast Alaska, and to a lesser degree, off Washington and Oregon. Nehalem, Siletz, and Siuslaw stocks are subject to the PSC ISBM harvest limitations.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 3 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p>COLUMBIA RIVER BASIN - All pertinent fall, summer, and spring stocks of the Columbia River and its tributaries. Stocks within this complex are noted by area of origin: lower river (below Bonneville Dam), mid-river (Bonneville to McNary Dams), and upper river (above McNary Dam). Spawner escapement goals for these stocks are set through procedures of the U.S. District Court in <i>U.S. v. Oregon</i> and subsequent court orders. These goals are set forth in the Columbia River Fishery Management Plan and are recognized in the Council's conservation objectives. Annual inside fishery management planning activities are conducted within the Columbia River Compact and other state and tribal management forums. The Columbia River Compact, initially established by Oregon and Washington to jointly administer commercial fisheries within the Columbia River, takes into account the impacts from other state and tribal fisheries (e.g., recreational, ceremonial, subsistence, etc.) authorized under <i>U.S. v. Oregon</i>. The majority of ocean Chinook harvest north of Cape Falcon is provided by Columbia River salmon stocks, primarily hatchery production of tule fall Chinook from the Bonneville Pool (Spring Creek) and lower river hatcheries, smaller numbers of upper river bright hatchery and natural fall Chinook, and some lower river hatchery spring Chinook (Cowlitz). Hatchery objectives are based on long-range production programs and/or mitigation requirements associated with displaced natural stocks. Threatened Snake River fall Chinook, which suffer from severe dam passage mortalities and extreme loss of freshwater habitat, are of prime concern in limiting ocean exploitation rates in all ocean fisheries north of Pigeon Pt., California. These limits act to provide considerable protection to other weak natural stocks subject to ocean fishery impacts. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.</p>			
<p>North Lewis River Fall Threatened (1999)</p>	<p>NMFS ESA consultation standard/recovery plan (not established at time of printing). Mclsaac (1990) stock-recruit analysis supports MSY objective of 5,700 natural adult spawners.</p>	<p>No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program. Base period Council-area ocean fishery impacts around 7%.</p>	<p>Below average abundance in 2008. Present in ocean fisheries north of Cape Falcon to SE Alaska. Subject to the PSC ISBM harvest limitations.</p>
<p>Lower River Hatchery Fall</p>	<p>15,400 adults to meet egg-take goal or as determined by management entities. 41.0% total RER in 2008 for ESA listed lower Columbia River natural tule fall Chinook estimated from Cowlitz, Washougal, Kalama and Big Creek hatchery fall Chinook.</p>	<p>No (hatchery exception or listed stock). NMFS ESA consultation standard provides interim rebuilding program.</p>	<p>Below average abundance in 2008. Major contributor to ocean fisheries north of Cape Falcon to central British Columbia.</p>
<p>Lower River Hatchery (Spring)</p>	<p>2,700 adults to meet Cowlitz, Kalama, and Lewis Rivers broodstock needs.</p>	<p>No (hatchery exception).</p>	<p>Below average abundance in 2008. Present in ocean fisheries north of Cape Falcon to southeast Alaska.</p>
<p>Upper Willamette (Spring) Threatened (1999)</p>	<p>NMFS ESA consultation standard/recovery plan (ODFW FMFP). Willamette River Management Plan provides an MSY proxy of 30,000 to 45,000 hatchery and natural adults over Willamette River falls, depending on run size.</p>	<p>No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program. Base period Council-area ocean fishery exploitation rate of <5% prevents effective Council fishery management and rebuilding.</p>	<p>Below average abundance in 2008. Present in fisheries north of Cape Falcon to southeast Alaska.</p>
<p>Mid-Columbia Bright Hatchery (Fall)</p>	<p>None for ocean fishery management.</p>	<p>No (hatchery exception).</p>	<p>Below average abundance in 2008. Contributor to ocean fisheries off Washington, British Columbia, and southeast Alaska.</p>
<p>Spring Creek Hatchery (Fall)</p>	<p>7,000 adults to meet hatchery egg-take goal.</p>	<p>No (hatchery exception).</p>	<p>Average abundance in 2008. Major contributor to ocean fisheries north of Cape Falcon to southern British Columbia.</p>

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 4 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
COLUMBIA RIVER BASIN (continued)			
Klickitat, Deschutes, John Day, and Yakima Rivers (Spring)	Hold ocean fishery impacts at or below base period (<1%) and recognize CRFMP objective - MSY proxy of 115,000 adults above Bonneville Dam, including upper and mid-Columbia and Snake River stocks (state and tribal management entities considering separate conservation objectives for these stocks).	Limited. Base period Council-area ocean fishery exploitation rate of <1% prevents effective Council fishery management and rebuilding. Major habitat restoration addressing water withdrawals and dam passage and blockages is necessary for rebuilding.	Below average abundance in 2008. No significance to ocean fisheries, infrequent occurrence in fisheries north of Cape Falcon to Alaska.
Snake River Fall Threatened (1992)	NMFS ESA consultation/recovery standard. Since 1995, Council has met a standard of limiting its fisheries so that the total exploitation rate on age-3 and age-4 Lyons Ferry Hatchery fall Chinook (representing Snake River fall Chinook) for all ocean fisheries (including Canada) has been ≤70.0% of the 1988-1993 average adult equivalent exploitation rate. Prior to listing, managed within objectives for upper Columbia River bright fall Chinook.	No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim rebuilding program. Recovering historic abundance unlikely, as dams block former primary spawning area.	Depressed. Present in ocean fisheries from central California to southeast Alaska with greatest contribution to Canadian fisheries. Primary impacts in Council fisheries north of Cape Falcon, but also extending to Pigeon Pt., California.
Snake River Spring/Summer Threatened (1992)	Not applicable for ocean fisheries.	No. Listed stock. Base period Council-area ocean fishery impacts rare (unmeasurable). Dam passage mortality must be reduced to allow stock recovery.	Depressed, recent upward trend. Rare occurrence in ocean fisheries from Washington to southeast Alaska.
Upper River Bright (Fall)	40,000 natural bright adults above McNary Dam (MSY proxy) adopted in 1984 based on CRFMP. The management goal was increased to 45,000 by Columbia River managers between 1986 and 1993. Since 1994, inriver fisheries management based on a NMFS ESA consultation standard exploitation rate to protect Snake River wild fall Chinook	Limited. Base period Council-area ocean fishery exploitation rate <4% prevents effective Council fishery management and rebuilding.	Below average abundance in 2008. Major contributor to ocean fisheries off Canada, and to a lesser extent, Washington and Oregon. Primary impact area north of Cape Falcon. Subject to the PSC ISBM harvest limitations.
Upper River Summer	Hold ocean fishery impacts at or below base period (<2%); recognize <i>U.S. v. Oregon</i> objective - MSY proxy of 29,300 adults to river mouth destined to for areas above Priest Rapids Dam (excludes Snake River stocks).	Limited. Base period Council-area ocean fishery exploitation rate <2% prevents effective Council fishery management and rebuilding. Dam passage mortalities must be reduced to allow rebuilding.	Long-term depressed abundance, significant upward trend in the last few years. Present in ocean fisheries north of Cape Falcon to southeast Alaska. Subject to the PSC ISBM harvest limitations.
Upper Columbia River Spring Endangered (1999)	None applicable to ocean fisheries. Ensure ocean fishery impacts remain rare and recognize CRFMP objective - MSY proxy of 115,000 adults above Bonneville Dam, including upper and mid-Columbia and Snake River stocks (state/tribal management entities considering separate objectives for these stocks).	No. Listed stock. Base period Council-area ocean fishery impacts rare (not measurable), making Council management and rebuilding ineffective. Reduce dam passage mortalities to allow rebuilding.	Long-term depressed abundance, recent upward trend. Captive broodstock programs started in 1997. No significance to ocean fisheries. Rare occurrence in ocean fisheries north of Cape Falcon to Canada.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 5 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
WASHINGTON COAST - All pertinent fall, summer and spring stocks from coastal streams north of the Columbia River through the western Strait of Juan de Fuca (west of the Elwha River). This stock complex consists of several natural stocks, generally of small to medium sized populations, and some hatchery production (Willapa Bay and the Quinault River). Stocks in this complex tend to range further north than most Columbia River stocks and, while present in fisheries from Cape Falcon to southeast Alaska, are not significantly impacted by Council-area ocean fisheries. Preseason abundance estimates are generally not available for Council management. These stocks qualify as exceptions to the Council's overfishing criteria, due to very low fishery impacts. Spawning escapement goals for stocks managed within this complex, established in U.S. District Court by WDFW and the treaty tribes, are recognized in the Council's conservation objectives below. Objectives for Grays Harbor and the north coast river systems have been established pursuant to the U.S. District Court order in <u>Hoh v. Baldrige</u> . However, annual natural spawning escapement targets may vary from the conservation objectives below if agreed to by WDFW and the treaty tribes under the provisions of <u>Hoh v. Baldrige</u> and subsequent U.S. District Court orders. After agreement is reached on the annual targets, ocean fishery escapement objectives are established for each river, or region of origin, which include provisions for treaty allocation and inside, non-Indian fishery needs. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.			
Willapa Bay Fall (Natural)	No FMP objective. WDFW goal of 4,400 natural spawners.	Limited (exploitation rate exception).	
Willapa Bay Fall (Hatchery)	9,800 adult return to hatchery.	No (hatchery exception).	
Grays Harbor Fall	14,600 natural adult spawners--MSP based on full seeding of spawning and rearing habitat (WDF 1979). 1,400 natural adult spawners.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Grays Harbor Spring		Limited (exploitation rate exception).	
Quinault Fall	Hatchery production.	No (hatchery exception).	
Queets Fall	Manage terminal fisheries for 40% harvest rate, but no less than 2,500 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Queets Spring/Summer	Manage terminal fisheries for 30% harvest rate, but no less than 700 natural adult spawners.	Limited (exploitation rate exception).	
Hoh Fall	Manage terminal fisheries for 40% harvest rate, but no less than 1,200 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Hoh Spring/Summer	Manage terminal fisheries for 31% harvest rate, but no less than 900 natural adult spawners.	Limited (exploitation rate exception).	
Quillayute Fall	Manage terminal fisheries for 40% harvest rate, but no less than 3,000 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Quillayute Spring/Summer	1,200 natural adult spawners for summer component (MSY).	Limited (exploitation rate exception).	
Hoko Summer/Fall (Western Strait of Juan de Fuca)	850 natural adult spawners, the MSP level estimated by Ames and Phinney (1977). May include adults used for supplementation program.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 6 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p>PUGET SOUND - All fall, summer, and spring stocks originating from U.S. tributaries to Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek). This stock complex consists of numerous natural Chinook stocks of small to medium sized populations and significant hatchery production. Puget Sound stocks contribute to fisheries off British Columbia and are present into southeast Alaska, but are impacted to a minor degree by Council-area ocean fisheries. Base period, Council-area ocean fishery exploitation rates (adult equivalent) of 2% or less are below a management threshold which allows effective Council management of these stocks and they qualify as exceptions to the Council's overfishing criteria. The naturally spawning stocks within this complex are listed as threatened under the ESA. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border. Management objectives for hatchery stocks are based on hatchery escapement needs. Fisheries in Puget Sound conducted under a Resource Management Plan (RMP) are exempted from ESA Section 9 take prohibitions under Limit 6 of the 4(d) rule. This RMP will expire on May 1 of this year. A new RMP is currently under review by NOAA Fisheries but this review will not be completed prior to the March Council meeting.</p>			
Eastern Strait of Juan de Fuca Summer/Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
Skokomish Summer/Fall (Hood Canal) Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
Nooksack Spring (early) Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Skagit Summer/Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Skagit Spring Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Stillaguamish Summer/Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Snohomish Summer/Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Cedar River Summer/Fall (Lake Washington) Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). The preliminary 2004 consultation standard is an RER constraint total mortality in all fisheries not to exceed 31%.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 7 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
PUGET SOUND (continued)			
White River Spring Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
Puyallup Summer/Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
Green River Summer/Fall Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
Nisqually River Summer/Fall (South Puget Sound) Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
Mid Hood Canal Fall Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
SOUTHERN BRITISH COLUMBIA - Fall and spring stocks of British Columbia coastal streams and the Fraser River. Management based primarily on natural and hatchery fall Chinook. Base period, Council-area ocean fishery exploitation rates (adult equivalent) on the coastal stocks of 1% or less are below a management threshold which allows effective Council management of these stocks, and they qualify as exceptions to the Council's overfishing criteria.			
Coastal Stocks	Undefined for Council fisheries. Manage consistent with the Pacific Salmon Treaty.	No. Under Canadian authority and would also be an exploitation rate exception.	Medium abundance. Major contributors to ocean fisheries off British Columbia; significant contributors north into southeast Alaska and present off northern Washington.
Fraser River	Undefined for Council fisheries. Manage consistent with the Pacific Salmon Treaty.	No. Under Canadian authority.	Medium abundance. Major contributors to ocean fisheries off British Columbia; contributors off northern Washington; and present north into southeast Alaska. Harrison River stock subject to the PSC ISBM harvest limitations.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 8 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
<p>OREGON PRODUCTION INDEX AREA - All Washington, Oregon, and California natural and hatchery coho stocks from streams south of Leadbetter Pt. Significant production from Columbia River and Oregon coastal hatcheries provide harvest in ocean fisheries throughout the Council management area. Ocean fisheries are usually limited primarily to meet natural escapement objectives. Treaty Indian obligations, non-Indian harvest opportunity, and hatchery requirements must also be factored in for the Columbia River stocks. Natural components have been severely depressed for several yeas due to a combination of previously high fishery impacts, major losses or degradation of freshwater habitat, and long-term marine conditions unfavorable to coho survival.</p>			
<p>Central California Coast Threatened (1996)</p>	<p>NMFS ESA consultation standard/recovery plan. Since 1998, no retention of coho in commercial and recreational fisheries off California in conjunction with total marine fishery impacts of no more than 13% on Rogue/Klamath hatchery coho (surrogate stock). Objective undefined prior to listing.</p>	<p>No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim protection of productive capacity. Recovery limited by deterioration of significant portions of freshwater habitat, distribution at southern edge of coho range, and ongoing unfavorable marine conditions.</p>	<p>Very minor component of OPI area fisheries, limited potential for significant contribution to ocean and inland fisheries. Current impacts incidental in ocean fisheries off California. Development of monitoring and assessment program considered for Ten Mile River, Noyo River, Gualala River, Lagunitas Creek, and Scott Creek. Rogue/Klamath coho are believed to have a similar, but more northerly distribution.</p>
<p>Northern California Threatened (1997)</p>	<p>NMFS ESA consultation standard/recovery plan. Since 1998, total marine fishery impacts limited to no more than 13.0% on Rogue/Klamath hatchery coho (surrogate stock) and no retention of coho in California ocean fisheries. Objective undefined prior to listing.</p>	<p>No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim protection of productive capacity. Recovery may last more than 10 years even with no fishery impacts, due to loss or deterioration of significant portions of freshwater habitat and ongoing unfavorable marine conditions.</p>	<p>Depressed and listed. Very minor natural component of OPI area fisheries, potential for minor contribution to ocean fisheries off California and southern Oregon, and inland California fisheries. Current impacts incidental in ocean and inland fisheries (total non-retention south of Cape Falcon since 1994). CDFG considering monitoring to provide data for the Smith, Trinity, Eel, Mattole, and Klamath Rivers.</p>
<p>Oregon Coastal Natural Comprised of Southern, South-Central, North-Central, and Northern Oregon stocks.</p>	<p>An allowable marine and freshwater exploitation rate of no more than 13% to 35%, depending on parent escapement and ocean survival trends, based on Amendment 13 of the Salmon FMP, or no more than 8% to 45% based on the OCN workgroup review of Amendment 13. Standard is ≤8.0% in 2008</p>	<p>No. Listed stock, rebuilding program initiated in 1998. The annual conservation objective should allow component stocks to rebuild when environmental conditions are favorable. Recovery for some components may last more than 10 years even with no fishery impacts, due to loss or deterioration of significant portions of freshwater habitat and ongoing unfavorable marine conditions.</p>	<p>Decline in 2007 after recent increases in abundance. Major natural component of OPI area and freshwater fisheries in Oregon coastal streams. Current impacts are primarily incidental in ocean fisheries under a total nonretention regulation south of Cape Falcon since 1994 (except 2007).</p>

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 9 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
OREGON PRODUCTION INDEX (continued)			
Columbia River Late (Hatchery)	Hatchery rack return goal of 17,200 adults.	No (hatchery exception).	Major component of ocean fisheries north of Cape Falcon. When abundant, significant contributors to ocean fisheries off Oregon north into Canada and Columbia River fisheries.
Columbia River Early (Hatchery)	Hatchery rack return goal of 18,800 adults.	No (hatchery exception).	Major component of OPI area fisheries. When abundant, significant contributors to ocean fisheries off California and north to Leadbetter Pt., Washington and to Columbia River fisheries. Current ocean fishery impacts from very limited retention fisheries north of Cape Falcon and incidental hook-and-release mortality in fisheries south of Cape Falcon.
Columbia River (Natural) Threatened, 2005	NMFS ESA consultation standard for 2008 is $\leq 8.0\%$ in total exploitation rate in marine and mainstem Columbia River fisheries.	No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program.	Extinct above the Dalles Dam, small populations in Clackamas, and Sandy rivers in Oregon, and Cedar Creek (Lewis River) Washington. Lower river coho are also listed under the Oregon State ESA.
<p>WASHINGTON COASTAL - All pertinent natural and hatchery stocks originating in Washington coastal streams north of the Columbia River through the western Strait of Juan de Fuca (West of the Elwha River). Management goals for Grays Harbor and Olympic Peninsula coho stocks include achieving natural spawning escapement objectives and treaty allocation requirements, although Grays Harbor also contains a significant amount of hatchery production. The conservation objectives for these stocks are based on MSY spawner escapements established pursuant to the U.S. District Court order in <u>Hoh v. Baldrige</u>. Annual natural spawning escapement targets and total escapement objectives are established by the WDFW and treaty tribes under the provisions of <u>U.S. v. Washington</u> and subsequent U.S. District Court orders. After agreement to annual targets is reached by the parties in this litigation, ocean fishery escapement objectives are established for each river, or region of origin, which include provisions for providing treaty allocation requirements and inside, non-Indian fishery needs. The conservation objectives for the Queets, Hoh, and Quillayute Rivers were developed as ranges intended to bracket the current best estimates of MSY escapement. The range of each objective reflects the degree of uncertainty inherent by using the high estimate of recruits-per-spawner and low estimate of carrying capacity for the lower bound and the low estimate of recruits-per-spawner with the high estimate of smolt carrying capacity for the upper end of the range. The ranges were subsequently adjusted upward for risk aversion and again for habitat considerations by 26% to 184% (Lestelle <i>et al.</i> 1984). These stocks are also subject to provisions of the 2002 PSC Coho Management Plan, which requires the United States and Canada to constrain total fishery exploitation rates to levels associated with the categorical status (low, moderate, and abundant) and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries are established through formulas specified in the PSC Management Plan. However, the salmon FMP management objectives determine the criteria for triggering a conservation alert or an overfishing concern; annual management objectives established pursuant to U.S. District Court orders and the PSC Coho Management Plan do not.</p>			
Willapa Bay (Hatchery)	Meet WDFW program objectives.	No (hatchery exception).	Contributes to ocean fisheries off northern Oregon north into Canada. Significant contributor to inside non-Indian commercial net and recreational fisheries. WDFW critically reviewing current management to determine if objectives for natural stocks are warranted.
Grays Harbor	35,400 natural adult spawners (MSP based on WDF [1979]) or annual target agreed to by WDFW and the Quinault Indian Nation.	Yes. Conservation alert or overfishing concern based on fewer than 35,400 natural spawners.	Ocean distribution from Oregon to northern British Columbia. Harvested by treaty Indian, non-Indian commercial, and recreational fisheries in Grays Harbor and tributary rivers.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 10 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
WASHINGTON COAST (continued)			
Queets	MSY range of 5,800 to 14,500 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and the Quinault Indian Nation.	Yes. Conservation alert or overfishing concern based on fewer than 5,800 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver. Coho supplementation project conducted since the late 1970s.
Hoh	MSY range of 2,000 to 5,000 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and Hoh Tribe.	Yes. Conservation alert or overfishing concern based on fewer than 2,000 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver.
Quillayute Fall	MSY range of 6,300 to 15,800 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and the Quillayute Tribe.	Yes. Conservation alert or overfishing concern based on fewer than 6,300 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver.
Quillayute Summer (Hatchery)	Meet hatchery program objectives.	No (hatchery exception).	Early river entry timing. Contributor to ocean fisheries off Washington north into British Columbia; present south to central Oregon.
Western Strait of Juan de Fuca (Sekiu, Hoko, Clallam, Pysht, East and West, and Lyre Rivers and miscellaneous streams west of the Elwha River)	11,900 natural adult spawners PSC 2007 annual management objective: 40% (low status) exploitation rate.	Yes. Overfishing concern based on fewer than 11,900 natural spawners.	Little information on ocean distribution.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 11 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
---COHO---			
<p>PUGET SOUND - All pertinent natural and hatchery stocks originating from U.S. tributaries to Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek). The Puget Sound Salmon Management Plan defines management objectives and long-term goals for these stocks as developed by representatives from federal, state, and tribal agencies. Conservation objectives for specific stocks are currently based on either MSP principles for stocks managed primarily for natural production or upon hatchery escapement needs for stocks managed for artificial production. Puget Sound management procedures are outlined in a "Memorandum Adopting Salmon Management Plan" (<u>U.S. v. Washington</u>, 626 F. Supp. 1405 [1985]). The original conservation objectives were developed by a State/Tribal Management Plan Development Team following the Boldt Decision with the goal for natural spawning stocks defined as "the adult spawning population that will, on the average, maximize biomass of juvenile outmigrants subsequent to incubation and freshwater rearing under average environmental conditions." The methodology used to develop the objectives was based on assessment of the quantity and quality of rearing habitat and the number of adult spawners required to fully seed the habitat (Zillges 1977). Some objectives have subsequently been modified under fixed procedures set by the U.S. District Court and its Fisheries Advisory Board (Clark 1983 and PSSSRG 1997) and later determinations of the WDFW/Tribal Technical Committee. These natural stocks are also subject to provisions of the 2002 PSC Coho Management Plan, which requires the United States and Canada to constrain total fishery exploitation rates to levels associated with the categorical status (low, moderate, and abundant) and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries are established through formulas specified in the PSC Management Plan. However, the salmon FMP management objectives determine criteria for triggering a conservation alert or an overfishing concern; annual management objectives established pursuant to U.S. District Court orders and the PSC Coho Management Plan do not.</p>			
Eastern Strait of Juan de Fuca (Streams east of Salt Creek through Chimacum Creek))	FMP: MSP objective of 950 natural adult spawners (Clark 1983 modified by habitat apportionment of WDFW/Tribal Technical Committee in 1998) or annual target agreed to in U.S. District Court procedures. The Elwha and Dungeness rivers are not included in this objective but are managed on a harvest rate basis. PSC 2008: 40% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 950 natural spawners.	Little information on ocean distribution.
Hood Canal	FMP: MSP objective of 21,500 natural adult spawners (Clark 1983 modified since 1994 by WDFW/Tribal Technical Committee) or annual target agreed to in U.S. District Court procedures. PSC 2008: 45% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 21,500 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
Skagit	FMP: MSP objective of 30,000 natural adult spawners (Zillges 1977 and Clark 1983) or annual target agreed to in U.S. District Court procedures. (The spawner assessment methodology is currently being revised and may result in an objective significantly different from 30,000.) PSC 2008: 35% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 30,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
Stillaguamish	FMP: MSP objective of 17,000 natural adult spawners (Zillges 1977) or annual target agreed to in U.S. District Court procedures. PSC 2008: 50% (normal status) total exploitation rate.	Yes. Overfishing concern based on fewer than 17,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
Snomish	FMP: MSP objective of 70,000 natural adult spawners (Zillges 1977 as modified by WDFW/Tribal Technical Committee) or annual target agreed to in U.S. District Court procedures. PSC 2008: 40% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 70,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
South Puget Sound (Hatchery)	Hatchery rack return goal of 52,000 adults. Natural production goals under development.	No (hatchery exception).	Ocean distribution from Cape Falcon, Oregon to British Columbia.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 12 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
SOUTHERN BRITISH COLUMBIA COAST - Stocks of southern British Columbia coastal streams (including Vancouver Island) and the Fraser River.			
Coastal Stocks	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty.	No. Not under Council management authority.	Contributes to ocean fisheries off British Columbia, north into southeast Alaska and present off northern Washington.
Fraser River	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty. For 2008, southern U.S. fisheries total exploitation rate of ≤10.0%.	No. Not under Council management authority.	Contributes to ocean fisheries off British Columbia and Washington, and to Strait of Juan de Fuca and Puget Sound fisheries.
--- PINK (odd-numbered years) ---			
The Fraser River Panel of the PSC manages fisheries for pink salmon in the Fraser River Panel Area (U.S.) north of 48° N latitude to meet Fraser River natural spawning escapement and U.S./Canada allocation requirements. The Council manages pink salmon harvests in that portion of the EEZ, which is not in the Fraser River Panel Area (U.S.) waters consistent with Fraser River Panel management intent. Pink salmon management objectives must address meeting natural spawning escapement objectives, allowing ocean pink harvest within fixed constraints of coho and Chinook harvest ceilings and providing for treaty allocation requirements.			
Puget Sound	900,000 natural spawners or consistent with provisions of the Pacific Salmon Treaty (Fraser River Panel).	No. Minor impacts in Council fisheries and not under Council management authority.	Contributes to ocean fisheries off British Columbia and in Puget Sound. Present south into Oregon. Rare off California.
Fraser River	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty (Fraser River Panel).	No. Minor impacts in Council fisheries and not under Council management authority.	Contributes to ocean fisheries off British Columbia; present into southeast Alaska and off Washington and northern Oregon. Rare off California.

TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13. (Page 1 of 1)

PARENT SPAWNER STATUS		MARINE SURVIVAL INDEX (based on return of jacks per hatchery smolt)			
		Low (<0.0009)	Medium (0.0009 to 0.0034)	High (>0.0034)	
		Allowable Total Fishery Impact Rate			
High:	Parent spawners achieved Level #2 rebuilding criteria, grandparent spawners achieved Level #1	≤15%	≤30% ^{a/}	≤35% ^{a/}	
Medium:	Parent spawners achieved Level #1 or greater rebuilding criteria	≤15%	≤20% ^{a/}	≤25% ^{a/}	
Low:	Parent spawners less than Level #1 rebuilding criteria	≤15% ≤10-13% ^{b/}	≤15%	≤15%	
OCN Coho Spawners by Stock Component					
Rebuilding Criteria	Northern	North-Central	South-Central	Southern	Total
Full Seeding at Low Marine Survival:	21,700	55,000	50,000	5,400	132,100
Level #2 (75% of full seeding):	16,400	41,300	37,500	4,100	99,300
Level #1 (50% of full seeding):	10,900	27,500	25,000	2,700	66,100
38% of Level #1 (19% of full seeding):	4,100	10,500	9,500	1,000	25,100
Stock Component (Boundaries)	Full Seeding of Major Basins at Low Marine Survival (Number of Adult Spawners)				
Northern: (Necanicum River to Neskowin Creek)	Nehalem	Tillamook	Nestucca	Ocean Tribs.	
	17,500	2,000	1,800	400	
North-Central: (Salmon River to Siuslaw River)	Siletz	Yaquina	Alsea	Siuslaw	Ocean Tribs.
	4,300	7,100	15,100	22,800	5,700
South-Central: (Siltcoos River to Sixes River)	Umpqua	Coos	Coquille	Coastal Lakes	
	29,400	7,200	5,400	8,000	
Southern: (Elk River to Winchuck River)	Rogue				
	5,400				

a/ When a stock component achieves a medium or high parent spawner status under a medium or high marine survival index, but a major basin within the stock component is less than 10% of full seeding, (1) the parent spawner status will be downgraded one level to establish the allowable fishery impact rate for that component, and (2) no coho-directed harvest impacts will be allowed within that particular basin.

b/ This exploitation rate criteria applies when (1) parent spawners are less than 38% of the Level #1 rebuilding criteria, or (2) marine survival conditions are projected to be at an extreme low as in 1994-1996 (<0.0006 jack per hatchery smolt). If parent spawners decline to lower levels than observed through 1998, rates of less than 10% would be considered, recognizing that there is a limit to further bycatch reduction opportunities.

TABLE A-3. Fishery **impact** rate criteria for **OCN coho** stock components based on the harvest matrix resulting from the **OCN work group** 2000 review of Amendment 13. (Page 1 of 1).

Parent Spawner Status ^{a/}	Marine Survival Index (based on return of jacks per hatchery smolt)						
	Extremely Low (<0.0008)	Low (0.0008 to 0.0014)	Medium (>0.0014 to 0.0040)	High (>0.0040)			
High Parent Spawners > 75% of full seeding	E ≤ 8%	J ≤ 15%	O ≤ 30%	T ≤ 45%			
Medium Parent Spawners > 50% & ≤ 75% of full seeding	D ≤ 8%	I ≤ 15%	N ≤ 20%	S ≤ 38%			
Low Parent Spawners > 19% & ≤ 50% of full seeding	C ≤ 8%	H ≤ 15%	M ≤ 15%	R ≤ 25%			
Very Low Parent Spawners > 4 fish per mile & ≤ 19% of full seeding	B ≤ 8%	G ≤ 11%	L ≤ 11%	Q ≤ 11%			
Critical^{b/} Parental Spawners ≤ 4 fish per mile	A 0 - 8%	F 0 - 8%	K 0 - 8%	P 0 - 8%			
Sub-aggregate and Basin Specific Spawner Criteria Data							
Sub-aggregate	Miles of Available Spawning Habitat	100% of Full Seeding	"Critical"		Very Low, Low, Medium & High		
			4 Fish per Mile	12% of Full Seeding	19% of Full Seeding	50% of Full Seeding	75% of full Seeding
Northern	899	21,700	3,596	NA	4,123	10,850	16,275
North - Central	1,163	55,000	4,652	NA	10,450	27,500	41,250
South - Central	1,685	50,000	6,740	NA	9,500	25,000	37,500
Southern	450	5,400	NA	648	1,026	2,700	4,050
Coastwide Total	4,197	132,100	15,636		25,099	66,050	99,075

a/ Parental spawner abundance status for the OCN aggregate assumes the status of the weakest sub-aggregate.

b/ "Critical" parental spawner status is defined as 4 fish per mile for the Northern, North-Central, and South-Central subaggregates. Because the ratio of high quality spawning habitat to total spawning habitat in the Rogue River Basin differs significantly from the rest of the basins on the coast, the spawner density of 4 fish per mile does not represent "Critical" status for that basin. Instead, "Critical" status for the Rogue Basin (Southern Sub-aggregate) is estimated as 12% of full seeding of high quality

**APPENDIX B
OREGON PRODUCTION INDEX DATA**

LIST OF TABLES

	<u>Page</u>
TABLE B-1. Millions of coho smolts released annually into the OPI area by geographic area and rearing agency	81
TABLE B-2. Data set used in predicting 2007 Oregon production index hatchery (OPIH) adult coho with Stratified Random Sampling accounting. Adults and jacks shown in thousands of fish and smolts in millions of fish	82
TABLE B-3. Estimated coho salmon natural spawner abundance (SRS accounting) in Oregon coastal basins for each OCN coho management component	83
TABLE B-4. Data set used in predicting 2007 Oregon coastal natural river (OCNR) coho ocean recruits with Stratified Random Sampling (SRS) accounting	84

TABLE B-1. Millions of coho smolts^{a/} released annually into the OPI area by geographic area and rearing agency. (Page 1 of 1)

Year or Average	Columbia River						Oregon Coast				
	Oregon	Washington			Federal	Total	ODFW ^{b/}	Private		California	Total OPI
		Early	Late	Combined				Yearlings	Total		
1960-1965	5.6	-	-	6.1	4.5	16.2	2.0	-	2.0	0.4	18.6
1966-1970	6.0	10.2	4.9	15.1	6.5	27.6	2.9	0.0	2.9	1.3	31.8
1971-1975	6.8	10.7	6.8	17.5	4.5	28.8	3.9	0.0	3.9	1.2	33.9
1976-1980	8.0	7.3	10.1	17.4	4.7	30.1	3.8	1.4	5.2	0.7	36.0
1981-1985	7.1	4.3	14.4	18.7	3.2	29.0	3.9	3.3	7.2	0.7	36.9
1986-1990	7.3	3.1	15.6	18.7	4.1	30.1	5.2	1.9	7.1	1.4	38.6
1991	10.4	3.7	15.3	19.0	5.9	35.2	5.3	-	5.3	1.5	42.0
1992	11.5	4.3	14.3	18.6	2.7	32.8	6.2	-	6.2	0.7	39.7
1993	11.1	4.3	14.8	19.1	4.1	34.3	4.3	-	4.3	0.8	39.4
1994	9.1	2.5	12.0	14.5	3.0	26.6	5.2	-	5.2	0.6	32.4
1995	7.1	3.4	12.9	16.3	1.7	25.1	3.7	-	3.7	0.7	29.5
1996	8.4	3.4	12.9	16.3	3.4	28.1	3.3	-	3.3	0.3	31.7
1997	6.1	3.2	7.8	11.0	3.9	21.0	2.9	-	2.9	0.7	24.6
1998	6.1	5.8	11.4	17.2	3.6	26.8	1.7	-	1.7	0.6	29.1
1999	7.6	4.0	11.5	15.5	4.8	27.9	1.0	-	1.0	0.7	29.6
2000	7.8	6.2	10.8	17.0	5.9	30.7	0.9	-	0.9	0.6	32.2
2001	7.6	4.2	9.7	13.9	3.7	25.2	0.9	-	0.9	0.6	26.7
2002	7.5	3.3	8.6	11.9	4.3	23.7	1.0	-	1.0	0.6	25.3
2003	8.2	3.3	8.7	12.0	3.1	23.3	0.8	-	0.8	0.5	24.6
2004	6.7	3.0	8.8	11.8	3.6	22.1	0.8	-	0.8	0.6	23.5
2005	6.1	2.5	9.1	11.6	2.8	20.6	0.8	-	0.8	0.6	22.0
2006	6.1	2.8	9.0	11.7	2.6	20.4	0.8	-	0.8	0.6	21.8
2007 ^{c/}	6.2	3.1	9.0	12.1	3.1	21.4	0.7	-	0.7	0.6	22.7

a/ Defined here as 30 fish per pound or larger and released in February or later.

b/ Beginning in 1989, does not include minor releases from STEP projects.

c/ Preliminary.

TABLE B-2. Data set used in predicting 2008 Oregon production index hatchery (OPIH) adult coho. Adults and jacks shown in thousands of fish and smolts in millions of fish. (Page 1 of 1)

Year	Adult OPIH ^{a/}	Columbia River Jacks ^{b/}	Oregon Coast/ California Jacks ^{c/}	Columbia River Smolts ^{d/}	Columbia River Delayed Smolts ^{e/}
1970	2,765.1	148.6	13.6	27.6	0.0
1971	3,365.0	172.8	6.6	24.0	0.0
1972	1,924.8	100.8	2.9	28.3	0.0
1973	1,817.0	85.7	5.7	29.9	1.8
1974	3,071.1	132.1	12.1	28.5	2.9
1975	1,652.8	75.1	1.1	27.8	1.8
1976	3,885.3	146.2	25.3	29.0	2.0
1977	987.5	46.2	7.5	28.9	0.2
1978	1,824.1	99.2	4.0	31.4	0.0
1979	1,476.7	64.1	8.4	32.6	5.0
1980	1,224.0	51.6	6.0	28.9	6.7
1981	1,064.5	40.6	8.1	28.1	5.6
1982	1,266.8	55.0	6.3	32.4	6.8
1983 ^{f/}	599.2	61.0	7.2	27.7	5.0
1984	691.3	28.1	3.6	27.0	5.1
1985	717.5	18.2	7.8	29.2	9.1
1986	2,416.6	64.6	12.9	28.8	12.2
1987	761.9	24.2	8.7	32.9	9.0
1988	1,479.9	72.3	12.9	28.8	7.7
1989	1,878.9	55.0	5.8	29.5	7.2
1990	673.5	37.1	9.6	29.6	8.5
1991	1,753.6	60.8	7.9	30.3	7.1
1992	482.9	19.9	5.7	35.3	6.0
1993	223.3	19.6	7.5	32.8	5.5
1994	214.3	3.9	1.3	34.4	6.0
1995	139.4	8.8	2.7	26.6	3.1
1996	176.5	14.1	3.2	25.2	4.2
1997	195.6	15.8	4.6	28.0	3.4
1998	228.7	6.8	3.0	21.0	2.5
1999	372.0	23.3	5.9	26.8	3.0
2000	617.7	31.2	3.5	27.9	4.1
2001	1,480.1	71.1	15.7	30.6	2.0
2002	688.9	18.9	6.3	23.5	1.4
2003	1,010.6	42.2	8.2	23.7	0.3
2004	692.2	29.4	6.0	23.2	2.0
2005	415.6	21.2	4.7	22.0	0.8
2006	431.5	20.9	5.4	20.6	0.4
2007	476.6	34.1	2.5	20.4	0.1
2008	216.1 ^{g/}	14.0	1.4	22.0	0.6

a/ Adult OPIH = Harvest impacts plus escapement for public hatchery stocks originating in the Columbia River, Oregon coastal rivers, and the Klamath River, California (1970-1985 with Stratified Random Sampling accounting; 1986-2007 with MSM abundance used for 2008).

b/ Jack CR = Columbia River jack returns corrected for small adults.

c/ Jack OC = Oregon coastal and California hatchery jack returns corrected for small adults.

d/ Sm CR = Columbia River smolt release from the previous year expected to return as adults in the year listed.

e/ Sm D = Columbia River delayed smolt releases from the previous year expected to return as adults in the year listed.

f/ Data not used in subsequent predictions due to El Niño impacts.

g/ Preseason predicted adults.

TABLE B-3. Estimated coho salmon natural spawner abundance (SRS accounting) in Oregon coastal basins for each OCN coho management component. Estimates adjusted for visual observation bias by multiplying observed count by 1.33. (Page 1 of 1)

Component and Basin ^{a/}	Miles	Adjusted SRS Natural Coho Spawner Estimates															1993-2007
		1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Avg.
NORTHERN																	
Nehalem	386	2,265	2,007	1,463	1,057	1,173	1,190	3,713	14,285	22,310	20,903	33,059	21,479	10,451	11,614	9,887	10,457
Tillamook	249	860	652	289	661	388	271	2,175	1,983	1,883	15,715	14,584	2,290	1,995	8,774	3,167	3,712
Nestucca	167	401	313	1,811	519	271	169	2,201	1,171	3,940	13,003	8,929	6,152	695	1,876	1,377	2,855
Ind. Tribs.	97	983	485	319	1,043	314	946	728	474	5,247	2,912	3,068	3,142	1,218	750	457	1,472
TOTAL	899	4,508	3,457	3,882	3,280	2,146	2,576	8,842	17,913	33,380	52,515	59,563	33,063	16,475	24,135	15,143	18,725
NORTH CENTRAL																	
Siletz	118	400	1,200	607	763	336	394	706	3,553	1,437	2,252	9,736	6,399	14,567	5,205	1,750	3,287
Yaquina	109	549	2,448	5,668	5,127	384	365	2,588	647	3,039	23,981	13,254	4,989	3,441	4,247	2,887	4,908
Alea	221	1,071	1,279	681	1,637	680	213	2,050	2,465	3,339	6,170	8,957	6,005	13,907	1,972	1,384	3,454
Siuslaw	514	4,428	3,205	6,089	7,625	668	1,089	2,724	6,767	11,024	57,129	29,257	8,443	16,907	5,869	2,743	10,931
Ind. Tribs.	201	1,331	1,683	560	2,975	774	1,222	3,691	817	5,636	10,371	7,664	14,558	2,589	3,931	1,195	3,933
TOTAL	1,163	7,779	9,815	13,605	18,127	2,842	3,283	11,442	14,261	25,239	99,506	66,550	40,393	51,411	21,224	9,959	26,362
SOUTH CENTRAL																	
Umpqua	1,083	10,244	5,336	12,809	10,824	2,960	9,153	7,685	12,233	35,702	37,591	29,607	31,346	42,676	18,154	11,253	18,505
Coos	208	15,284	14,685	10,351	12,128	1,127	3,167	4,945	5,386	43,301	35,688	29,559	24,116	17,048	11,266	1,414	15,298
Coquille	331	7,384	5,035	2,116	16,169	5,720	2,466	3,001	6,130	13,310	8,610	23,909	22,276	11,806	28,577	4,879	10,759
Coastal Lakes	-	10,145	5,841	11,216	13,493	8,603	11,107	12,710	12,747	19,669	22,162	16,688	18,687	14,724	24,378	8,885	14,070
Ind. Tribs.	83	-	-	-	-	-	-	-	-	-	-	-	-	-	1,104	342	723
TOTAL	1,622	43,057	30,897	36,492	52,614	18,410	25,893	28,341	36,496	111,982	104,051	99,763	96,425	86,254	83,479	26,773	58,728
SOUTH																	
Rogue ^{b/}	-	361	5,439	3,761	4,622	8,282	2,316	1,438	10,966	12,213	7,800	6,754	24,481	9,953	3,937	5,242	7,171
COASTWIDE	-	55,705	49,608	57,740	78,643	31,680	34,068	50,063	79,636	182,814	263,872	232,630	194,362	164,093	132,775	57,116	110,987

a/ The sum of the individual basins may not equal the aggregate totals, due to the use of independent estimates at different geographic scales.

b/ Mark recapture estimate based on seining at Huntley Park in the lower Rogue River.

TABLE B-4. Data set used in predicting Oregon coastal natural river (OCNR) coho ocean recruits with Stratified Random Sampling (SRS) accounting prior to 2008. Recruits shown in thousands of fish. (Page 1 of 1)

Year	Recruits to Ocean			
	SRS	Ln SRS	JanAnom ^{a/}	UpAnom (t-1) ^{b/}
1970	183.1	5.21003	0.307	-16.92
1971	416.3	6.03141	-1.293	30.08
1972	185.5	5.22305	-1.393	10.08
1973	235.0	5.45959	-0.493	23.08
1974	196.4	5.28015	-0.693	47.08
1975	208.4	5.33946	-0.493	48.08
1976	451.7	6.11302	-0.893	65.08
1977	161.2	5.08265	-0.193	32.08
1978	111.6	4.71492	1.207	17.08
1979	188.8	5.24069	-1.193	-2.92
1980	108.3	4.68491	0.507	17.08
1981	174.5	5.16192	1.607	-1.92
1982	185.7	5.22413	-0.093	-8.92
1983	96.0	4.56435	1.007	14.08
1984	94.7	4.55071	0.607	-24.92
1985	124.9	4.82751	0.007	-24.92
1986	97.9	4.58395	0.107	-24.92
1987	70.1	4.24992	0.507	-39.92
1988	124.4	4.82350	-0.093	-21.92
1989	103.8	4.64247	-0.493	-43.92
1990	60.4	4.10099	-0.007	-21.92
1991	68.8	4.23120	-0.893	-37.92
1992	86.9	4.46476	0.107	43.08
1993	81.1	4.39568	-0.593	7.08
1994	40.6	3.70377	1.107	-50.92
1995	47.6	3.86283	0.707	-3.92
1996	65.5	4.18205	1.807	-1.92
1997	16.3	2.79117	0.907	9.08
1998	21.7	3.07731	2.407	-24.92
1999	37.8	3.63231	-0.393	18.08
2000	58.9	4.07584	0.107	84.08
2001	161.4	5.08389	0.707	9.08
2002	266.5	5.58537	0.207	65.08
2003	249.4	5.51906	1.107	54.08
2004	175.2	5.16593	0.407	53.08
2005	134.4	4.90082	0.317	3.08
2006	116.4	4.75703	1.607	-34.92
2007	49.6	3.90399	-1.153	16.08
2008			-0.933	24.08

a/ JanAnom = The annual deviation from mean (1969-1996) January sea surface temperature (degrees Centigrade) at Charleston, Oregon.

b/ UpAnom = Annual deviation from mean (1946-1996) April-June Bakun upwelling index at 42° N latitude.

c/ Preseason adult prediction.

**APPENDIX C
SALMON HARVEST ALLOCATION SCHEDULES**

TABLE OF CONTENTS

	<u>Page</u>
HARVEST ALLOCATION -- SECTION 5.3 OF THE PACIFIC COAST SALMON PLAN.....	85
5.3 ALLOCATION.....	85
5.3.1 Commercial (Non-Tribal) and Recreational Fisheries North of Cape Falcon	85
5.3.1.1 Goal, Objectives, and Priorities.....	85
5.3.1.2 Allocation Schedule Between Gear Types	86
5.3.1.3 Recreational Subarea Allocations	87
5.3.2 Commercial and Recreational Fisheries South of Cape Falcon.....	89
 SELECTIVE FISHERY GUIDELINES -- SECTION 6.5 OF THE PACIFIC COAST SALMON PLAN	 92
6.5 SEASONS AND QUOTAS.....	92
6.5.3 Species-Specific and Other Selective Fisheries	92
6.5.3.1 Guidelines.....	92
6.5.3.2 Selective Fisheries Which May Change Allocation Percentages North of Cape Falcon	 92

HARVEST ALLOCATION -- SECTION 5.3 OF THE PACIFIC COAST SALMON PLAN

5.3 ALLOCATION

“Conservation and management measures shall not discriminate between residents of different states. If it becomes necessary to allocate or assign fishing privileges among various United States fishermen, such allocation shall be (A) fair and equitable to all such fishermen; (B) reasonably calculated to promote conservation; and (C) carried out in such manner that no particular individual, corporation, or other entity acquires an excessive share of such privileges.”

Magnuson-Stevens Act, National Standard 4

Harvest allocation is required when the number of fish is not adequate to satisfy the perceived needs of the various fishing industry groups and communities, to divide the catch between (non-Indian) ocean and inside fisheries and among ocean fisheries, and to provide treaty Indian fishing opportunity. In allocating the resource between ocean and inside fisheries, the Council considers both inriver harvest and spawner escapement needs. The magnitude of inriver harvest is determined by the states in a variety of ways, depending upon the management area. Some levels of inriver harvests are designed to accommodate federally recognized inriver Indian fishing rights, while others are established to allow for non-Indian harvests of historic magnitudes. Several fora exist to assist this process on an annual basis. The North of Cape Falcon Forum, a state and tribal sponsored forum, convenes the pertinent parties during the Council's pre-season process to determine allocation and conservation recommendations for fisheries north of Cape Falcon. The Klamath Fishery Management Council fulfills much the same roll with regard to Klamath River salmon stocks. The individual states also convene fishery industry meetings to coordinate their input to the Council.

5.3.1 Commercial (Non-Tribal) and Recreational Fisheries North of Cape Falcon

5.3.1.1 Goal, Objectives, and Priorities

Harvest allocations will be made from a total allowable ocean harvest which is maximized to the largest extent possible but still consistent with treaty obligations, state fishery needs and spawning escapement requirements, including jeopardy standards for stocks listed under the ESA. The Council shall make every effort to establish seasons and gear requirements which provide troll and recreational fleets a reasonable opportunity to catch the available harvest. These may include single-species directed fisheries with landing restrictions for other species.

The goal of allocating ocean harvest north of Cape Falcon is to achieve, to the greatest degree possible, the objectives for the commercial and recreational fisheries as follows:

- Provide recreational opportunity by maximizing the duration of the fishing season while minimizing daily and area closures and restrictions on gear and daily limits.
- Maximize the value of the commercial harvest while providing fisheries of reasonable duration.

The priorities listed below will be used to help guide establishment of the final harvest allocation while meeting the overall commercial and recreational fishery objectives.

At total allowable harvest levels up to 300,000 coho and 100,000 Chinook:

- Provide coho to the recreational fishery for a late June through early September all-species season. Provide Chinook to allow (1) access to coho and, if possible, (2) a minimal Chinook-only fishery prior to the all-species season. Adjust days per week and/or institute area restrictions to stabilize season duration.
- Provide Chinook to the troll fishery for a May and early June Chinook season and provide coho to (1) meet coho hooking mortality in June where needed and (2) access a pink salmon fishery in odd years. Attempt to ensure that part of the Chinook season will occur after June 1.

At total allowable harvest levels above 300,000 coho and above 100,000 Chinook:

- Relax any restrictions in the recreational all-species fishery and/or extend the all-species season beyond Labor Day as coho quota allows. Provide Chinook to the recreational fishery for a Memorial Day through late June Chinook-only fishery. Adjust days per week to ensure continuity with the all-species season.
- Provide coho for an all-salmon troll season in late summer and/or access to a pink fishery. Leave adequate Chinook from the May through June season to allow access to coho.

5.3.1.2 Allocation Schedule Between Gear Types

Initial commercial and recreational allocation will be determined by the schedule of percentages of total allowable harvest as follows:

TABLE 5-1. Initial commercial/recreational harvest allocation schedule north of Cape Falcon.

Coho			Chinook		
Harvest (thousands of fish)	Percentage ^{a/}		Harvest (thousands of fish)	Percentage ^{a/}	
	Troll	Recreational		Troll	Recreational
0-300	25	75	0-100	50	50
>300	60	40	>100-150	60	40
			>150	70	30

a/ The allocation must be calculated in additive steps when the harvest level exceeds the initial tier.

This allocation schedule should, on average, allow for meeting the specific fishery allocation priorities described above. The initial allocation may be modified annually by preseason and inseason trades to better achieve (1) the commercial and recreational fishery objectives and (2) the specific fishery allocation priorities. The final preseason allocation adopted by the Council will be expressed in terms of quotas which are neither guaranteed catches nor inflexible ceilings. Only the total ocean harvest quota is a maximum allowable catch.

To provide flexibility to meet the dynamic nature of the fisheries and to assure achievement of the allocation objectives and fishery priorities, deviations from the allocation schedule will be allowed as provided below and as described in Section 6.5.3.2 for certain selective fisheries.

1. Preseason species trades (Chinook and coho) which vary from the allocation schedule may be made by the Council based upon the recommendation of the pertinent recreational and commercial SAS representatives north of Cape Falcon. The Council will compare the socioeconomic impacts of any such recommendation to those of the standard allocation schedule before adopting the allocation which best meets FMP management objectives.

2. Inseason transfers, including species trades of Chinook and coho, may be permitted in either direction between recreational and commercial fishery quotas to allow for uncatchable fish in one fishery to be reallocated to the other. Fish will be deemed "uncatchable" by a respective commercial or recreational fishery only after considering all possible annual management actions to allow for their harvest which meet framework harvest management objectives, including single species or exclusive registration fisheries. Implementation of inseason transfers will require (a) consultation with the pertinent recreational and commercial SAS members and the STT and (b) a clear establishment of available fish and impacts from the transfer.
3. An exchange ratio of four coho to one Chinook shall be considered a desirable guideline for preseason trades. Deviations from this guideline should be clearly justified. Inseason trades and transfers may vary to meet overall fishery objectives. (The exchange ratio of four coho to one Chinook approximately equalizes the species trade in terms of average ex-vessel values of the two salmon species in the commercial fishery. It also represents an average species catch ratio in the recreational fishery.)
4. Any increase or decrease in the recreational or commercial total allowable catch (TAC), resulting from an inseason restructuring of a fishery or other inseason management action, does not require reallocation of the overall north of Cape Falcon non-Indian TAC.
5. The commercial TACs of Chinook and coho derived during the preseason allocation process may be varied by major subareas (i.e., north of Leadbetter Point and south of Leadbetter Point) if there is a need to do so to decrease impacts on weak stocks. Deviations in each major subarea will generally not exceed 50% of the TAC of each species that would have been established without a geographic deviation in the distribution of the TAC. Deviation of more than 50% will be based on a conservation need to protect the weak stocks and will provide larger overall harvest for the entire fishery north of Cape Falcon than would have been possible without the deviation. In addition, the actual harvest of coho may deviate from the initial allocation as provided in Section 6.5.3.2 for certain selective fisheries.
6. The recreational TACs of Chinook and coho derived during the preseason allocation process will be distributed among four major recreational port areas as described in the coho and Chinook distribution sections below. Additionally, based on the recommendations of the SAS members representing the ocean sport fishery north of Cape Falcon, the Council will include criteria in its preseason salmon management recommendations to guide any inseason transfer of coho among the recreational subareas to meet recreational season duration objectives. Inseason redistributions of quotas within the recreational fishery or the distribution of allowable coho catch transfers from the commercial fishery may deviate from the preseason distribution. The Council may also deviate from subarea quotas to (1) meet recreational season objectives based on agreement of representatives of the affected ports and (2) in accordance with Section 6.5.3.2 with regard to certain selective fisheries.

5.3.1.3 Recreational Subarea Allocations

Coho

The north of Cape Falcon preseason recreational TAC of coho will be distributed to provide 50% to the area north of Leadbetter Point and 50% to the area south of Leadbetter Point. The distribution of the

allocation north of Leadbetter point will vary, depending on the existence and magnitude of an inside fishery in Area 4B which is served by Neah Bay.

In years with no Area 4B fishery, the distribution of coho north of Leadbetter Point (50% of the total recreational TAC) will be divided to provide 74% to the area between Leadbetter Point and the Queets River (Westport), 5.2% to the area between Queets River and Cape Flattery (La Push), and 20.8% to the area north of the Queets River (Neah Bay). In years when there is an Area 4B (Neah Bay) fishery under state management, the allocation percentages north of Leadbetter Point will be modified to maintain more equitable fishing opportunity among the ports by decreasing the ocean harvest share for Neah Bay. This will be accomplished by adding 25% of the numerical value of the Area 4B fishery to the recreational TAC north of Leadbetter Point prior to calculating the shares for Westport and La Push. The increase to Westport and La Push will be subtracted from the Neah Bay ocean share to maintain the same total harvest allocation north of Leadbetter Point. Table 5-2 displays the resulting percentage allocation of the total recreational coho catch north of Cape Falcon among the four recreational port areas (each port area allocation will be rounded to the nearest hundred fish, with the largest quotas rounded downward if necessary to sum to the TAC).

TABLE 5-2. Percentage allocation of total allowable coho harvest among the four recreational port areas north of Cape Falcon.

Port Area	Without Area 4B Add-on		With Area 4B Add-on	
Columbia River	50.0%	50.0%		
Westport	37.0%	37.0%	plus 17.3% of the Area 4B add-on	
La Push	2.6%	2.6%	plus 1.2% of the Area 4B add-on	
Neah Bay	10.4%	10.4%	minus 18.5% of the Area 4B add-on	

Example distributions of the recreational coho TAC north of Leadbetter Point would be as follows:

Sport TAC North of Cape Falcon	Without Area 4B Add-On					With Area 4B Add-On ^{a/}					
	Columbia River	Westport	La Push	Neah Bay		Columbia River	Westport	La Push	Neah Bay		
									Ocean	Add-on	Total
50,000	25,000	18,500	1,300	5,200	25,000	19,900	1,400	3,700	8,000	11,700	
150,000	75,000	55,500	3,900	15,600	75,000	57,600	4,000	13,600	12,000	25,600	
300,000	150,000	111,000	7,800	31,200	150,000	114,500	8,000	27,500	20,000	47,500	

a/ The add-on levels are merely examples. The actual numbers in any year would depend on the particular mix of stock abundances and season determinations.

Chinook

Subarea distributions of Chinook will be managed as guidelines and shall be calculated by the STT with the primary objective of achieving all-species fisheries without imposing Chinook restrictions (i.e., area closures or bag limit reductions). Chinook in excess of all-species fisheries needs may be utilized by directed Chinook fisheries north of Cape Falcon or by negotiating a Chinook/coho trade with another fishery participant group.

Inseason management actions may be taken by NMFS Regional Director to assure that the primary objective of the Chinook harvest guidelines for each of the three recreational subareas north of Cape Falcon are met. Such actions might include: closure from 0 to 3, or 0 to 6, or 3 to 200, or 5 to 200 nautical miles from shore; closure from a point extending due west from Tatoosh Island for 5 miles, then south to a point due west of Umatilla Reef Buoy, then due east to shore; closure from North Head at the

Columbia River mouth north to Leadbetter Point; change species which may be landed; or other actions as prescribed in the annual regulations.

5.3.2 Commercial and Recreational Fisheries South of Cape Falcon

The allocation of allowable ocean harvest of coho salmon south of Cape Falcon has been developed to provide a more stable recreational season and increased economic benefits of the ocean salmon fisheries at varying stock abundance levels. When coupled with various recreational harvest reduction measures or the timely transfer of unused recreational allocation to the commercial fishery, the allocation schedule is designed to help secure recreational seasons extending at least from Memorial Day through Labor Day, assist in maintaining commercial markets even at relatively low stock sizes, and fully utilize available harvest. Total ocean catch of coho south of Cape Falcon will be treated as a quota to be allocated between troll and recreational fisheries as provided in Table 5-3.

(Note: The allocation schedule provides guidance only when coho abundance permits a directed coho harvest, not when the allowable impacts are insufficient to allow coho retention south of Cape Falcon. At such low levels, allocation of the allowable impacts will be accomplished during the Council's preseason process.)

TABLE 5-3. Allocation of allowable ocean harvest of coho salmon (thousands of fish) south of Cape Falcon.^{a/}

Total Allowable Ocean Harvest	Recreational Allocation		Commercial Allocation	
	Number	Percentage	Number	Percentage
≤100	≤100 ^{b/c/}	100 ^{b/}	b/	b/
200	167 ^{b/c/}	84 ^{b/}	33 ^{b/}	17 ^{b/}
300	200	67	100	33
350	217	62	133	38
400	224	56	176	44
500	238	48	262	52
600	252	42	348	58
700	266	38	434	62
800	280	35	520	65
900	290	32	610	68
1,000	300	30	700	70
1,100	310	28	790	72
1,200	320	27	880	73
1,300	330	25	970	75
1,400	340	24	1,060	76
1,500	350	23	1,150	77
1,600	360	23	1,240	78
1,700	370	22	1,330	78
1,800	380	21	1,420	79
1,900	390	21	1,510	79
2,000	400	20	1,600	80
2,500	450	18	2,050	82
3,000	500	17	2,500	83

a/ The allocation schedule is based on the following formula: first 150,000 coho to the recreational base (this amount may be reduced as provided in footnote b); over 150,000 to 350,000 fish, share at 2:1, 0.667 to troll and 0.333 to recreational; over 350,000 to 800,000 the recreational share is 217,000 plus 14% of the available fish over 350,000; above 800,000 the recreational share is 280,000 plus 10% of the available fish over 800,000.

Note: The allocation schedule provides guidance only when coho abundance permits a directed coho harvest, not when the allowable impacts are insufficient to allow general coho retention south of Cape Falcon. At such low levels, allocation of the allowable impacts will be determined in the Council's preseason process. Deviations from the allocation may also be allowed to meet consultation standards for ESA listed stocks (e.g., the 1998 biological opinion for California coastal coho requires no retention of coho in fisheries off California).

b/ If the commercial allocation is insufficient to meet the projected hook-and-release mortality associated with the commercial all-salmon-except-coho season, the recreational allocation will be reduced by the number needed to eliminate the deficit.

c/ When the recreational allocation is 167,000 coho or less, special allocation provisions apply to the recreational harvest distribution by geographic area (unless superseded by requirements to meet a consultation standard for ESA listed stocks); see text of FMP as modified by Amendment 11 allocation provisions.

The allocation schedule is designed to give sufficient coho to the recreational fishery to increase the probability of attaining no less than a Memorial Day to Labor Day season as stock sizes increase. This increased allocation means that, in many years, actual catch in the recreational fishery may fall short of its allowance. In such situations, managers will make an inseason reallocation of unneeded recreational coho to the south of Cape Falcon troll fishery. The reallocation should be structured and timed to allow the commercial fishery sufficient opportunity to harvest any available reallocation prior to September 1, while still assuring completion of the scheduled recreational season (usually near mid-September) and, in any event, the continuation of a recreational fishery through Labor Day. This reallocation process will

occur no later than August 15 and will involve projecting the recreational fishery needs for the remainder of the summer season. The remaining projected recreational catch needed to extend the season to its scheduled closing date will be a harvest guideline rather than a quota. If the guideline is met prior to Labor Day, the season may be allowed to continue if further fishing is not expected to result in any significant danger of impacting the allocation of another fishery or of failing to meet an escapement goal.

The allocation schedule is also designed to assure there are sufficient coho allocated to the troll fishery at low stock levels to ensure a full Chinook troll fishery. This hooking mortality allowance will have first priority within the troll allocation. If the troll allocation is insufficient for this purpose, the remaining number of coho needed for the estimated incidental coho mortality will be deducted from the recreational share. At higher stock sizes, directed coho harvest will be allocated to the troll fishery after hooking mortality needs for Chinook troll fishing have been satisfied.

The allowable harvest south of Cape Falcon may be further partitioned into subareas to meet management objectives of the FMP. Allowable harvests for subareas south of Cape Falcon will be determined by an annual blend of management considerations including:

1. abundance of contributing stocks
2. allocation considerations of concern to the Council
3. relative abundance in the fishery between Chinook and coho
4. escapement goals
5. maximizing harvest potential

Troll coho quotas may be developed for subareas south of Cape Falcon consistent with the above criteria. California recreational catches of coho, including projections of the total catch to the end of the season, would be included in the recreational allocation south of Cape Falcon, but the area south of the Oregon-California border would not close when the allocation is met; except as provided below when the recreational allocation is at 167,000 or fewer fish.

When the south of Cape Falcon recreational allocation is equal to or less than 167,000 coho:

1. The recreational fisheries will be divided into two major subareas, as listed in #2 below, with independent quotas (i.e., if one quota is not achieved or is exceeded, the underage or overage will not be added to or deducted from the other quota; except as provided under #3 below).
2. The two major recreational subareas will be managed within the constraints of the following impact quotas, expressed as a percentage of the total recreational allocation (percentages based on avoiding large deviations from the historical harvest shares):
 - a. Central Oregon (Cape Falcon to Humbug Mountain) - 70%
 - b. South of Humbug Mountain - 30%

In addition,

- (1) Horse Mountain to Point Arena will be managed for an impact guideline of 3 percent of the south of Cape Falcon recreational allocation, and

- (2) there will be no coho harvest constraints south of Point Arena. However, the projected harvest in this area (which averaged 1,800 coho from 1986-1990) will be included in the south of Humbug Mountain impact quota.
3. Coho quota transfers can occur on a one-for-one basis between subareas if Chinook constraints preclude access to coho.

SELECTIVE FISHERY GUIDELINES – SECTION 6.5 OF THE PACIFIC COAST SALMON PLAN

6.5 SEASONS AND QUOTAS

* * * * *

6.5.3 Species-Specific and Other Selective Fisheries

6.5.3.1 Guidelines

In addition to the all-species and single or limited species seasons established for the commercial and recreational fisheries, other species-limited fisheries, such as "ratio" fisheries and fisheries selective for marked or hatchery fish, may be adopted by the Council during the preseason regulatory process. In adopting such a fishery, the Council will consider the following guidelines:

Harvestable fish of the target species are available.

Harvest impacts on incidental species will not exceed allowable levels determined in the management plan.

Proven, documented, selective gear exists (if not, only an experimental fishery should be considered).

Significant wastage of incidental species will not occur or a written economic analysis demonstrates the landed value of the target species exceeds the potential landed value of the wasted species.

The species specific or ratio fishery will occur in an acceptable time and area where wastage can be minimized and target stocks are maximally available.

Implementation of selective fisheries for marked or hatchery fish must be in accordance with U.S. v. Washington stipulation and order concerning co-management and mass marking (Case No. 9213, Subproceeding No. 96-3) and any subsequent stipulations or orders of the U.S. District Court, and consistent with international objectives under the Pacific Salmon Treaty (e.g., to ensure the integrity of the coded-wire tag program).

6.5.3.2 Selective Fisheries Which May Change Allocation Percentages North of Cape Falcon

As a tool to increase management flexibility to respond to changing harvest opportunities, the Council may implement deviations from the specified port area allocations and/or gear allocations to increase harvest opportunity through fisheries that are selective for marked salmon stocks (e.g., marked hatchery salmon). The benefits of any selective fishery will vary from year to year and fishery to fishery depending on stock abundance, the mix of marked and unmarked fish, projected hook-and-release mortality rates, and public acceptance. These factors should be considered on an annual and case-by-case basis when utilizing selective fisheries. The deviations for selective fisheries are subordinate to the allocation priorities in Section 5.3.1.1 and may be allowed under the following management constraints:

Selective fisheries will first be considered during the months of August and/or September. However, the Council may consider selective fisheries at other times, depending on year to year circumstances identified in the preceding paragraph.

The total impacts within each port area or gear group on the critical natural stocks of management concern are not greater than those under the original allocation without the selective fisheries. Other allocation objectives (i.e., treaty Indian, or ocean and inside allocations) are satisfied during negotiations in the North of Cape Falcon Forum.

The selective fishery is assessed against the guidelines in Section 6.5.3.1.

Selective fishery proposals need to be made in a timely manner in order to allow sufficient time for analysis and public comment on the proposal before the Council finalizes its fishery recommendations.

If the Council chooses to deviate from the specified port and/or gear allocations, the process for establishing a selective fishery would be as follows:

Allocate the TAC among the gear groups and port areas according to the basic FMP allocation process described in Section 5.3.1 without the selective fishery.

Each gear group or port area may utilize the critical natural stock impacts allocated to its portion of the TAC to access additional harvestable, marked fish, over and above the harvest share established in step one, within the limits of the management constraints listed in the preceding paragraph.

APPENDIX D
CVI PREDICTOR: EXCLUSION OF 2005 DATA POINT

LIST OF FIGURES

	<u>Page</u>
FIGURE D-1. Regression diagnostics for the CVI predictor including the 2005 data point.	100
FIGURE D-2. Regression diagnostics for the CVI predictor excluding the 2005 data point.....	101

CVI Predictor: Exclusion of 2005 Data Point

The 2005 data point for the CVI predictor (see Figure II-1) was identified as an “outlier”. Its x -value is nearly 50% greater than the next highest x -value, while its y -value is near the mid-range of the rest of the y -values. This suggests that the 2005 data point may have considerable influence on the linear predictor, and if so, would significantly influence the predictor not only at the high end of x , but at the low end of x as well. This is a particular concern given the situation in 2008, where the jack return (x -value) being used to predict the CVI is the lowest on record (more than 50% lower than the previous low) and near the origin.

Figure D-I displays several regression diagnostics for the predictor including the 2005 data point. By all of these measures it is clear that the 2005 data point is not only an outlier, but has excessive leverage on the resulting predictor. Figure D-II displays the same diagnostics for the predictor excluding the 2005 data point. These measures are more consistent with a linear predictor model, although it is suggested that the variance of y increases with x and that a weighted regression model may be a more efficient alternative.

Based on this information, the STT has decided to exclude the 2005 data point from the 2008 CVI predictor, but not to alter the prediction methodology.







FIGURE D-1. Regression diagnostics for the CVI predictor including the 2005 data point.







FIGURE D-2. Regression diagnostics for the CVI predictor excluding the 2005 data point.

