

Agenda Item F.6.b
Supplemental GMT Report
April 2009

Projected mortality impacts (mt) of overfished groundfish species updated with most recent research estimates and fishery projections through April 2009.

| Fishery | Bocaccio b/ | Canary | Cowcod | Dkbl | POP | Widow | Yelloweye |
|---|---|------------|------------|------------|------------|------------|------------|
| Limited Entry Trawl- Non-whiting | 15.1 | 16.2 | 1.3 | 214.4 | 82.1 | 18.1 | 0.3 |
| Limited Entry Trawl- Whiting | | | | | | | |
| At-sea whiting motherships a/ | | 4.3 | | 6.0 | 0.5 | 60.0 | 0.0 |
| At-sea whiting cat-proc a/ | | 6.1 | | 8.5 | 0.5 | 85.0 | 0.0 |
| Shoreside whiting a/ | | 7.6 | | 10.5 | 0.1 | 105.0 | 0.0 |
| Tribal whiting | | 1.4 | | 0.0 | 0.7 | 3.7 | 0.0 |
| Tribal | | | | | | | |
| Midwater Trawl | | 3.6 | | 0.0 | 0.0 | 40.0 | 0.0 |
| Bottom Trawl | | 0.8 | | 0.0 | 3.7 | 0.0 | 0.0 |
| Troll | | 0.5 | | 0.0 | 0.0 | | 0.0 |
| Fixed gear | | 0.3 | | 0.0 | 0.0 | 0.0 | 2.3 |
| Fixed Gear Sablefish | 0.0 | 0.3 | 0.0 | 1.0 | 0.2 | 0.3 | 1.1 |
| Fixed Gear Nearshore | 0.3 | 3.3 | 0.0 | 0.0 | 0.0 | 0.3 | 1.2 |
| Fixed Gear Other | 5.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.7 | 0.0 |
| Open Access: Incidental Groundfish | 2.0 | 0.9 | 0.0 | 0.0 | 0.0 | 4.0 | 0.3 |
| Recreational Groundfish c/ | | | | | | | |
| WA | | 20.9 | | | | | 5.2 |
| OR | | | | | | 1.0 | |
| CA | 67.3 | 22.9 | 0.1 | | | 6.2 | 2.8 |
| EFPs | 13.7 | 2.7 | 0.3 | 1.3 | 0.0 | 5.5 | 0.3 |
| Research: Includes NMFS trawl shelf-slope surveys, the IPHC halibut survey, and expected impacts from SRPs and LOAs. | | | | | | | |
| | 2.0 | 8.0 | 0.2 | 2.0 | 2.0 | 5.7 | 2.4 |
| TOTAL | 105.4 | 99.8 | 1.9 | 252.7 | 89.8 | 335.5 | 15.9 |
| 2009 OY d/ | 288 | 105 | 4.0 | 285 | 189 | 522 | 17 |
| Difference | 182.6 | 5.2 | 2.1 | 32.3 | 99.2 | 186.5 | 1.1 |
| Percent of OY | 36.6% | 95.1% | 47.5% | 88.7% | 47.5% | 64.3% | 93.3% |
| Key | = either not applicable; trace amount (<0.01 mt); or not reported in available data | | | | | | |
| a/ Non-tribal whiting values for canary, darkblotched, and widow reflect bycatch limits for the non-tribal whiting sectors. | | | | | | | |
| b/ South of 40°10' N. lat. | | | | | | | |
| c/ Values in scorecard represent projected impacts for all species except canary and yelloweye rockfish, which are the prescribed harvest guidelines. | | | | | | | |
| d/ 2009 and 2010 OYs are the same except for darkblotched (291 mt in 2010), POP (200 mt in 2010), and widow (509 mt in 2010). | | | | | | | |

PFMC
04/09/09