

PRESEASON REPORT I
STOCK ABUNDANCE ANALYSIS
AND
ENVIRONMENTAL ASSESSMENT PART 1
FOR 2012 OCEAN SALMON FISHERY
REGULATIONS

REGULATION IDENTIFIER NUMBER 0648-XA921



Pacific Fishery Management Council
7700 NE Ambassador Place, Suite 101
Portland, OR 97220-1384
(503) 820-2280
www.pcouncil.org

FEBRUARY 2012

ACKNOWLEDGMENTS

SALMON TECHNICAL TEAM

DR. ROBERT KOPE, CHAIR

National Marine Fisheries Service, Seattle, Washington

DR. MICHAEL O'FARRELL, VICE-CHAIR

National Marine Fisheries Service, Santa Cruz, California

MR. CRAIG FOSTER

Oregon Department of Fish and Wildlife, Clackamas, Oregon

MR. LARRIE LAVOY

National Marine Fisheries Service, Seattle, Washington

MR. KEITH LUTZ

Northwest Indian Fisheries Commission, Olympia, Washington

MR. DOUG MILWARD

Washington Department of Fish and Wildlife, Olympia, Washington

MS. JENNIFER SIMON

California Department of Fish and Game, Santa Rosa, California

MR. HENRY YUEN

U.S. Fish and Wildlife Service, Vancouver, Washington

PACIFIC FISHERY MANAGEMENT COUNCIL STAFF

MR. CHUCK TRACY

MS. RENEE DORVAL

MS. KIM MERYDITH

MR. KRIS KLEINSCHMIDT

The Salmon Technical Team and the Council staff express their thanks for the expert assistance provided by Mr. Jeff Haymes, Ms. Cindy LeFleur, and Ms. Robin Ehlke, Washington Department of Fish and Wildlife; Mr. Eric Schindler, Oregon Department of Fish and Wildlife; Ms. Melodie Palmer-Zwahlen, California Department of Fish and Game; Ms. Sandy Zeiner, Northwest Indian Fisheries Commission; Dr. Pete Lawson, National Marine Fisheries Service, Northwest Fisheries Science Center, and numerous other agency and tribal personnel in completing this report.

This document may be cited in the following manner:

Pacific Fishery Management Council. 2012. *Preseason Report I: Stock Abundance Analysis and Environmental Assessment Part 1 for 2012 Ocean Salmon Fishery Regulations*. (Document prepared for the Council and its advisory entities.) Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, Oregon 97220-1384.



A report of the Pacific Fishery Management Council pursuant to National Oceanic and Atmospheric Administration Award Number NA10NMF4410014.

TABLE OF CONTENTS

| | <u>Page</u> |
|---|-------------|
| LIST OF TABLES | v |
| LIST OF FIGURES | vi |
| LIST OF ACRONYMS AND ABBREVIATIONS | vii |
| INTRODUCTION | 1 |
| Purpose and Need | 1 |
| STT Concerns | 3 |
| CHAPTER I: DESCRIPTION OF THE AFFECTED ENVIRONMENT | 4 |
| ABUNDANCE FORECASTS | 5 |
| ACCEPTABLE BIOLOGICAL CATCH, ANNUAL CATCH LIMITS, AND OVERFISHING LIMITS | 5 |
| Overfishing Limit | 5 |
| Acceptable Biological Catch | 6 |
| Annual Catch Limit | 6 |
| STATUS DETERMINATION CRITERIA | 6 |
| CHAPTER II: AFFECTED ENVIRONMENT - CHINOOK SALMON ASSESSMENT | 14 |
| CHINOOK STOCKS SOUTH OF CAPE FALCON | 14 |
| Sacramento River Fall Chinook | 14 |
| Predictor Description and Performance | 14 |
| Stock Forecast and Status | 14 |
| OFL, ABC, and ACL | 14 |
| Klamath River Fall Chinook | 15 |
| Predictor Description | 15 |
| Predictor Performance | 15 |
| Stock Forecast and Status | 15 |
| OFL, ABC, and ACL | 16 |
| Other California Coastal Chinook Stocks | 16 |
| Oregon Coast Chinook Stocks | 16 |
| Far-North and North Migrating Chinook (NOC and MOC groups) | 16 |
| South/Local Migrating Chinook (SOC group) | 17 |
| CHINOOK STOCKS NORTH OF CAPE FALCON | 19 |
| Columbia River Chinook | 19 |
| Predictor Description | 19 |
| Predictor Performance | 20 |
| Stock Forecasts and Status | 20 |
| Washington Coast Chinook | 21 |
| Predictor Description and Past Performance | 21 |
| Stock Forecasts and Status | 21 |
| Puget Sound Chinook | 21 |
| Predictor Description | 21 |
| Predictor Performance | 22 |
| Stock Forecasts and Status | 22 |
| STOCK STATUS DETERMINATION UPDATES | 22 |
| SELECTIVE FISHERY CONSIDERATIONS FOR CHINOOK | 23 |

TABLE OF CONTENTS (continued)

| | <u>Page</u> |
|--|-------------|
| CHAPTER III - COHO SALMON ASSESMENT..... | 46 |
| COLUMBIA RIVER AND OREGON/CALIFORNIA COAST COHO | 46 |
| (OREGON PRODUCTION INDEX AREA) | 46 |
| Public Hatchery Coho | 46 |
| Predictor Description | 46 |
| Predictor Performance | 47 |
| Stock Forecast and Status | 47 |
| Oregon Coastal Natural Coho | 47 |
| Predictor Description | 48 |
| Predictor Performance | 49 |
| Stock Forecasts and Status | 49 |
| Lower Columbia River Natural..... | 49 |
| Predictor Description | 49 |
| Predictor Performance | 49 |
| Stock Forecast and Status | 50 |
| Oregon Production Index Area Summary of 2012 Stock Forecasts | 50 |
| WASHINGTON COAST COHO | 50 |
| Willapa Bay | 50 |
| Predictor Description | 50 |
| Predictor Performance | 50 |
| Stock Forecasts and Status..... | 51 |
| Grays Harbor..... | 51 |
| Predictor Description | 51 |
| Predictor Performance | 51 |
| Stock Forecasts and Status..... | 51 |
| OFL | 51 |
| Quinault River..... | 51 |
| Predictor Description | 51 |
| Predictor Performance | 51 |
| Stock Forecasts and Status..... | 52 |
| Queets River..... | 52 |
| Predictor Description | 52 |
| Predictor Performance | 52 |
| Stock Forecasts and Status..... | 52 |
| OFL | 52 |
| Hoh River..... | 52 |
| Predictor Description | 52 |
| Predictor Performance | 53 |
| Stock Forecasts and Status..... | 53 |
| OFL | 53 |
| Quillayute River..... | 53 |
| Predictor Description | 53 |
| Predictor Performance | 54 |
| Stock Forecasts and Status..... | 54 |
| North Washington Coast Independent Tributaries..... | 54 |
| Predictor Description | 54 |
| Predictor Performance | 55 |
| Stock Forecasts and Status..... | 55 |

TABLE OF CONTENTS (continued)

| | <u>Page</u> |
|--|-------------|
| PUGET SOUND COHO STOCKS | 55 |
| Strait of Juan de Fuca..... | 55 |
| Predictor Description | 55 |
| Predictor Performance | 56 |
| Stock Forecasts and Status..... | 56 |
| OFL | 56 |
| Nooksack-Samish | 56 |
| Predictor Description | 56 |
| Predictor Performance | 56 |
| Stock Forecasts and Status..... | 56 |
| Skagit | 57 |
| Predictor Description | 57 |
| Predictor Performance | 57 |
| Stock Forecasts and Status..... | 57 |
| OFL | 57 |
| Stillaguamish..... | 57 |
| Predictor Description | 57 |
| Predictor Performance | 57 |
| Stock Forecasts and Status..... | 57 |
| OFL | 58 |
| Snohomish..... | 58 |
| Predictor Performance | 58 |
| Stock Forecasts and Status..... | 58 |
| OFL | 58 |
| Hood Canal | 58 |
| Predictor Description | 58 |
| Predictor Performance | 59 |
| Stock Forecasts and Status..... | 59 |
| OFL | 59 |
| South Sound | 59 |
| Predictor Description | 59 |
| Stock Forecasts and Status..... | 59 |
| STOCK STATUS DETERMINATION UPDATES | 59 |
| SELECTIVE FISHERY CONSIDERATIONS FOR COHO | 60 |
| CHAPTER IV: AFFECTED ENVIRONMENT - PINK SALMON ASSESSMENT..... | 72 |

TABLE OF CONTENTS (continued)

| | <u>Page</u> |
|--|-------------|
| CHAPTER V: DESCRIPTION AND ANALYSIS OF THE NO ACTION ALTERNATIVE..... | 73 |
| DESCRIPTION OF THE NO-ACTION ALTERNATIVE..... | 73 |
| ANALYSIS OF EFFECTS ON THE ENVIRONMENT OF THE NO-ACTION ALTERNATIVE..... | 73 |
| Overview | 73 |
| Sacramento River Fall Chinook | 73 |
| Klamath River Fall Chinook | 74 |
| California Coastal Chinook Stocks | 74 |
| Oregon Coast Chinook Stocks | 74 |
| Columbia River Chinook Stocks..... | 74 |
| Washington Coast and Puget Sound Chinook Stocks..... | 74 |
| Oregon Production Index Area Coho Stocks | 74 |
| Washington Coast, Puget Sound, and Canadian Coho Stocks | 75 |
| Summary | 75 |
| Conclusion | 76 |
| CHAPTER VI: REFERENCES | 91 |
| APPENDIX A SUMMARY OF COUNCIL STOCK MANAGEMENT GOALS | 93 |
| APPENDIX B SALMON HARVEST ALLOCATION SCHEDULES | 105 |
| APPENDIX C OREGON PRODUCTION INDEX DATA | 119 |

LIST OF TABLES

| | <u>Page</u> |
|---|-------------|
| TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish | 8 |
| TABLE I-2. Preseason adult coho salmon stock forecasts in thousands of fish | 12 |
| TABLE II-1. Harvest and abundance indices for Sacramento River fall Chinook (SRFC) in thousands of fish. | 24 |
| TABLE II-2. Klamath River fall Chinook ocean abundance (thousands), harvest rate, and river run size estimates (thousands) by age | 25 |
| TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook | 26 |
| TABLE II-4. Summary of management objectives and predictor performance for Klamath River fall Chinook | 30 |
| TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook | 31 |
| TABLE II-6. Rogue River fall Chinook inriver run and ocean population indices..... | 35 |
| TABLE II-7. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish | 36 |
| TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish..... | 39 |
| TABLE III-1. Preliminary 1996-2011 preseason and postseason coho stock abundance estimates for Oregon production index area stocks in thousands of fish..... | 61 |
| TABLE III-2. Oregon production index (OPI) area coho harvest impacts, spawning, abundance, and exploitation rate estimates in thousands of fish. | 64 |
| TABLE III-3. Preseason forecasts and postseason estimates of ocean escapements for selected Washington coastal adult natural coho stocks in thousands of fish..... | 65 |
| TABLE III-4. Preseason forecasts and postseason estimates of ocean escapements for selected Puget Sound adult natural coho stocks in thousands of fish | 66 |
| TABLE III-5. Status categories and constraints for Puget Sound and Washington Coast coho under the FMP and PST Southern Coho Management Plan. | 68 |
| TABLE III-6. Projected coho mark rates for 2012 fisheries under base period fishing patterns (percent marked) | 69 |
| TABLE IV-1. Estimated annual (odd-numbered years) run sizes and forecasts for Fraser River and Puget Sound pink salmon in millions of fish. | 72 |
| TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011..... | 77 |
| TABLE V-2. Recreational management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011 | 82 |
| TABLE V-3. Treaty Indian ocean troll management measures adopted by the Council for ocean salmon fisheries, 2011 | 86 |
| TABLE V-4. Stock status relative to overfished and overfishing criteria. | 87 |
| TABLE V-5. Estimated ocean escapements and exploitation rates for critical natural and Columbia River hatchery coho stocks (thousands of fish) based on preliminary 2012 preseason abundance forecasts and 2011 Council management measures | 88 |

LIST OF TABLES (continued)

| | | <u>Page</u> |
|------------|--|-------------|
| TABLE V-6. | Comparison of Lower Columbia natural (LCN), Oregon coastal natural (OCN), and Rogue/Klamath (RK) coho projected harvest mortality and exploitation rates by fishery under Council-adopted 2011 management measures and preliminary 2012 preseason abundance estimates..... | 89 |
| TABLE V-7 | Maximum allowable fishery impact rate for OCN coho under Amendment 13 matrix and the OCN work group matrix based on parent escapement levels by stock component and marine survival category..... | 90 |

LIST OF FIGURES

| | | <u>Page</u> |
|----------------|---|-------------|
| FIGURE II-1. | The Sacramento Index (SI) and relative levels of its components. The Sacramento River fall Chinook escapement goal range of 122,000-180,000 adult spawners is noted on the vertical axis..... | 43 |
| FIGURE II-2. | Regression estimator for the SI based on previous year's escapement of Sacramento River fall Chinook jacks, 1990-2010. Years shown are SI years..... | 43 |
| FIGURE II-3. | Regression estimators for Klamath River fall Chinook ocean abundance (September 1) based on that year's river return of same cohort | 44 |
| FIGURE II-4. | Selected preseason vs. postseason forecasts for Chinook stocks with significant contribution to Council area fisheries. | 45 |
| FIGURE III-1a. | Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries. | 70 |
| FIGURE III-1b. | Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries | 71 |

LIST OF ACRONYMS AND ABBREVIATIONS

| | |
|-----------|--|
| ABC | acceptable biological catch |
| ACL | annual catch limit |
| BY | brood year |
| CDFG | California Department of Fish and Game |
| CoTC | Coho Technical Committee (of the PSC) |
| Council | Pacific Fishery Management Council |
| CRFMP | Columbia River Fishery Management Plan |
| CVI | Central Valley Index |
| CWT | coded-wire tag |
| EA | Environmental Assessment |
| EEZ | exclusive economic zone (from 3-200 miles from shore) |
| EIS | Environmental Impact Statement |
| EMAP | Environmental Monitoring and Assessment Program |
| ESA | Endangered Species Act |
| ESU | evolutionarily significant unit |
| F_{ABC} | exploitation rate associated with ABC |
| F_{ACL} | exploitation rate associated with ACL ($= F_{ABC}$) |
| FMP | fishery management plan |
| F_{MSY} | MSY exploitation rate |
| F_{OFL} | exploitation rate associated with the overfishing limit ($= F_{MSY}$, MFMT) |
| FONSI | Finding of No Significant Impacts |
| FRAM | Fishery Regulatory Assessment Model |
| GAM | generalized additive models |
| ISBM | individual stock-based management |
| Jack CR | Columbia River jacks (coho) |
| Jack OC | Oregon coastal and Klamath River Basin jacks (coho) |
| Jack OPI | Jack CR + Jack OC (coho) |
| KMZ | Klamath management zone (ocean zone between Humbug Mountain and Horse Mountain where management emphasis is on Klamath River fall Chinook) |
| KOHM | Klamath Ocean Harvest Model |
| KRFC | Klamath River fall Chinook |
| KRTT | Klamath River Technical Team |
| LCN | lower Columbia River natural (coho) |
| LCR | lower Columbia River (natural tule Chinook) |
| LRB | lower Columbia River bright (Chinook) |
| LRH | lower Columbia River hatchery (tule fall Chinook returning to hatcheries below Bonneville Dam) |
| LRW | lower Columbia River wild (bright fall Chinook spawning naturally in tributaries below Bonneville Dam) |
| MCB | mid-Columbia River brights (bright hatchery fall Chinook released below McNary Dam) |
| MFMT | maximum fishery mortality threshold |
| MOC | mid-Oregon coast |
| MSST | minimum stock size threshold |
| MSM | mixed stock model |
| MSA | Magnuson-Stevens Act |
| MSY | maximum sustainable yield |
| NA | not available |

LIST OF ACRONYMS AND ABBREVIATIONS (continued)

| | |
|-----------|--|
| NEPA | National Environmental Policy Act |
| NMFS | National Marine Fisheries Service |
| NOC | north Oregon coast |
| OCN | Oregon coast natural (coho) |
| OCNL | Oregon coast natural lake (coho) |
| OCNR | Oregon coast natural river (coho) |
| ODFW | Oregon Department of Fish and Wildlife |
| OFL | overfishing limit |
| OPI | Oregon Production Index (coho salmon stock index south of Leadbetter Point) |
| OPIH | Oregon Production Index public hatchery |
| OPITT | Oregon Production Index Technical Team |
| OY | Optimum Yield |
| PDO | Pacific Decadal Oscillation |
| PFMC | Pacific Fishery Management Council (Council) |
| PRIH | Private hatchery |
| PSC | Pacific Salmon Commission |
| PST | Pacific Salmon Treaty |
| RER | rebuilding exploitation rate |
| RK | Rogue/Klamath (coho) |
| RMP | Resource Management Plan (for exemption from ESA section 9 take prohibitions under limit 6 of the 4(d) rule) |
| ROPI | Rogue Ocean Production Index (Chinook) |
| SAB | Select Area brights |
| S_{ABC} | spawning escapement associated with ABC |
| S_{ACL} | spawning escapement associated with ACL ($= F_{ABC}$) |
| SCH | Spring Creek Hatchery (tule fall Chinook returning to Spring Creek Hatchery) |
| SHM | Sacramento Harvest Model |
| SI | Sacramento Index |
| SJF | Strait of Juan de Fuca |
| S_{MSY} | MSY spawning escapement |
| S_{OFL} | spawning escapement associated with the overfishing limit ($= S_{MSY}$) |
| SOC | south Oregon Coast |
| SRFC | Sacramento River fall Chinook |
| SRS | Stratified Random Sampling |
| STEP | Salmon Trout Enhancement Program |
| STT | Salmon Technical Team (formerly the Salmon Plan Development Team) |
| TAC | Technical Advisory Committee (<i>U.S. v. Oregon</i>) |
| URB | upper river brights (naturally spawning bright fall Chinook normally migrating past McNary Dam) |
| VSI | visual stock identification |
| WCVI | West Coast Vancouver Island |
| WDFW | Washington Department of Fish and Wildlife |

INTRODUCTION

This is the second report in an annual series of four reports prepared by the Salmon Technical Team (STT) of the Pacific Fishery Management Council (Council) to document and help guide salmon fishery management off the coasts of Washington, Oregon, and California. The report focuses on Chinook, coho, and pink salmon stocks that have been important in determining Council fisheries in recent years, and on stocks listed under the Endangered Species Act (ESA) with established National Marine Fisheries Service (NMFS) ESA consultation standards. This report will be formally reviewed at the Council's March 2012 meeting.

This report provides 2012 salmon stock abundance forecasts, and an analysis of the impacts of 2011 management measures, or regulatory procedures, on the projected 2012 abundance. This analysis is intended to give perspective in developing 2012 management measures. This report also constitutes the first part of an Environmental Assessment (EA) to comply with National Environmental Policy Act (NEPA) requirements for the 2012 ocean salmon management measures. An EA is used to determine whether an action being considered by a Federal agency has significant impacts. This part of the EA includes a statement of the purpose and need, a summary description of the affected environment, a description of the No-Action Alternative, and an analysis of the No-Action Alternative effects on the salmon stocks included in the Council's Salmon Fishery management Plan (FMP).

The STT and Council staff will provide two additional reports prior to the beginning of the ocean salmon season to help guide the Council's selection of annual fishery management measures: Preseason Report II and Preseason Report III. These reports will analyze the impacts of the Council's proposed alternatives and adopted fishery management recommendations. Preseason Report II will constitute the second part of the EA, and will include additional description of the affected environment relevant to the alternative management measures considered for 2012 ocean salmon fisheries, a description of the alternatives, and an analysis of the environmental consequences of the alternatives. Preseason Report II will analyze the potential impacts of a reasonable range of alternatives, which will inform the final fishery management measures included in Preseason Report III. Preseason Report III will describe and analyze the effects of the Council's final proposed action. Together, these parts of the EA will provide the necessary components to determine if a finding of no significant impact (FONSI) or Environmental Impact Statement (EIS) is warranted.

Chapter I provides a summary of stock abundance forecasts. Chapters II and III provide detailed stock-by-stock analyses of abundance, a description of prediction methodologies, and accuracy of past abundance forecasts for Chinook and coho salmon, respectively. Chapter IV summarizes abundance and forecast information for pink salmon. Chapter V provides an assessment of 2011 regulations applied to 2012 abundance forecasts. Three appendices provide supplementary information as follows: Appendix A provides a summary of Council stocks and their management objectives; Appendix B contains the Council's current harvest allocation schedules, and; Appendix C contains pertinent data for Oregon production index (OPI) area coho. For NEPA purposes, Chapters I-IV of this document describe the affected environment and Chapter V provides a description and analysis of the No-Action Alternative.

Purpose and Need

The purpose of this action, implementation of the 2012 ocean salmon fishery management measures, is to allow fisheries to harvest surplus production of healthy natural and hatchery salmon stocks within the constraints specified under the Salmon FMP, the Pacific Salmon Treaty (PST), and consultation standards established for ESA listed salmon stocks. In achieving this purpose, management measures must take into account the allocation of harvest among different user groups and port areas. Without this action, 2011 management measures would be in effect, which do not consider changes in abundance of stocks in the mixed stock ocean salmon fisheries. Therefore, this action is needed to ensure constraining stocks are

not overharvested and that harvest of abundant stocks can be optimized and achieve the most overall benefit to the nation.

This action will also establish a rebuilding plan for Sacramento River fall Chinook (SRFC), which were determined to be overfished in 2010. This is needed to comply with the MSA requirement for adopting and implementing a rebuilding plan for an overfished stock within two years of an overfished status determination.

The Salmon FMP also establishes nine more general harvest-related objectives:

1. Establish ocean exploitation rates for commercial and recreational salmon fisheries that are consistent with requirements for stock conservation objectives, specified ESA consultation standards, or Council adopted rebuilding plans.
2. Fulfill obligations to provide for Indian harvest opportunity as provided in treaties with the United States, as mandated by applicable decisions of the Federal courts, and as specified in the October 4, 1993, opinion of the Solicitor, Department of Interior, with regard to Federally-recognized Indian fishing rights of Klamath River Tribes.
3. Maintain ocean salmon fishing seasons that support established recreational and commercial fisheries, while meeting salmon harvest allocation objectives among ocean and inside recreational and commercial fisheries. These allocations will be fair and equitable, and fishing interests shall equitably share the obligations of fulfilling any treaty or other legal requirements for harvest opportunities.
4. Minimize fishery mortalities for those fish not landed from all ocean salmon fisheries as consistent with achieving optimum yield (OY) and bycatch management specifications.
5. Manage and regulate fisheries, so the OY encompasses the quantity and value of food produced, the recreational value, and the social and economic values of the fisheries.
6. Develop fair and creative approaches to managing fishing effort and evaluate and apply effort management systems as appropriate to achieve these management objectives.
7. Support the enhancement of salmon stock abundance in conjunction with fishing effort management programs to facilitate economically viable and socially acceptable commercial, recreational, and tribal seasons.
8. Achieve long-term coordination with the member states of the Council, Indian tribes with Federally recognized fishing rights, Canada, the North Pacific Fishery Management Council, Alaska, and other management entities which are responsible for salmon habitat or production. Manage consistent with the Pacific Salmon Treaty and other international treaty obligations.
9. In recommending seasons, to the extent practicable, promote the safety of human life at sea.

These objectives, along with the conservation objectives established under the ESA, provide "sideboards" for setting management measures necessary to implement the Salmon FMP, which conforms to the terms and requirements of the Magnuson-Stevens Act (MSA) and the National Standards Guidelines.

Implementation of 2012 management measures will allow fisheries to harvest surplus production of healthy natural and hatchery salmon stocks within the constraints specified under the Salmon FMP and consultation standards established for ESA-listed salmon stocks.

The reauthorization of the MSA in 2006 established new requirements to end and prevent overfishing through specification of overfishing limits (OFL) acceptable biological catch (ABC), annual catch limits (ACLs) and accountability measures (AMs). Because OFLs, ABCs, and ACLs are based on annual abundance forecasts, Preseason Report I also specifies OFLs, ABCs, and ACLs for 2012 fisheries.

STT Concerns

The Sacramento Index (SI) forecast has exceeded its postseason estimate for three consecutive years (2009-2011). In response to these over forecasts and the markedly different pattern in the jack escapement to SI relationship since 2009, the STT based the 2012 forecast on data from 2009-2011 rather than the longer data range that has been used previously.

As with the SI forecasts made for 2009-2011, the 2012 SI forecast is being made under conditions where the most recent jack escapement estimate, the largest on record for SRFC, exceeds the jack escapement estimate from the previous year by a large margin. Under such conditions, there has been a tendency to over predict the SI. The 2012 modification to the data range used for the SI forecast is intended to account for this and other factors that have likely contributed to recent forecast errors.

Age-specific escapement and river harvest data can enable the development of age-specific abundance forecasts, which would likely reduce the errors associated with forecasting a combined-age index (SI) with information from a single year class (jack escapement). The STT encourages the continued development of the scale ageing program and continuation of coded-wire tag (CWT) collection programs in the Sacramento Basin, which will help address some of these concerns.

CHAPTER I: DESCRIPTION OF THE AFFECTED ENVIRONMENT

The affected environment relevant to establishing the 2012 ocean salmon fishery management measures consists of the following components:

- Target Species – Chinook, coho, and pink salmon
- ESA-listed salmon stocks
- Socioeconomic aspects of coastal communities, federally recognized Tribes, and states

A description of the historical baseline for these components of the affected environment is presented in the Review of 2011 Ocean Salmon Fisheries (PFMC 2012). The current status (2012 ocean abundance forecasts) of the environmental components expected to be affected by the 2012 ocean salmon fisheries regulation alternatives (FMP salmon stocks) are described in this report (Part 1 of the 2012 salmon EA); the Review of 2011 Ocean Salmon Fisheries (PFMC 2012) provides an historical description of the salmon fishery-affected environment, including stock status and socioeconomic impacts, and represents the current status of the socioeconomic component of the affected environment.

The No-Action alternative was assessed in the 2011 NEPA process for ocean salmon regulations (Preseason Reports II and III; PFMC 2011a and 2011b). In those analyses, several components of the affected environment were determined to have no significant impacts. These components included:

- Non-target species – Pacific Halibut, groundfish (NMFS 2003; PFMC 2006, 2011a)
- Marine mammals – pinnipeds, killer whales (NMFS 2003, 2008; PFMC 2006, 2011a)
- Seabirds (NMFS 2003; PFMC 2006, 2011a)
- Ocean and coastal habitats, ESA critical habitat, and essential fish habitat (NMFS 2003; PFMC 2006, 2011a)
- Biodiversity and ecosystem function (NMFS 2003; PFMC 2006, 2011a)
- Unique characteristics of the geographic area (NMFS 2003; PFMC 2006, 2011a)
- Cultural, scientific, or historical resources such as those eligible for listing in the National Register of Historic Places (NMFS 2003; PFMC 2006, 2011a)
- Public health or safety (NMFS 2003; PFMC 2006, 2011a)

The 2012 No-Action alternative is not expected to differ from the 2011 action in any ways that would change the effects of the action on these elements of the environment.

The component of the affected environment that is analyzed in this document consists only of the salmon stocks identified in the FMP (Appendix A). The 2012 forecast abundance of the FMP salmon stocks represents this component of the affected environment. The surviving stock after fishery-related mortality is generally referred to as spawning escapement (S), and the proportion of the stock that succumbs to fishing related mortality is generally referred to as the exploitation rate (F); these are the metrics that constitute conservation objectives for FMP stocks, and by which effects of the alternatives to this part of the affected environment are evaluated. Thus, application of management measures (alternatives) to the abundance forecasts (affected environment) results in projected exploitation rates and spawning escapements (effects).

A description of the other components of the affected environment considered for 2012 ocean salmon fishery regulation alternatives, including socioeconomic components and updated additional information on the biological components of the environment, will be presented in the Preseason Report II, to be issued after the March Council meeting.

ABUNDANCE FORECASTS

Abundance forecasts in 2012 are summarized for key Chinook and coho salmon stocks in Tables I-1 and I-2, respectively. A cursory comparison of preseason forecast and postseason abundance estimates for selected stocks is presented in Figures I-1 and I-2. More detailed analyses of this subject are covered in Chapters II (Chinook) and III (coho). Information on pink salmon abundance and forecasts, which are only significant in odd-numbered years, is contained in Chapter IV. Council Salmon Fishery Management Plan (FMP) conservation objectives are presented in Appendix A; allocation objectives are presented in Appendix B.

In addition to the key stocks with abundance forecasts listed in Tables I-1 and I-2, Council management decisions for the 2012 ocean salmon fishing seasons may be constrained by other stocks, such as those listed under the ESA or subject to PSC agreements, which may not have abundance forecasts made, or do not have abundance forecasts available in time for inclusion in this report. These include the following Evolutionarily Significant Units (ESUs): Sacramento River Winter, Central Valley Spring, California Coastal, Lower Columbia River (LCR) natural tule, and Snake River Fall Chinook; and Central California and Southern Oregon/Northern California coho, as well as Interior Fraser (including Thompson River) coho.

ACCEPTABLE BIOLOGICAL CATCH, ANNUAL CATCH LIMITS, AND OVERFISHING LIMITS

Amendment 16 to the Salmon FMP was approved in December 2011 to comply with the requirements of the 2006 MSA reauthorization, including specification of acceptable biological catch (ABC) and annual catch limits (ACLs), overfishing limits (OFLs), and Scientific and Statistical Committee (SSC) recommendations for ABC. Amendment 16 established that ABC and ACLs were required for two stocks, Sacramento River fall Chinook (SRFC) and Klamath River fall Chinook (KRFC), which serve as indicator stocks for the Central Valley Fall and Southern Oregon/Northern California Chinook complexes, respectively. Other stocks in the FMP were not required to have ACLs either because they were components of these two stock complexes, or they were ESA-listed, hatchery stocks, or managed under an international agreement.

ABC and ACLs are not specified for stocks that are managed under an international agreement as there is a statutory exception in the MSA to the requirement for ACLs, and the NSIGs state that an ABC is not required if stocks meet this international exception. The NSIGs allow the flexibility to consider alternative approaches for specifying ACLs for stocks with unusual life history characteristics like Pacific salmon, and particularly for species listed under the ESA and hatchery stocks. For hatchery stocks, broodstock goals serve as conservation objectives rather than specifying ACLs. For ESA stocks, biological opinions and associated consultation standards provide necessary controls to ensure their long-term conservation.

Preseason OFLs are determined for all non-ESA-listed and non-hatchery stocks with an estimate of F_{MSY} (or MFMT) and sufficient information available to make abundance forecasts.

Overfishing Limit

For salmon, OFL is defined in terms of spawner escapement (S_{OFL}), which is consistent with the common practice of using spawner escapement to assess stock status for salmon. S_{OFL} is determined annually based on stock abundance, in spawner equivalent units (N) and the exploitation rate F_{OFL} .

F_{OFL} is defined as being equal to F_{MSY} (or MFMT) and

$$S_{OFL} = N \times (1 - F_{MSY}).$$

Acceptable Biological Catch

For salmon, ABC is defined in terms of spawner escapement (S_{ABC}), which is determined annually based on stock abundance, in spawner equivalent units (N) and the exploitation rate F_{ABC} .

$$S_{ABC} = N \times (1 - F_{ABC}).$$

The ABC control rule defines F_{ABC} as a fixed exploitation rate reduced from F_{MSY} to account for scientific uncertainty. The degree of the reduction in F between F_{ABC} and F_{MSY} depends on whether F_{MSY} is directly estimated (tier 1 stock) or a proxy value is used (tier 2 stock). For tier 1 stocks, F_{ABC} equals F_{MSY} reduced by five percent. For tier 2 stocks, F_{ABC} equals F_{MSY} reduced by ten percent.

Tier-1: $F_{ABC} = F_{MSY} \times 0.95$.

Tier-2: $F_{ABC} = F_{MSY} \times 0.90$.

Annual Catch Limit

ACLs are also defined in terms of spawner escapement (S_{ACL}) based on N and the corresponding exploitation rate (F_{ACL}), where the exploitation rate is a fixed value that does not change on an annual basis.

F_{ACL} is equivalent to F_{ABC} and

$$S_{ACL} = N \times (1 - F_{ACL}),$$

which results in $S_{ACL} = S_{ABC}$ for each management year.

During the annual preseason salmon management process, S_{ACL} is estimated using the fixed F_{ACL} exploitation rate and the preseason forecast of N. Thus, fishery management measures must result in an expected spawning escapement greater than or equal to this estimate of S_{ACL} .

STATUS DETERMINATION CRITERIA

In 2011 the Council also adopted new status determination criteria (SDC) for overfishing, approaching an overfished condition, overfished, not overfished/rebuilding, and rebuilt under Salmon Fishery Management Plan (FMP) Amendment 16. These criteria, approved and implemented in December 2011, were:

- Overfishing occurs when a single year exploitation rate exceeds the maximum fishing mortality threshold (MFMT), which is based on the maximum sustainable yield exploitation rate (F_{MSY});
- Approaching an overfished condition occurs when the geometric mean of the two most recent postseason estimates of spawning escapement, and the current preseason forecast of spawning escapement, is less than the minimum stock size threshold (MSST);
- Overfished status occurs when the most recent 3-year geometric mean spawning escapement is less than the MSST;
- Not overfished/rebuilding status occurs when a stock has been classified as overfished and has not yet been rebuilt, and the most recent 3-year geometric mean spawning escapement is greater than the MSST but less than S_{MSY} ;
- A stock is rebuilt when the most recent 3-year geometric mean spawning escapement exceeds S_{MSY} .

Status determinations for overfishing, overfished, not overfished/rebuilding, and rebuilt were reported in the annual SAFE document, Review of 2011 Ocean Salmon Fisheries (PFMC 2012). Because approaching an overfished condition relies on a preseason forecast and proposed fishing regulations, that status determination is reported in Chapter V of this document. All SDC rely on the most recent estimates available, which in some cases may be a year or more in the past because of incomplete broods or data availability; however, some status determinations reported in the SAFE document may be updated if more recent spawning escapement or exploitation rate estimates become available between the time the SAFE document and this document are published.

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 1 of 4)

| Production Source and Stock or Stock Group | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | Methodology for 2012 Prediction and Source |
|--|-------|---------------------|-------|-------|--------------------|-------|-------|-------|---------|---|
| Sacramento Index | | | | | | | | | | |
| Fall | - | - | - | - | 54.6 ^{a/} | 122.2 | 245.5 | 729.9 | 819.4 | Linear regression analysis of jack escapement on SI of the following year using 2009-2011 data. STT |
| Klamath River (Ocean Abundance) | | | | | | | | | | |
| Fall | 216.3 | 239.8 | 110.0 | 546.2 | 190.7 | 505.7 | 331.5 | 371.1 | 1,651.8 | Linear regression analysis of age-specific ocean abundance estimates on river runs of same cohort. STT. |
| Oregon Coast | | | | | | | | | | |
| North and South/Local Migrating | | | | | | | | | | None. |
| Columbia River (Ocean Escapement) | | | | | | | | | | |
| Upriver Spring | 360.7 | 254.1 ^{b/} | 88.4 | 78.5 | 269.3 | 298.9 | 470.0 | 198.4 | 314.2 | Log-normal sibling regressions of cohort returns in previous run years. WDFW staff. |
| Willamette Spring | 109.4 | 116.9 | 46.5 | 52.0 | 34.0 | 37.6 | 62.7 | 104.1 | 83.4 | Age-specific linear regressions of cohort returns in previous run years. ODFW staff. |
| Sandy Spring | 5.2 | 7.4 | 8.2 | 7.9 | 6.8 | 5.2 | 3.7 | 5.5 | 4.8 | Recent year average. ODFW staff. |
| Cowlitz Spring | 15.9 | 12.7 | 3.0 | 6.4 | 5.2 | 4.1 | 12.5 | 6.6 | 8.7 | Age-specific linear regressions of cohort returns in previous run years. WDFW. |
| Kalama Spring | 6.0 | 4.5 | 1.5 | 4.0 | 3.7 | 0.9 | 0.9 | 0.6 | 0.7 | Age-specific linear regressions of cohort returns in previous run years. WDFW. |
| Lewis Spring | 5.4 | 7.6 | 1.8 | 5.9 | 3.5 | 2.2 | 6.0 | 3.4 | 2.7 | Age-specific linear regressions of cohort returns in previous run years. WDFW. |
| Upriver Summer | 102.8 | 62.4 ^{b/} | 49.0 | 45.6 | 52.0 | 70.7 | 88.8 | 91.9 | 91.2 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |
| URB Fall | 292.2 | 352.2 | 253.9 | 182.4 | 162.5 | 259.9 | 310.8 | 398.2 | 353.5 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |
| SCH Fall | 138.0 | 114.1 | 50.0 | 21.8 | 87.2 | 59.3 | 169.0 | 116.4 | 63.8 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |
| LRW Fall | 24.1 | 20.2 | 16.6 | 10.1 | 3.8 | 8.5 | 9.7 | 12.5 | 16.2 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |
| LRH Fall | 77.1 | 74.1 | 55.8 | 54.9 | 59.0 | 88.8 | 90.6 | 133.5 | 127.0 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |
| MCB Fall | 90.4 | 89.4 | 88.3 | 68.0 | 54.0 | 94.5 | 72.6 | 100.0 | 90.8 | Age-specific average cohort ratios/cohort regressions. Columbia River TAC subgroup and WDFW |

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 2 of 4)

| Production Source and Stock or Stock | | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | Methodology for 2012 Prediction and Source |
|---|----------|------|------|------|------|-------------------|-----------------|-------------------|-------------------|-------------------|--|
| Washington Coast (Ocean Escapement) | | | | | | | | | | | |
| Willapa Bay Fall | Natural | 4.1 | 3.2 | 2.0 | 2.0 | 2.5 | 2.0 | 2.0 | 2.0 | 5.2 | Based on average 1999-2007 returns/spawner applied to Brood Years 2005-2008. WDFW |
| | Hatchery | 14.7 | 17.4 | 29.8 | 29.8 | 27.0 | 34.8 | 31.1 | 31.1 | 40.5 | Based on average 1998-2007 returns/release applied to Brood Years 2005-2008, adjusted by model performance. WDFW |
| Quinalt Fall | Natural | 2.2 | 3.9 | 8.7 | 7.3 | 3.7 | 6.9 | 7.6 | 5.9 | 7.7 | Return per spawner by age with a 5 year adjusted average adjusted with brood year sibling return. |
| | Hatchery | 2.9 | 6.2 | 7.3 | 8.7 | 1.3 | 7.8 | 5.5 | 4.7 | 3.8 | Recent 5 year average return per spawner |
| Queets Spring/Sum | Natural | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | Recent 5 year average |
| Queets Fall | Natural | 4.4 | 4.3 | 3.5 | 2.6 | 3.5 | 4.5 | 4.1 | 2.7 | 5.8 | Return per spawner by age with a 5 year adjusted average adjusted with brood year sibling return. |
| Hoh Spring/Summer | Hatchery | 0.7 | 1.2 | 1.4 | 1.5 | 7.0 | 1.2 | 9.8 | 1.9 | 1.8 | Recent 5 year average return per spawner |
| | Natural | 1.5 | 1.5 | 1.4 | 1.6 | 0.9 | 1.1 | 0.8 | 1.0 | 1.0 | Forecast from returns per spawner using recent 5 year mean. |
| Hoh Fall | Natural | 4.2 | 3.8 | 4.0 | 2.7 | 2.9 | 2.6 | 3.3 | 2.9 | 2.7 | Forecast from returns per spawner using recent 5 year mean. |
| Quillayute Spring | Hatchery | 1.4 | 1.2 | 1.7 | 1.3 | 1.7 | 2.0 | 1.5 | 1.4 | 1.5 | Mean return per release using most recent 4 years, 5 year adjusted means for age-5 and age-6. |
| Quillayute Sum/Fall | Natural | 7.8 | 6.7 | 6.8 | 7.7 | 6.0 | 6.8 | 7.5 | 8.8 | 7.4 | Summer: Recent 5 year mean return per spawner. Fall: Returns per spawner mean recent 5 years. |
| Hoko | Natural | - | - | - | - | 1.1 ^{e/} | 1 ^{e/} | 1.8 ^{e/} | 0.6 ^{e/} | 1.9 ^{e/} | Sibling regressions. |
| North Coast Totals | | | | | | | | | | | |
| Spring/Summer | Natural | 1.9 | 2.0 | 1.9 | 2.0 | 1.3 | 1.5 | 1.2 | 1.4 | NA | |
| Fall | Natural | 18.6 | 18.7 | 23.0 | 20.3 | 16.1 | 20.8 | 22.5 | 20.3 | NA | |
| Spring/Summer | Hatchery | 1.4 | 1.2 | 1.7 | 1.3 | 1.7 | 2.0 | 1.5 | 1.4 | 1.5 | |
| Fall | Hatchery | 3.6 | 7.4 | 8.7 | 10.2 | 8.3 | 9.0 | 15.3 | 6.6 | NA | |

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 3 of 4)

| Production Source and Stock or Stock | | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | Methodology for 2012 Prediction and Source |
|---|----------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--|
| Stillaguamish | Natural | 3.3 ^{e/} | 2.0 ^{e/} | 1.6 ^{e/} | 1.9 ^{e/} | 1.1 ^{e/} | 1.7 ^{e/} | 1.4 ^{e/} | 1.8 ^{e/} | 0.9 ^{e/} | Natural plus supplemental production from average of FRAM CWT reconstruction and an independent environmental model to link to return rates of specific age classes. FRAM CWT reconstruction uses BY 1993-2003 tagged fish survival rates for supplemental forecast, and BY 1986-1993 recruits/spawner for the natural return. |
| Snohomish | Natural | 15.7 ^{e/} | 14.2 ^{e/} | 8.7 ^{e/} | 12.3 ^{e/} | 6.5 ^{e/} | 8.4 ^{e/} | 9.9 ^{e/} | 7.4 ^{e/} | 2.8 ^{e/} | Recent year average brood recruits/spawner applied to the 2006-2010 parent escapements. Hatchery forecasts based on average CWT survival rates (yearlings: BY 1996-97; fingerlings: BY 2000-2003) from Wallace Hatchery applied to releases. |
| | Hatchery | 10.1 ^{e/} | 9.9 ^{e/} | 9.6 ^{e/} | 8.7 ^{e/} | 8.8 ^{e/} | 4.9 ^{e/} | 5.6 ^{e/} | 5.2 ^{e/} | 3.9 ^{e/} | Yearlings based on CWT groups for Wallace Hatchery (BYs 1987 and 1992-1996). Fingerlings based on survival estimate from Tulalip Hatchery 1998-2003. |
| Tulalip | Hatchery | 7.6 ^{e/} | 9.2 ^{e/} | 10.0 ^{e/} | 8.1 ^{e/} | 4.1 ^{e/} | 4.0 ^{e/} | 3.4 ^{e/} | 3.5 ^{e/} | 5.9 ^{e/} | CWT survival rates (1998-2003) multiplied by release numbers for brood years 2006-2009. |
| South Puget Sound | Natural | 17.5 | 17.7 | 21.3 | 17.0 | 21.1 | 17.2 | 12.7 | 8.9 | 8.9 | Puyallup R. recent five year average return per spawner applied to brood years contributing ages 3-6. For Nisqually, recent 5 year average (2004-2010 return years) of runsizes. Green R. spawning escapement in terms of natural origin adults. |
| | Hatchery | 86.5 | 83.1 | 85.8 | 92.1 | 101.3 | 93.0 | 97.4 | 118.6 | 95.8 | Average return at age multiplied by cohort release for Green, Carr Inlet, and Area 10E. Nisqually based on return rates/realease for age-3 -5. |
| Hood Canal | Natural | 2.4 ^{d/} | 3.1 ^{d/} | 2.5 ^{d/} | 3.8 ^{d/} | 2.6 ^{d/} | 2.5 ^{d/} | 2.4 ^{d/} | 2.2 ^{d/} | 2.9 ^{d/} | Natural fish based on the Hood Canal terminal run reconstruction-based relative contribution of the individual Hood Canal management units in the 2008-2011 return years. |
| | Hatchery | 27.2 ^{d/} | 27.5 ^{d/} | 27.7 ^{d/} | 43.6 ^{d/} | 34.2 ^{d/} | 40.1 ^{d/} | 42.6 ^{d/} | 38.4 ^{d/} | 43.9 ^{d/} | Brood 2008 fingerling lbs released from WDFW facilities in 2009, multiplied by the average of postseason estimated terminal area return rates (total terminal run / hatchery fingerling lbs released three years previous) for the last four return years (2008-2011). |
| Strait of Juan de Fuca Including Dungeness spring run | Natural | 3.6 ^{d/} | 4.2 ^{d/} | 4.2 ^{d/} | 4.4 ^{d/} | 3.2 ^{d/} | 2.4 ^{d/} | 1.9 ^{d/} | 2.5 ^{d/} | 2.9 ^{d/} | Dungeness and Elwha hatchery estimated by four-year average releases times average return rates. Dungeness wild estimated by smolts times average hatchery return rate. Elwha estimate separates hatchery and wild fish based on otolith sampling. |
| | Hatchery | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | Hatchery production included in naturals. |

TABLE I-1. Preseason adult Chinook salmon stock forecasts in thousands of fish. (Page 4 of 4)

a/ Does not include the river harvest component. SI forecasts after 2008 include river harvest.

b/ Beginning in 2005, the upriver spring/summer designation was changed, with stream type Snake Basin summer fish being combined with the spring stock.

c/ Unless otherwise noted, forecasts are for Puget Sound run size (4B) available to U.S. net fisheries. Does not include fish caught in troll and recreational fisheries.

d/ Terminal run forecast.

e/ Expected spawning escapement without fishing.

TABLE I-2. Preseason adult coho salmon stock forecasts in thousands of fish. (Page 1 of 2)

| Production Source and Stock or Stock Group | | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | Methodology for 2012 Prediction and Source |
|---|----------------------------|-------|-------|-------|-------|-------|---------|-------|-------|-------|--|
| OPI Area (Total Abundance) (California and Oregon Coasts and Columbia River) | | 777.9 | 542.9 | 460.2 | 849.2 | 276.1 | 1,284.7 | 556.0 | 624.5 | 632.7 | Abundance of all OPI components based on cohort reconstruction including all fishery impacts using Mixed Stock Model (MSM); prior to 2008 only fishery impacts south of Leadbetter Point were used (traditional OPI accounting). OPITT, see Chapter III for details. |
| OPI Public | Hatchery | 623.9 | 389.9 | 398.8 | 593.6 | 216.1 | 1,073.1 | 408.0 | 375.1 | 341.7 | OPIH: 1969-2010 Columbia River jacks adjusted for delayed smolt releases and total OPI jacks regressed on 1970-2011 adults. Columbia/Coastal proportions based on jacks; Columbia early/late proportions based on jacks; Coastal N/S proportions based on smolts. |
| Columbia River Early | | 313.6 | 284.6 | 245.8 | 424.9 | 110.3 | 672.7 | 245.3 | 216.0 | 229.8 | |
| Columbia River Late | | 274.7 | 78.0 | 113.8 | 139.5 | 86.4 | 369.7 | 144.2 | 146.5 | 87.4 | |
| Coastal N. of Cape Blanco | | 16.6 | 11.5 | 8.6 | 7.0 | 1.7 | 7.3 | 4.4 | 3.6 | 6.4 | |
| Coastal S. of Cape Blanco | | 19.0 | 15.8 | 30.6 | 22.2 | 17.7 | 23.4 | 14.1 | 9.0 | 18.1 | |
| Lower Columbia River | Natural | NA | NA | NA | 21.5 | 13.4 | 32.7 | 15.1 | 22.7 | 30.1 | Oregon: recent three year average; Washington: natural smolt production multiplied by 2009 brood marine survival rate. Abundance is subset of early/late hatchery abundance above. |
| Oregon Coast (OCN) | Natural | 150.9 | 152.0 | 60.8 | 255.4 | 60.0 | 211.6 | 148.0 | 249.4 | 291.0 | Rivers: Generalized additive model (GAM) relating ocean recruits to parental spawners and marine environmental variables. See text in Chapter III for details. Lakes: recent three year average return. |
| STEP ^{a/} | Hatchery | 3.1 | 1.0 | 0.6 | 0.2 | - | - | - | - | - | No forecast since 2007; releases discontinued. |
| Washington Coast | | | | | | | | | | | A variety of methods were used for 2012, primarily based on smolt production and survival. See text in Chapter III for details. |
| Willapa | Natural | 36.7 | 35.9 | 30.3 | 24.4 | 35.1 | 33.5 | 20.4 | 47.8 | 81.3 | |
| | Hatchery | 55.0 | 56.4 | 37.7 | 37.2 | 25.5 | 59.4 | 78.7 | 64.7 | 88.8 | |
| Grays Harbor | Natural | 117.9 | 91.1 | 67.3 | 59.4 | 42.7 | 59.2 | 67.9 | 89.1 | 150.2 | |
| | Hatchery | 67.8 | 54.4 | 52.4 | 74.0 | 53.1 | 63.5 | 33.3 | 44.0 | 47.8 | |
| Quinalt | Natural | 50.5 | 44.9 | 28.8 | 18.6 | 17.4 | 16.3 | 16.7 | 22.9 | 27.3 | |
| | Hatchery | 18.2 | 33.6 | 34.5 | 22.7 | 24.5 | 26.2 | 26.6 | 35.5 | 35.4 | |
| Queets | Natural | 18.5 | 17.1 | 8.3 | 13.6 | 10.2 | 31.4 | 21.8 | 13.3 | 37.2 | |
| | Hatchery | 17.1 | 17.4 | 11.9 | 19.1 | 10.3 | 13.5 | 11.9 | 16.3 | 25.3 | |
| | Supplemental ^{b/} | 2.5 | 2.4 | - | - | - | - | - | - | - | |
| Hoh | Natural | 8.1 | 7.6 | 6.4 | 5.4 | 4.3 | 9.5 | 7.6 | 11.6 | 14.3 | |

TABLE I-2. Preseason adult coho salmon stock forecasts in thousands of fish. (Page 2 of 2)

TABLE 12. 1

a/ Program ended in 2005.

b/ Strait of Juan de Fuca and Hood Canal Hatchery numbers in 2002-2005 include natural coho from secondary (hatchery) management zones.

CHAPTER II: AFFECTED ENVIRONMENT - CHINOOK SALMON ASSESSMENT

CHINOOK STOCKS SOUTH OF CAPE FALCON

Sacramento River Fall Chinook

The Council's Salmon FMP sets the escapement goal for SRFC as a range from 122,000 to 180,000 hatchery and natural area adults. This stock comprises a large proportion of the Chinook spawners returning to Central Valley streams and hatcheries. SRFC are designated as the indicator stock for the Central Valley Fall Chinook stock complex, which was established under FMP Amendment 16 to facilitate setting and assessing compliance with ABC and ACLs, as required by the 2006 revision of the MSA. SRFC are currently in an overfished status, and will be managed in accordance with a rebuilding plan to be adopted during the 2012 preseason process.

Predictor Description and Performance

The Sacramento Index (SI) is the sum of (1) SRFC ocean fishery harvest south of Cape Falcon between September 1 and August 31, (2) SRFC impacts from non-retention ocean fisheries when they occur, (3) the recreational harvest of SRFC in the Sacramento River Basin, and (4) the SRFC adult spawner escapement (Table II-1, Figure II-1).

In 2011, the STT based the forecast of the SI on a zero-intercept linear model relating the previous year ($t-1$) SRFC jack escapement to the SI in year t , for $t = 1990-2010$. In 2011, the SI preseason forecast of 729,893 was 3.7 times its postseason value of 199,308.

The SI forecast has exceeded its postseason estimate for three consecutive years (2009-2011). Each of these years has been characterized by the most recent jack escapement estimate (year $t-1$) exceeding the jack escapement estimate from the previous year (year $t-2$) by a large margin. This is the case again for the 2012 SI forecast, where the 2011 jack escapement estimate is the largest on record.

For a variety of potential reasons, including the increasing trend in jack escapement, the relationship between jack escapement and the SI for years 2009-2011 exhibits a markedly different pattern than what existed for years prior to 2009 (Figure II-2). As a result, the 2012 SI forecast is based on data from 2009-2011. For reference, the SI forecast based on data from 1990-2011 is presented as well.

Stock Forecast and Status

A total of 85,719 SRFC jacks were estimated to have escaped to Sacramento River basin hatcheries and natural spawning areas in 2011. The resulting 2012 SI forecast is 819,400 (Figure II-2). For comparison, the SI forecast that would result from using data from 1990-2011 is 2,199,565.

In 2012, invoking *de minimis* fishing rates under Amendment 16 will be unnecessary because SRFC potential spawner abundance is projected to be greater than 162,667 hatchery and natural area adults; therefore, the S_{MSY} conservation objective of 122,000 should be exceeded with an AEQ exploitation rate greater than 0.25.

OFL, ABC, and ACL

The OFL, ABC, and ACL OFL are defined in terms of spawner escapement (S_{OFL} , S_{ABC} , and S_{ACL}). For SRFC $F_{MSY} = 0.78$, the proxy value for Tier-2 Chinook stocks that do not have estimates of this rate derived from a stock-specific spawner-recruit analysis. The OFL for SRFC is $S_{OFL} = 819,400 \times (1-0.78) = 180,268$. Because SRFC is a Tier-2 stock, $F_{ABC} = F_{MSY} \times 0.90 = 0.70$, and $F_{ACL} = F_{ABC}$. The 2012 preseason ABC for SRFC is: $S_{ABC} = 819,400 \times (1-0.70) = 245,820$, with $S_{ACL} = S_{ABC}$. Therefore, fisheries

impacting SRFC must be crafted to achieve, in expectation, a minimum of 245,820 hatchery and natural-area adult spawners in 2012. These preseason estimates will be recalculated with postseason abundance estimates (when available) to assess ACL and OFL compliance.

Klamath River Fall Chinook

Predictor Description

For Klamath River fall Chinook, linear regressions are used to relate September 1 ocean abundance estimates of age-3, age-4, and age-5 fish to that year's river run size estimates of age-2, age-3, and age-4 fish, respectively (Table II-2). Historical abundance estimates were derived from a cohort analysis of CWT information (brood years 1979-2007). The y-intercept of the regressions is constrained to zero, which gives the biologically reasonable expectation that a river run size of zero predicts an ocean abundance remainder of zero for the same cohort. The abundance of age-2 fish is not forecasted because no precursor to age-2 fish of that brood is available. Ocean fisheries harvest small numbers of age-2 KRFC.

Predictor Performance

Since 1985, the preseason ocean abundance forecasts for age-3 fish have ranged from 0.33 to 2.72 times the postseason estimates; for age-4 fish from 0.47 to 2.60 times the postseason estimates; and for the adult stock as a whole from 0.34 to 2.03 times the postseason estimates (Table II-3). The September 1, 2010 age-3 forecast (304,600) was 1.31 times its postseason estimate (232,749). The age-4 forecast (61,600) was 0.94 times its postseason estimate (65,714); and the age-5 forecast (5,000) was 1.8 times its postseason estimate (2,772) (Table II-3). The preseason forecast of the adult stock as a whole was 1.23 times the postseason estimate.

Management of KRFC harvest since 1986 has attempted to achieve specific harvest rates on fully-vulnerable age-4 and age-5 fish in ocean and river fisheries (Table II-4). The Council has used a combination of quotas and time/area restrictions in ocean fisheries in an attempt to meet the harvest rate objective set each year. Since 1992, fisheries have been managed to achieve 50/50 allocation between tribal and non-tribal fisheries. Tribal and recreational river fisheries have been managed on the basis of adult Chinook quotas.

The Council's FMP conservation objective for KRFC (Amendment 16) permits an average natural spawner reduction rate via fisheries of no more than 0.68, with a minimum escapement of 40,700 natural spawning adults. The FMP allows for any ocean and river harvest allocation that meets the spawner reduction rate constraint, provided it also meets the minimum escapement goal. The regulations adopted in 2011 were expected to result in 35,000 natural area spawning adults and an age-4 ocean harvest rate of 16.0 percent. Postseason estimates of these quantities were 47,754 natural area adult spawners and an age-4 ocean harvest rate of 7.8 percent (Table II-5).

Stock Forecast and Status

The 2012 forecast for the ocean abundance of KRFC as of September 1, 2011 (preseason) is 1,567,600 age-3 fish, the age-4 forecast is 79,600, and the age-5 forecast is 4,600 fish.

Late-season ocean fisheries in 2011 (September through November) were estimated to have harvested 143 adult KRFC, including 70 age-4 (0.1 percent ocean harvest rate), which will be deducted from the ocean fishery's allocation in determining the 2012 allowable ocean harvest.

In 2012, invoking *de minimis* fishing rates under Amendment 16 will be unnecessary because KRFC potential spawner abundance is projected to be substantially greater than 54,267 natural area adults.

Therefore, the S_{MSY} conservation objective of 40,700 should be achieved with an exploitation rate greater than 0.25.

OFL, ABC, and ACL

The OFL, ABC, and ACL OFL are defined in terms of spawner escapement (S_{OFL} , S_{ABC} , and S_{ACL}). For KRFC $F_{MSY} = 0.71$, the value estimated from a stock-specific spawner-recruit analysis (STT 2005). The OFL for KRFC is $S_{OFL} = 269,649 \times (1-0.71) = 78,198$. Because KRFC is a Tier-1 stock, $F_{ABC} = F_{MSY} \times 0.95 = 0.68$, and $F_{ACL} = F_{ABC}$. For KRFC, the preseason forecast of potential natural area adult spawners is 269,649, which results in $S_{ABC} = 269,649 \times (1-0.68) = 86,288$, with $S_{ACL} = S_{ABC}$. Therefore, fisheries impacting KRFC must be crafted to achieve, in expectation, a minimum of 86,288 natural-area adult spawners in 2012. These preseason estimates will be recalculated with postseason abundance estimates (when available) to assess ACL and OFL compliance.

Other California Coastal Chinook Stocks

Other California coastal streams that support fall Chinook stocks which contribute to ocean fisheries off Oregon and California, include the Smith, Little, Mad, Eel, and Mattole rivers, and Redwood Creek. Except for the Smith River, these stocks are included in the California coastal Chinook ESU, which is listed as threatened under the ESA. Current information is insufficient to forecast the ocean abundance of these stocks, however, the NMFS ESA consultation standard restricts the KRFC age-4 ocean harvest rate to no more than 16.0 percent to limit impacts on these stocks. In 2011 the age-4 ocean harvest rate was 7.8 percent. The Klamath River spring, Smith River, Rogue River, Umpqua River, and other Oregon Chinook stocks south of the Elk River are components of the Southern Oregon/Northern California (SONC) Chinook complex, and as such, specification of ACLs is deferred to KRFC, the indicator stock for the SONC complex.

Oregon Coast Chinook Stocks

Oregon coast Chinook stocks are categorized into three major subgroups based on ocean migration patterns; the North Oregon Coast (NOC) Chinook aggregate, the Mid Oregon Coast (MOC) Chinook aggregate, and the South Oregon Coast (SOC) Chinook aggregate. Although their ocean harvest distributions overlap somewhat, they have been labeled as far-north, north, or south/local migrating, respectively.

Far-North and North Migrating Chinook (NOC and MOC groups)

Far-north and north migrating Chinook stocks include spring and fall stocks north of and including the Elk River, with the exception of Umpqua River spring Chinook. Based on CWT analysis, the populations from ten major NOC river systems from the Nehalem through the Siuslaw Rivers are harvested primarily in ocean fisheries off British Columbia and Southeast Alaska, and to a much lesser degree in Council area and terminal area (state waters) fisheries off Washington and Oregon. CWT analysis indicates populations from five major MOC systems, from the Coos through the Elk Rivers, are harvested primarily in ocean fisheries off British Columbia, Canada, Washington, Oregon, and in terminal area fisheries. Minor catches occur in California fisheries, and variable catches have been observed in southeast Alaska troll fisheries.

NOC and MOC Chinook stocks are components of the Far-North-Migrating Coastal (FNMC) Chinook complex, which is an exception to the ACL requirements of the MSA because they are managed under an international agreement (the PST); therefore, specification of ACLs is not necessary for stocks in the FNMC complex.

Predictor Description

Quantitative abundance predictions are made for all three of the coastal Chinook groups (NOC, MOC, and SOC), but are not used in annual development of Council area fishery regulations. Quantitative forecasts of abundance are based on sibling regression analyses from individual basins' escapement assessment data and scale sampling, which occurs coast-wide. Forecast data for the NOC are used in the PSC management process in addition to terminal area management actions.

Natural spawner escapement is assessed yearly from the Nehalem through Sixes rivers. Peak spawning counts of adults are obtained from standard index areas on these rivers and monitored to assess stock trends (PFMC 2012, Chapter II, Table II-4 and Figure II-3). Natural fall Chinook stocks from both the NOC and MOC dominate production from this subgroup. Also present in lesser numbers are naturally-produced spring Chinook stocks from several rivers, and hatchery fall and/or spring Chinook released in the Trask, Nestucca, Salmon, Alsea, and Elk rivers.

Basin-specific forecasts constitute the overall aggregate forecasts and are derived in conjunction with annual PSC Chinook model input and calibration activities; however, they were not available at publication time.

Predictor Performance

There was no information available to evaluate performance of predictors for NOC and MOC stocks.

Stock Forecast and Status

North Oregon Coast

Since 1977, the Salmon River Hatchery production has been tagged for use primarily as a PSC indicator stock for the NOC stock component. Because these fish are primarily harvested in fisheries north of the Council management area, the STT has not reviewed the procedure by which this indicator stock is used in estimating annual stock status. The annual spawner counts have been gradually increasing since 2007. The 2011 spawner counts were a 5 percent increase from 2010 (PFMC 2012, Appendix B, Table B-11).

Based on the density index of total spawners, the generalized expectation for NOC stocks in 2012 is above recent years' average abundance. Specifically, the 2011 spawner density in standard survey areas for the NOC averaged 91 spawners per mile.

Mid Oregon Coast

Since 1977, the Elk River Hatchery production has been tagged for potential use as a PSC indicator stock for the MOC stock aggregate. Age-specific ocean abundance forecasts for 2012 are not currently available, but are being developed. The STT has not undertaken a review of the methods used by Oregon Department of Fish and Wildlife (ODFW) staff in developing these abundance forecasts.

The 2011 MOC density from standard survey areas was 106 adult spawners per mile, the highest since 2003 (PFMC 2012, Appendix B, Table B-11). Fall Chinook escapement goals are currently under development for the South Umpqua and Coquille basins of the MOC.

South/Local Migrating Chinook (SOC group)

South/local migrating Chinook stocks include Rogue River spring and fall Chinook, fall Chinook from smaller rivers south of the Elk River, and Umpqua River spring Chinook. These stocks are important contributors to ocean fisheries off Oregon and northern California. Umpqua River spring Chinook contribute to a lesser degree to fisheries off Washington, British Columbia, and southeast Alaska.

SOC stocks are components of the Southern Oregon/Northern California (SONC) Chinook complex, and as such, specification of ACLs is deferred to KRFC, the indicator stock for the SONC complex.

Rogue River Fall Chinook

Rogue River fall Chinook contribute to ocean fisheries principally as age-3 through age-5 fish. Mature fish enter the river each year from mid-July through October, with the peak of the run occurring during August and September.

Predictor Description

Carcass recoveries in Rogue River index surveys covering a large proportion of the total spawning area were available for 1977-2004. Using Klamath Ocean Harvest Model (KOHM) methodology, these carcass numbers, allocated into age-classes from scale data, were used to estimate the Rogue Ocean Population Index (ROPI) for age-3 to age-5 fish. A linear regression was developed using the escapement estimates (all ages) in year t based on seining at Huntley Park (1976-2004) to predict the ROPI in year $t+1$ (1977-2005). The 2011 Huntley Park escapement estimate and the resulting 2012 ROPI forecast was then scaled to the historical carcass survey-based ROPI. The 2012 ROPI forecast (45,000) consisting of age-3 (25,400), age-4 (16,800) and age-5 (2,700) are based on the average annual age-class strengths of the carcass-based ROPIs from 1991-2004. This data set was truncated at 1991 because significant harvest restrictions that could affect age structure began that year.

Predictor Performance

The ROPI is based on cohort reconstruction methods with index values predicted from regression equations. Because postseason estimates of the ROPI are not available, it is not possible to assess predictor performance.

Stock Forecast and Status

The 2012 ROPI is three times higher than the recent three-year average of 14,900, and the highest on record since 1988 (Table II-6).

Other SOC Stocks

Umpqua and Rogue spring Chinook contribute to ocean fisheries primarily as age-3 fish. Mature Chinook enter the rivers primarily during April and May and generally prior to annual ocean fisheries. Quantitative abundance predictions are not made for these stocks.

Natural fall Chinook stocks from river systems south of the Elk River and spring Chinook stocks from the Rogue and Umpqua rivers dominate production from this subgroup. Substantial releases of hatchery spring Chinook occur in both the Rogue and Umpqua rivers, although also present in lesser numbers are hatchery fall Chinook, primarily from the Chetco River.

Fall Chinook escapement goals and forecasts are currently under development for stocks south of the Elk River. These stocks are minor contributors to general season mixed stock ocean fisheries. Standard fall Chinook spawning index escapement data were available for the smaller SOC rivers (Winchuck, Chetco, and Pistol rivers). The 2011 average density from standard survey areas was 35 adult spawners per mile (PFMC 2012, Appendix B, Table B-8).

Quantitative abundance predictions are not made for these stocks, although general trends in stock abundance for SOC Chinook stocks are assessed through escapement indices (PFMC 2012, Chapter II, Table II-4 and Figure II-3).

CHINOOK STOCKS NORTH OF CAPE FALCON

Columbia River Chinook

Columbia River fall Chinook stocks typically form the largest contributing stock group to Council Chinook fisheries north of Cape Falcon. Abundance of these stocks is a major factor in determining impacts of fisheries on weak natural stocks critical to Council area management, particularly ESA-listed Lower Columbia River (LCR) natural tule Chinook. Abundance predictions are made for five major fall stock units characterized as being hatchery or natural production, and originating above or below Bonneville Dam. The upriver brights (URB) and lower river wild (LRW) are primarily naturally-produced stocks, although the upriver brights do have a significant hatchery component. The lower river hatchery (LRH) tule, Spring Creek Hatchery (SCH) tule, and mid-Columbia brights (MCB) are primarily hatchery-produced stocks. The MCB include the lower river bright (LRB) stock as a small naturally-produced component. LRB spawn in the mainstem Columbia River near Beacon Rock and are believed to have originated from MCB hatchery strays. The tule stocks generally mature at an earlier age than the bright fall stocks and do not migrate as far north. Minor fall stocks include the Select Area brights (SAB), a stock originally from the Rogue River.

Columbia Upper River summer Chinook also contribute to Council area fisheries, although like URB and LRW, most ocean impacts occur in B.C. and SEAK fisheries. Columbia River summer Chinook have both natural and hatchery components, and originate in areas upstream from Rock Island Dam.

URB and Columbia summer Chinook are exceptions to the ACL requirements of the MSA because they are managed under an international agreement (the PST); therefore, specification of ACLs is not necessary for URB and Columbia summer Chinook. ESA consultation standards serve the purpose of ACLs for ESA-listed stocks like LRW Chinook, and are deferred to ESA consultation standards. Broodstock goals serve the purpose of ACLs for hatchery origin stocks like LRH, SCH and MCB.

Predictor Description

Preseason forecasts of Columbia River fall and summer Chinook stock abundance, used by the STT to assess the Council's adopted fishery regulations, are based on age-specific and stock-specific forecasts of annual ocean escapement (return to the Columbia River). These forecasts are developed by WDFW and a subgroup of the *U.S. v Oregon* Technical Advisory Committee (TAC). Columbia River return forecast methodologies used for Council management are identical to those used for planning Columbia River fall season fisheries, although minor updates to Council estimates of inriver run size may occur prior to finalization of the inriver fishery plans, based on results of planned ocean fisheries.

The 2012 return of summer and each fall Chinook stock group is forecasted using relationships between successive age groups within a cohort. The database for these relationships was constructed by combining age-specific estimates of escapement and inriver fishery catches for years since 1964 (except for MCB, which started in the 1980's). Typically, only the more recent broods are used in the current predictions. Fall Chinook stock identification in the Columbia River mixed stock fisheries is determined by sampling catch and escapement for CWTs and visual stock identification (VSI). Age composition estimates are based on CWT data and scale reading of fishery and escapement samples, where available. These stock and age data for Columbia River fall Chinook are the basis for the return data presented in the *Review of 2011 Ocean Salmon Fisheries* (Appendix B, Tables B-15 through B-20). The 2011 returns for summer Chinook and the five fall Chinook stocks listed in this report may differ somewhat from those provided in the *Review of 2011 Ocean Salmon Fisheries*, since ocean escapement estimates may have been updated after that report was printed.

Summer and fall Chinook ocean escapement forecasts developed for the March Council meeting do not take into account variations in marine harvest. The STT combines the initial inriver run size (ocean escapement; Table II-7) with expected Council area fishery harvest levels and stock distribution patterns to produce adjusted ocean escapement forecasts based on the proposed ocean fishing regulations. These revised forecasts are available at the end of the Council preseason planning process in April and are used for preseason fishery modeling in the Columbia River.

Predictor Performance

Performance of the preliminary inriver run size estimation methodology can be assessed, in part, by examining the differences between preseason forecasts and postseason estimates (Table II-7;

Figure II-4). The recent 10-year average March preliminary preseason forecasts as a percentage of the postseason estimates for the URB, LRW, LRH, SCH, and MCB are 1.08, 1.06, 1.02, 1.19, and 1.00 respectively. None of the fall Chinook stocks had a notable bias in the recent time series of March preliminary forecasts. The recent 4-year average March preliminary preseason forecasts as a percentage of the postseason estimates for SUM is 1.15 with a bias toward over-forecasting.

Stock Forecasts and Status

The preliminary forecast for 2012 URB fall Chinook ocean escapement is 383,500 adults, about 109 percent of last year's return and about 131 percent of the recent 10-year average of 268,860. This escapement is well above the FMP S_{MSY} conservation objective of 39,625 natural area spawners in the Hanford Reach, Yakima River, and areas above Priest Rapids Dam, and should allow opportunity for both ocean and in-river fisheries.

The preliminary forecast for 2012 ocean escapement of ESA-listed Snake River wild fall Chinook is 15,100, about 101 percent of last year's preliminary return estimate of 14,911.

Ocean escapement of LRW fall Chinook in 2012 is forecast at 16,200 adults, about 107 percent of last year's forecast, and about 106 percent of the recent 10-year average return of 15,310. The forecast is greater than last year's actual return, and the spawning escapement goal of 5,700 in the North Fork Lewis River should be achieved this year.

The preliminary forecast for 2012 ocean escapement of LRH fall Chinook is for a return of 127,000 adults, about 116 percent of last year's return and 135 percent of the recent 10-year average of 93,890. Based on this abundance forecast, the total allowable LCR natural tule exploitation rate for 2012 fisheries is no greater than 41.0 percent under the matrix developed by the Tule Chinook Workgroup in 2011, which the Council recommended NMFS use in developing ESA guidance for 2012 fisheries (Appendix A Table A-5). This is the highest exploitation rate allowed under the recommended matrix.

The preliminary ocean escapement forecast of SCH fall Chinook in 2012 is 63,800 adults, about 92 percent of last year's return and 64 percent of the 10-year average of 99,360.

The preliminary forecast for the 2012 ocean escapement of MCB fall Chinook is 90,800 adults, about 107 percent of last year's return and about 99 percent of the recent 10-year average of 91,400.

The preliminary forecast for summer Chinook in 2012 is 91,200 adults, about 113 percent of last year's return and about 139 percent of the recent 4-year average of 68,583. This escapement is well above the FMP S_{MSY} conservation objective of 12,143 escapement above Rock Island Dam, and should allow opportunity for both ocean and in-river fisheries.

Washington Coast Chinook

Washington Coast Chinook consist of spring, summer, and fall stocks from Willapa Bay through the Hoko River. Based on limited CWT analysis, these populations are harvested primarily in ocean fisheries off British Columbia and Southeast Alaska, and to a lesser degree in Council-area fisheries off Washington and Oregon.

Washington Coast Chinook stocks are components of the FNMC Chinook complex, which is an exception to the ACL requirements of the MSA because it is managed under an international agreement (the PST); therefore, specification of ACLs is not necessary for stocks in the FNMC complex.

Predictor Description and Past Performance

Council fisheries have negligible impacts on Washington coast Chinook stocks, and except for Willapa Bay fall Chinook, Queets River fall Chinook, Hoh River Chinook, and Quillayute River Chinook, forecast data is unavailable in time for publication of this report; therefore, preseason abundance estimates are not presented. However, abundance estimates are provided for Washington Coastal fall stocks in subsequent preseason fishery impact assessment reports prepared by the STT (e.g., Preseason Report III).

Stock Forecasts and Status

The 2012 Willapa Bay hatchery fall Chinook ocean escapement forecast is 40,518, which is higher than the 2011 prediction of 32,476. The 2012 natural fall Chinook ocean escapement forecast is 5,222, which is higher than last year's prediction of 4,341, and is above the FMP S_{MSY} conservation objective of 3,393.

The 2012 Queets River natural fall Chinook forecast is for an ocean escapement of 5,800, which is higher than the 2011 forecast of 2,700. The ocean escapement is greater than the 2,500 FMP S_{MSY} conservation objective, which should allow flexibility in structuring 2012 ocean and river fisheries. The 2012 Queets River hatchery fall Chinook forecast is for an ocean escapement of 1,835, which is slightly less than the 2011 forecast of 1,900.

For the Hoh River, the 2012 natural spring/summer Chinook ocean escapement forecast is 997, above the FMP conservation objective of 900. The natural fall Chinook forecast is 2,683, above the FMP S_{MSY} conservation objective of 1,200.

The 2012 Quillayute hatchery spring Chinook ocean escapement forecast is 1,453 and the natural summer/fall Chinook forecast is 7,359 (767 summer and 6,592 fall). The FMP S_{MSY} conservation objectives are spawning escapements of 1,200 summer Chinook and 3,000 fall Chinook.

Puget Sound Chinook

Puget Sound Chinook stocks include all fall, summer, and spring stocks originating from U.S. tributaries in Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek, inclusive). Puget Sound Chinook consists of numerous natural Chinook stocks of small to medium-sized populations and significant hatchery production. The Puget Sound ESU was listed under the ESA as threatened in March 1999.

Southern U.S. fisheries that impact Puget Sound Chinook are constrained by terms of a Resource Management Plan (RMP), and are exempted from ESA Section 9 take prohibitions under Limit 6 of the 4(d) rule. Puget Sound stocks contribute to fisheries off B.C., are present to a lesser degree off SEAK, and are impacted to a minor degree by Council-area ocean fisheries. Because Council-area fishery impacts to Puget Sound Chinook stocks are negligible, ocean regulations are not generally used to manage these stocks.

Predictor Description

Methodologies for estimates are described in the annual Puget Sound management reports (starting in 1993, reports are available by Puget Sound management unit, not by individual species). Forecasts for Puget Sound stocks generally assume production is dominated by age-4 adults. The STT has not undertaken a review of the methods employed by state and tribal staffs in preparing these abundance forecasts. Run-size expectations for various Puget Sound stock management units are listed in Table I-1.

Predictor Performance

There was no information available to evaluate performance of predictors for Puget Sound Chinook stocks.

Stock Forecasts and Status

ACLs are undefined in the FMP for ESA-listed stocks like Puget Sound Chinook, and are deferred to ESA consultation standards.

Spring Chinook

Spring Chinook originating in Puget Sound are expected to remain depressed. Runs in the Nooksack, Skagit, White, and Dungeness rivers are of particular concern.

Summer/Fall Chinook

The 2012 preliminary forecast for Puget Sound summer/fall stocks is for a return of 229,989 Chinook, slightly lower than the 2011 preseason forecast of 244,377. The 2012 natural Chinook return forecast of 25,643 (includes supplemental category forecasts) is lower than the 2011 forecast of 39,333.

Since ESA listing and development of the RMP, fishery management for Puget Sound Chinook has changed from an escapement goal basis to the use of stock-specific exploitation rates and “critical abundance thresholds.” This new approach is evaluated on an annual basis through the RMP.

STOCK STATUS DETERMINATION UPDATES

There were several updates and additions to the spawning escapement estimates for Chinook stocks in the SAFE document. Previously unavailable 2011 natural spawning escapements are now available for Columbia River URB, Willapa Bay fall, Grays Harbor spring, Queets spring/summer and fall, and Hoh spring/summer and fall Chinook. Updates to 2010 and previously unavailable 2011 exploitation rate estimates were available for SRFC and KRFC.

Preliminary 2011 Feather River natural area escapement estimates have been changed to 10,443 jacks and 32,531 adults. Because of this change, the total SRFC escapement values used for 2011 stock status determination and abundance forecast modeling was 85,719 jacks and 114,741 adults. Using these updated escapement estimates, the most recent three-year (2009-2011) geometric mean of SRFC hatchery and natural adult escapement is 83,530. This value is below the MSST of 91,500, which results in an overfished status for this stock (Table V-4).

The SAFE document reported a 2010 exploitation rate for SRFC of 0.18. Updated information resulted in a 2010 exploitation rate of 0.17, well below the MFMT of 0.78. The preliminary 2011 SRFC exploitation rate estimate was 0.42, also less than the MFMT. Hence, SRFC were not subject to overfishing in 2010 or 2011 (Table V-4).

The SAFE document reported a 2010 exploitation rate (spawner reduction rate) for KRFC of 0.42, well below the MFMT of 0.71. Additional information resulted in a preliminary exploitation rate estimate for

2011 of 0.38, also less than the MFMT. Therefore KRFC were not subject to overfishing in 2010 or 2011 (Table V-4).

The most recent 3-year (2008-2010) geometric mean spawning escapement estimated for Queets spring/summer Chinook reported in the SAFE document was 339, less than the MSST of 350; however, a preliminary 2011 spawning estimate of 373 results in 3-year geometric mean of 363. Therefore, Queets spring/summer Chinook should not be considered overfished (Table V-4).

Other than Queets spring/summer Chinook, the updated 2010 and 2011 estimates did not change the status (e.g., overfished, rebuilt, etc.) for any of these stocks.

SELECTIVE FISHERY CONSIDERATIONS FOR CHINOOK

As the North of Falcon region has moved forward with mass marking of hatchery Chinook salmon stocks, the first mark selective fishery for Chinook salmon in Council waters was implemented in June, 2010 in the recreational fishery north of Cape Falcon. In 2011, the mark selective Chinook quota season of 4,800 occurred from June 18-25 (8 days). Selective fishing options for non-Indian fisheries are likely to be under consideration again in the ocean area from Cape Falcon, Oregon to the U.S./Canada border. Observed mark rates on Chinook in 2011 ocean fisheries in this area ranged from 57 to 70 percent. Based on preseason abundance forecasts, the expected mark rate for Chinook in this area for 2012 should be similar to those observed in 2011.

TABLE II-1. Harvest and abundance indices for Sacramento River fall Chinook (SRFC) in thousands of fish.

| Year | SRFC Ocean Harvest | | | | River Harvest | Spawning Escapement | | | Sacramento Index (SI) ^{c/} | Exploitation Rate (%) ^{d/} |
|--------------------|------------------------------------|-------|-----------------------|---------|--------------------|---------------------|----------|-------|-------------------------------------|-------------------------------------|
| | South of Cape Falcon ^{a/} | | | | | Natural | Hatchery | Total | | |
| | Troll | Sport | Non-Ret ^{b/} | Total | | | | | | |
| 1984 | 266.8 | 87.1 | 0.0 | 353.9 | 26.1 | 119.5 | 39.5 | 159.0 | 539.0 | 71 |
| 1985 | 359.0 | 159.3 | 0.0 | 518.4 | 39.3 | 209.5 | 29.9 | 239.3 | 796.9 | 70 |
| 1986 | 620.1 | 137.5 | 0.0 | 757.6 | 39.4 | 216.3 | 23.8 | 240.1 | 1,037.1 | 77 |
| 1987 | 686.6 | 173.8 | 0.0 | 860.4 | 32.0 | 174.8 | 20.3 | 195.1 | 1,087.5 | 82 |
| 1988 | 1,163.0 | 188.3 | 0.0 | 1,351.3 | 37.3 | 198.0 | 29.5 | 227.5 | 1,616.1 | 86 |
| 1989 | 605.9 | 158.9 | 0.0 | 764.8 | 25.0 | 126.7 | 25.9 | 152.6 | 942.4 | 84 |
| 1990 | 507.5 | 150.8 | 0.0 | 658.3 | 17.2 | 83.2 | 21.9 | 105.1 | 780.6 | 87 |
| 1991 | 301.0 | 90.7 | 0.0 | 391.7 | 26.0 ^{e/} | 91.4 | 27.5 | 118.9 | 536.6 | 78 |
| 1992 | 233.3 | 70.2 | 0.0 | 303.5 | 13.3 ^{e/} | 59.5 | 22.1 | 81.5 | 398.3 | 80 |
| 1993 | 342.8 | 115.5 | 0.0 | 458.3 | 27.7 ^{e/} | 110.6 | 26.8 | 137.4 | 623.4 | 78 |
| 1994 | 303.3 | 164.8 | 0.0 | 468.1 | 28.9 ^{e/} | 133.0 | 32.6 | 165.6 | 662.5 | 75 |
| 1995 | 730.4 | 387.9 | 0.0 | 1,118.3 | 48.5 | 253.5 | 41.8 | 295.3 | 1,462.1 | 80 |
| 1996 | 426.8 | 157.0 | 0.0 | 583.8 | 49.5 | 267.1 | 34.6 | 301.6 | 934.9 | 68 |
| 1997 | 579.7 | 210.3 | 0.0 | 790.0 | 56.6 | 279.6 | 65.2 | 344.8 | 1,191.5 | 71 |
| 1998 | 292.8 | 113.9 | 0.0 | 406.7 | 69.8 ^{e/} | 168.1 | 77.8 | 245.9 | 722.5 | 66 |
| 1999 | 308.1 | 76.7 | 0.0 | 384.8 | 68.9 ^{e/} | 353.7 | 46.1 | 399.8 | 853.5 | 53 |
| 2000 | 432.7 | 153.2 | 0.0 | 585.8 | 59.5 ^{e/} | 369.2 | 48.3 | 417.5 | 1,062.8 | 61 |
| 2001 | 285.2 | 94.3 | 0.0 | 379.5 | 97.9 | 537.4 | 59.4 | 596.8 | 1,074.2 | 44 |
| 2002 | 454.2 | 185.2 | 0.0 | 639.4 | 89.2 ^{e/} | 682.7 | 87.2 | 769.9 | 1,498.5 | 49 |
| 2003 | 506.5 | 106.9 | 0.0 | 613.4 | 85.8 | 413.4 | 109.6 | 523.0 | 1,222.2 | 57 |
| 2004 | 622.0 | 213.0 | 0.0 | 835.0 | 47.1 | 203.5 | 83.4 | 286.9 | 1,169.0 | 75 |
| 2005 | 370.3 | 127.7 | 0.0 | 498.0 | 65.0 | 210.7 | 185.3 | 396.0 | 959.0 | 59 |
| 2006 | 149.9 | 107.8 | 0.0 | 257.7 | 45.1 | 195.1 | 79.9 | 275.0 | 577.8 | 52 |
| 2007 | 120.0 | 32.2 | 0.0 | 152.2 | 14.3 ^{e/} | 70.0 | 21.4 | 91.4 | 257.9 | 65 |
| 2008 | 3.2 | 0.9 | 0.0 | 4.1 | 0.1 ^{e/} | 46.9 | 18.5 | 65.4 | 69.6 | 6 |
| 2009 | 0.0 | 0.2 | 0.1 | 0.3 | 0.0 ^{e/} | 23.3 | 17.5 | 40.9 | 41.1 | 1 |
| 2010 | 11.8 | 11.4 | 0.3 | 23.6 | 2.5 ^{e/} | 84.6 | 39.7 | 124.3 | 150.3 | 17 |
| 2011 ^{f/} | 45.7 | 21.5 | 0.0 | 67.2 | 17.4 ^{e/} | 71.9 | 42.9 | 114.7 | 199.3 | 42 |

a/ Ocean harvest for the period September 1 (t-1) through August 31 (t).

b/ Mortalities estimated from non-retention ocean fisheries (e.g., coho-only fisheries, non-retention GSI sampling).

c/ The SI is the sum of (1) SRFC ocean fishery harvest south of Cape Falcon between September 1 and August 31, (2) SRFC impacts from non-retention ocean fisheries when they occur, (3) the recreational harvest of SRFC in the Sacramento River Basin, and (4) the SRFC adult spawner escapement.

d/ Total ocean harvest, non-retention ocean fishery mortalities, and river harvest of SRFC as a percentage of the SI.

e/ Estimates derived from CDFG Sacramento River Basin angler survey. Estimates not marked with a footnote are inferred from escapement data and the mean river harvest rate estimate.

f/ Preliminary.

TABLE II-2. Klamath River fall Chinook ocean abundance (thousands), harvest rate, and river run size estimates (thousands) by age.

| Year (t) | Ocean Abundance Sept. 1 (t-1) | | | Annual Ocean Harvest Rate Sept. 1 (t-1) - Aug. 31 (t) | | Klamath Basin River Run (t) | | | | |
|----------|-------------------------------|--------------------|---------|--|--------------------|-----------------------------|-------|-------|-------|--------------|
| | Age-3 | Age-4 | Total | Age-3 | Age-4 | Age-2 | Age-3 | Age-4 | Age-5 | Total Adults |
| 1981 | 493.2 | 57.0 | 550.2 | 0.21 | 0.53 | 28.2 | 64.1 | 14.4 | 1.8 | 80.3 |
| 1982 | 561.1 | 133.4 | 694.5 | 0.30 | 0.52 | 39.4 | 30.1 | 33.9 | 2.6 | 66.6 |
| 1983 | 313.3 | 114.2 | 427.5 | 0.19 | 0.60 | 3.8 | 35.9 | 20.7 | 0.9 | 57.5 |
| 1984 | 157.3 | 82.8 | 240.1 | 0.08 | 0.38 | 8.3 | 21.7 | 24.4 | 1.1 | 47.2 |
| 1985 | 374.8 | 56.9 | 431.7 | 0.11 | 0.24 | 69.4 | 32.9 | 25.7 | 5.8 | 64.4 |
| 1986 | 1,304.4 | 140.8 | 1,445.2 | 0.18 | 0.46 | 44.6 | 162.9 | 29.8 | 2.3 | 195.0 |
| 1987 | 781.2 | 341.9 | 1,123.1 | 0.16 | 0.43 | 19.1 | 89.7 | 112.6 | 6.8 | 209.1 |
| 1988 | 756.3 | 234.8 | 991.0 | 0.20 | 0.39 | 24.1 | 101.2 | 86.5 | 3.9 | 191.6 |
| 1989 | 369.8 | 177.2 | 547.1 | 0.15 | 0.36 | 9.1 | 50.4 | 69.6 | 4.3 | 124.3 |
| 1990 | 176.1 | 104.0 | 280.1 | 0.30 | 0.55 | 4.4 | 11.6 | 22.9 | 1.3 | 35.9 |
| 1991 | 69.4 | 37.2 | 106.6 | 0.03 | 0.18 | 1.8 | 10.0 | 21.6 | 1.1 | 32.7 |
| 1992 | 39.5 | 28.2 | 67.7 | 0.02 | 0.07 | 13.7 | 6.9 | 18.8 | 1.0 | 26.7 |
| 1993 | 168.5 | 15.0 | 183.5 | 0.05 | 0.16 | 7.6 | 48.3 | 8.2 | 0.7 | 57.2 |
| 1994 | 119.9 | 41.7 | 161.6 | 0.03 | 0.09 | 14.4 | 37.0 | 26.0 | 1.0 | 64.0 |
| 1995 | 784.3 | 28.7 | 813.0 | 0.04 | 0.14 | 22.8 | 201.9 | 18.3 | 2.6 | 222.8 |
| 1996 | 192.3 | 225.5 | 417.8 | 0.05 | 0.16 | 9.5 | 38.8 | 136.7 | 0.3 | 175.8 |
| 1997 | 140.2 | 62.8 | 203.0 | 0.01 | 0.06 | 8.0 | 35.0 | 44.2 | 4.6 | 83.7 |
| 1998 | 154.8 | 44.7 | 199.5 | 0.00 | 0.09 | 4.6 | 59.2 | 29.7 | 1.7 | 90.6 |
| 1999 | 129.1 | 30.5 | 159.5 | 0.02 | 0.09 | 19.2 | 29.2 | 20.5 | 1.3 | 51.0 |
| 2000 | 617.1 | 44.2 | 661.3 | 0.06 | 0.10 | 10.2 | 187.1 | 30.5 | 0.5 | 218.1 |
| 2001 | 356.1 | 133.8 | 489.9 | 0.03 | 0.09 | 11.3 | 99.1 | 88.2 | 0.2 | 187.4 |
| 2002 | 513.6 | 98.9 | 612.5 | 0.02 | 0.15 | 9.2 | 94.6 | 62.5 | 3.7 | 160.8 |
| 2003 | 400.2 | 192.2 | 592.4 | 0.08 | 0.21 | 3.8 | 94.3 | 96.8 | 0.9 | 191.9 |
| 2004 | 159.6 | 105.1 | 264.6 | 0.12 | 0.34 | 9.7 | 33.2 | 40.7 | 5.3 | 79.2 |
| 2005 | 190.0 | 38.1 | 228.1 | 0.02 | 0.20 | 2.3 | 43.8 | 17.5 | 3.9 | 65.2 |
| 2006 | 90.6 | 63.4 | 154.0 | 0.01 | 0.10 | 26.9 | 18.5 | 41.6 | 1.3 | 61.4 |
| 2007 | 376.8 | 33.6 | 410.5 | 0.06 | 0.21 | 1.7 | 113.7 | 16.8 | 1.6 | 132.1 |
| 2008 | 68.0 | 81.4 | 149.4 | 0.00 | 0.10 | 25.2 | 18.6 | 50.2 | 1.7 | 70.6 |
| 2009 | 240.8 | 21.1 | 261.9 | 0.00 | 0.00 | 11.9 | 78.6 | 16.4 | 5.6 | 100.6 |
| 2010 | 194.7 ^{a/} | 62.1 | 256.8 | 0.01 ^{a/} | 0.04 | 16.6 | 46.1 | 44.3 | 0.4 | 90.9 |
| 2011 | 232.7 ^{b/} | 65.7 ^{a/} | 298.5 | NA ^{c/} | 0.08 ^{a/} | 85.9 | 59.7 | 41.3 | 2.0 | 103.0 |

a/ Preliminary: incomplete cohort data (age-5 unavailable).

b/ Preliminary: incomplete cohort data (age-4 and age-5 unavailable).

c/ Not estimated: incomplete cohort data (age-4 and age-5 unavailable).

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 1 of 4)

| | Preseason Forecast ^{a/} | Postseason Estimate | |
|--------------------|----------------------------------|---------------------|----------------|
| Year (t) | Sept. 1 (t-1) | Sept. 1 (t-1) | Pre/Postseason |
| | | Age-3 | |
| 1985 | 113,000 | 276,000 | 0.41 |
| 1986 | 426,000 ^{b/} | 1,304,409 | 0.33 |
| 1987 | 511,800 | 781,198 | 0.66 |
| 1988 | 370,800 | 756,261 | 0.49 |
| 1989 | 450,600 | 369,828 | 1.22 |
| 1990 | 479,000 | 176,133 | 2.72 |
| 1991 | 176,200 | 69,424 | 2.54 |
| 1992 | 50,000 | 39,502 | 1.27 |
| 1993 | 294,400 | 168,473 | 1.75 |
| 1994 | 138,000 | 119,913 | 1.15 |
| 1995 | 269,000 | 784,260 | 0.34 |
| 1996 | 479,800 | 192,272 | 2.50 |
| 1997 | 224,600 | 140,153 | 1.60 |
| 1998 | 176,000 | 154,799 | 1.14 |
| 1999 | 84,800 | 129,066 | 0.66 |
| 2000 | 349,600 | 617,098 | 0.57 |
| 2001 | 187,200 | 356,128 | 0.53 |
| 2002 | 209,000 | 513,561 | 0.41 |
| 2003 | 171,300 | 400,242 | 0.43 |
| 2004 | 72,100 | 159,560 | 0.45 |
| 2005 | 185,700 | 189,976 | 0.98 |
| 2006 | 44,100 | 90,606 | 0.49 |
| 2007 | 515,400 | 376,841 | 1.37 |
| 2008 | 31,600 | 68,003 | 0.46 |
| 2009 | 474,900 | 240,760 | 1.97 |
| 2010 | 223,400 | 194,655 | 1.15 |
| 2011 ^{c/} | 304,600 | 232,749 | 1.31 |
| 2012 | 1,567,600 | -- | -- |

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 2 of 4)

| Year (t) | Preseason Forecast ^{a/} | Postseason Estimate | Pre/Postseason |
|--------------------|----------------------------------|---------------------|----------------|
| | Sept. 1 (t-1) | Sept. 1 (t-1) | |
| | Age-4 | | |
| 1985 | 56,875 | 57,500 | 0.99 |
| 1986 | 66,250 | 140,823 | 0.47 |
| 1987 | 206,125 | 341,875 | 0.60 |
| 1988 | 186,375 | 234,772 | 0.79 |
| 1989 | 215,500 | 177,245 | 1.22 |
| 1990 | 50,125 | 103,951 | 0.48 |
| 1991 | 44,625 | 37,172 | 1.20 |
| 1992 | 44,750 | 28,169 | 1.59 |
| 1993 | 39,125 | 15,037 | 2.60 |
| 1994 | 86,125 | 41,736 | 2.06 |
| 1995 | 47,000 | 28,725 | 1.64 |
| 1996 | 268,500 | 225,521 | 1.19 |
| 1997 | 53,875 | 62,820 | 0.86 |
| 1998 | 46,000 | 44,733 | 1.03 |
| 1999 | 78,750 | 30,456 | 2.59 |
| 2000 | 38,875 | 44,176 | 0.88 |
| 2001 | 247,000 | 133,801 | 1.85 |
| 2002 | 143,800 | 98,928 | 1.45 |
| 2003 | 132,400 | 192,156 | 0.69 |
| 2004 | 134,500 | 105,051 | 1.28 |
| 2005 | 48,900 | 38,079 | 1.28 |
| 2006 | 63,700 | 63,383 | 1.00 |
| 2007 | 26,100 | 33,615 | 0.78 |
| 2008 | 157,200 | 81,366 | 1.93 |
| 2009 | 25,200 | 21,124 | 1.19 |
| 2010 | 106,300 | 62,119 | 1.71 |
| 2011 ^{c/} | 61,600 | 65,714 | 0.94 |
| 2012 | 79,600 | -- | -- |

TABLE II-3. Comparisons of preseason forecasts and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 3 of 4)

| Year (t) | Preseason Forecast ^{a/} | Postseason Estimate | Pre/Postseason |
|--------------------|----------------------------------|---------------------|----------------|
| | Sept. 1 (t-1) | Sept. 1 (t-1) | |
| | Age-5 | | |
| 1985 | NA | 11,113 | NA |
| 1986 | NA | 6,376 | NA |
| 1987 | 5,250 | 19,414 | 0.27 |
| 1988 | 13,250 | 14,632 | 0.91 |
| 1989 | 10,125 | 9,612 | 1.05 |
| 1990 | 7,625 | 7,767 | 0.98 |
| 1991 | 1,500 | 2,774 | 0.54 |
| 1992 | 1,250 | 1,444 | 0.87 |
| 1993 | 1,125 | 1,759 | 0.64 |
| 1994 | 500 | 1,468 | 0.34 |
| 1995 | 2,000 | 3,805 | 0.53 |
| 1996 | 1,125 | 787 | 1.43 |
| 1997 | 7,875 | 8,859 | 0.89 |
| 1998 | 3,250 | 2,382 | 1.36 |
| 1999 | 2,000 | 2,106 | 0.95 |
| 2000 | 1,375 | 1,051 | 1.31 |
| 2001 | 1,250 | 258 | 4.84 |
| 2002 | 9,700 | 6,933 | 1.40 |
| 2003 | 6,500 | 1,915 | 3.39 |
| 2004 | 9,700 | 17,170 | 0.56 |
| 2005 | 5,200 | 6,857 | 0.76 |
| 2006 | 2,200 | 5,236 | 0.42 |
| 2007 | 4,700 | 2,911 | 1.61 |
| 2008 | 1,900 | 2,900 | 0.66 |
| 2009 | 5,600 | 7,059 | 0.79 |
| 2010 | 1,800 | 518 | 3.47 |
| 2011 ^{c/} | 5,000 | 2,772 | 1.80 |
| 2012 | 4,600 | -- | -- |

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 4 of 4)

| Year (t) | Preseason Forecast ^{a/} | Postseason Estimate | Pre/Postseason |
|--------------------|----------------------------------|---------------------|----------------|
| | Sept. 1 (t-1) | Sept. 1 (t-1) | |
| | Total Adults | | |
| 1985 | 169,875 ^{d/} | 344,613 | 0.49 |
| 1986 | 492,250 ^{d/} | 1,451,608 | 0.34 |
| 1987 | 723,175 | 1,142,487 | 0.63 |
| 1988 | 570,425 | 1,005,665 | 0.57 |
| 1989 | 676,225 | 556,685 | 1.21 |
| 1990 | 536,750 | 287,851 | 1.86 |
| 1991 | 222,325 | 109,370 | 2.03 |
| 1992 | 96,000 | 69,115 | 1.39 |
| 1993 | 334,650 | 185,269 | 1.81 |
| 1994 | 224,625 | 163,117 | 1.38 |
| 1995 | 318,000 | 816,790 | 0.39 |
| 1996 | 749,425 | 418,580 | 1.79 |
| 1997 | 286,350 | 211,832 | 1.35 |
| 1998 | 225,250 | 201,914 | 1.12 |
| 1999 | 165,550 | 161,628 | 1.02 |
| 2000 | 389,850 | 662,325 | 0.59 |
| 2001 | 435,450 | 490,187 | 0.89 |
| 2002 | 362,500 | 619,422 | 0.59 |
| 2003 | 310,200 | 594,313 | 0.52 |
| 2004 | 216,300 | 281,781 | 0.77 |
| 2005 | 239,800 | 234,912 | 1.02 |
| 2006 | 110,000 | 159,225 | 0.69 |
| 2007 | 546,200 | 413,367 | 1.32 |
| 2008 | 190,700 | 152,269 | 1.25 |
| 2009 | 505,700 | 268,943 | 1.88 |
| 2010 | 331,500 | 257,292 | 1.29 |
| 2011 ^{c/} | 371,200 | 301,235 | 1.23 |
| 2012 | 1,651,800 | -- | -- |

a/ Original preseason forecasts for years 1985-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the assumed May 1 (t) number by the Sept. 1 (t-1) through May 1 (t) survival rate in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ A scalar of 0.75 was applied to the jack count to produce the forecast because, (1) most jacks returned to the Trinity River, and (2) the jack count was outside the database range.

c/ Postseason estimates are preliminary.

d/ Does not include age-5 adults.

TABLE II-4. Summary of management objectives and predictor performance for Klamath River fall Chinook.

| Year(t) | Preseason Ocean Abundance Forecast ^{a/} Sept. 1 (t-1) | | Postseason Ocean Abundance Estimate Sept. 1 (t-1) | | Preseason Age-4 Harvest Rate Forecast ^{b/} | | Postseason Age-4 Harvest Rate Estimate ^{c/} | | Preseason Adult Harvest Forecast | | Postseason Adult Harvest Estimate | |
|--------------------|---|---------|--|---------|---|-------|--|-------|----------------------------------|---------|-----------------------------------|--------|
| | Age-3 | Age-4 | Age-3 | Age-4 | Ocean | River | Ocean | River | Ocean | River | Ocean | River |
| 1987 | 511,800 | 206,125 | 781,198 | 341,875 | 0.28 | 0.53 | 0.43 | 0.44 | 121,200 | 78,200 | 277,224 | 73,265 |
| 1988 | 370,800 | 186,375 | 756,261 | 234,772 | 0.31 | 0.53 | 0.39 | 0.52 | 114,100 | 65,400 | 253,905 | 73,854 |
| 1989 | 450,600 | 215,500 | 369,828 | 177,245 | 0.30 | 0.49 | 0.36 | 0.70 | 128,100 | 67,600 | 125,117 | 54,340 |
| 1990 | 479,000 | 50,125 | 176,133 | 103,951 | 0.30 | 0.49 | 0.55 | 0.36 | 85,100 | 31,200 | 114,786 | 11,459 |
| 1991 | 176,200 | 44,625 | 69,424 | 37,172 | 0.13 | 0.28 | 0.18 | 0.45 | 16,700 | 12,800 | 9,872 | 13,581 |
| 1992 | 50,000 | 44,750 | 39,502 | 28,169 | 0.06 | 0.15 | 0.07 | 0.27 | 4,200 | 4,200 | 3,142 | 6,787 |
| 1993 | 294,400 | 39,125 | 168,473 | 15,037 | 0.12 | 0.43 | 0.16 | 0.49 | 20,100 | 22,500 | 11,355 | 12,808 |
| 1994 | 138,000 | 86,125 | 119,913 | 41,736 | 0.07 | 0.20 | 0.09 | 0.29 | 10,400 | 14,300 | 7,961 | 13,524 |
| 1995 | 269,000 | 47,000 | 784,260 | 28,725 | 0.07 | 0.32 | 0.14 | 0.19 | 13,500 | 18,500 | 32,233 | 21,637 |
| 1996 | 479,800 | 268,500 | 192,272 | 225,521 | 0.17 | 0.66 | 0.16 | 0.39 | 88,400 | 129,100 | 45,155 | 69,241 |
| 1997 | 224,600 | 53,875 | 140,153 | 62,820 | 0.10 | 0.43 | 0.06 | 0.26 | 17,600 | 26,500 | 8,656 | 17,764 |
| 1998 | 176,000 | 46,000 | 154,799 | 44,733 | 0.07 | 0.29 | 0.09 | 0.30 | 10,200 | 14,800 | 4,891 | 17,897 |
| 1999 | 84,800 | 78,750 | 129,066 | 30,456 | 0.10 | 0.28 | 0.09 | 0.45 | 12,300 | 18,100 | 5,116 | 16,942 |
| 2000 | 349,600 | 38,875 | 617,098 | 44,176 | 0.11 | 0.53 | 0.10 | 0.25 | 24,000 | 32,400 | 42,050 | 35,066 |
| 2001 | 187,200 | 247,000 | 356,128 | 133,801 | 0.14 | 0.61 | 0.09 | 0.29 | 45,600 | 105,300 | 21,747 | 50,780 |
| 2002 | 209,000 | 143,800 | 513,561 | 98,928 | 0.13 | 0.57 | 0.15 | 0.26 | 30,000 | 70,900 | 28,895 | 35,069 |
| 2003 | 171,300 | 132,400 | 400,242 | 192,156 | 0.16 | 0.50 | 0.21 | 0.28 | 30,600 | 52,200 | 70,684 | 39,715 |
| 2004 | 72,100 | 134,500 | 159,560 | 105,051 | 0.15 | 0.38 | 0.34 | 0.48 | 26,500 | 35,800 | 63,885 | 29,807 |
| 2005 | 185,700 | 48,900 | 189,976 | 38,079 | 0.08 | 0.16 | 0.20 | 0.19 | 7,100 | 9,600 | 12,826 | 10,001 |
| 2006 | 44,100 | 63,700 | 90,606 | 63,383 | 0.11 | 0.23 | 0.10 | 0.18 | 10,000 | 10,000 | 10,401 | 10,345 |
| 2007 | 515,400 | 26,100 | 376,841 | 33,615 | 0.16 | 0.63 | 0.21 | 0.56 | 30,200 | 51,400 | 30,244 | 33,884 |
| 2008 | 31,600 | 157,200 | 68,003 | 81,366 | 0.02 | 0.43 | 0.10 | 0.38 | 4,500 | 49,500 | 8,679 | 24,180 |
| 2009 | 474,900 | 25,200 | 240,760 | 21,124 | 0.00 | 0.57 | 0.00 | 0.40 | 100 | 61,700 | 51 | 34,040 |
| 2010 | 223,400 | 106,300 | 194,655 | 62,119 | 0.12 | 0.49 | 0.04 | 0.40 | 22,600 | 46,600 | 4,467 | 32,920 |
| 2011 ^{d/} | 304,600 | 61,600 | 232,749 | 65,714 | 0.16 | 0.54 | 0.08 | 0.34 | 26,900 | 42,700 | 10,151 | 30,518 |
| 2012 | 1,567,600 | 79,600 | - | - | - | - | - | - | - | - | - | - |

a/ Original preseason forecasts for years 1986-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the May 1 (t) number by the assumed Sept. 1 (t-1) through May 1 (t) survival rate assumed in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ Ocean harvest rate forecast is the fraction of the predicted ocean abundance expected to be harvested Sept. 1 (t-1) through August 31(t). River harvest rate forecast is the fraction of the predicted river run expected to be harvested in river fisheries. Original ocean harvest rate forecasts for year (t), 1986-2001, were based on a May 1 (t) ocean abundance denominator; converted to Sept. 1 (t-1) abundance denominator by multiplying former values by 0.8 (the assumed age-4 survival rate between Sept. 1 (t-1) and May 1 (t) in those years).

c/ Ocean harvest rate is the fraction of the postseason ocean abundance harvested Sept. 1 (t-1) through August 31 (t). River harvest rate is the fraction of the river run harvested by river fisheries.

d/ Postseason estimates are preliminary.

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 1 of 4)

| Harvest Levels and Rates of Age-3 and Age-4 Atlantic River-Hair Shinnock (Page 1 of 4) | | | | | | | | | | |
|--|--|--------|----------|----------|----------|----------|-------------|---------------------|--------|--------|
| Year (t) | Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t)) | | | | | | | River Fisheries (t) | | |
| | KMZ | | | North of | South of | Subtotal | Ocean Total | Net | Sport | Total |
| | Troll | Sport | Subtotal | KMZ | KMZ | | | | | |
| HARVEST (numbers of fish) | | | | | | | | | | |
| Age-3 | | | | | | | | | | |
| 1986 | 35,632 | 4,876 | 40,508 | 73,777 | 122,913 | 196,690 | 237,198 | 8,100 | 18,100 | 26,200 |
| 1987 | 17,240 | 5,083 | 22,323 | 43,439 | 56,378 | 99,817 | 122,140 | 11,400 | 11,400 | 22,800 |
| 1988 | 15,999 | 5,165 | 21,164 | 24,317 | 107,971 | 132,288 | 153,452 | 12,500 | 15,600 | 28,100 |
| 1989 | 6,456 | 11,783 | 18,239 | 15,315 | 23,729 | 39,044 | 57,283 | 2,700 | 900 | 3,600 |
| 1990 | 81 | 4,357 | 4,438 | 36,579 | 11,006 | 47,585 | 52,023 | 1,300 | 1,400 | 2,700 |
| 1991 | 0 | 1,022 | 1,022 | 344 | 810 | 1,154 | 2,176 | 2,123 | 1,277 | 3,400 |
| 1992 | 0 | 0 | 0 | 972 | 0 | 972 | 972 | 970 | 251 | 1,221 |
| 1993 | 0 | 822 | 822 | 833 | 6,424 | 7,257 | 8,079 | 5,426 | 2,917 | 8,343 |
| 1994 | 42 | 604 | 646 | 0 | 3,387 | 3,387 | 4,033 | 4,543 | 965 | 5,508 |
| 1995 | 0 | 999 | 999 | 12,213 | 14,810 | 27,023 | 28,022 | 11,840 | 5,536 | 17,376 |
| 1996 | 0 | 0 | 0 | 0 | 9,314 | 9,314 | 9,314 | 12,363 | 3,661 | 16,024 |
| 1997 | 0 | 232 | 232 | 620 | 1,215 | 1,835 | 2,067 | 2,166 | 2,736 | 4,902 |
| 1998 | 0 | 6 | 6 | 298 | 466 | 764 | 770 | 2,231 | 5,781 | 8,012 |
| 1999 | 63 | 180 | 243 | 1,262 | 433 | 1,695 | 1,938 | 4,981 | 1,748 | 6,729 |
| 2000 | 404 | 3,282 | 3,686 | 8,604 | 25,203 | 33,807 | 37,493 | 22,458 | 4,893 | 27,351 |
| 2001 | 113 | 105 | 218 | 2,749 | 6,082 | 8,831 | 9,049 | 17,885 | 7,294 | 25,179 |
| 2002 | 220 | 784 | 1,004 | 1,501 | 9,915 | 11,416 | 12,420 | 11,734 | 6,258 | 17,992 |
| 2003 | 173 | 679 | 852 | 1,885 | 27,309 | 29,194 | 30,046 | 6,996 | 5,061 | 12,057 |
| 2004 | 402 | 971 | 1,373 | 9,719 | 7,331 | 17,050 | 18,423 | 4,679 | 2,051 | 6,730 |
| 2005 | 0 | 568 | 568 | 619 | 2,381 | 3,000 | 3,568 | 4,394 | 1,641 | 6,035 |
| 2006 | 0 | 477 | 477 | 32 | 341 | 373 | 850 | 2,388 | 13 | 2,401 |
| 2007 | 770 | 8,099 | 8,869 | 4,193 | 9,365 | 13,558 | 22,427 | 17,543 | 5,734 | 23,277 |
| 2008 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3,225 | 608 | 3,833 |
| 2009 | 0 | 51 | 51 | 0 | 0 | 0 | 51 | 19,820 | 4,715 | 24,535 |
| 2010 ^{a/} | 104 | 28 | 132 | 0 | 1,638 | 1,638 | 1,770 | 13,132 | 1,884 | 15,016 |
| 2011 ^{a/} | 245 | 845 | 1,090 | 25 | 3,620 | 3,645 | 4,735 | 13,286 | 2,637 | 15,923 |

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 2 of 4)

| Year (t) | Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t)) | | | | | | | River Fisheries (t) | | |
|---------------------------|--|-------|----------|----------|----------|----------|-------------|---------------------|-------|--------|
| | KMZ | | | North of | South of | Subtotal | Ocean Total | Net | Sport | Total |
| | Troll | Sport | Subtotal | KMZ | KMZ | | | | | |
| HARVEST (numbers of fish) | | | | | | | | | | |
| Age-4 | | | | | | | | | | |
| 1986 | 7,745 | 1,113 | 8,858 | 23,486 | 31,913 | 55,399 | 64,257 | 17,000 | 2,900 | 19,900 |
| 1987 | 21,736 | 4,427 | 26,163 | 70,645 | 48,832 | 119,477 | 145,640 | 41,000 | 8,500 | 49,500 |
| 1988 | 11,870 | 3,596 | 15,466 | 26,381 | 50,296 | 76,677 | 92,143 | 38,600 | 6,200 | 44,800 |
| 1989 | 6,064 | 9,735 | 15,799 | 32,116 | 16,608 | 48,724 | 64,523 | 41,000 | 7,700 | 48,700 |
| 1990 | 3,997 | 2,919 | 6,916 | 39,627 | 10,624 | 50,251 | 57,167 | 6,000 | 2,200 | 8,200 |
| 1991 | 0 | 1,001 | 1,001 | 1,513 | 4,135 | 5,648 | 6,649 | 7,593 | 2,016 | 9,609 |
| 1992 | 171 | 55 | 226 | 1,783 | 12 | 1,795 | 2,021 | 4,360 | 723 | 5,083 |
| 1993 | 0 | 0 | 0 | 849 | 1,616 | 2,465 | 2,465 | 3,786 | 243 | 4,029 |
| 1994 | 0 | 1,124 | 1,124 | 1,168 | 1,499 | 2,667 | 3,791 | 6,666 | 818 | 7,484 |
| 1995 | 0 | 242 | 242 | 1,879 | 1,771 | 3,650 | 3,892 | 2,957 | 480 | 3,437 |
| 1996 | 773 | 3,464 | 4,237 | 10,337 | 20,741 | 31,078 | 35,315 | 43,959 | 9,080 | 53,039 |
| 1997 | 3 | 172 | 175 | 463 | 2,994 | 3,457 | 3,632 | 8,734 | 2,586 | 11,320 |
| 1998 | 0 | 105 | 105 | 3,942 | 0 | 3,942 | 4,047 | 7,164 | 1,822 | 8,986 |
| 1999 | 15 | 381 | 396 | 1,657 | 696 | 2,353 | 2,749 | 8,789 | 494 | 9,283 |
| 2000 | 117 | 895 | 1,012 | 2,327 | 1,076 | 3,403 | 4,415 | 6,733 | 756 | 7,489 |
| 2001 | 1,312 | 1,604 | 2,916 | 5,819 | 3,926 | 9,745 | 12,661 | 20,759 | 4,819 | 25,578 |
| 2002 | 1,938 | 827 | 2,765 | 2,811 | 9,416 | 12,227 | 14,992 | 11,929 | 4,063 | 15,992 |
| 2003 | 834 | 918 | 1,752 | 7,855 | 30,007 | 37,862 | 39,614 | 22,754 | 4,592 | 27,346 |
| 2004 | 1,421 | 1,215 | 2,636 | 11,504 | 21,949 | 33,453 | 36,089 | 17,623 | 1,751 | 19,374 |
| 2005 | 247 | 317 | 564 | 5,243 | 1,909 | 7,152 | 7,716 | 3,048 | 304 | 3,352 |
| 2006 | 196 | 725 | 921 | 4,192 | 985 | 5,177 | 6,098 | 7,569 | 42 | 7,611 |
| 2007 | 270 | 2,336 | 2,606 | 1,991 | 2,472 | 4,463 | 7,069 | 8,987 | 502 | 9,489 |
| 2008 | 6,376 | 1,105 | 7,481 | 546 | 113 | 659 | 8,140 | 17,891 | 1,260 | 19,151 |
| 2009 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5,831 | 706 | 6,537 |
| 2010 | 36 | 111 | 147 | 892 | 1,487 | 2,379 | 2,526 | 16,630 | 1,134 | 17,764 |
| 2011 ^{a/} | 397 | 166 | 563 | 992 | 3,592 | 4,584 | 5,147 | 12,587 | 1,475 | 14,062 |

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 3 of 4)

| TABLE 11.3: Harvest totals and rates of Age-3 and Age-4 Atlantic Kingfisher Chinook (Page 3 of 7) | | | | | | | | | | |
|---|--|-------|----------|----------|----------|-------------|---------------------|-------|-------|----------|
| Year (t) | Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t)) | | | | | | River Fisheries (t) | | | |
| | KMZ | | | North of | South of | Ocean Total | Net | Sport | Total | |
| | Troll | Sport | Subtotal | KMZ | KMZ | | | | | Subtotal |
| HARVEST RATE ^{b/} | | | | | | | | | | |
| Age-3 | | | | | | | | | | |
| 1986 | 0.03 | 0.00 | 0.03 | 0.06 | 0.09 | 0.15 | 0.18 | 0.05 | 0.11 | 0.16 |
| 1987 | 0.02 | 0.01 | 0.03 | 0.06 | 0.07 | 0.13 | 0.16 | 0.13 | 0.13 | 0.25 |
| 1988 | 0.02 | 0.01 | 0.03 | 0.03 | 0.14 | 0.17 | 0.20 | 0.12 | 0.15 | 0.28 |
| 1989 | 0.02 | 0.03 | 0.05 | 0.04 | 0.06 | 0.11 | 0.15 | 0.05 | 0.02 | 0.07 |
| 1990 | 0.00 | 0.02 | 0.03 | 0.21 | 0.06 | 0.27 | 0.30 | 0.11 | 0.12 | 0.23 |
| 1991 | 0.00 | 0.01 | 0.01 | 0.00 | 0.01 | 0.02 | 0.03 | 0.21 | 0.13 | 0.34 |
| 1992 | 0.00 | 0.00 | 0.00 | 0.02 | 0.00 | 0.02 | 0.02 | 0.14 | 0.04 | 0.18 |
| 1993 | 0.00 | 0.00 | 0.00 | 0.00 | 0.04 | 0.04 | 0.05 | 0.11 | 0.06 | 0.17 |
| 1994 | 0.00 | 0.01 | 0.01 | 0.00 | 0.03 | 0.03 | 0.03 | 0.12 | 0.03 | 0.15 |
| 1995 | 0.00 | 0.00 | 0.00 | 0.02 | 0.02 | 0.03 | 0.04 | 0.06 | 0.03 | 0.09 |
| 1996 | 0.00 | 0.00 | 0.00 | 0.00 | 0.05 | 0.05 | 0.05 | 0.32 | 0.09 | 0.41 |
| 1997 | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.01 | 0.06 | 0.08 | 0.14 |
| 1998 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.04 | 0.10 | 0.14 |
| 1999 | 0.00 | 0.00 | 0.00 | 0.01 | 0.00 | 0.01 | 0.02 | 0.17 | 0.06 | 0.23 |
| 2000 | 0.00 | 0.01 | 0.01 | 0.01 | 0.04 | 0.05 | 0.06 | 0.12 | 0.03 | 0.15 |
| 2001 | 0.00 | 0.00 | 0.00 | 0.01 | 0.02 | 0.02 | 0.03 | 0.18 | 0.07 | 0.25 |
| 2002 | 0.00 | 0.00 | 0.00 | 0.00 | 0.02 | 0.02 | 0.02 | 0.12 | 0.07 | 0.19 |
| 2003 | 0.00 | 0.00 | 0.00 | 0.00 | 0.07 | 0.07 | 0.08 | 0.07 | 0.05 | 0.13 |
| 2004 | 0.00 | 0.01 | 0.01 | 0.06 | 0.05 | 0.11 | 0.12 | 0.14 | 0.06 | 0.20 |
| 2005 | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 | 0.02 | 0.02 | 0.10 | 0.04 | 0.14 |
| 2006 | 0.00 | 0.01 | 0.01 | 0.00 | 0.00 | 0.00 | 0.01 | 0.13 | 0.00 | 0.13 |
| 2007 | 0.00 | 0.02 | 0.02 | 0.01 | 0.02 | 0.04 | 0.06 | 0.15 | 0.05 | 0.20 |
| 2008 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.17 | 0.03 | 0.21 |
| 2009 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.25 | 0.06 | 0.31 |
| 2010 ^{a/} | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.01 | 0.28 | 0.04 | 0.33 |
| 2011 ^{a/} | 0.00 | 0.00 | 0.00 | 0.00 | 0.02 | 0.02 | 0.02 | 0.22 | 0.04 | 0.27 |

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 4 of 4)

| Year (t) | Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t)) | | | | | | River Fisheries (t) | | | |
|----------------------------|--|-------|----------|----------|----------|-------------|---------------------|-------|-------|----------|
| | KMZ | | | North of | South of | Ocean Total | Net | Sport | Total | |
| | Troll | Sport | Subtotal | KMZ | KMZ | | | | | Subtotal |
| HARVEST RATE ^{b/} | | | | | | | | | | |
| Age-4 | | | | | | | | | | |
| 1986 | 0.05 | 0.01 | 0.06 | 0.17 | 0.23 | 0.39 | 0.46 | 0.57 | 0.10 | 0.67 |
| 1987 | 0.06 | 0.01 | 0.08 | 0.21 | 0.14 | 0.35 | 0.43 | 0.36 | 0.08 | 0.44 |
| 1988 | 0.05 | 0.02 | 0.07 | 0.11 | 0.21 | 0.33 | 0.39 | 0.45 | 0.07 | 0.52 |
| 1989 | 0.03 | 0.05 | 0.09 | 0.18 | 0.09 | 0.27 | 0.36 | 0.59 | 0.11 | 0.70 |
| 1990 | 0.04 | 0.03 | 0.07 | 0.38 | 0.10 | 0.48 | 0.55 | 0.26 | 0.10 | 0.36 |
| 1991 | 0.00 | 0.03 | 0.03 | 0.04 | 0.11 | 0.15 | 0.18 | 0.35 | 0.09 | 0.45 |
| 1992 | 0.01 | 0.00 | 0.01 | 0.06 | 0.00 | 0.06 | 0.07 | 0.23 | 0.04 | 0.27 |
| 1993 | 0.00 | 0.00 | 0.00 | 0.06 | 0.11 | 0.16 | 0.16 | 0.46 | 0.03 | 0.49 |
| 1994 | 0.00 | 0.03 | 0.03 | 0.03 | 0.04 | 0.06 | 0.09 | 0.26 | 0.03 | 0.29 |
| 1995 | 0.00 | 0.01 | 0.01 | 0.07 | 0.06 | 0.13 | 0.14 | 0.16 | 0.03 | 0.19 |
| 1996 | 0.00 | 0.02 | 0.02 | 0.05 | 0.09 | 0.14 | 0.16 | 0.32 | 0.07 | 0.39 |
| 1997 | 0.00 | 0.00 | 0.00 | 0.01 | 0.05 | 0.06 | 0.06 | 0.20 | 0.06 | 0.26 |
| 1998 | 0.00 | 0.00 | 0.00 | 0.09 | 0.00 | 0.09 | 0.09 | 0.24 | 0.06 | 0.30 |
| 1999 | 0.00 | 0.01 | 0.01 | 0.05 | 0.02 | 0.08 | 0.09 | 0.43 | 0.02 | 0.45 |
| 2000 | 0.00 | 0.02 | 0.02 | 0.05 | 0.02 | 0.08 | 0.10 | 0.22 | 0.02 | 0.25 |
| 2001 | 0.01 | 0.01 | 0.02 | 0.04 | 0.03 | 0.07 | 0.09 | 0.24 | 0.05 | 0.29 |
| 2002 | 0.02 | 0.01 | 0.03 | 0.03 | 0.10 | 0.12 | 0.15 | 0.19 | 0.06 | 0.26 |
| 2003 | 0.00 | 0.00 | 0.01 | 0.04 | 0.16 | 0.20 | 0.21 | 0.24 | 0.05 | 0.28 |
| 2004 | 0.01 | 0.01 | 0.03 | 0.11 | 0.21 | 0.32 | 0.34 | 0.43 | 0.04 | 0.48 |
| 2005 | 0.01 | 0.01 | 0.01 | 0.14 | 0.05 | 0.19 | 0.20 | 0.17 | 0.02 | 0.19 |
| 2006 | 0.00 | 0.01 | 0.01 | 0.07 | 0.02 | 0.08 | 0.10 | 0.18 | 0.00 | 0.18 |
| 2007 | 0.01 | 0.07 | 0.08 | 0.06 | 0.07 | 0.13 | 0.21 | 0.53 | 0.03 | 0.56 |
| 2008 | 0.08 | 0.01 | 0.09 | 0.01 | 0.00 | 0.01 | 0.10 | 0.36 | 0.03 | 0.38 |
| 2009 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.36 | 0.04 | 0.40 |
| 2010 | 0.00 | 0.00 | 0.00 | 0.01 | 0.02 | 0.04 | 0.04 | 0.37 | 0.03 | 0.40 |
| 2011 ^{a/} | 0.01 | 0.00 | 0.01 | 0.02 | 0.05 | 0.07 | 0.08 | 0.30 | 0.04 | 0.34 |

a/ Preliminary (incomplete cohort).

b/ Ocean harvest rates are the fraction of Sept. 1 (t-1) ocean abundance harvested in these fisheries. River harvest rates are the fraction of the river run (t) harvested in these fisheries.

TABLE II-6. Rogue River fall Chinook inriver run and ocean population indices.

| Return Year | Inriver Run Index in Thousands of Fish ^{a/} | | | | | Ocean Impact Rate by Age ^{b/} | | Rogue Ocean Population Index (ROPI) in Thousands of Fish ^{c/} | | | |
|--------------------|--|-------|-------|-------|---------------------|--|---------|--|--------------------|-------------------|--------------------|
| | Age-2 | Age-3 | Age-4 | Age-5 | Total ^{d/} | Age-3 | Age-4-5 | Age-3 | Age-4 | Age-5 | Total |
| 1982 | 0.7 | 1.3 | 1.3 | 0.1 | 3.4 | 0.30 | 0.52 | 9.8 | 2.9 | 0.3 | 13.0 |
| 1983 | 0.3 | 1.1 | 1.5 | 0.0 | 2.9 | 0.19 | 0.60 | 8.6 | 4.4 | 0.1 | 13.1 |
| 1984 | 0.4 | 1.2 | 1.8 | 0.1 | 3.5 | 0.08 | 0.38 | 9.9 | 4.7 | 0.2 | 14.8 |
| 1985 | 2.5 | 1.3 | 3.5 | 0.6 | 7.9 | 0.11 | 0.25 | 9.7 | 6.3 | 0.9 | 16.9 |
| 1986 | 3.1 | 12.5 | 2.3 | 0.5 | 18.4 | 0.18 | 0.46 | 71.3 | 5.9 | 1.0 | 78.2 |
| 1987 | 2.6 | 7.8 | 18.1 | 0.4 | 28.9 | 0.16 | 0.43 | 80.3 | 36.3 | 0.6 | 117.2 |
| 1988 | 1.4 | 4.8 | 25.2 | 1.5 | 32.9 | 0.20 | 0.39 | 17.3 | 47.9 | 2.5 | 67.7 |
| 1989 | 0.5 | 1.3 | 4.0 | 2.0 | 7.8 | 0.15 | 0.36 | 8.4 | 7.2 | 3.2 | 18.8 |
| 1990 | 0.0 | 0.3 | 1.4 | 0.2 | 1.9 | 0.30 | 0.55 | 6.0 | 4.7 | 0.5 | 11.2 |
| 1991 | 0.2 | 0.4 | 1.9 | 0.5 | 3.0 | 0.03 | 0.18 | 3.5 | 3.2 | 0.6 | 7.3 |
| 1992 | 0.5 | 0.3 | 1.5 | 0.5 | 2.8 | 0.02 | 0.07 | 4.4 | 2.4 | 0.6 | 7.4 |
| 1993 | 0.3 | 3.5 | 1.5 | 0.5 | 5.8 | 0.05 | 0.16 | 16.1 | 3.2 | 0.6 | 19.9 |
| 1994 | 0.5 | 0.8 | 5.8 | 0.9 | 8.0 | 0.03 | 0.09 | 3.0 | 9.5 | 0.9 | 13.4 |
| 1995 | 0.2 | 0.6 | 1.4 | 2.0 | 4.2 | 0.04 | 0.13 | 4.3 | 1.7 | 2.3 | 8.3 |
| 1996 | 0.1 | 0.4 | 1.8 | 0.1 | 2.4 | 0.05 | 0.16 | 2.4 | 2.8 | 0.1 | 5.3 |
| 1997 | 0.1 | 0.3 | 1.0 | 0.3 | 1.7 | 0.01 | 0.06 | 5.2 | 1.5 | 0.3 | 7.0 |
| 1998 | 0.0 | 0.5 | 2.8 | 0.3 | 3.6 | 0.00 | 0.09 | 3.8 | 3.9 | 0.3 | 8.0 |
| 1999 | 0.2 | 0.3 | 1.6 | 0.5 | 2.6 | 0.01 | 0.09 | 1.5 | 2.7 | 0.6 | 4.8 |
| 2000 | 0.2 | 2.0 | 0.8 | 0.6 | 3.6 | 0.06 | 0.10 | 9.9 | 0.9 | 0.6 | 11.4 |
| 2001 | 0.8 | 2.3 | 4.2 | 0.0 | 7.3 | 0.03 | 0.09 | 14.1 | 5.9 | 0.0 | 20.0 |
| 2002 | 0.9 | 4.0 | 7.1 | 0.8 | 12.7 | 0.02 | 0.15 | 32.2 | 9.1 | 0.9 | 42.2 |
| 2003 | 0.9 | 2.3 | 12.0 | 0.4 | 15.6 | 0.08 | 0.21 | 14.4 | 22.1 | 0.5 | 37.0 |
| 2004 | 0.4 | 0.6 | 4.9 | 2.9 | 8.8 | 0.12 | 0.34 | 3.9 | 9.7 | 4.4 | 18.0 |
| 2005 ^{f/} | NA | NA | NA | NA | NA | 0.02 | 0.20 | 7.6 | 5.0 | 0.8 | 13.4 |
| 2006 ^{f/} | NA | NA | NA | NA | NA | 0.01 | 0.10 | 4.9 | 3.2 | 0.5 | 8.6 |
| 2007 ^{f/} | NA | NA | NA | NA | NA | 0.06 | 0.21 | 5.8 | 3.8 | 0.6 | 10.2 |
| 2008 ^{f/} | NA | NA | NA | NA | NA | 0.00 | 0.10 | 6.9 | 4.6 | 0.7 | 12.2 |
| 2009 ^{f/} | NA | NA | NA | NA | NA | 0.00 | 0.00 | 6.1 | 4.0 | 0.7 | 10.7 |
| 2010 ^{f/} | NA | NA | NA | NA | NA | 0.01 | 0.04 | 9.8 ^{e/} | 6.5 | 1.1 | 17.3 ^{e/} |
| 2011 ^{f/} | NA | NA | NA | NA | NA | NA | 0.08 | 9.5 ^{g/} | 6.3 ^{g/} | 1.0 | 16.8 ^{g/} |
| 2012 ^{f/} | NA | NA | NA | NA | NA | - | - | 25.4 ^{g/} | 16.8 ^{g/} | 2.7 ^{g/} | 45.0 ^{g/} |

a/ Index based on carcass counts in spawning survey index areas. Carcass counts in 1978, 1979, and 1980 adjusted for prespawning mortality. Age composition developed from carcass scale sampling.

b/ Exploitation rates since 1981 are based on Klamath River fall Chinook cohort analysis, 1977-1980 based on 1981-1983 average.

c/ Based on cohort reconstruction methods. Index values for 2011 predicted from regression equations; postseason estimates are not available.

d/ Excludes age-6 fish.

e/ Preliminary, complete cohort not available, mean maturity rate used to derive estimate.

f/ Spawning surveys were discontinued 2005.

g/ Preseason forecast.

TABLE II-7. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 1 of 3)

| Year | March Preseason Forecast ^{a/} | April STT Modeled Forecast ^{b/} | Postseason Return | March Pre/Postseason | April Pre/Postseason |
|--------------------|---|---|-------------------|-------------------------|-------------------------|
| URB | | | | | |
| 1986 | 285.90 | 286.10 | 281.60 | 1.02 | 1.02 |
| 1987 | 436.40 | 436.40 | 420.70 | 1.04 | 1.04 |
| 1988 | 450.70 | 446.50 | 339.90 | 1.33 | 1.31 |
| 1989 | 234.00 | 231.80 | 261.30 | 0.90 | 0.89 |
| 1990 | 127.20 | 126.90 | 153.60 | 0.83 | 0.83 |
| 1991 | 88.80 | 88.90 | 103.30 | 0.86 | 0.86 |
| 1992 | 68.40 | 66.30 | 81.00 | 0.84 | 0.82 |
| 1993 | 84.50 | 82.70 | 102.90 | 0.82 | 0.80 |
| 1994 | 85.40 | 94.70 | 132.80 | 0.64 | 0.71 |
| 1995 | 103.70 | 125.00 | 106.50 | 0.97 | 1.17 |
| 1996 | 88.90 | 94.20 | 143.20 | 0.62 | 0.66 |
| 1997 | 166.40 | 158.00 | 161.70 | 1.03 | 0.98 |
| 1998 | 150.80 | 141.80 | 142.30 | 1.06 | 1.00 |
| 1999 | 147.50 | 102.10 | 166.10 | 0.89 | 0.61 |
| 2000 | 171.10 | 208.20 | 155.70 | 1.10 | 1.34 |
| 2001 | 127.20 | 132.70 | 232.60 | 0.55 | 0.57 |
| 2002 | 281.00 | 273.80 | 276.90 | 1.01 | 0.99 |
| 2003 | 280.40 | 253.20 | 373.20 | 0.75 | 0.68 |
| 2004 | 292.20 | 287.00 | 367.90 | 0.79 | 0.78 |
| 2005 | 352.20 | 354.60 | 268.70 | 1.31 | 1.32 |
| 2006 | 253.90 | 249.10 | 230.40 | 1.10 | 1.08 |
| 2007 | 182.40 | 185.20 | 112.60 | 1.62 | 1.64 |
| 2008 | 162.50 | 165.90 | 196.90 | 0.83 | 0.84 |
| 2009 | 259.90 | 269.80 | 212.00 | 1.23 | 1.27 |
| 2010 | 310.80 | 319.10 | 324.90 | 0.96 | 0.98 |
| 2011 ^{c/} | 398.20 | 399.50 | 324.10 | 1.23 | 1.23 |
| 2012 | 353.50 | - | - | - | - |
| LRW | | | | | |
| 1986 | 15.70 | NA | 24.50 | 0.64 | NA |
| 1987 | 29.20 | NA | 37.90 | 0.77 | NA |
| 1988 | 43.30 | 42.10 | 41.70 | 1.04 | 1.01 |
| 1989 | 27.30 | 26.90 | 38.60 | 0.71 | 0.70 |
| 1990 | 23.70 | 23.40 | 20.30 | 1.17 | 1.15 |
| 1991 | 12.70 | 12.70 | 19.80 | 0.64 | 0.64 |
| 1992 | 17.40 | 16.70 | 12.50 | 1.39 | 1.34 |
| 1993 | 12.50 | 11.90 | 13.30 | 0.94 | 0.89 |
| 1994 | 14.70 | 13.20 | 12.20 | 1.20 | 1.08 |
| 1995 | 12.40 | 11.50 | 16.00 | 0.78 | 0.72 |
| 1996 | 8.80 | 8.10 | 14.60 | 0.60 | 0.55 |
| 1997 | 7.50 | 7.20 | 12.30 | 0.61 | 0.59 |
| 1998 | 8.10 | 7.00 | 7.30 | 1.11 | 0.96 |
| 1999 | 2.60 | 2.50 | 3.30 | 0.79 | 0.76 |
| 2000 | 3.50 | 2.70 | 10.20 | 0.34 | 0.26 |
| 2001 | 16.70 | 18.50 | 15.70 | 1.06 | 1.18 |
| 2002 | 18.70 | 18.30 | 24.90 | 0.75 | 0.73 |
| 2003 | 24.60 | 23.40 | 26.00 | 0.95 | 0.90 |
| 2004 | 24.10 | 24.20 | 22.30 | 1.08 | 1.09 |
| 2005 | 20.20 | 21.40 | 16.80 | 1.20 | 1.27 |
| 2006 | 16.60 | 16.60 | 18.10 | 0.92 | 0.92 |
| 2007 | 10.10 | 10.00 | 4.30 | 2.35 | 2.33 |
| 2008 | 3.80 | 3.80 | 7.10 | 0.54 | 0.54 |
| 2009 | 8.50 | 8.60 | 7.50 | 1.13 | 1.15 |
| 2010 | 9.70 | 10.00 | 10.90 | 0.89 | 0.92 |
| 2011 ^{c/} | 12.50 | 13.10 | 15.20 | 0.82 | 0.86 |
| 2012 | 16.20 | - | - | - | - |

TABLE II-7. Predicted and postseason returns of Columbia River adult summer and fall Chinook in thousands of fish.
(Page 2 of 3)

| Year | March Preseason Forecast ^{a/} | April STT Modeled Forecast ^{b/} | Postseason Return | March Pre/Postseason | April Pre/Postseason |
|--------------------|---|---|-------------------|-------------------------|-------------------------|
| LRH | | | | | |
| 1986 | 171.60 | 173.90 | 154.80 | 1.11 | 1.12 |
| 1987 | 294.90 | 298.70 | 344.10 | 0.86 | 0.87 |
| 1988 | 267.70 | 246.50 | 309.90 | 0.86 | 0.80 |
| 1989 | 104.90 | 97.50 | 130.90 | 0.80 | 0.74 |
| 1990 | 68.50 | 65.50 | 60.00 | 1.14 | 1.09 |
| 1991 | 71.40 | 73.10 | 62.70 | 1.14 | 1.17 |
| 1992 | 113.20 | 121.50 | 62.60 | 1.81 | 1.94 |
| 1993 | 79.30 | 77.70 | 52.30 | 1.52 | 1.49 |
| 1994 | 36.10 | 46.50 | 53.60 | 0.67 | 0.87 |
| 1995 | 35.80 | 42.40 | 46.40 | 0.77 | 0.91 |
| 1996 | 37.70 | 48.30 | 75.50 | 0.50 | 0.64 |
| 1997 | 54.20 | 68.70 | 57.40 | 0.94 | 1.20 |
| 1998 | 19.20 | 22.50 | 45.30 | 0.42 | 0.50 |
| 1999 | 34.80 | 38.20 | 40.00 | 0.87 | 0.96 |
| 2000 | 23.70 | 26.40 | 27.00 | 0.88 | 0.98 |
| 2001 | 32.20 | 30.50 | 94.30 | 0.34 | 0.32 |
| 2002 | 137.60 | 133.00 | 156.40 | 0.88 | 0.85 |
| 2003 | 115.90 | 116.90 | 155.00 | 0.75 | 0.75 |
| 2004 | 77.10 | 79.00 | 108.90 | 0.71 | 0.73 |
| 2005 | 74.10 | 78.44 | 78.30 | 0.95 | 1.00 |
| 2006 | 55.80 | 57.50 | 58.30 | 0.96 | 0.99 |
| 2007 | 54.90 | 54.40 | 32.70 | 1.68 | 1.66 |
| 2008 | 59.00 | 55.90 | 60.30 | 0.98 | 0.93 |
| 2009 | 88.80 | 88.20 | 76.70 | 1.16 | 1.15 |
| 2010 | 90.60 | 85.60 | 103.00 | 0.88 | 0.83 |
| 2011 ^{c/} | 133.50 | 128.90 | 109.00 | 1.22 | 1.18 |
| 2012 | 127.00 | - | - | - | - |
| SCH | | | | | |
| 1986 | 16.00 | 16.20 | 16.60 | 0.96 | 0.98 |
| 1987 | 9.10 | 9.20 | 9.10 | 1.00 | 1.01 |
| 1988 | 6.50 | 5.90 | 12.00 | 0.54 | 0.49 |
| 1989 | 29.50 | 23.00 | 26.80 | 1.10 | 0.86 |
| 1990 | 27.30 | 23.70 | 18.90 | 1.44 | 1.25 |
| 1991 | 56.30 | 61.40 | 52.40 | 1.07 | 1.17 |
| 1992 | 40.90 | 41.30 | 29.50 | 1.39 | 1.40 |
| 1993 | 19.90 | 18.20 | 16.80 | 1.18 | 1.08 |
| 1994 | 20.20 | 28.90 | 18.50 | 1.09 | 1.56 |
| 1995 | 17.50 | 22.50 | 33.80 | 0.52 | 0.67 |
| 1996 | 27.60 | 35.40 | 33.10 | 0.83 | 1.07 |
| 1997 | 21.90 | 25.70 | 27.40 | 0.80 | 0.94 |
| 1998 | 14.20 | 14.20 | 20.20 | 0.70 | 0.70 |
| 1999 | 65.80 | 61.00 | 50.20 | 1.31 | 1.22 |
| 2000 | 21.90 | 26.90 | 20.50 | 1.07 | 1.31 |
| 2001 | 56.60 | 61.90 | 125.00 | 0.45 | 0.50 |
| 2002 | 144.40 | 136.00 | 160.80 | 0.90 | 0.85 |
| 2003 | 96.90 | 101.90 | 180.60 | 0.54 | 0.56 |
| 2004 | 138.00 | 150.00 | 175.30 | 0.79 | 0.86 |
| 2005 | 114.10 | 115.79 | 93.10 | 1.23 | 1.24 |
| 2006 | 50.00 | 51.80 | 27.90 | 1.79 | 1.86 |
| 2007 | 21.80 | 21.30 | 14.60 | 1.49 | 1.46 |
| 2008 | 87.20 | 86.20 | 91.90 | 0.95 | 0.94 |
| 2009 | 59.30 | 56.50 | 49.00 | 1.21 | 1.15 |
| 2010 | 169.00 | 162.90 | 130.80 | 1.29 | 1.25 |
| 2011 ^{c/} | 116.40 | 116.70 | 70.10 | 1.66 | 1.66 |
| 2012 | 63.80 | - | - | - | - |

TABLE II-7. Predicted and postseason returns of Columbia River adult summer and fall Chinook in thousands of fish.
(Page 3 of 3)

| Year | March Preseason Forecast ^{a/} | April STT Modeled Forecast ^{b/} | Postseason Return | March Pre/Postseason | April Pre/Postseason |
|--------------------|---|---|-------------------|-------------------------|-------------------------|
| MCB | | | | | |
| 1990 | 69.50 | 69.30 | 58.90 | 1.18 | 1.18 |
| 1991 | 48.40 | 48.50 | 35.40 | 1.37 | 1.37 |
| 1992 | 42.50 | 40.70 | 31.10 | 1.37 | 1.31 |
| 1993 | 33.00 | 32.30 | 27.50 | 1.20 | 1.17 |
| 1994 | 23.90 | 26.70 | 33.70 | 0.71 | 0.79 |
| 1995 | 25.00 | 30.00 | 34.20 | 0.73 | 0.88 |
| 1996 | 40.80 | 43.20 | 59.70 | 0.68 | 0.72 |
| 1997 | 72.10 | 61.90 | 59.00 | 1.22 | 1.05 |
| 1998 | 47.80 | 44.90 | 36.80 | 1.30 | 1.22 |
| 1999 | 38.30 | 27.70 | 50.70 | 0.76 | 0.55 |
| 2000 | 50.60 | 61.60 | 36.80 | 1.38 | 1.67 |
| 2001 | 43.50 | 45.30 | 76.40 | 0.57 | 0.59 |
| 2002 | 96.20 | 91.80 | 108.40 | 0.89 | 0.85 |
| 2003 | 104.80 | 94.60 | 150.20 | 0.70 | 0.63 |
| 2004 | 90.40 | 88.80 | 117.60 | 0.77 | 0.76 |
| 2005 | 89.40 | 89.73 | 98.00 | 0.91 | 0.92 |
| 2006 | 88.30 | 86.60 | 80.40 | 1.10 | 1.08 |
| 2007 | 68.00 | 69.10 | 46.90 | 1.45 | 1.47 |
| 2008 | 54.00 | 55.10 | 75.50 | 0.72 | 0.73 |
| 2009 | 94.40 | 97.90 | 73.10 | 1.29 | 1.34 |
| 2010 | 79.00 | 74.60 | 79.00 | 1.00 | 0.94 |
| 2011 ^{c/} | 100.00 | 100.40 | 85.40 | 1.17 | 1.18 |
| 2012 | 90.80 | - | - | - | - |
| SUMMER | | | | | |
| 2008 | 52.00 | | 55.53 | 0.94 | |
| 2009 | 70.70 | | 53.88 | 1.31 | |
| 2010 | 88.80 | | 72.35 | 1.23 | |
| 2011 ^{c/} | 91.10 | | 80.57 | 1.13 | |
| 2012 | 91.20 | | - | - | |

a/ March preseason forecasts are ocean escapements based on terminal run size and stock-specific cohort relationships affected by the historical "normal" ocean fisheries, generally between 1979 and the most recent complete broods.

b/ STT-modeled forecasts adjust March preseason forecasts for Council-adopted ocean regulations each year, and should provide a more accurate estimate of expected ocean escapement.

c/ Postseason estimates are preliminary.

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 1 of 4)

| Year | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season |
|--------------------|---|----------------------|---------------------|------------------------------------|----------------------|---------------------|----------------------------|----------------------|---------------------|---------------------------|----------------------|---------------------|
| | Nooksack-Samish Hatchery and Natural | | | East Sound Bay Hatchery | | | Skagit Hatchery | | | Skagit Natural | | |
| 1993 | 50.4 | 32.3 | 1.53 | 3.2 | 3.8 | 0.84 | 1.0 | 1.4 | 0.71 | 14.0 | 6.9 | 2.00 |
| 1994 | 46.6 | 28.1 | 1.66 | 3.2 | 0.7 | 4.00 | 1.3 | 5.5 | 0.30 | 8.4 | 5.9 | 1.27 |
| 1995 | 38.5 | 22.3 | 1.73 | 3.5 | 0.2 | 17.50 | 1.6 | 3.4 | 0.48 | 5.0 | 9.2 | 0.52 |
| 1996 | 27.0 | 29.2 | 0.92 | 1.7 | 0.5 | 2.43 | 1.0 | 1.2 | 0.83 | 7.1 | 10.9 | 0.58 |
| 1997 | 34.0 | 41.7 | 0.99 | 1.2 | 1.2 | 1.00 | 0.1 | 0.0 | - | 6.4 | 6.1 | 1.03 |
| 1998 | 28.0 | 31.5 | 0.95 | 0.5 | 0.3 | 1.67 | 0.0 | 0.0 | - | 6.6 | 15.0 | 0.44 |
| 1999 | 27.0 | 42.1 | 0.66 | 2.3 | 0.3 | 7.67 | 0.0 | 0.0 | - | 7.6 | 5.3 | 1.46 |
| 2000 | 19.0 | 32.6 | 0.57 | 5.0 | 0.1 | 50.00 | 0.0 | 0.0 | - | 7.3 | 17.3 | 0.42 |
| 2001 | 34.9 | 65.6 | 0.55 | 1.6 | 0.9 | 16.00 | 0.0 | 0.0 | - | 9.1 | 14.1 | 0.65 |
| 2002 | 52.8 | 57.0 | 0.99 | 1.6 | 0.9 | 2.29 | 0.0 | 0.1 | - | 13.8 | 20.0 | 0.69 |
| 2003 | 45.8 | 30.0 | 1.51 | 1.6 | 0.2 | 8.00 | 0.0 | 0.3 | - | 13.7 | 10.3 | 1.38 |
| 2004 | 34.2 | 18.1 | 1.83 | 0.8 | 0.0 | 200.00 | 0.5 | 0.0 | - | 20.3 | 24.3 | 0.83 |
| 2005 | 19.5 | 16.5 | 1.07 | 0.4 | 0.0 | 13.33 | 0.7 | 0.4 | 3.50 | 23.4 | 23.4 | 0.99 |
| 2006 | 16.9 | 31.9 | 0.53 | 0.4 | 0.0 | 25.00 | 0.6 | 0.4 | 1.51 | 24.1 | 22.5 | 1.07 |
| 2007 | 18.8 | 26.5 | 0.71 | 0.4 | 0.0 | 66.67 | 1.1 | 0.4 | 2.75 | 15.0 | 13.0 | 1.15 |
| 2008 | 35.3 | 29.1 | 1.21 | 0.8 | 0.0 | 0.00 | 0.7 | 0.2 | 3.50 | 23.8 | 15.0 | 1.59 |
| 2009 | 23.0 | 20.9 | 1.10 | 0.1 | 0.0 | 25.00 | 0.6 | 0.1 | 6.00 | 23.4 | 12.5 | 1.87 |
| 2010 ^{b/} | 30.3 | 41.2 | 0.74 | 2.3 | NA | NA | 0.9 | 0.1 | 11.25 | 13.0 | 10.0 | 1.30 |
| 2011 | 37.5 | NA | NA | 0.4 | NA | NA | 1.5 | NA | NA | 14.3 | NA | NA |
| 2012 | 42.8 | - | - | 1.1 | - | - | 1.3 | - | - | 8.3 | - | - |

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 2 of 4)

| Year | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season |
|--------------------|---|----------------------|---------------------|--|----------------------|---------------------|---|----------------------|---------------------|--|----------------------|---------------------|
| | Stillaguamish^{c/} Natural | | | Snohomish^{c/} Hatchery | | | Snohomish^{c/} Natural | | | Tulalip^{c/} Hatchery | | |
| 1993 | NA | 1.3 | - | 1.6 | 2.7 | 0.58 | 4.9 | 5.5 | 0.89 | 2.8 | 1.4 | 2.03 |
| 1994 | NA | 1.3 | - | 1.8 | 5.4 | 0.33 | 4.5 | 5.0 | 0.90 | 2.8 | 1.8 | 1.59 |
| 1995 | 1.8 | 0.9 | 1.92 | 2.2 | 4.0 | 0.54 | 4.3 | 4.0 | 1.08 | 2.3 | 8.5 | 0.27 |
| 1996 | 1.3 | 1.2 | 1.04 | 6.7 | 4.6 | 1.47 | 4.2 | 5.9 | 0.71 | 2.7 | 11.5 | 0.24 |
| 1997 | 1.6 | 1.2 | 1.36 | 7.7 | 12.0 | 0.64 | 5.2 | 4.4 | 1.19 | 4.0 | 8.7 | 0.46 |
| 1998 | 1.6 | 1.6 | 1.03 | 6.5 | 4.7 | 1.37 | 5.6 | 6.4 | 0.88 | 2.5 | 7.2 | 0.35 |
| 1999 | 1.5 | 1.1 | 1.36 | 7.8 | 4.7 | 1.65 | 5.6 | 4.8 | 1.16 | 4.5 | 15.2 | 0.30 |
| 2000 | 2.0 | 1.7 | 1.21 | 6.2 | 1.9 | 3.20 | 6.0 | 6.1 | 0.98 | 5.0 | 8.3 | 0.60 |
| 2001 | 1.7 | 1.4 | 1.22 | 4.1 | 0.9 | 4.57 | 5.8 | 8.4 | 0.69 | 5.5 | 5.1 | 1.08 |
| 2002 | 2.0 | 1.6 | 1.25 | 6.8 | 2.6 | 2.66 | 6.7 | 7.3 | 0.92 | 5.8 | 5.2 | 1.12 |
| 2003 | 2.0 | 1.0 | 1.98 | 9.4 | 5.8 | 1.63 | 5.5 | 5.6 | 0.99 | 6.0 | 8.7 | 0.69 |
| 2004 | 3.3 | 1.6 | 1.19 | 10.1 | 6.4 | 1.58 | 15.7 | 11.2 | 1.40 | 6.8 | 6.5 | 1.05 |
| 2005 | 2.0 | 1.2 | 1.42 | 9.9 | 4.0 | 2.48 | 14.2 | 5.0 | 2.84 | 6.4 | 7.4 | 0.86 |
| 2006 | 1.6 | 1.3 | 1.26 | 9.6 | 4.3 | 2.23 | 8.7 | 8.8 | 0.99 | 9.3 | 5.8 | 1.60 |
| 2007 | 1.9 | 0.8 | 2.38 | 8.7 | 6.6 | 1.32 | 12.3 | 4.0 | 3.08 | 8.4 | 6.1 | 1.38 |
| 2008 | 1.1 | 1.8 | 0.61 | 8.8 | 6.3 | 1.40 | 6.5 | 8.7 | 0.75 | 2.7 | 3.2 | 0.84 |
| 2009 | 1.7 | 1.2 | 1.42 | 4.9 | 2.2 | 2.23 | 8.4 | 2.3 | 3.65 | 4.0 | 1.7 | 2.35 |
| 2010 ^{b/} | 1.4 | 1.0 | 1.40 | 5.6 | 2.7 | 2.07 | 9.9 | 4.8 | 2.06 | 3.4 | 3.2 | 1.06 |
| 2011 | 1.8 | NA | NA | 5.2 | NA | NA | 7.4 | NA | NA | 3.5 | NA | NA |
| 2012 | 0.4 | - | - | 3.9 | - | - | 2.8 | - | - | 3.5 | - | - |

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 3 of 4)

| Year | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season |
|--------------------|---------------------------------------|----------------------|---------------------|--------------------------------------|----------------------|---------------------|--|----------------------|---------------------|---|----------------------|---------------------|
| | South Puget Sound Hatchery | | | South Puget Sound Natural | | | Strait of Juan de Fuca Hatchery | | | Strait of Juan de Fuca Natural | | |
| 1993 | 61.8 | 43.1 | 1.68 | 26.5 | 9.6 | 1.34 | 0.7 | 1.0 | 3.50 | 3.1 | 1.6 | 1.29 |
| 1994 | 52.7 | 49.9 | 1.08 | 18.0 | 10.5 | 0.60 | 3.9 | 1.2 | 2.44 | 1.0 | 1.0 | 2.00 |
| 1995 | 49.6 | 75.4 | 0.67 | 21.7 | 24.9 | 0.63 | 3.0 | 0.7 | 30.00 | 0.9 | 2.3 | 0.33 |
| 1996 | 51.9 | 53.2 | 0.89 | 19.0 | 16.5 | 0.53 | 2.8 | 1.4 | 14.00 | 0.9 | 2.0 | 0.29 |
| 1997 | 65.1 | 38.3 | 1.40 | 18.2 | 15.9 | 0.88 | 2.2 | 1.0 | 7.33 | 0.8 | 2.9 | 0.23 |
| 1998 | 67.8 | 49.6 | 1.24 | 21.8 | 14.6 | 0.79 | 1.7 | 1.7 | 1.00 | 0.9 | 2.1 | 0.47 |
| 1999 | 59.4 | 67.3 | 0.71 | 19.6 | 33.5 | 1.15 | 1.9 | 0.7 | 2.71 | 0.9 | 2.7 | 0.33 |
| 2000 | 77.5 | 47.4 | 1.39 | 17.5 | 39.5 | 1.26 | 2.0 | 1.2 | 1.67 | 1.1 | 1.7 | 0.65 |
| 2001 | 73.7 | 76.6 | 0.76 | 16.2 | 60.6 | 0.80 | 0.0 | 1.7 | 0.00 | 3.5 | 2.0 | 1.75 |
| 2002 | 90.8 | 69.3 | 1.07 | 16.9 | 57.0 | 0.79 | 0.0 | 1.6 | 0.00 | 3.6 | 2.2 | 0.97 |
| 2003 | 86.6 | 57.2 | 1.14 | 19.6 | 38.6 | 1.28 | 0.0 | 1.3 | 0.00 | 3.4 | 2.8 | 0.72 |
| 2004 | 86.5 | 66.6 | 1.16 | 17.5 | 42.3 | 0.61 | 0.0 | 1.4 | 0.00 | 3.6 | 4.1 | 0.85 |
| 2005 | 83.1 | 73.9 | 0.95 | 17.7 | 19.0 | 0.46 | 0.0 | 1.4 | 0.00 | 4.2 | 2.1 | 2.00 |
| 2006 | 85.8 | 104.1 | 0.82 | 21.3 | 37.0 | 0.58 | 0.0 | 1.2 | 0.00 | 4.2 | 3.2 | 1.31 |
| 2007 | 83.0 | 140.3 | 0.59 | 17.0 | 30.1 | 0.56 | 0.0 | 0.8 | 0.00 | 4.4 | 1.3 | 3.38 |
| 2008 | 101.6 | 90.6 | 1.12 | 21.1 | 32.2 | 0.65 | 0.0 | 0.7 | 0.00 | 3.2 | 1.2 | 2.67 |
| 2009 | 93.0 | 72.7 | 1.28 | 17.2 | 13.3 | 1.29 | 0.0 | 1.5 | 0.00 | 2.4 | 1.3 | 1.85 |
| 2010 ^{b/} | 97.4 | 82.9 | 1.17 | 12.7 | 13.9 | 0.91 | 0.0 | 0.7 | 0.00 | 1.9 | 2.6 | 0.73 |
| 2011 | 118.6 | NA | NA | 8.9 | NA | NA | 0.0 | NA | NA | 2.5 | NA | NA |
| 2012 | 95.8 | - | - | 8.9 | - | - | 2.7 | - | - | 2.1 | - | - |

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 4 of 4)

| Year | Preseason Forecast | Postseason Return | Pre/Post-season |
|--------------------|--|-------------------|-----------------|
| | Hood Canal Hatchery and Natural | | |
| 1993 | NA | 9.2 | - |
| 1994 | 11.7 | 8.1 | 1.44 |
| 1995 | 11.5 | 7.8 | 1.47 |
| 1996 | 3.9 | 16.2 | 0.24 |
| 1997 | 9.0 | 30.2 | 0.30 |
| 1998 | 2.7 | 20.9 | 0.13 |
| 1999 | 6.7 | 30.4 | 0.22 |
| 2000 | 14.0 | 34.4 | 0.41 |
| 2001 | 19.2 | 26.1 | 0.74 |
| 2002 | 25.3 | 30.2 | 0.84 |
| 2003 | 24.0 | 33.0 | 0.73 |
| 2004 | 29.6 | 34.3 | 0.86 |
| 2005 | 30.6 | 54.7 | 0.56 |
| 2006 | 30.2 | 40.7 | 0.74 |
| 2007 | 47.5 | 32.5 | 1.46 |
| 2008 | 36.8 | 33.1 | 1.11 |
| 2009 | 42.6 | 38.0 | 1.12 |
| 2010 ^{b/} | 45.0 | 43.7 | 1.03 |
| 2011 | 40.6 | NA | NA |
| 2012 | 46.8 | - | - |

a/ Puget Sound run size is defined as the run available to Puget Sound net fisheries. Does not include fish caught by troll and recreational fisheries inside Puget Sound.

b/ Postseason returns are preliminary.

c/ These numbers are in terms of terminal run of Chinook returning to area 8A. This includes all adult Chinook harvested in the net fisheries in Areas 8A, 8D, the Stillaguamish and Snohomish Rivers; harvest in sport fisheries in Area 8D and the Stillaguamish and Snohomish Rivers; and escapement.




FIGURE II-1. The Sacramento Index (SI) and relative levels of its components. The Sacramento River fall Chinook escapement goal range of 122,000-180,000 adult spawners is noted on the vertical axis.




FIGURE II-2. Regression estimator for the SI based on previous year's escapement of Sacramento River fall Chinook jacks. Years shown are SI years. The dashed line represents the 2012 SI predictor using 2009-2011 data, which are denoted by circles. The solid line represents the predictor using data from 1990-2011.






FIGURE II-3. Regression estimators for Klamath River fall Chinook ocean abundance (September 1) based on that year's river return of same cohort. Numbers in plots denote brood years.




FIGURE II-4. Selected preseason vs. postseason forecasts for Chinook stocks with significant contribution to Council area fisheries.

CHAPTER III - COHO SALMON ASSESMENT

COLUMBIA RIVER AND OREGON/CALIFORNIA COAST COHO

(OREGON PRODUCTION INDEX AREA)

The majority of coho harvested in the OPI area originate from stocks produced in rivers located within the OPI area (Leadbetter Point, Washington, to the U.S./Mexico border). These stocks include hatchery and natural production from the Columbia River, Oregon Coast, and northern California, and are divided into the following components: (1) public hatchery (OPIH), (2) Oregon coastal natural (OCN), including river and lake components, (3) Lower Columbia natural (LCN), and (4) natural and hatchery stocks south of Cape Blanco, Oregon, which include the Rogue, Klamath, and Northern California coastal stocks. Direct comparisons of 2012 abundance forecasts with recent year preseason abundance forecasts and postseason estimates, are reported in Table III-1.

Beginning in 2008, a new method was developed to estimate coho abundances for both the natural and hatchery components of the Columbia River and the Oregon coast. The traditional method of stock abundance estimation used only catch data from Leadbetter Point, Washington, to the U.S./Mexico border. The assumption prior to 2008 was that OPI stocks that were caught north of the OPI area were balanced by northern stocks that were caught inside the OPI area. This assumption was valid as long as fisheries north and south were balanced. However, in recent years, fisheries to the south have been more restrictive than those to the north, leading to underestimation of harvest of OPI area stocks. In addition, the estimation technique was not consistent with the methods used in Coho FRAM. The Mixed Stock Model (MSM) used for constructing the FRAM base period data was used to estimate the contribution of various coho stocks, including the OPI area stocks, to ocean fisheries and was based on CWT recoveries and associated tag rates. The MSM includes all fisheries that impact a particular stock, and therefore should provide a better overall accounting of total harvest and mortality of both Columbia River and Oregon coast coho stocks. The new run size estimates are based on the 1986-1997 base period and backwards FRAM run reconstructions for more recent years. The Oregon Production Index Technical Team (OPITT) decided to use the MSM run reconstruction database for future accounting and forecasts. The MSM estimates were refined for use in 2009, with particular attention to the base period reconstruction for OCN coho. In 2010, the relationship between the MSM and previous time series was reconsidered. The changes in fishery effort patterns that resulted in biased harvest estimates began in the mid- to late-1990s, so the first few years of the MSM time series should be equivalent to the previous time series. This was used as justification to use the MSM data set as a continuation of the previous time series starting in 1986. In 2012, the OPI hatchery and OCN predictors used the longer, merged time series. This results in a higher level of statistical significance for the predictors and lower residuals in most recent years.

Public Hatchery Coho

OPI area public hatchery coho smolt production occurs primarily in Columbia River facilities and net pens. Several facilities located in Oregon coastal rivers and in the Klamath River Basin, California, collectively produce fewer coho. OPI area smolt releases since 1960 are reported by geographic area in Appendix C, Table C-1.

Predictor Description

Prior to 2008, the OPIH stock predictor was a multiple linear regression with the following variables: (1) Columbia River jacks (Jack CR), (2) Oregon coastal and Klamath River Basin jacks (Jack OC), and (3) a correction term for the proportion of delayed smolts released from Columbia River hatcheries (Jack CR * [SmD/SmCR]).

In 2008, the stock predictor was modified slightly from that used in previous years. Because of the shorter data set (1986-2007 vs. 1970-2007) and the near-total phase-out of coastal coho salmon hatcheries, the factor for Oregon and California jacks (Jack OC) was not significant in the regression. A simplified model with all OPI jacks combined into one term (Jack OPI) was used, and all parameters were significant. In 2011, the longer (1970-2010) time series was used with the simplified model.

The OPIH stock predictor is partitioned into Columbia River early and late stocks based on the proportion of the 2011 jack returns of each stock adjusted for stock-specific maturation rates. The coastal hatchery stock is partitioned into northern and southern coastal stock components. The northern OPIH coastal stock is comprised of hatchery production from the central Oregon Coast. The southern OPIH coastal stock is comprised of hatchery production from the Rogue River basin in southern Oregon and the Klamath and Trinity basins in northern California. The 2012 partition was based on the proportion of the smolt releases in 2011.

For the 2012 abundance forecast, the data base includes 1970-2011 recruits and 1969-2010 jack returns (in thousands of fish). The model was:

$$\text{OPIH}(t) = a + b (\text{Jack OPI}(t-1) + c ((\text{Jack CR}(t-1) ([\text{SmD}(t-1)/\text{SmCR}(t-1)]))$$

Where:

$$\begin{aligned} a &= -65.13 \\ b &= 18.96 \\ c &= 26.96 \\ \text{adjusted } r^2 &= 0.95 \end{aligned}$$

The OPIH stock data set and a definition of the above terms are presented in Appendix C, Table C-2.

Predictor Performance

Recent year OPIH stock preseason abundance forecasts, partitioned by production area, stock, and as a total, are compared with postseason estimates in Table III-1. The 2011 preseason abundance prediction of 375,100 OPIH coho was 85 percent of the preliminary postseason estimate of 442,300 coho.

Since 1983, the OPIH predictor has performed well (Figure III-1a). The years with the highest variations were due principally to high interannual variability in the jack-to-adult ratios.

Stock Forecast and Status

Using the appropriate values from Appendix C, Table C-2, the OPIH abundance forecast for 2012 is 341,700 coho, 91 percent of the 2011 prediction and 85 percent of the preliminary 2011 postseason estimate.

Oregon Coastal Natural Coho

The OCN stock is composed of natural production north of Cape Blanco, Oregon from river (OCNR) and lake (OCNL) systems, which are forecasted independently.

ACLs are undefined in the FMP for ESA-listed stocks like OCN (and SONCC and CCC) coho, and are deferred to ESA consultation standards.

Predictor Description

Oregon Coastal Natural Rivers

Prior to 2010 a variety of methods were used to forecast OCNR coho abundance. Beginning in 2011, generalized additive models (GAMs) were used to relate OCNR recruitment to ocean environment indices. Nine variables were evaluated, ranging from indices of large-scale ocean patterns (e.g., Pacific Decadal Oscillation (PDO)) to local ecosystem variables (e.g., sea surface temperature at Charleston, OR). It was found that high explanatory power and promising forecast skill could be achieved when the mean May-July PDO averaged over the four years prior to the return year was used in combination with two other variables in a GAM. The multi-year average of the PDO, in essence, explains the lower frequency (multi-year) variability in recruitment and can be viewed as a replacement of the Regime Index used previously. A final set of six models using six different environmental indices plus parent spawner abundance was chosen from the possible model combinations. When averaging the predictions from the set of models (the ensemble mean), a higher skill (in terms of variance explained or cross-validation) was achieved than by selecting any single model. Making multiple forecasts from a set of models also provides a range of possible outcomes that reflects, to some degree, the uncertainty in understanding how salmon productivity is driven by ocean conditions.

The GAM with 3 predictor variables can be expressed in the following general form:

$$\hat{Y} = f(X_1) + f(X_2) + f(X_3) + \varepsilon$$

Where \hat{Y} is the prediction, X_1 through X_3 are the predictor variables, and ε is the deviation of \hat{Y} from the observation Y . For the prediction, Y was the log-transformation of annual recruit abundance. The term f represents a smooth function, which in this case is a cubic spline.

GAM Model Predictor used for 2012 forecast was:

Ensemble Mean of six forecasts based on environmental conditions and spawners.

| Variables | | | Prediction | r^2 | OCV ^{a/} |
|---|--|--|------------------------------|-------|-------------------|
| PDO | Spring Transition (Julian date; t-1) | Log Spawners (t-3) | 302,300 | 0.79 | 0.72 |
| PDO | Multivariate ENSO Index (Oct-Dec; t-1) | Upwelling (July-Sept; t-1) | 281,400 | 0.79 | 0.70 |
| PDO | Spring Transition (Julian date; t-1) | Multivariate ENSO Index (Oct-Dec; t-1) | 242,200 | 0.77 | 0.70 |
| PDO | Upwelling (July-Sept; t-1) | Sea Surface Temperature (May-Jul; t-1) | 298,400 | 0.78 | 0.69 |
| PDO | Sea Surface Height (Apr-June; t-1) | Upwelling (July-Sept; t-1) | 212,500 | 0.77 | 0.67 |
| PDO | Upwelling (Sept-Nov; t-1) | Sea Surface Temperature (Jan; t) | 250,100 | 0.76 | 0.67 |
| Ensemble Mean (90% prediction intervals) | | | 262,400 (137,600-496,300) | 0.81 | 0.74 |

a/ OCV – ordinary cross-validation score

The OCNR stock data set and a definition of the above terms are presented in Appendix C, Table C-4.

Oregon Coastal Natural Lakes

Since 1988, except for 2008, the abundance of OCNL index coho has been predicted using the most recent three-year average adult stock abundance. OCNL coho production occurs from three lake systems (Tennile, Siltcoos, and Tahkenitch). Production from these systems has declined substantially from the levels observed during 1950-1973, but has steadily increased in recent years.

For 2012, OPITT chose to use the most recent three-year average adult stock abundance which predicts 28,600 coho.

Predictor Performance

Recent year OCN preseason abundance predictions are compared to postseason estimates in Table III-1. Since 2000, the OCN predictor has underestimated abundance except for 2005 and 2007. The 2011 preseason abundance prediction of 294,000 OCN coho was 80 percent of the preliminary postseason estimate of 311,600 coho.

Stock Forecasts and Status

The 2012 preseason prediction for OCN (river and lake systems combined) is 291,000 coho, 117 percent of the 2011 preseason prediction and 93 percent of the 2011 postseason estimate (Table III-1). The 2012 preseason prediction for OCNR and OCNL components are 262,400 and 28,600 coho, respectively.

Based on parent escapement levels and observed OPI smolt-to-jack survival for 2009 brood OPI smolts, the total allowable OCN coho exploitation rate for 2012 fisheries is no greater than 20.0 percent under the Salmon FMP (Amendment 13) and no greater than 15.0 percent under the matrix developed by the OCN Coho Work Group during their review of Amendment 13 (Table V-7; Appendix A, Tables A-2 and A-3, respectively). The work group recommendation was accepted by the Council as expert biological advice in November 2000.

Private Hatchery Coho

There have been no Oregon coastal PRIH coho smolt releases since 1990.

Salmon Trout Enhancement Hatchery Coho Smolt Program

STEP program releases were discontinued after the 2004 brood.

Lower Columbia River Natural

LCN coho consist of naturally produced coho mostly from Columbia River tributaries below Bonneville dam; however, coho produced in the upper Willamette are not part of the ESA-listed ESU and are not included in the LCN coho forecast. LCN coho were listed as endangered under the Oregon State ESA in 2002, and as threatened under the Federal ESA on June 28, 2005. ACLs are undefined in the FMP for ESA-listed stocks like LCN coho, and are deferred to ESA consultation standards.

Predictor Description

The 2012 prediction for the Clackamas River is based on the recent 3-year cohort average counts at North Fork dam. The Clackamas ocean abundance forecast for 2012 is 5,100. The forecast for other Oregon lower Columbia natural (LCN) populations, including the Sandy River, are 3-year averages of recent year abundances based on spawning ground counts. The 2012 LCN coho ocean abundance forecast for all Oregon areas combined is 11,100 coho.

The 2012 prediction for the Washington LCN coho populations are derived by combining estimates of the 2009 brood year natural smolt production based on watershed area and the 5-year average ocean survival rate of 3.3 percent. The 2012 adult ocean abundance forecast for Washington LCN coho is 19,000 coho.

Predictor Performance

The LCN stock predictor methodology was developed in 2007. The preseason abundance compared to the postseason estimate is presented in Table III-1. The 2011 preseason abundance prediction of 22,700 LCN coho was 97 percent of the preliminary postseason estimate of 23,400 coho.

Stock Forecast and Status

The 2012 prediction for LCN coho is 30,100 coho (Table III-1). This ocean abundance estimate includes both Oregon and Washington LCN components.

NMFS ESA guidance for harvest of LCN coho in marine and mainstem Columbia River fisheries in recent years has been based on the allowable marine exploitation rate in a matrix developed by ODFW, similar to the OCN matrix. Based on parent escapement levels in the Sandy and Clackamas and observed OPI smolt-to-jack survival for 2009 brood OPI smolts, the allowable LCN coho marine exploitation rate in the ODFW matrix for 2012 fisheries is no greater than 15.0 percent; therefore, if the NMFS guidance is consistent with recent years, the total allowable marine and mainstem Columbia River exploitation rate for LCN coho in 2012 fisheries would be no more than 15.0 percent.

Oregon Production Index Area Summary of 2012 Stock Forecasts

The 2012 combined OPI area stock abundance is predicted to be 632,700 coho, which is 101 percent of the 2011 preseason prediction of 624,500 coho and 84 percent of the 2011 preliminary postseason estimate of 753,900 coho. The historical OPI abundances are reported in Table III-2.

WASHINGTON COAST COHO

Washington coastal coho stocks include all natural and hatchery stocks originating in Washington coastal streams north of the Columbia River to the western Strait of Juan de Fuca (west of the Sekiu River). The stocks in this group most pertinent to ocean salmon fishery management are Willapa Bay (hatchery), Grays Harbor, Quinalt (hatchery), Queets, Hoh, and Quillayute coho. These stocks contribute primarily to ocean fisheries off Washington and B.C.

A variety of preseason abundance estimators currently are employed for Washington coast and Puget Sound coho stocks, primarily based on smolt production and survival (Table I-2). These estimators are used to forecast preseason abundance of adult ocean (age-3) recruits.

A comparison was made of preseason ocean age-3 forecasts with postseason estimates derived from run reconstructions using FRAM ("Backwards" mode) to expand observed escapements to ocean abundance from CWT recovery data. It should be noted that forecast methodology has changed over time, and the overall trends and biases may not reflect the current methods.

Washington Coast coho are exceptions to the ACL requirements of the MSA because they are managed under an international agreement (the PST); therefore, specification of ACLs is not necessary for these stocks.

Willapa Bay

Predictor Description

The hatchery forecast is based on the marine survival rate of 5.3 percent from Zimmerman et al. 2012 applied to the 2009 brood year smolts. The natural forecast is based on a 3-year average terminal runsize (2009-2011) expanded to ocean age-3 recruits using an average pre-terminal catch (2005-2008) of Willapa Bay double index tag groups as a surrogate for natural harvest.

Predictor Performance

There was no information available to evaluate performance of predictors for Willapa coho stocks.

Stock Forecasts and Status

The 2012 Willapa Bay hatchery coho abundance forecast is 88,774 ocean recruits compared to a 2011 preseason forecast of 64,658. The 2012 natural coho forecast is 81,325 ocean recruits, compared to a 2011 preseason forecast of 47,788.

Grays Harbor

Preseason abundance forecasts are made for natural fish throughout the system and for hatchery fish returning to three freshwater rearing complexes and three saltwater net-pen sites. The forecasts include fish originating from numerous volunteer production projects.

Predictor Description

The natural coho forecast consists of an estimate of smolt production in the Humptulips and Chehalis basins multiplied by a PDO-based marine survival rate.

The 2012 hatchery coho forecast is an estimate of smolt releases from on- and off-station sites, multiplied by the average return per release for four years (2005-2008 BY) and then expanded to ocean recruit abundance based on CWT recoveries for 2003-2007 return years.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates for Grays Harbor natural coho derived from Backwards FRAM run reconstruction indicated no notable bias (Table III-3, Figure III-1).

Stock Forecasts and Status

The abundance forecast for Grays Harbor natural stock coho for 2012 is 150,200 ocean age-3 recruits. This ocean abundance results in an allowable exploitation rate of 65 percent under the FMP and the 2002 PST Southern Coho Management Plan (Table III-5).

The forecast for hatchery stock ocean abundance is 47,804 ocean age-3 recruits.

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Grays Harbor coho $MFMT = 0.65$ and the OFL is $S_{OFL} = 150,200 \times (1 - 0.65) = 52,570$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Quinault River

Predictor Description

The Quinault River natural coho forecast is based on the mean estimate of recent ocean recruits for 2004 through 2010. All natural coho are unmarked.

The Quinault River hatchery coho forecast is based on an estimated release of 657,993 smolts, multiplied by the recent 5-year average smolt return rate of 5.38 percent for the Quinault National Fish Hatchery.

Predictor Performance

There was no information available to evaluate performance of predictors for these stocks.

Stock Forecasts and Status

The 2012 forecast for Quinault natural coho is 27,278 age-3 ocean recruits, an increase of 19 percent from the 2011 forecast of 22,947.

The Quinault hatchery coho forecast is 35,421 age-3 ocean recruits, including 30,785 marked coho and 4,636 unmarked coho.

Queets River

Predictor Description

The natural coho forecast represents the estimated smolt production (412,722) multiplied by an expected survival rate of 9.02 percent. The survival rate estimate is based on a binomial logistic regression model developed by Quinault Fisheries Department. This model consists of a regression of Queets survival rates from return years 1992-2009 as estimated using backward FRAM run reconstructions, and the standardized monthly mean Pacific Decadal Oscillation (PDO) values from January through August for the corresponding years the smolts entered salt water.

The 2012 hatchery coho forecast is based on a smolt release of 696,482 multiplied by the recent 3 year average marine survival rate (3.64 percent). Approximately 88 percent of the fish released from the Salmon River facility were marked with an adipose fin clip.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated no persistent tendency to under- or over- predict abundance. The 2010 forecast was slightly higher than the postseason estimate (Table III-3; Figure III-1).

Stock Forecasts and Status

The 2012 Queets natural coho forecast is 37,228 ocean recruits, an increase of 180 percent compared to the 2011 forecast level of 13,279. This ocean abundance results in an allowable exploitation rate of 65 percent under the FMP and the 2002 PST Southern Coho Management Plan (Table III-5).

The 2011 Queets hatchery (Salmon River) coho forecast is 25,327 ocean recruits, an increase of 55 percent compared to the 2011 forecast of 16,331.

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Queets River coho $MFMT = 0.65$, and the OFL is $S_{OFL} = 37,228 \times (1-0.65) = 13,030$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Hoh River

Predictor Description

The natural coho forecast is based on estimated smolt production per square mile of watershed from the Clearwater tributary to the Queets River (958 smolts/square mile), multiplied by the size of the Hoh watershed (299 square miles), for a total of 286,442 smolts. The total natural smolt production estimate was then multiplied by an expected survival rate of 5.0 percent. To the north, the Strait of Juan de Fuca wild coho survival was estimated at 4.14 percent, and to the south the Chehalis was estimated at 4.8 percent. The Queets survival rate was not available, but will likely be between 6.5 percent and 9 percent.

The estimated survival rate for Hoh wild coho was selected as a rate within the range of these other forecasts, and 1 percentage point lower than the rate adopted for the Quillayute system just to the north.

No hatchery production is projected for the Hoh system for 2012.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated a tendency to under-predict actual run-size (Table III-3; Figure III-1). In 2010, the preseason forecast was lower than the postseason return.

Stock Forecasts and Status

The 2012 Hoh River natural coho forecast is 14,322 ocean recruits, an increase of 23 percent compared to the 2011 forecast of 11,625. This ocean abundance results in an allowable exploitation rate of 65 percent under the FMP and the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Hoh River coho $MFMT = 0.65$, and the OFL is $S_{OFL} = 14,322 \times (1 - 0.65) = 9,309$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Quillayute River

Quillayute River coho consist of a summer run that is managed primarily for hatchery production, and a fall run that is managed primarily for natural production. Quillayute River coho have both natural and hatchery components to both runs.

Predictor Description

The basin total coho smolt production estimate (summer and fall stocks) was derived using the estimated coho smolt production in the Clearwater Basin of 134,052, which was 2.12 times its average production during the years a smolt trap was operated on the Bogachiel River (1987, 1988 and 1990), and 2.20 times its average production during the years a trap was operated on the Dickey River (1992-1994). Using 2.12 as a multiplier of the estimated average smolt production of the Quillayute system excluding the Dickey (217,257) yields an estimated production of 459,851 coho smolts. The Dickey production yields an additional 194,142 smolts to the system. The total freshwater production for the basin is estimated to be 653,993 smolts. Smolt production was apportioned according to brood year natural spawning escapements of summer and fall coho to yield the smolt estimates for each natural population.

Summer Coho

The summer natural coho forecast is based on the estimated total summer coho smolt production (95,042) and a projected ocean survival rate of 6.0 percent. This is a lower ocean survival rate than the 7.0 percent used in 2011.

An examination of the return rates of both hatchery releases and natural smolts indicates that hatchery return rates are 1.5 to 2.0 percent below natural returns. Thus, for the hatchery component, an ocean survival rate of 4.0 percent was selected. The survival rate of 4.0 percent was multiplied by a release of 106,580 smolts.

Fall Coho

The forecast for the natural component was based on the estimated total fall coho smolt production (558,951) multiplied by an expected marine survival rate of 6.0 percent, which was the same as used for the summer natural returns.

The fall hatchery production forecast was based on the same prediction of marine survival (4.0 percent) used for the summer hatchery coho forecast, multiplied by a release of 422,612 smolts.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates for fall natural coho derived from Backwards FRAM run reconstruction indicated no notable bias (Table III-3; Figure III-1). The 2010 preseason forecast exceeded the postseason estimate by a factor of 1.29.

Stock Forecasts and Status

The 2012 Quillayute River summer natural and hatchery coho forecasts are 5,702 and 4,263 ocean recruits, respectively. Approximately 99 percent of the hatchery smolts were marked with an adipose fin clip; an additional 635 unmarked smolts were released. The 2012 forecast abundance of natural summer coho is 104 percent higher than the 2011 forecast, while the hatchery forecast is 21 percent lower than the 2011 forecast.

The 2012 Quillayute River fall natural and hatchery coho forecasts are 33,537 and 16,904 ocean recruits, respectively. The 2012 forecast abundance of natural Quillayute fall coho is 18 percent higher, and the hatchery forecast is 46 percent lower, than their respective 2011 forecasts. The hatchery smolts were marked as follows: 263,437 with adipose fin-clip only; 78,763 with adipose fin-clip and CWT; 78,098 with CWT only; 2,314 with no mark or tag.

The ocean abundance forecast for Quillayute fall natural coho results in an allowable exploitation rate of 65 percent under the 2002 PST Southern Coho Management Plan (Table III-5). The MFMT for Quillayute coho is currently undefined in the FMP.

North Washington Coast Independent Tributaries

Predictor Description

Production from several smaller rivers and streams along the North Washington Coast (Waatch River, Sooes River, Ozette River, Goodman Creek, Mosquito Creek, Cedar Creek, Kalaloch Creek, Raft River, Camp Creek, Duck Creek, Moclips River, Joe Creek, Copalis River, Conner Creek), which flow directly into the Pacific Ocean, is forecast as an aggregate. Generally, stock assessment programs on these systems are minimal.

The 2012 forecast of natural coho production for these independent streams is based on a prediction of 700 smolts per square mile of watershed drainage, 424 square miles of watershed, and an expected marine survival rate of 5.3 percent. This rate was the average of the jack-based and the PDO models.

The hatchery forecast is based on the relationship between the log-transformed jack return rate to Makah National Fish Hatchery and the log-transformed marine survival rate from smolt to January age-3. The predicted marine survival of 6.55 percent for the brood year 2009 was multiplied by the 2009 brood year smolt release (215,035) from the Makah National Fish Hatchery.

Predictor Performance

There was no information available to evaluate performance of predictors for these stocks.

Stock Forecasts and Status

The 2012 forecast of natural coho production for these independent streams is 15,730 age-3 ocean recruits. The hatchery forecast is 11,430 age-3 ocean recruits, and approximately 81 percent of the smolts released were marked with an adipose fin clip.

PUGET SOUND COHO STOCKS

Puget Sound coho salmon stocks include natural and hatchery stocks originating from U.S. tributaries in Puget Sound and the Strait of Juan de Fuca. The primary stocks in this group that are most pertinent to ocean salmon fishery management were Strait of Juan de Fuca, Hood Canal, Skagit, Stillaguamish, Snohomish, and South Puget Sound (hatchery) coho. These stocks contribute primarily to ocean fisheries off Washington and B.C.

A variety of preseason abundance estimators currently are employed for Puget Sound coho stocks, primarily based on smolt production and survival (Table I-2). These estimators are used to forecast preseason abundance of adult ocean age-3 recruits. Forecasts for natural Puget Sound coho stocks were generally derived by measured or predicted smolt production from each major watershed or region, multiplied by stock-specific marine survival rate predictions based on a jack return model from the WDFW Big Beef Creek Research Station in Hood Canal, adult recruits/smolt data generated from the WDFW Deschutes River Research Station, and a natural coho CWT tagging program at Baker Lake (Skagit River basin), or other information. Puget Sound hatchery forecasts were generally the product of 2009 brood year (BY) smolt releases from each facility, and a predicted marine survival rate for each program. Marine survival rates were typically based on recent year average survival rates derived from CWT recovery information and/or run reconstructions, and review of relationships between jack returns and adult marine survival rates at selected hatcheries.

The 2012 total hatchery and natural coho ocean recruit forecast for the Puget Sound region of is 731,000, compared to a 2011 forecast of 981,000. The hatchery coho forecast is 371,800 compared to the 2011 forecast of 380,900, and the natural coho forecast for 2012 of 359,100 is much lower than the 2011 forecast of 600,100.

A comparison was made of preseason ocean age-3 forecasts with postseason estimates derived from run reconstructions using FRAM ("Backwards" mode) to expand observed escapements to ocean abundance from CWT recovery data. It should be noted that forecast methodology has changed over time, and the overall trends and biases may not reflect the current methods.

Puget Sound coho are exceptions to the ACL requirements of the MSA because they are managed under an international agreement (the PST); therefore, specification of ACLs is not necessary for these stocks.

Strait of Juan de Fuca

Predictor Description

As in past years, the natural and hatchery coho forecasts include both Eastern and Western Strait of Juan de Fuca drainages. The natural coho forecast was derived by multiplying the estimated 2009 brood natural smolt production for the region by a predicted ocean marine survival rate that is the weighted mean of four predictions of marine survival from four regression models. Those models were:

- the log-transformed jack return rate to the Lower Elwha Hatchery
- the coho CPUE from the NOAA September trawl survey
- the copepod species-richness anomaly, generated by NOAA and
- a multiple regression model with the independent variables of $\ln(\text{jack return rate})$ and copepod species-richness.

The hatchery forecasts were based on applying hatchery-specific marine survival rate predictions (2.28 percent for Dungeness, 2.07 percent for Elwha) to the 2009 BY smolt releases for each hatchery. The marine survival rate predictions for the hatchery stocks were based on 2-year averages of estimated return rates of adults in 2009 and 2010.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated a tendency to under predict actual run-size (Table III-4). The 2010 postseason estimate exceeded the preseason forecast by a factor of 2.5.

Stock Forecasts and Status

The 2012 forecasts for Strait of Juan de Fuca natural and hatchery coho age-3 ocean recruits are 12,628 and 18,647, respectively.

The preliminary preseason forecast of 12,628 age-3 ocean recruits places Strait of Juan de Fuca natural coho in the low abundance based status category, which results in an allowable total exploitation rate of no more than 40 percent under both the Council adopted exploitation rate matrix (Appendix A, Table A-4) the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Strait of Juan de Fuca coho $MFMT = 0.60$, and the OFL is $S_{OFL} = 12,628 \times (1 - 0.60) = 5,051$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Nooksack-Samish

Predictor Description

The natural coho forecast is the product of projected natural smolt production from each stream basin in the region, multiplied by a marine survival rate expectation of 5.07 percent.

The hatchery forecasts are based on the 2005-2007 BY average recruits/smolt rate of 1.02 percent (Lummi Bay Hatchery) or 5.18 percent (Skookum Hatchery) multiplied by the number of smolts released.

Predictor Performance

There was no information available to evaluate performance of predictors for Nooksack-Samish coho stocks.

Stock Forecasts and Status

The 2012 forecasts for Nooksack-Samish natural and hatchery coho ocean recruits are 62,833 and 25,188 respectively.

Skagit

Predictor Description

The natural coho forecast is the product of measured smolt production from the Skagit basin multiplied by a marine survival rate expectation of 5.07 percent. The natural coho marine survival rate was based on the median of 2004-2008 brood year recruits/smolt of CWT Baker River natural coho.

The hatchery forecasts are based on Marblemount Hatchery CWT recoveries from the 2003-2007 brood years. Marine survival rates were calculated separately for adipose-marked and non-marked returns, and the median marine survival of the two groups was averaged resulting in a recruits/smolt rate of 4.3 percent, which was multiplied by the total number of smolts released from all regional hatcheries.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated a tendency to over-predict actual run-size, especially early in the time series (Table III-4; Figure III-1b). However, the 2009 postseason estimate exceeded the preseason forecast by a factor of 2.2.

Stock Forecasts and Status

The 2012 forecasts for Skagit River natural and hatchery coho ocean recruits are 48,310 and 14,922 (13,632 from in-river hatchery production, 1,289 from Oak Harbor net-pens), respectively.

The preliminary preseason forecast of 48,310 age-3 ocean recruits places Skagit natural coho in the low abundance based status category, which results in an allowable total exploitation rate of no more than 35 percent under both the Council adopted exploitation rate matrix (Appendix A, Table A-4) and the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Skagit River coho $MFMT = 0.60$, and the OFL is $S_{OFL} = 48,310 \times (1 - 0.60) = 19,324$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Stillaguamish

Predictor Description

The natural coho forecast was based upon a smolt trap catch per unit effort (CPUE) regressed against adult terminal returns, for brood years 1999-2008. This terminal runsize estimate was then expanded by a pre-terminal Puget Sound exploitation rate.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated a tendency to under-predict actual run-size (Table III-4; Figure III-1b). The 2009 postseason estimate exceeded the preseason forecast by a factor of 2.

Stock Forecasts and Status

The 2012 forecast for Stillaguamish River natural coho age-3 ocean recruits is 47,510.

The preliminary preseason forecast of 47,507 age-3 ocean recruits places Stillaguamish natural coho in the normal abundance based status category, which results in an allowable total exploitation rate of no

more than 50 percent under both the Council adopted exploitation rate matrix (Appendix A, Table A-4) and the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Stillaguamish coho $MFMT = 0.50$, and the OFL is $S_{OFL} = 47,507 \times (1-0.60) = 23,754$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Snohomish

The natural coho forecast used the estimated 2009 brood year smolt production from multiplied by a 10.9 percent marine survival rate expectation, which is based on average South Fork Skykomish coho marine survival (return years 1998-2010).

The hatchery forecasts were based on brood year 2009 releases multiplied by a 6.0 percent marine survival rate of Wallace Hatchery CWT releases (2000-2007 brood year average).

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated no persistent tendency to under- or over- predict abundance. The 2010 forecast was higher than the postseason estimate by a factor of 1.85 (Table III-4).

Stock Forecasts and Status

The 2012 forecast for Snohomish River natural coho ocean recruits is 109,000. The Snohomish regional hatchery coho forecast is 49,837; 8,460 for Skykomish River/Wallace River Hatchery facility releases and 36,628 for the Tulalip Bay facility.

The preliminary preseason forecast of 109,000 age-3 ocean recruits places Snohomish natural coho in the low abundance based status category, which results in an allowable total exploitation rate of no more than 40 percent under both the Council adopted exploitation rate matrix (Appendix A, Table A-4) and the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Snohomish coho $MFMT = 0.65$, and the OFL is $S_{OFL} = 109,000 \times (1-0.60) = 43,600$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

Hood Canal

Predictor Description

The natural coho forecast is based on a regression of CWT natural Big Beef Creek jacks on Hood Canal natural coho run sizes, using brood years 1983-1998 and 2002-2007. The 1999-2001 broods were excluded because of the unusually high recruit per tagged jack ratio, which is not expected to occur this year.

The hatchery coho forecasts are based on average cohort reconstruction-based recruits/smolt for the 2002-2007 broods from each facility, applied to the 2009 brood smolt releases for each facility. The marine survival rates used for these forecasts were 9.0 percent for George Adams Hatchery, 2.9 percent for Port Gamble Net Pens, 9.7 percent for the Quilcene National Fish Hatchery, and 3.9 percent for the Quilcene Bay Net Pens.

Predictor Performance

A comparison of preseason ocean age-3 forecasts with postseason estimates derived from Backwards FRAM run reconstruction indicated no persistent tendency to under- or over- predict abundance in recent years. The 2010 forecast was slightly higher than the postseason estimate (Table III-4; Figure III-1b).

Stock Forecasts and Status

The 2012 forecasts for Hood Canal region natural and hatchery coho ocean recruits are 73,415 and 62,641 respectively.

The preliminary preseason forecast of 73,415 age-3 ocean recruits places Hood Canal natural coho in the normal abundance based status category, which results in an allowable total exploitation rate of no more than 65 percent under both the Council adopted exploitation rate matrix (Appendix A, Table A-4) and the 2002 PST Southern Coho Management Plan (Table III-5).

OFL

The OFL is defined in terms of spawner escapement (S_{OFL}). For Hood Canal coho $MFMT = 0.65$, and the OFL is $S_{OFL} = 73,415 \times (1 - 0.65) = 25,695$. The preseason S_{OFL} value will be recalculated with postseason abundance estimates (when available) to assess OFL compliance.

South Sound

Predictor Description

The natural coho forecast is the product of projected smolt production from each of the stream basins in the region multiplied by variable marine survival rate expectations of 4.0 to 4.6 percent for natural coho in the region. The upper South Sound natural stocks' marine survival rate (4.6 percent) was based upon a five year average rate of return (return years 2006-2010) of Lake Washington natural smolts. The deep South Sound stocks' marine survival prediction (4.0 percent) was selected from results of regressions of Deschutes River natural-origin coho against various ocean physical and biological indicators. The hatchery coho forecasts were typically based on the 2005-2007 brood year average CWT-based recruits/smolt rate for each facility, applied to the 2009 brood smolt releases. The expected survival rates range from 0.6 to 5.2 percent.

Stock Forecasts and Status

The 2012 forecasts for South Sound region natural and hatchery coho ocean recruits are 43,055 and 162,916 respectively.

STOCK STATUS DETERMINATION UPDATES

There were several updates and additions to the spawning escapement estimates for Puget Sound and Washington Coast coho in the SAFE document. Previously unavailable 2011 spawning escapements are now available for Willapa, Grays Harbor, and Hoh, Strait of Juan de Fuca, and Skagit coho.

The most recent 3-year geometric mean spawning escapement estimated for Strait of Juan de Fuca coho reported in the SAFE document was 9,875, less than the S_{MSY} objective of 11,000. In the SAFE document the three years were erroneously reported as 2009-2011, however, the three years were actually 2008-2010. New information in the form of a preliminary 2011 spawning estimate indicates a 3-year geometric mean (2009-2011) of 11,295, which meets the default rebuilt criterion in the Salmon FMP for an overfished stock (Table V-4). Other than Strait of Juan de Fuca coho, the 2011 estimates did not change the status (e.g., overfished, rebuilt, etc.) for any of these stocks.

SELECTIVE FISHERY CONSIDERATIONS FOR COHO

As the region has moved forward with mass marking of hatchery coho salmon stocks, selective fishing options have become an important consideration for fishery managers. Projected coho mark rates in Canadian, Puget Sound and north Washington Coast fisheries are generally higher than 2011 projections, but lower in fisheries to the south. Table III-6 summarizes projected 2012 mark rates for coho fisheries by month from Southern British Columbia, Canada to the Oregon Coast, based on preseason abundance forecasts.

TABLE III-1. Preliminary 1996-2011 preseason and postseason coho stock abundance estimates for Oregon production index area stocks in thousands of fish. (Page 1 of 2)

| Stock | Year | Preseason | Postseason ^{a/} | Preseason/Postseason ^{a/} |
|---|------|-----------|--------------------------|------------------------------------|
| Oregon Production Index Area Hatchery Total^{b/} | 1996 | 309.2 | 182.6 | 1.69 |
| | 1997 | 376.1 | 215.3 | 1.75 |
| | 1998 | 118.4 | 203.6 | 0.58 |
| | 1999 | 559.2 | 319.6 | 1.75 |
| | 2000 | 671.4 | 677.1 | 0.99 |
| | 2001 | 1,707.6 | 1,454.2 | 1.17 |
| | 2002 | 361.7 | 660.1 | 0.55 |
| | 2003 | 863.1 | 952.5 | 0.91 |
| | 2004 | 623.9 | 634.6 | 0.98 |
| | 2005 | 389.9 | 443.1 | 0.88 |
| | 2006 | 398.8 | 440.6 | 0.91 |
| | 2007 | 593.6 | 476.5 | 1.25 |
| | 2008 | 216.1 | 565.4 | 0.38 |
| | 2009 | 1,073.1 | 1,066.2 | 1.01 |
| Columbia River Early | 2010 | 408.0 | 551.3 | 0.74 |
| | 2011 | 375.1 | 442.3 | 0.85 |
| | 2012 | 341.7 | - | - |
| | 1996 | 142.2 | 98.0 | 1.45 |
| | 1997 | 206.9 | 129.8 | 1.59 |
| | 1998 | 63.8 | 126.4 | 0.50 |
| | 1999 | 325.5 | 174.9 | 1.86 |
| | 2000 | 326.3 | 378.0 | 0.86 |
| | 2001 | 1,036.5 | 873.0 | 1.19 |
| | 2002 | 161.6 | 324.7 | 0.50 |
| | 2003 | 440.0 | 645.7 | 0.68 |
| | 2004 | 313.6 | 389.0 | 0.81 |
| | 2005 | 284.6 | 282.7 | 1.01 |
| | 2006 | 245.8 | 251.4 | 0.98 |
| | 2007 | 424.9 | 291.0 | 1.46 |
| Columbia River Late | 2008 | 110.3 | 333.9 | 0.33 |
| | 2009 | 672.7 | 681.4 | 0.99 |
| | 2010 | 245.3 | 274.3 | 0.89 |
| | 2011 | 216.0 | 288.5 | 0.75 |
| | 2012 | 229.8 | - | - |
| | 1996 | 114.4 | 30.8 | 3.71 |
| | 1997 | 86.5 | 53.7 | 1.61 |
| | 1998 | 24.9 | 47.3 | 0.53 |
| | 1999 | 140.9 | 120.7 | 1.17 |
| | 2000 | 278.0 | 260.1 | 1.07 |
| | 2001 | 491.8 | 488.3 | 1.01 |
| | 2002 | 143.5 | 271.8 | 0.53 |
| | 2003 | 377.9 | 248.0 | 1.52 |
| | 2004 | 274.7 | 203.0 | 1.35 |
| | 2005 | 78.0 | 111.6 | 0.70 |
| | 2006 | 113.8 | 156.3 | 0.73 |
| | 2007 | 139.5 | 171.0 | 0.82 |
| | 2008 | 86.4 | 207.6 | 0.42 |
| | 2009 | 369.7 | 374.1 | 0.99 |
| | 2010 | 144.2 | 263.6 | 0.55 |
| | 2011 | 146.5 | 141.2 | 1.04 |
| | 2012 | 87.4 | - | - |

TABLE III-1. Preliminary 1996-2012 preseason and postseason coho stock abundance estimates for Oregon production index area stocks in thousands of fish. (Page 2 of 3)

| Stock | Year | Preseason | Postseason ^{a/} | Preseason/Postseason ^{a/} |
|--|------|-----------|--------------------------|------------------------------------|
| Oregon Coast North of Cape Blanco | 1996 | 38.5 | 28.0 | 1.38 |
| | 1997 | 60.4 | 19.0 | 3.18 |
| | 1998 | 21.6 | 19.7 | 1.10 |
| | 1999 | 59.4 | 14.4 | 4.13 |
| | 2000 | 48.5 | 23.4 | 2.07 |
| | 2001 | 127.3 | 46.9 | 2.71 |
| | 2002 | 36.6 | 41.6 | 0.88 |
| | 2003 | 29.3 | 34.5 | 0.85 |
| | 2004 | 16.6 | 21.7 | 0.76 |
| | 2005 | 11.5 | 10.7 | 1.07 |
| | 2006 | 8.6 | 7.9 | 1.09 |
| | 2007 | 7.0 | 1.3 | 5.38 |
| | 2008 | 1.7 | 7.1 | 0.24 |
| | 2009 | 7.3 | 7.5 | 0.97 |
| | 2010 | 4.4 | 8.6 | 0.51 |
| Oregon and California Coast South of Cape Blanco | 2011 | 3.6 | 3.6 | 1.00 |
| | 2012 | 6.4 | - | - |
| | 1996 | 14.2 | 25.8 | 0.55 |
| | 1997 | 22.3 | 12.8 | 1.74 |
| | 1998 | 8.1 | 10.2 | 0.79 |
| | 1999 | 33.4 | 9.6 | 3.48 |
| | 2000 | 18.6 | 15.6 | 1.19 |
| | 2001 | 52.0 | 46.0 | 1.13 |
| | 2002 | 20.0 | 22.0 | 0.91 |
| | 2003 | 15.9 | 24.3 | 0.65 |
| | 2004 | 19.0 | 29.9 | 0.64 |
| | 2005 | 15.8 | 38.1 | 0.41 |
| | 2006 | 30.6 | 25.0 | 1.22 |
| | 2007 | 22.2 | 13.2 | 1.68 |
| | 2008 | 17.7 | 16.8 | 1.05 |
| | 2009 | 23.4 | 3.1 | 7.55 |
| Lower Columbia River Natural | 2010 | 14.1 | 4.8 | 2.94 |
| | 2011 | 9.0 | 9.0 | 1.00 |
| | 2012 | 18.1 | - | - |
| | 2007 | 21.5 | 19.4 | 1.11 |
| | 2008 | 13.4 | 27.2 | 0.49 |
| | 2009 | 32.7 | 40.4 | 0.81 |
| | 2010 | 15.1 | 30.8 | 0.49 |
| | 2011 | 22.7 | 23.4 | 0.97 |
| | 2012 | 30.1 | - | - |

TABLE III-1. Preliminary 1996-2012 preseason and postseason coho stock abundance estimates for Oregon production index area stocks in thousands of fish. (Page 3 of 3)

| Stock | Year | Preseason | Postseason ^{a/} | Preseason/Postseason ^{a/} |
|--|------|-----------|--------------------------|------------------------------------|
| Oregon Coast Natural (Rivers and Lakes) | 1996 | 63.2 | 86.1 | 0.73 |
| | 1997 | 86.4 | 27.8 | 3.11 |
| | 1998 | 47.2 | 29.2 | 1.62 |
| | 1999 | 60.7 | 51.9 | 1.17 |
| | 2000 | 55.9 | 69.0 | 0.81 |
| | 2001 | 50.1 | 163.2 | 0.31 |
| | 2002 | 71.8 | 304.5 | 0.24 |
| | 2003 | 117.9 | 278.8 | 0.42 |
| | 2004 | 150.9 | 197.0 | 0.77 |
| | 2005 | 152.0 | 150.1 | 1.01 |
| | 2006 | 60.8 | 116.4 | 0.52 |
| | 2007 | 255.4 | 60.0 | 4.26 |
| | 2008 | 60.0 | 170.9 | 0.35 |
| | 2009 | 211.6 | 257.0 | 0.82 |
| | 2010 | 148.0 | 266.8 | 0.55 |
| | 2011 | 249.4 | 311.6 | 0.80 |
| | 2012 | 291.0 | - | - |
| Salmon Trout Enhancement Program^{c/} | 1996 | 0.4 | 1.2 | 0.33 |
| | 1997 | 1.3 | 0.3 | 4.33 |
| | 1998 | 0.2 | 0.3 | 0.67 |
| | 1999 | 0.7 | 0.4 | 1.75 |
| | 2000 | 0.6 | 0.5 | 1.20 |
| | 2001 | 1.0 | 1.4 | 0.71 |
| | 2002 | 0.6 | 3.0 | 0.20 |
| | 2003 | 3.6 | 3.6 | 1.00 |
| | 2004 | 3.1 | 1.0 | 3.10 |
| | 2005 | 1.0 | 0.4 | 2.50 |
| | 2006 | 0.6 | 0.1 | 6.00 |
| | 2007 | 0.2 | 0.0 | - |
| | 2008 | - | - | - |
| | 2009 | - | - | - |
| | 2010 | - | - | - |
| | 2011 | - | - | - |

a/ Postseason estimates are based on preliminary data, and not all stocks have been updated with final estimates.

b/ LCN abundance is included as a subset of early/late hatchery abundance beginning in 2007. STEP estimates not included

c/ Program was discontinued in 2005.

TABLE III-2. Oregon production index (OPI) area coho harvest impacts, spawning, abundance, and exploitation rate estimates in thousands of fish.^{a/}

| Year or Avg. | Oregon and California Coastal Returns | | | | | | | Ocean | OCN |
|--------------------|---------------------------------------|-------|-------------------------------------|------------------------|------------|----------------|-------------------------|-------------------------|-------------------|
| | Ocean Fisheries ^{b/} | | Hatcheries and | OCN | Private | Columbia River | Abundance ^{e/} | Exploitation Rate | Exploitation Rate |
| | Troll | Sport | Freshwater Harvest ^{c/} | Spawners ^{d/} | Hatcheries | Returns | | Based on OPI | Based on |
| | | | | | | | | Abundance ^{f/} | Postseason |
| | | | | | | | | | FRAM |
| 1970-1975 | 1,629.6 | 558.4 | 45.8 | 55.2 | - | 460.4 | 2,749.3 | 0.80 | - |
| 1976-1980 | 1,253.6 | 555.0 | 31.2 | 31.1 | 26.1 | 263.3 | 2,154.2 | 0.83 | - |
| 1981-1985 | 451.2 | 274.0 | 37.2 | 56.0 | 176.8 | 305.3 | 1,328.6 | 0.60 | - |
| 1986 | 638.9 | 320.6 | 79.3 | 70.0 | 332.0 | 1,549.1 | 3,195.4 | 0.37 | 0.44 |
| 1987 | 468.2 | 296.2 | 45.1 | 30.1 | 453.7 | 316.5 | 1,272.4 | 0.83 | 0.65 |
| 1988 | 844.7 | 297.2 | 61.1 | 56.8 | 119.3 | 670.9 | 1,918.9 | 0.69 | 0.66 |
| 1989 | 645.1 | 425.5 | 61.1 | 46.4 | 116.1 | 709.0 | 2,176.5 | 0.52 | 0.62 |
| 1990 | 275.9 | 357.1 | 28.7 | 22.5 | 46.9 | 196.7 | 987.4 | 0.78 | 0.73 |
| 1991 | 448.4 | 469.9 | 77.8 | 38.1 | 35.6 | 955.1 | 2,040.4 | 0.48 | 0.64 |
| 1992 | 67.4 | 256.5 | 51.0 | 44.2 | - | 216.1 | 629.6 | 0.51 | 0.63 |
| 1993 | 13.1 | 140.8 | 38.6 | 56.1 | - | 114.2 | 315.9 | 0.49 | 0.40 |
| 1994 | 2.7 | 3.0 | 28.2 | 48.5 | - | 169.2 | 267.5 | 0.02 | 0.06 |
| 1995 | 5.4 | 43.5 | 37.5 | 57.3 | - | 74.8 | 204.1 | 0.24 | 0.11 |
| 1996 | 7.0 | 31.8 | 45.7 | 79.3 | - | 113.0 | 260.3 | 0.15 | 0.06 |
| 1997 | 5.5 | 22.4 | 26.9 | 31.6 | - | 149.1 | 230.5 | 0.12 | 0.09 |
| 1998 | 3.5 | 12.8 | 29.4 | 34.3 | - | 168.4 | 270.8 | 0.06 | 0.08 |
| 1999 | 3.6 | 36.5 | 22.6 | 51.2 | - | 274.1 | 432.0 | 0.09 | 0.07 |
| 2000 | 25.2 | 74.6 | 33.2 | 81.1 | - | 548.2 | 762.4 | 0.13 | 0.04 |
| 2001 | 38.1 | 216.8 | 75.8 | 185.2 | - | 1,108.3 | 1,673.2 | 0.15 | 0.04 |
| 2002 | 15.0 | 118.7 | 54.0 | 269.0 | - | 499.9 | 972.2 | 0.14 | 0.05 |
| 2003 | 28.8 | 252.4 | 45.1 | 235.3 | - | 677.7 | 1,266.9 | 0.22 | 0.08 |
| 2004 | 26.2 | 159.3 | 38.1 | 197.2 | - | 442.6 | 904.5 | 0.21 | 0.08 |
| 2005 | 10.5 | 58.2 | 42.8 | 164.6 | - | 341.0 | 629.9 | 0.11 | 0.04 |
| 2006 | 4.5 | 47.5 | 29.6 | 132.7 | - | 386.2 | 674.1 | 0.08 | 0.08 |
| 2007 | 26.2 | 128.5 | 10.9 | 71.4 | - | 336.9 | 631.3 | 0.25 | 0.12 |
| 2008 | 0.6 | 26.4 | 16.0 | 180.1 | - | 494.3 | 769.8 | 0.04 | 0.02 |
| 2009 | 27.7 | 201.2 | 16.7 | 265.3 | - | 729.8 | 1,341.3 | 0.17 | 0.07 |
| 2010 | 5.8 | 48.8 | 19.6 | 286.5 | - | 440.4 | 848.4 | 0.06 | 0.04 |
| 2011 ^{g/} | 4.2 | 54.7 | 19.3 | 295.3 | - | 352.0 | 760.7 | 0.08 | 0.08 |

a/ The OPI area includes ocean and inside harvest impacts and escapement to streams and lakes south of Leadbetter Pt., Washington.

b/ Includes estimated nonretention mortality: troll fishery--hook-and-release mortality for 1982-2005 and drop-off mortality for all years; sport fishery--hook-and-release mortality for 1994-2005 and drop-off mortality for all years.

c/ Includes returns from Salmon-Trout Enhancement Program (STEP) smolt releases through the 2007 return year, after which the program was terminated.

d/ Includes Rogue River.

e/ FRAM post season runs used after 1985 and includes OPI origin stock catches in all fisheries.

f/ Ocean fishery impacts on private hatchery stock and returns to private hatcheries are excluded in calculating the OPI area stock aggregate ocean exploitation

g/ Preliminary.

TABLE III-3. Preseason forecasts and postseason estimates of ocean escapements for selected Washington coastal adult natural coho stocks in thousands of fish.

| Year | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season | Preseason Forecast | Postseason Return | Pre/Post- season |
|------------------------------|-----------------------|----------------------|---------------------|-----------------------|----------------------|---------------------|-----------------------|----------------------|---------------------|----------------------------------|----------------------|---------------------|
| Quillayute River Fall | | | | Hoh River | | | Queets River | | | Grays Harbor^{a/} | | |
| 1986 | 11.6 | 36.3 | 0.32 | 4.1 | 18.1 | 0.23 | 9.8 | 24.6 | 0.40 | 93.8 | 123.3 | 0.76 |
| 1987 | 27.3 | 33.8 | 0.81 | 13.0 | 14.2 | 0.91 | 20.6 | 15.9 | 1.29 | 218.6 | 66.3 | 3.30 |
| 1988 | 23.0 | 13.5 | 1.70 | 4.4 | 19.4 | 0.23 | 10.3 | 17.9 | 0.57 | 55.7 | 96.8 | 0.58 |
| 1989 | 28.2 | 18.8 | 1.50 | 11.0 | 9.2 | 1.19 | 13.6 | 12.0 | 1.13 | 82.3 | 156.5 | 0.53 |
| 1990 | 45.5 | 11.7 | 3.91 | 8.1 | 8.7 | 0.93 | 13.6 | 27.3 | 0.50 | 81.2 | 96.1 | 0.84 |
| 1991 | 16.3 | 26.4 | 0.62 | 6.3 | 11.6 | 0.55 | 16.1 | 26.6 | 0.60 | 244.6 | 139.1 | 1.76 |
| 1992 | 22.8 | 15.8 | 1.44 | 8.9 | 15.4 | 0.58 | 11.7 | 17.7 | 0.66 | 60.4 | 58.0 | 1.04 |
| 1993 | 13.2 | 10.5 | 1.26 | 8.3 | 3.4 | 2.47 | 12.9 | 12.7 | 1.01 | 144-153 | 58.5 | 2.46-2.62 |
| 1994 | 11.6 | 8.4 | 1.38 | 5.0 | 2.2 | 2.31 | 6.9 | 2.5 | 2.78 | 53.8-60.2 | 14.0 | 3.84-4.30 |
| 1995 | 13.1 | 19.8 | 0.66 | 6.8 | 9.7 | 0.70 | 12.1 | 10.7 | 1.13 | 103.4 | 70.2 | 1.47 |
| 1996 | 13.0 | 20.3 | 0.64 | 4.2 | 7.7 | 0.54 | 8.3 | 22.6 | 0.37 | 121.4 | 89.7 | 1.35 |
| 1997 | 8.9 | 5.8 | 1.53 | 2.8 | 4.1 | 0.68 | 4.3 | 2.2 | 1.92 | 26.1 | 20.2 | 1.29 |
| 1998 | 8.0 | 17.4 | 0.46 | 3.4 | 5.6 | 0.61 | 4.2 | 6.3 | 0.66 | 30.1 | 46.4 | 0.65 |
| 1999 | 14.5 | 16.1 | 0.90 | 3.2 | 6.8 | 0.47 | 4.3 | 8.6 | 0.50 | 57.7 | 42.7 | 1.35 |
| 2000 | 8.7 | 16.5 | 0.53 | 3.5 | 9.3 | 0.38 | 2.7 | 12.1 | 0.22 | 47.8 | 51.9 | 0.92 |
| 2001 | 23.0 | 28.4 | 0.81 | 8.5 | 16.2 | 0.52 | 12.0 | 35.8 | 0.33 | 51.3 | 103.2 | 0.50 |
| 2002 | 22.3 | 33.2 | 0.67 | 8.5 | 13.2 | 0.64 | 12.5 | 26.3 | 0.47 | 55.4 | 142.0 | 0.39 |
| 2003 | 24.9 | 22.5 | 1.11 | 12.5 | 8.7 | 1.44 | 24.0 | 15.7 | 1.52 | 58.0 | 108.4 | 0.54 |
| 2004 | 21.2 | 20.7 | 1.02 | 8.1 | 6.9 | 1.17 | 18.5 | 13.3 | 1.39 | 117.9 | 90.8 | 1.30 |
| 2005 | 18.6 | 20.9 | 0.89 | 7.6 | 8.2 | 0.93 | 17.1 | 11.9 | 1.43 | 91.1 | 65.9 | 1.38 |
| 2006 | 14.6 | 9.9 | 1.48 | 6.4 | 2.7 | 2.36 | 8.3 | 9.2 | 0.90 | 67.3 | 30.6 | 2.20 |
| 2007 | 10.8 | 10.7 | 1.01 | 5.4 | 5.8 | 0.93 | 13.6 | 7.1 | 1.92 | 59.4 | 34.6 | 1.72 |
| 2008 | 10.5 | 11.1 | 0.95 | 4.3 | 4.3 | 1.00 | 10.2 | 7.4 | 1.39 | 42.7 | 49.0 | 0.87 |
| 2009 | 19.3 | 15.5 | 1.24 | 9.5 | 9.5 | 1.00 | 31.4 | 16.0 | 1.97 | 59.2 | 104.6 | 0.57 |
| 2010 | 22.0 | 17.0 | 1.29 | 7.6 | 11.6 | 0.65 | 21.8 | 19.1 | 1.14 | 67.9 | 130.7 | 0.52 |
| 2011 | 28.2 | NA | NA | 11.6 | NA | NA | 13.3 | NA | NA | 89.1 | NA | NA |
| 2012 | 33.5 | - | - | 14.3 | - | - | 37.2 | - | - | 150.2 | - | - |

a/ Coho FRAM was used to estimate post season ocean abundance.

TABLE III-4. Preseason forecasts and postseason estimates of ocean escapements for selected Puget Sound adult natural coho stocks in thousands of fish. (Page 1 of 2)

| Year | Preseason Forecast | Postseason Return | Pre/Postseason | Preseason Forecast | Postseason Return | Pre/Postseason | Preseason Forecast | Postseason Return | Pre/Postseason |
|--------------------|-----------------------|----------------------|----------------|----------------------------|----------------------|----------------|-----------------------|----------------------|----------------|
| | Skagit River | | | Stilliguamish River | | | Hood Canal | | |
| 1986 | 76.8 | 69.7 | 1.10 | NA | 49.9 | NA | 110.8 | 82.2 | 1.35 |
| 1987 | 70.5 | 39.4 | 1.79 | NA | 46.3 | NA | 96.5 | 71.7 | 1.35 |
| 1988 | 81.8 | 28.4 | 2.88 | NA | 35.4 | NA | 39.6 | 15.5 | 2.55 |
| 1989 | 80.3 | 24.4 | 3.29 | NA | 13.5 | NA | 77.4 | 25.5 | 3.04 |
| 1990 | 98.9 | 24.3 | 4.07 | 75.8 | 34.1 | 2.22 | 94.2 | 14.2 | 6.63 |
| 1991 | 95.3 | 10.3 | 9.25 | 71.5 | 11.3 | 6.33 | 38.1 | 15.3 | 2.49 |
| 1992 | 80.1 | 9.4 | 8.52 | 42.4 | 18.0 | 2.36 | 23.2 | 19.9 | 1.17 |
| 1993 | 70.7 | 14.2 | 4.98 | 61.8 | 10.6 | 5.83 | 89.6 | 16.7 | 5.37 |
| 1994 | 39.0 | 30.3 | 1.29 | 21.9 | 30.3 | 0.72 | 25.4 | 57.0 | 0.45 |
| 1995 | 64.7 | 15.8 | 4.09 | 70.3 | 20.4 | 3.45 | 36.4 | 41.1 | 0.89 |
| 1996 | 44.8 | 8.6 | 5.19 | 51.6 | 12.5 | 4.13 | 25.1 | 37.2 | 0.67 |
| 1997 | 70.9 | 45.7 | 1.55 | 36.0 | 14.1 | 2.56 | 78.4 | 101.8 | 0.77 |
| 1998 | 55.0 | 85.2 | 0.65 | 47.8 | 31.1 | 1.54 | 108.0 | 118.5 | 0.91 |
| 1999 | 75.7 | 38.3 | 1.98 | 35.7 | 7.5 | 4.77 | 65.1 | 17.6 | 3.70 |
| 2000 | 30.2 | 75.1 | 0.40 | 17.7 | 31.2 | 0.57 | 61.0 | 39.7 | 1.54 |
| 2001 | 87.2 | 115.6 | 0.75 | 24.4 | 81.8 | 0.30 | 62.0 | 110.0 | 0.56 |
| 2002 | 98.5 | 70.8 | 1.39 | 19.7 | 30.4 | 0.65 | 34.9 | 81.0 | 0.43 |
| 2003 | 116.6 | 114.4 | 1.02 | 37.8 | 49.8 | 0.76 | 33.4 | 199.9 | 0.17 |
| 2004 | 155.8 | 151.0 | 1.03 | 38.0 | 73.9 | 0.51 | 98.7 | 219.7 | 0.45 |
| 2005 | 61.8 | 53.1 | 1.16 | 56.7 | 29.1 | 1.95 | 98.4 | 68.3 | 1.44 |
| 2006 | 106.6 | 12.8 | 8.33 | 45.0 | 11.8 | 3.81 | 59.4 | 49.7 | 1.20 |
| 2007 ^{b/} | 26.8 | 71.2 | 0.38 | 69.2 | 45.2 | 1.53 | 42.4 | 78.6 | 0.54 |
| 2008 ^{b/} | 61.4 | 32.1 | 1.91 | 31.0 | 15.3 | 2.03 | 30.4 | 25.8 | 1.18 |
| 2009 ^{b/} | 33.4 | 72.7 | 0.46 | 13.4 | 27.4 | 0.49 | 48.6 | 45.7 | 1.06 |
| 2010 | 95.9 | NA | - | 25.9 | NA | - | 33.2 | NA | - |
| 2011 | 138.1 | NA | - | 66.6 | NA | - | 74.7 | NA | - |
| 2012 | 48.3 | - | - | 47.5 | - | - | 73.4 | - | - |

TABLE III-4. Preseason and postseason estimates of ocean escapements^{a/} for selected Puget Sound adult natural coho stocks in thousands of fish. (Page 2 of 2)

| Year | Preseason Forecast | Postseason Return | Pre/Postseason | Preseason Forecast | Postseason Return | Pre/Postseason |
|--------------------|-----------------------|----------------------|-------------------------------|-----------------------|----------------------|----------------|
| Snohomish | | | Strait of Juan de Fuca | | | |
| 1986 | NA | 49.9 | - | 24.7 | 48.9 | 0.51 |
| 1987 | NA | 46.3 | - | 17.8 | 23.9 | 0.74 |
| 1988 | NA | 35.4 | - | 19.5 | 25.6 | 0.76 |
| 1989 | NA | 13.5 | - | 17.0 | 28.7 | 0.59 |
| 1990 | 308.8 | 276.5 | 1.12 | 25.8 | 28.5 | 0.91 |
| 1991 | 308.8 | 163.4 | 1.89 | 24.1 | 21.5 | 1.12 |
| 1992 | 389.7 | 192.5 | 2.02 | 25.7 | 27.8 | 0.93 |
| 1993 | 394.4 | 142.3 | 2.77 | 20.8 | 11.5 | 1.81 |
| 1994 | 256.7 | 293.6 | 0.87 | 20.8 | 11.3 | 1.84 |
| 1995 | 358.3 | 211.3 | 1.70 | 11.4 | 22.6 | 0.51 |
| 1996 | 338.1 | 132.3 | 2.55 | 10.7 | 19.1 | 0.56 |
| 1997 | 186.6 | 106.4 | 1.75 | 6.5 | 20.1 | 0.32 |
| 1998 | 165.3 | 193.9 | 0.85 | 16.8 | 20.9 | 0.80 |
| 1999 | 141.6 | 82.2 | 1.72 | 14.7 | 9.9 | 1.49 |
| 2000 | 53.0 | 154.6 | 0.34 | 13.5 | 28.5 | 0.47 |
| 2001 | 129.6 | 360.1 | 0.36 | 21.4 | 43.8 | 0.49 |
| 2002 | 123.1 | 185.5 | 0.66 | 21.3 | 26.3 | 0.81 |
| 2003 | 203.0 | 198.0 | 1.03 | 25.6 | 22.9 | 1.12 |
| 2004 | 192.1 | 287.9 | 0.67 | 35.7 | 23.7 | 1.51 |
| 2005 | 241.6 | 133.4 | 1.81 | 20.7 | 12.5 | 1.66 |
| 2006 | 139.5 | 94.2 | 1.48 | 26.1 | 4.6 | 5.67 |
| 2007 ^{b/} | 98.9 | 156.4 | 0.63 | 29.9 | 10.2 | 2.94 |
| 2008 ^{b/} | 92.0 | 49.5 | 1.86 | 24.1 | 3.8 | 6.27 |
| 2009 ^{b/} | 67.0 | 133.4 | 0.50 | 20.5 | 24.6 | 0.83 |
| 2010 | 99.4 | 53.8 | 1.85 | 8.5 | 21.5 | 0.40 |
| 2011 | 180.0 | NA | - | 12.3 | NA | - |
| 2012 | 109.0 | - | - | 12.6 | - | - |

a/ Coho FRAM was used to estimate post season ocean abundance.

b/ Preliminary postseason return.

TABLE III-5. Status categories and constraints for Puget Sound and Washington Coast coho under the FMP and PST Southern Coho Management Plan.

| FMP | | |
|------------------------|--|----------------------------------|
| FMP Stock | Total Exploitation Rate Constraint ^{a/} | Categorical Status ^{a/} |
| Skagit | 35% | low |
| Stillaguamish | 50% | normal |
| Snohomish | 40% | low |
| Hood Canal | 65% | normal |
| Strait of Juan de Fuca | 40% | low |
| Quillayute Fall | Undefined | |
| Hoh | 65% | |
| Queets | 65% | |
| Grays Harbor | 65% | |

| PST Southern Coho Management Plan | | |
|--|--|----------------------------------|
| U.S. Management Unit | Total Exploitation Rate Constraint ^{b/} | Categorical Status ^{c/} |
| Skagit | 35% | Moderate |
| Stillaguamish | 50% | Abundant |
| Snohomish | 40% | Moderate |
| Hood Canal | 65% | Abundant |
| Strait of Juan de Fuca | 40% | Moderate |
| Quillayute Fall ^{c/} | 65% | Abundant |
| Hoh ^{c/} | 65% | Abundant |
| Queets ^{c/} | 65% | Abundant |
| Grays Harbor | 65% | Abundant |

a/ Preliminary. For Puget Sound stocks, the exploitation rate constraints and categorical status (normal, low, critical) reflect application of Comprehensive Coho Agreement rules, as adopted in the FMP. For Washington Coast stocks, exploitation rate constraints represent MFMT. Note that under *U.S. v. Washington* and *Hoh v. Baldrige* case law, the management objectives can differ from FMP objectives provided there is an annual agreement among the state and tribal comanagers; therefore, the exploitation rates used to report categorical status do not necessarily represent maximum allowable rates for these stocks.

b/ Preliminary. For Puget Sound and Washington Coast management units, the exploitation rate constraints reflect application of the 2002 PST Southern Coho Management Plan.

c/ Categories (abundant, moderate, low) correspond to the general exploitation rate ranges depicted in paragraph 3(a) of the 2002 PST Southern Coho Management Plan. For Washington Coast stocks, categorical status is determined by taking the midpoint of the range of exploitation rates associated with achieving the escapement goal ranges. The exploitation rate ranges are based on preseason abundance forecasts and the upper and lower ends of the escapement goal ranges. Maximum exploitation rates are computed using the lower end of the escapement range; minimum exploitation rates are computed using the upper end of the escapement range.

TABLE III-6. Projected coho mark rates for 2012 fisheries under base period fishing patterns (percent marked).

| Area | Fishery | June | July | August | Sept |
|----------------------------------|--------------|------|------|--------|------|
| Canada | | | | | |
| Johnstone Strait | Recreational | - | 34% | 35% | - |
| West Coast Vancouver Island | Recreational | 39% | 21% | 14% | 13% |
| North Georgia Strait | Recreational | 55% | 55% | 55% | 49% |
| South Georgia Strait | Recreational | 52% | 57% | 50% | 53% |
| Juan de Fuca Strait | Recreational | 49% | 50% | 47% | 46% |
| Johnstone Strait | Troll | 63% | 52% | 38% | 49% |
| NW Vancouver Island | Troll | 30% | 27% | 27% | 28% |
| SW Vancouver Island | Troll | 43% | 39% | 40% | 42% |
| Georgia Strait | Troll | 62% | 62% | 62% | 58% |
| Puget Sound | | | | | |
| Strait of Juan de Fuca (Area 5) | Recreational | 50% | 49% | 47% | 48% |
| Strait of Juan de Fuca (Area 6) | Recreational | 51% | 48% | 48% | 46% |
| San Juan Island (Area 7) | Recreational | 54% | 55% | 57% | 42% |
| North Puget Sound (Areas 6 & 7A) | Net | - | 45% | 53% | 51% |
| Council Area | | | | | |
| Neah Bay (Area 4/4B) | Recreational | 42% | 49% | 47% | 51% |
| LaPush (Area 3) | Recreational | 50% | 48% | 50% | 39% |
| Westport (Area 2) | Recreational | 54% | 53% | 48% | 40% |
| Columbia River (Area 1) | Recreational | 62% | 59% | 54% | 57% |
| Tillamook | Recreational | 49% | 44% | 37% | 21% |
| Newport | Recreational | 44% | 38% | 35% | 21% |
| Coos Bay | Recreational | 31% | 27% | 18% | 8% |
| Brookings | Recreational | 24% | 16% | 14% | 7% |
| Neah Bay (Area 4/4B) | Troll | 47% | 47% | 46% | 41% |
| LaPush (Area 3) | Troll | 44% | 49% | 44% | 44% |
| Westport (Area 2) | Troll | 42% | 46% | 47% | 45% |
| Columbia River (Area 1) | Troll | 53% | 53% | 47% | 50% |
| Tillamook | Troll | 47% | 43% | 42% | 38% |
| Newport | Troll | 42% | 40% | 35% | 32% |
| Coos Bay | Troll | 30% | 28% | 23% | 13% |
| Brookings | Troll | 19% | 22% | 24% | 38% |
| Columbia River | | | | | |
| Buoy 10 | Recreational | - | - | - | 59% |




FIGURE III-1a. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.




FIGURE III-1b. Selected preseason vs. postseason forecasts for coho stocks with significant contribution to Council area fisheries.

CHAPTER IV: AFFECTED ENVIRONMENT - PINK SALMON ASSESSMENT

Two major runs comprise the pink salmon population available to Council fisheries during odd-numbered years: the Fraser River (British Columbia) run, which is more abundant, and the Puget Sound run. The 2011 run size forecast for Fraser pinks was 17.50 million fish; actual run size was estimated at 18.3 million. The 2011 Puget Sound pink salmon run size forecast was 5.98 million, with 5.97 million natural and 4,100 hatchery fish. The actual run size estimate for 2011 was not available.

Table IV-1 provides a summary of recent run sizes and forecasts.

TABLE IV-1. Estimated annual (odd-numbered years) run sizes and forecasts for Fraser River and Puget Sound pink salmon in millions of fish.

| Year | Puget Sound | | Fraser River ^{a/} | |
|--------------------|-------------|--------|----------------------------|--------|
| | Forecast | Actual | Forecast | Actual |
| 1977 | NA | 0.88 | NA | 8.21 |
| 1979 | NA | 1.32 | NA | 14.40 |
| 1981 | NA | 0.50 | NA | 18.69 |
| 1983 | NA | 1.01 | NA | 15.35 |
| 1985 | NA | 1.76 | NA | 19.10 |
| 1987 | NA | 1.57 | NA | 7.17 |
| 1989 | NA | 1.93 | NA | 16.63 |
| 1991 | NA | 1.09 | NA | 22.18 |
| 1993 | NA | 1.06 | NA | 16.98 |
| 1995 | 3.4 | 2.08 | NA | 12.90 |
| 1997 | NA | 0.44 | 11.40 | 8.18 |
| 1999 | NA | 0.96 | NA | 3.59 |
| 2001 | 2.92 | 3.56 | 5.47 | 21.17 |
| 2003 | 2.32 | 2.90 | 17.30 | 26.00 |
| 2005 | 1.98 | 1.23 | 16.30 | 10.00 |
| 2007 | 3.34 | 2.45 | 19.60 | 11.00 |
| 2009 | 5.16 | 9.84 | 17.54 | 19.50 |
| 2011 ^{b/} | 5.98 | NA | 17.50 | 18.30 |

a/ Total run size.

b/ Preliminary forecast.

CHAPTER V: DESCRIPTION AND ANALYSIS OF THE NO ACTION ALTERNATIVE

DESCRIPTION OF THE NO-ACTION ALTERNATIVE

The No-Action Alternative consists of the preseason management measures adopted by the Council and approved by the Secretary of Commerce for the 2011 ocean salmon management season between the U.S./Canada border and the U.S./Mexico border. The management measures relate to three fishery sectors: non-Indian commercial (Table V-1), recreational (Table V-2), and treaty Indian (Table V-3). A description of the 2011 preseason management measures and analyses of their projected effects on the biological and socioeconomic environment are presented in Preseason Report III (PFMC 2011b). A description of the 2011 management measures as implemented, including inseason modifications, and an analysis of their effects on the environment, including an historical perspective, is presented in the SAFE document - Review of 2011 Ocean Salmon Fisheries (PFMC 2012).

ANALYSIS OF EFFECTS ON THE ENVIRONMENT OF THE NO-ACTION ALTERNATIVE

Overview

Table V-4 provides a summary of Salmon FMP stock spawning escapement and exploitation rate projections for 2012 under the No-Action Alternative (2011 regulations), as well as postseason estimates of these quantities for earlier years, which are compared to FMP conservation objectives. For some stocks, postseason estimates of these metrics were either incomplete or unavailable when the Review of 2011 Ocean Salmon Fisheries was published. A preliminary determination of stock status under the FMP SDC was available for some of these stocks in time for this report; however, some estimates are still unavailable. The STT will report to the Council on stocks status at the March 2012 Council meeting, and may further update the status of stocks present in Table V-4 at that time.

Chinook escapements and fishery impacts were estimated using the Sacramento Harvest Model or Klamath Ocean Harvest Model for SRFC and KRFC, respectively. Assessment of effects under the No-Action Alternative for Oregon Coast Chinook are not available; for Columbia River Chinook stocks assessments were based on qualitative assessment of the magnitude of forecasts, if available, in relation to escapement goals.

Coho escapements and fishery impacts were estimated using Coho FRAM. Abundance forecasts for 2012 were updated for Washington and Oregon stocks, but forecasts for Canadian stocks are unchanged from those employed for 2011 planning. Updated forecasts for Canadian stocks are expected to become available in March 2012. To provide information on the effect of changes in abundance forecasts, the final 2011 pre-season regulatory package for ocean and inside fisheries was applied to 2012 projections of abundance.

Sacramento River Fall Chinook

A repeat of 2011 regulations would be expected to result in an escapement of 469,000 natural and hatchery SRFC adults, which is well above the 122,000 to 180,000 natural and hatchery adult escapement goal range, and exceeds the 2012 S_{ACL} of 245,820.

The geometric mean of the 2010 and 2011 spawning escapement estimates and the 2012 forecast spawning escapement under the No-Action Alternative is greater than S_{MSY} (Table V-4), which indicates that SRFC would meet the default rebuilt criterion if the forecast escapement was realized. The minimum 2012 spawning escapement necessary to reach that level would be 127,349.

Klamath River Fall Chinook

A repeat of 2011 fishery regulations, which included a river recreational harvest quota of 7,900 adults and a tribal allocation of 50 percent (of the overall adult harvest), would be expected to result in 179,000 natural area adult spawners. This projection exceeds the S_{MSY} of 40,700 natural area adults and the 2012 S_{ACL} of 86,288. If the ocean fisheries were closed from January through August 2012 between Cape Falcon and Point Sur, and the Klamath River fisheries (tribal and recreational) were closed in 2012, the expected number of natural area adult spawners would be 269,600.

California Coastal Chinook Stocks

The NMFS ESA consultation standard restricts the Klamath River fall Chinook age-4 ocean harvest rate to no more than 16.0 percent to limit impacts on these stocks. As indicated in the Chapter II, the postseason estimate of this rate for 2011 is 7.8 percent. Applying 2011 regulations to the 2012 abundance results in an age-4 ocean harvest rate forecast of 13.2 percent. If the ocean fisheries were closed from January through August 2012 between Cape Falcon and Point Sur, the expected age-4 ocean harvest rate for 2012 would be 0.1 percent (70 age-4 KRFC were harvested during the September through November 2011 period).

Oregon Coast Chinook Stocks

The FMP conservation objective for Oregon coast Chinook is 150,000 to 200,000 natural adult spawners, and attainment of this goal is assessed using peak spawner counts of 60 to 90 fish per mile observed in standard index reaches. In 2009, 2010, and 2011 the goal was achieved with 62, 79, and 78 fish per mile, respectively. No forecast is available for this stock, but given recent trends, it seems likely that it would meet its goal again in 2012 under 2011 fishing seasons.

Columbia River Chinook Stocks

The 2012 forecasts are lower than the 2011 forecast for all stocks except for LRW and summer Chinook, although the 2012 forecasts are higher than the 2011 actual returns for all stocks except SCH. Applying 2011 regulations to the forecasted 2012 abundance of Columbia River Chinook would result in ocean escapements meeting spawning escapement goals for all major fall Chinook stocks, including SCH, and summer Chinook (Table V-4).

Washington Coast and Puget Sound Chinook Stocks

Council fisheries north of Cape Falcon have only a minor impact on most stocks that originate in Washington coastal and Puget Sound rivers. These stocks have northerly marine distribution patterns, and are therefore impacted primarily by Canadian and Alaskan fisheries. An evaluation of 2011 Council area management measures on projected 2012 abundance would not provide a useful comparison of fishery impacts in relation to conservation objectives.

Oregon Production Index Area Coho Stocks

Ocean fisheries were modeled with 2011 Council regulations and 2011 expectations for non-Council area fisheries. Under this scenario, expected exploitation rates are 12.2 percent on OCN coho and 6.9 percent on Rogue/Klamath hatchery coho. Expected spawner escapement is 256,559 for OCN coho (Tables V-5 and V-6). For Columbia River hatchery coho stocks, the predicted ocean exploitation rate (excluding Buoy 10) is 22.2 percent on the Columbia River early stock and 35.7 percent on the Columbia River late stock. Predicted ocean escapements (after Buoy 10) into the Columbia River in 2012 under this exercise show that under 2011 ocean regulations, Columbia River early and late coho would be expected to meet egg take goals.

As noted in Chapter III, the total allowable OCN coho exploitation rate for 2011 fisheries is no greater than 15 percent under FMP Amendment 13 and no greater than 15 percent under the matrix developed by the OCN work group (Table V-7; Appendix A, Tables A-2 and A-3), and the total allowable RK hatchery coho marine exploitation rate is 13.0 percent (NMFS ESA consultation standard). Under 2011 fishery regulations and 2012 abundance forecasts, these exploitation rates are predicted to be 12.2 percent for OCN, and 6.9 percent for RK coho. The allowable LCN coho exploitation rate is 15.0 percent in marine area and mainstem Columbia River fisheries combined. Under 2011 fishery regulations and 2012 abundance forecasts, the exploitation rate is predicted to be 11.2 percent for marine fisheries (excluding the Buoy 10 fishery) using combined unmarked Columbia River hatchery stocks as the proxy. Given the 2011 inriver sharing arrangement, the total exploitation rate on LCN coho would be 15.2 percent.

Washington Coast, Puget Sound, and Canadian Coho Stocks

Exploitation rate and ocean escapement expectations in relation to management goals for selected naturally-spawning coho stocks, given 2012 preseason abundance forecasts and 2011 preseason projections for fishing patterns, are presented in Table V-5. The 2012 forecasts for Canadian coho stocks are not available, but are assumed to be at 2011 levels for this analysis. More detailed fishery management goals for Council area coho stocks are listed in Appendix A.

Under 2011 regulations, 2012 exploitation rates are expected to meet the allowable 2012 FMP conservation objectives for Puget Sound coho stocks with the exception of Skagit natural coho. Skagit coho would have a 38.1 percent exploitation rate with a conservation objective of 35 percent. Ocean escapements for Washington Coast natural coho stocks are expected to be at levels that would permit attainment of FMP spawning escapement conservation objectives. In addition, all annual management objectives for U.S. stocks subject to the PSC agreement would be met. The exploitation rate by U.S. fisheries south of the Canadian border on Interior Fraser (B.C.) coho is projected to be 10.3 percent, which is slightly over the anticipated 10.0 percent allowable exploitation rate under the 2002 PST Coho Agreement. The Council area fisheries portion would be 4.3 percent.

Coho bycatch during Puget Sound fisheries directed at chum and sockeye salmon will also be a consideration for preseason planning.

Summary

The effects of projected impacts (where available) under 2011 fishery regulations and 2012 abundance forecasts are as follows :

- All stocks would achieve S_{MSY} spawning escapement objectives.
- SRFC and KRFC would comply with 2012 preseason ACL requirements.
- All stocks would have projected exploitation rates less than MFMT or ESA consultation standards except LCN coho.
- All Puget Sound coho would have exploitation rates less than the annual rates allowed under the FMP harvest rate matrix and the PST 2002 Southern Coho Management Plan except Skagit natural coho and Interior Fraser (B.C.) natural coho.
- All Washington Coast coho would have exploitation rates less than the annual rates allowed under the PST 2002 Southern Coho Management Plan.
- No stocks would be approaching an overfished condition.
- SRFC would be projected to meet the FMP default rebuilt criterion of a 3-year geometric mean spawning escapement greater than S_{MSY} .

Conclusion

The No-Action Alternative would not meet the Purpose and Need for the proposed action because the 2012 ESA consultation standard of no more than 15.0 percent exploitation rate on LCN coho in marine and Columbia River mainstem fisheries would not be satisfied, Skagit coho would exceed the exploitation rate limit in the FMP, and the Southern U.S. exploitation rate limit on Interior Fraser coho would exceed the limit specified in the PST 2002 Southern Coho Management Plan. In addition, recreational opportunity and commercial value would not be optimized because surplus production of KRFC and SRFC would be forgone as a result of unnecessarily conservative management measures south of Cape Falcon.

The No-Action Alternative does not reflect consideration of changes in the status of salmon stocks from the previous year; therefore, over- or under- harvest of some salmon stocks would occur if this alternative was implemented. The analysis of the No-Action Alternative does, however, provide perspective that is useful in the planning process for 2012 ocean salmon fishery management measures. An understanding of stock shortfalls and surpluses under the No-Action Alternative helps managers, advisors, and constituents construct viable alternatives to the status-quo management measures.

| |
|--|
| TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011. (Page 1 of 5) |
| A. SEASON DESCRIPTIONS |
| North of Cape Falcon |
| Supplemental Management Information |
| <p>1. Overall non-Indian TAC: 64,600 (non-mark-selective equivalent of 61,800) Chinook and 80,000 coho marked with a healed adipose fin clip (marked).</p> <p>2. Non-Indian commercial troll TAC: 30,900 Chinook and 12,800 marked coho.</p> <p>3. No preseason trade of Chinook or coho between non-Indian commercial and recreational fisheries.</p> |
| <p>U.S./Canada Border to Cape Falcon</p> <ul style="list-style-type: none"> May 1 through earlier of June 30 or 20,600 Chinook quota. <p>Seven days per week (C.1). All salmon except coho (C.7). Cape Flattery, Mandatory Yelloweye Rockfish Conservation Area, and Columbia Control Zones closed (C.5). See gear restrictions and definitions (C.2, C.3). An inseason conference call will occur when it is projected that 13,700 Chinook have been landed to consider modifying the open period to five days per week and adding landing and possession limits to ensure the guideline is not exceeded.</p> |
| <p>U.S./Canada Border to Cape Falcon</p> <ul style="list-style-type: none"> July 1 through earlier of September 15 or 10,300 preseason Chinook guideline (C.8) or a 12,800 marked coho quota (C.8.d). <p>Friday through Tuesday; landing and possession limit of 50 Chinook and 50 coho per vessel per open period north of Leadbetter Point or 50 Chinook and 50 coho south of Leadbetter Point (C.1). All Salmon except no chum retention north of Cape Alava, Washington in August and September (C.7). All coho must be marked (C.8.d). See gear restrictions and definitions (C.2, C.3). Cape Flattery, Mandatory Yelloweye Rockfish Conservation Area, and Columbia Control Zones closed; Grays Harbor Control Zone closed in August and September (C.5).</p> |
| <p>Vessels must land and deliver their fish within 24 hours of any closure of this fishery. Under state law, vessels must report their catch on a state fish receiving ticket. Vessels fishing or in possession of salmon while fishing north of Leadbetter Point must land and deliver their fish within the area and north of Leadbetter Point. Vessels fishing or in possession of salmon while fishing south of Leadbetter Point must land and deliver their fish within the area and south of Leadbetter Point, except that Oregon permitted vessels may also land their fish in Garibaldi, Oregon. Oregon State regulations require all fishers landing salmon into Oregon from any fishery between Leadbetter Point, Washington and Cape Falcon, Oregon must notify ODFW within one hour of delivery or prior to transport away from the port of landing by either calling 541-867-0300 Ext. 271 or sending notification via e-mail to nfalcon.trollreport@state.or.us. Notification shall include vessel name and number, number of salmon by species, port of landing and location of delivery, and estimated time of delivery. Inseason actions may modify harvest guidelines in later fisheries to achieve or prevent exceeding the overall allowable troll harvest impacts (C.8).</p> |

TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 2 of 5)

| A. SEASON DESCRIPTIONS | |
|--|--|
| South of Cape Falcon | |
| Supplemental Management Information | |
| <ol style="list-style-type: none"> 1. Sacramento River Basin recreational fishery catch assumption: 61,400 adult Sacramento River fall Chinook. 2. Sacramento River fall Chinook spawning escapement of 377,000 adults. 3. Klamath River recreational fishery allocation: 7,900 adult Klamath River fall Chinook. 4. Klamath tribal allocation: 34,800 adult Klamath River fall Chinook. | |
| <p>Cape Falcon to Humbug Mt.</p> <ul style="list-style-type: none"> • April 15 through July 9, July 17 through August 31, October 1-31. (C.9). <p>Seven days per week. All salmon except coho; landing and possession limit of 50 Chinook per vessel per calendar week in October (C.7). All vessels fishing in the area must land their fish in the State of Oregon. See gear restrictions and definitions (C.2, C.3) and Oregon State regulations for a description of special regulations at the mouth of Tillamook Bay.</p> <p>In 2012, the season will open March 15 for all salmon except coho. This opening could be modified following Council review at its March 2012 meeting.</p> | |
| <p>Humbug Mt. to OR/CA Border (Oregon KMZ)</p> <ul style="list-style-type: none"> • May 1-31; • June 1 through earlier of June 30, or a 1,500 Chinook quota; • July 1 through earlier of July 31, or a 1,200 Chinook quota; • Aug. 1 through earlier of Aug. 31, or a 1,000 Chinook quota (C.9). <p>Seven days per week. All salmon except coho (C.7). Chinook 28 inch total length minimum size limit (B). June 1 through August 31, landing and possession limit of 30 Chinook per vessel per day. Any remaining portion of the June and/or July Chinook quotas may be transferred inseason on an impact neutral basis to the next open quota period (C.8). All vessels fishing in this area must land and deliver all fish within this area or Port Orford, within 24 hours of any closure in this fishery, and prior to fishing outside of this area (C.1, C.6). Oregon State regulations require all fishers landing salmon from any quota managed season within this area to notify Oregon Dept. of Fish and Wildlife (ODFW) within 1 hour of delivery or prior to transport away from the port of landing by either calling (541) 867-0300 ext. 252 or sending notification via e-mail to KMZOR.trollreport@state.or.us. Notification shall include vessel name and number, number of salmon by species, port of landing and location of delivery, and estimated time of delivery. See gear restrictions and definitions (C.2, C.3).</p> <p>In 2012, the season will open March 15 for all salmon except coho, with a 28 inch Chinook minimum size limit. This opening could be modified following Council review at its March 2012 meeting.</p> | |
| <p>OR/CA Border to Humboldt South Jetty (California KMZ)</p> <ul style="list-style-type: none"> • July 2 through the earlier of July 20 or a 1,400 Chinook quota, Saturday to Wednesday; • Aug. 1 through earlier of Aug. 15 or a 1,000 Chinook quota, seven days per week (C.9). <p>All salmon except coho (C.7). Chinook 27 inch total length minimum size limit (B). Landing and possession limit of 15 Chinook per vessel. Any remaining portion of the July Chinook quota may be transferred inseason on an impact neutral basis to the August quota (C.8). All vessels fishing in this area must land and deliver all fish within this area, within 24 hours of any closure in this fishery, and prior to fishing outside of this area (C.1, C.6). See compliance requirements (C.1) and gear restrictions and definitions (C.2, C.3). Klamath Control Zone closed (C.5.e). See California State regulations for additional closures adjacent to the Smith and Klamath rivers. When the fishery is closed between the OR/CA border and Humbug Mt. and open to the south, vessels with fish on board caught in the open area off California may seek temporary mooring in Brookings, Oregon prior to landing in California only if such vessels first notify the Chetco River Coast Guard Station via VHF channel 22A between the hours of 0500 and 2200 and provide the vessel name, number of fish on board, and estimated time of arrival.</p> | |
| <p>Humboldt South Jetty to Horse Mt. Closed.</p> | |
| <p>California State regulations require all salmon be made available to a CDFG representative for sampling immediately at port of landing. Any person in possession of a salmon with a missing adipose fin, upon request by an authorized agent or employee of the CDFG, shall immediately relinquish the head of the salmon to the state. (California Fish and Game Code §8226)</p> | |

TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 3 of 5)

| A. SEASON DESCRIPTIONS | | | | | |
|--|--|--|--|--|--|
| South of Cape Falcon | | | | | |
| Horse Mt. to Point Arena (Fort Bragg) <ul style="list-style-type: none"> July 23-27; July 29 through Aug. 29; Sept. 1-30 (C.9). <p>Seven days per week. All salmon except coho (C.7). Chinook 27 inch total length minimum size limit (B). All fish caught in the area when the KMZ quota fisheries are open must be landed south of Horse Mt.; all fish must be landed in California and offloaded within 24 hours of the August 29 closure (C.1, C.6). See gear restrictions and definitions (C.2, C.3).</p> | | | | | |
| Pt. Arena to Pigeon Pt. (San Francisco) <ul style="list-style-type: none"> May 1-31 seven days per week June 25 through July 5 seven days per week July 9-27 Saturday through Wednesday July 29 through Aug. 29 seven days per week September 1-30 seven days per week (C.9). <p>All salmon except coho (C.7). Chinook minimum size limit of 27 inches total length (B). All fish must be landed in California and offloaded within 24 hours of the August 29 closure. All fish caught in the area when the KMZ quota fisheries are open must be landed south of Horse Mt. (C.1, C.6). See gear restrictions and definitions (C.2, C.3).</p> <p>Pt. Reyes to Pt. San Pedro (Fall Area Target Zone)</p> <ul style="list-style-type: none"> October 3-14 <p>Monday through Friday. All salmon except coho (C.1). Chinook minimum size limit 27 inches total length (B). All vessels fishing in this area must land and deliver all fish between Point Arena and Pigeon Point (C.1, C.6). See gear restrictions and definitions (C.2, C.3).</p> | | | | | |
| Pigeon Pt. to Pt. Sur (Monterey) Same as Pt. Arena to Pigeon Pt. | | | | | |
| Pt. Sur to U.S./Mexico Border (Monterey south) <ul style="list-style-type: none"> May 1 through July 5 seven days per week July 9-27 Saturday through Wednesday July 29 through Aug. 29 seven days per week (C.9). <p>All salmon except coho (C.7). Chinook minimum size limit of 27 inches total length (B). All fish must be landed in California and offloaded within 24 hours of the August 29 closure; all fish caught in the area June 1-24 must be landed south of Pt. San Pedro; all fish caught in the area when the KMZ quota fisheries are open must be landed south of Horse Mt. (C.1, C.6). See gear restrictions and definitions (C.2, C.3).</p> | | | | | |
| California State regulations require all salmon be made available to a CDFG representative for sampling immediately at port of landing. Any person in possession of a salmon with a missing adipose fin, upon request by an authorized agent or employee of the CDFG, shall immediately relinquish the head of the salmon to the state. (California Fish and Game Code §8226) | | | | | |
| B. MINIMUM SIZE (Inches) (See C.1) | | | | | |

| Area (when open) | Chinook | | Coho | | Pink |
|------------------------------------|--------------|----------|--------------|----------|------|
| | Total Length | Head-off | Total Length | Head-off | |
| North of Cape Falcon | 28.0 | 21.5 | 16.0 | 12.0 | None |
| Cape Falcon to OR/CA Border | 28.0 | 21.5 | - | - | None |
| OR/CA Border to U.S./Mexico Border | 27.0 | 20.5 | - | - | None |

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS

C.1. Compliance with Minimum Size or Other Special Restrictions: All salmon on board a vessel must meet the minimum size, landing/possession limit, or other special requirements for the area being fished and the area in which they are landed if the area is open. Salmon may be landed in an area that has been closed more than 96 hours only if they meet the minimum size, landing/possession limit, or other special requirements for the area in which they were caught. Salmon may be landed in an area that has been closed less than 96 hours only if they meet the minimum size, landing/possession limit, or other special requirements for the areas in which they were caught and landed.

States may require fish landing/receiving tickets be kept on board the vessel for 90 days after landing to account for all previous salmon landings.

TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 4 of 5)

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS (continued)

C.2. Gear Restrictions:

- a. Salmon may be taken only by hook and line using single point, single shank, barbless hooks.
- b. Cape Falcon, Oregon, to the OR/CA border: No more than 4 spreads are allowed per line.
- c. OR/CA border to U.S./Mexico border: No more than 6 lines are allowed per vessel, and barbless circle hooks are required when fishing with bait by any means other than trolling.

C.3. Gear Definitions:

Trolling defined: Fishing from a boat or floating device that is making way by means of a source of power, other than drifting by means of the prevailing water current or weather conditions.

Troll fishing gear defined: One or more lines that drag hooks behind a moving fishing vessel. In that portion of the fishery management area (FMA) off Oregon and Washington, the line or lines must be affixed to the vessel and must not be intentionally disengaged from the vessel at any time during the fishing operation.

Spread defined: A single leader connected to an individual lure and/or bait.

Circle hook defined: A hook with a generally circular shape and a point which turns inward, pointing directly to the shank at a 90° angle.

C.4. Transit Through Closed Areas with Salmon on Board: It is unlawful for a vessel to have troll or recreational gear in the water while transiting any area closed to fishing for a certain species of salmon, while possessing that species of salmon; however, fishing for species other than salmon is not prohibited if the area is open for such species, and no salmon are in possession.

C.5. Control Zone Definitions:

- a. *Cape Flattery Control Zone* - The area from Cape Flattery (48°23'00" N. lat.) to the northern boundary of the U.S. EEZ; and the area from Cape Flattery south to Cape Alava (48°10'00" N. lat.) and east of 125°05'00" W. long.
- b. *Mandatory Yelloweye Rockfish Conservation Area* - The area in Washington Marine Catch Area 3 from 48°00.00' N. lat.; 125°14.00' W. long. to 48°02.00' N. lat.; 125°14.00' W. long. to 48°02.00' N. lat.; 125°16.50' W. long. to 48°00.00' N. lat.; 125°16.50' W. long. and connecting back to 48°00.00' N. lat.; 125°14.00' W. long.
- c. *Grays Harbor Control Zone* - The area defined by a line drawn from the Westport Lighthouse (46° 53'18" N. lat., 124° 07'01" W. long.) to Buoy #2 (46° 52'42" N. lat., 124°12'42" W. long.) to Buoy #3 (46° 55'00" N. lat., 124°14'48" W. long.) to the Grays Harbor north jetty (46° 36'00" N. lat., 124°10'51" W. long.).
- d. *Columbia Control Zone* - An area at the Columbia River mouth, bounded on the west by a line running northeast/southwest between the red lighted Buoy #4 (46°13'35" N. lat., 124°06'50" W. long.) and the green lighted Buoy #7 (46°15'09" N. lat., 124°06'16" W. long.); on the east, by the Buoy #10 line which bears north/south at 357° true from the south jetty at 46°14'00" N. lat., 124°03'07" W. long. to its intersection with the north jetty; on the north, by a line running northeast/southwest between the green lighted Buoy #7 to the tip of the north jetty (46°15'48" N. lat., 124°05'20" W. long.), and then along the north jetty to the point of intersection with the Buoy #10 line; and, on the south, by a line running northeast/southwest between the red lighted Buoy #4 and tip of the south jetty (46°14'03" N. lat., 124°04'05" W. long.), and then along the south jetty to the point of intersection with the Buoy #10 line.
- e. *Klamath Control Zone* - The ocean area at the Klamath River mouth bounded on the north by 41°38'48" N. lat. (approximately six nautical miles north of the Klamath River mouth); on the west, by 124°23'00" W. long. (approximately 12 nautical miles off shore); and on the south, by 41°26'48" N. lat. (approximately six nautical miles south of the Klamath River mouth).

C.6. Notification When Unsafe Conditions Prevent Compliance with Regulations: If prevented by unsafe weather conditions or mechanical problems from meeting special management area landing restrictions, vessels must notify the U.S. Coast Guard and receive acknowledgment of such notification prior to leaving the area. This notification shall include the name of the vessel, port where delivery will be made, approximate amount of salmon (by species) on board, the estimated time of arrival, and the specific reason the vessel is not able to meet special management area landing restrictions.

In addition to contacting the U.S. Coast Guard, vessels fishing south of the Oregon/California border must notify CDFG within one hour of leaving the management area by calling 800-889-8346 and providing the same information as reported to the U.S. Coast Guard. All salmon must be offloaded within 24 hours of reaching port.

TABLE V-1. Commercial troll management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 5 of 5)

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS (continued)

C.7. **Incidental Halibut Harvest:** During authorized periods, the operator of a vessel that has been issued an incidental halibut harvest license may retain Pacific halibut caught incidentally in Area 2A while trolling for salmon. Halibut retained must be no less than 32 inches in total length, measured from the tip of the lower jaw with the mouth closed to the extreme end of the middle of the tail, and must be landed with the head on. License applications for incidental harvest must be obtained from the International Pacific Halibut Commission (phone: 206-634-1838). Applicants must apply prior to April 1 of each year. Incidental harvest is authorized only during May and June troll seasons and after June 30 if quota remains and if announced on the NMFS hotline (phone: 800-662-9825). ODFW and Washington Department of Fish and Wildlife (WDFW) will monitor landings. If the landings are projected to exceed the 28,126 pound preseason allocation or the total Area 2A non-Indian commercial halibut allocation, NMFS will take inseason action to prohibit retention of halibut in the non-Indian salmon troll fishery.

Beginning May 1, license holders may land no more than one Pacific halibut per each 3 Chinook, except one Pacific halibut may be landed without meeting the ratio requirement, and no more than 35 halibut may be landed per trip. Pacific halibut retained must be no less than 32 inches in total length (with head on).

A "C-shaped" yelloweye rockfish conservation area is an area to be voluntarily avoided for salmon trolling. NMFS and the Council request salmon trollers voluntarily avoid this area in order to protect yelloweye rockfish. The area is defined in the Pacific Council Halibut Catch Sharing Plan in the North Coast subarea (Washington marine area 3), with the following coordinates in the order listed:

48°18' N. lat.; 125°18' W. long.;
48°18' N. lat.; 124°59' W. long.;
48°11' N. lat.; 124°59' W. long.;
48°11' N. lat.; 125°11' W. long.;
48°04' N. lat.; 125°11' W. long.;
48°04' N. lat.; 124°59' W. long.;
48°00' N. lat.; 124°59' W. long.;
48°00' N. lat.; 125°18' W. long.;
and connecting back to 48°18' N. lat.; 125°18' W. long.

C.8. **Inseason Management:** In addition to standard inseason actions or modifications already noted under the season description, the following inseason guidance is provided to NMFS:

- a. Chinook remaining from the May through June non-Indian commercial troll harvest guideline north of Cape Falcon may be transferred to the July through September harvest guideline on a fishery impact equivalent basis.
- b. Chinook remaining from the June and/or July non-Indian commercial troll quotas in the Oregon KMZ may be transferred to the Chinook quota for the next open period on a fishery impact equivalent basis.
- c. Chinook remaining from the July non-Indian commercial troll quota in the California KMZ area may be transferred to the August quota on a fishery impact equivalent basis.
- d. NMFS may transfer fish between the recreational and commercial fisheries north of Cape Falcon on a fishery impact neutral, fishery equivalent basis if there is agreement among the areas' representatives on the Salmon Advisory Subpanel (SAS).
- e. At the March 2012 meeting, the Council will consider inseason recommendations for special regulations for any experimental fisheries (proposals must meet Council protocol and be received in November 2011).
- f. If retention of unmarked coho is permitted by inseason action, the allowable coho quota will be adjusted to ensure preseason projected mortality of critical stocks is not exceeded.
- g. Landing limits may be modified inseason to sustain season length and keep harvest within overall quotas.

C.9. **State Waters Fisheries:** Consistent with Council management objectives:

- a. The State of Oregon may establish additional late-season fisheries in state waters.
 - b. The State of California may establish limited fisheries in selected state waters.
- Check state regulations for details.

C.10. For the purposes of California Department of Fish and Game (CDFG) Code, Section 8232.5, the definition of the Klamath Management Zone (KMZ) for the ocean salmon season shall be that area from Humbug Mt., Oregon, to Horse Mt., California.

TABLE V-2. Recreational management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 1 of 4)

| A. SEASON DESCRIPTIONS | |
|---|--|
| North of Cape Falcon | |
| Supplemental Management Information | |
| <p>1. Overall non-Indian TAC: 64,600 (non-mark-selective equivalent of 61,800) Chinook and 80,000 coho marked with a healed adipose fin clip (marked).</p> <p>2. Recreational TAC: 33,700 (non-mark selective equivalent of 30,900) Chinook and 67,200 marked coho; all retained coho must be marked.</p> <p>3. No preseason trade of Chinook or coho between non-Indian commercial and recreational fisheries.</p> <p>4. No Area 4B add-on fishery.</p> <p>5. Buoy 10 fishery opens Aug. 1 with an expected landed catch of 7,000 marked coho in August and September.</p> | |
| <p>U.S./Canada Border to Cape Falcon</p> <ul style="list-style-type: none"> June 18 through earlier of June 25 or a coastwide marked Chinook quota of 4,800 (C.5). Seven days per week. Two fish per day, all salmon except coho, all Chinook must be marked with a healed adipose fin clip (C.1). Chinook 24-inch total length minimum size limit (B). See gear restrictions (C.2). Inseason management may be used to sustain season length and keep harvest within the overall Chinook recreational TAC for north of Cape Falcon (C.5). | |
| <p>U.S./Canada Border to Cape Alava (Neah Bay)</p> <ul style="list-style-type: none"> June 26 through earlier of September 18 or 6,990 marked coho subarea quota with a subarea guideline of 3,200 Chinook. (C.5). Seven days per week. All salmon except no chum beginning August 1; two fish per day, no more than one of which can be a Chinook, plus one additional pink salmon. All coho must be marked (C.1). See gear restrictions (C.2). Beginning August 1, Chinook non-retention east of the Bonilla-Tatoosh line (C.4.a) during Council managed ocean fishery. Inseason management may be used to sustain season length and keep harvest within the overall Chinook and coho recreational TACs for north of Cape Falcon (C.5). | |
| <p>Cape Alava to Queets River (La Push Subarea)</p> <ul style="list-style-type: none"> June 26 through earlier of September 18 or 1,700 marked coho subarea quota with a subarea guideline of 1,350 Chinook. (C.5). September 24 through earlier of October 9 or 50 marked coho quota or 50 Chinook quota (C.5) in the area north of 47°50'00" N. lat. and south of 48°00'00" N. lat. <p>Seven days per week. All salmon; two fish per day, no more than one of which can be a Chinook, plus one additional pink salmon. All coho must be marked (C.1). See gear restrictions (C.2). Inseason management may be used to sustain season length and keep harvest within the overall Chinook and coho recreational TACs for north of Cape Falcon (C.5).</p> | |
| <p>Queets River to Leadbetter Point (Westport Subarea)</p> <ul style="list-style-type: none"> June 26 through earlier of September 18 or 24,860 marked coho subarea quota with a subarea guideline of 16,900 Chinook (C.5). <p>Sunday through Thursday. All salmon; two fish per day, no more than one of which can be a Chinook. All coho must be marked (C.1). See gear restrictions and definitions (C.2, C.3). Grays Harbor Control Zone closed beginning August 1 (C.4). Inseason management may be used to sustain season length and keep harvest within the overall Chinook and coho recreational TACs for north of Cape Falcon (C.5).</p> | |
| <p>Leadbetter Point to Cape Falcon (Columbia River Subarea)</p> <ul style="list-style-type: none"> June 26 through earlier of September 30 or 33,600 marked coho subarea quota with a subarea guideline of 7,400 Chinook (C.5). Seven days per week. All salmon; two fish per day, no more than one of which can be a Chinook. All coho must be marked (C.1). See gear restrictions and definitions (C.2, C.3). Columbia Control Zone closed (C.4.c). Inseason management may be used to sustain season length and keep harvest within the overall Chinook and coho recreational TACs for north of Cape Falcon (C.5). | |

TABLE V-2. Recreational management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 2 of 4)

| A. SEASON DESCRIPTIONS |
|--|
| South of Cape Falcon |
| Supplemental Management Information |
| <ol style="list-style-type: none"> 1. Sacramento River Basin recreational fishery catch assumption: 61,400 adult Sacramento River fall Chinook. 2. Sacramento River fall Chinook spawning escapement of 377,000 adults. 3. Klamath River recreational fishery allocation: 7,900 adult Klamath River fall Chinook. 4. Klamath tribal allocation: 34,800 adult Klamath River fall Chinook. 5. Overall recreational TAC: 15,000 marked coho and 3,000 unmarked coho. |
| <p>Cape Falcon to Humbug Mt.</p> <ul style="list-style-type: none"> • Except as provided below during the all-salmon mark-selective and non-mark-selective coho fisheries, the season will be March 15 through September 30 (C.6). <p>All salmon except coho; two fish per day (C.1). See gear restrictions and definitions (C.2, C.3).</p> <ul style="list-style-type: none"> • Cape Falcon to Humbug Mt. all-salmon mark-selective coho fishery: July 2 through earlier of August 13 or a landed catch of 15,000 marked coho. <p>Seven days per week. All salmon, two fish per day. All retained coho must be marked (C.1). Any remainder of the mark selective coho quota will be transferred on an impact neutral basis to the September non-selective coho quota listed below. The all salmon except coho season reopens the earlier of August 14 or attainment of the coho quota, through August 31.</p> <ul style="list-style-type: none"> • Cape Falcon to Humbug Mt. non-mark-selective coho fishery: September 1 through the earlier of September 10 or a landed catch of 3,000 non-mark-selective coho quota (C.5). <p>Thursday through Saturday all salmon, two fish per day; Sunday through Wednesday, all salmon except coho, two fish per day.</p> <p>The all salmon except coho season reopens the earlier of September 11 or attainment of the coho quota (C.5). Open days may be adjusted inseason to utilize the available coho quota (C.5).</p> <p>Fishing in the Stonewall Bank yelloweye rockfish conservation area restricted to trolling only on days the all depth recreational halibut fishery is open (call the halibut fishing hotline 1-800-662-9825 for specific dates) (C.3.b, C.4.d).</p> <p>In 2012, the season between Cape Falcon and Humbug Mt. will open March 15 for all salmon except coho, two fish per day (B, C.1, C.2, C.3).</p> |
| <p>Humbug Mt. to OR/CA Border. (Oregon KMZ)</p> <ul style="list-style-type: none"> • May 14 through September 5 (C.6). <p>Seven days per week. All salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B). See gear restrictions and definitions (C.2, C.3).</p> |
| <p>OR/CA Border to Horse Mt. (California KMZ)</p> <ul style="list-style-type: none"> • May 14 through September 5 (C.6). <p>Seven days per week. All salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B). See gear restrictions and definitions (C.2, C.3). Klamath Control Zone closed in August (C.4.e). See California State regulations for additional closures adjacent to the Smith, Eel, and Klamath rivers.</p> |
| <p>Horse Mt. to Point Arena (Fort Bragg)</p> <ul style="list-style-type: none"> • April 2 through October 30. <p>Seven days per week. All salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B). See gear restrictions and definitions (C.2, C.3).</p> <p>In 2012, season opens April 7 for all salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B); and the same gear restrictions as in 2011 (C.2, C.3).</p> |
| <p>Point Arena to Pigeon Pt. (San Francisco)</p> <ul style="list-style-type: none"> • April 2 through October 30. <p>Seven days per week. All salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B). See gear restrictions and definitions (C.2, C.3).</p> <p>In 2012, season opens April 7 for all salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B); and the same gear restrictions as in 2011 (C.2, C.3).</p> |

TABLE V-2. Recreational management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 3 of 4)

| A. SEASON DESCRIPTIONS | | | |
|---|--|--|--|
| South of Cape Falcon | | | |
| Pigeon Point to U.S./Mexico Border (Monterey South) | | | |
| <ul style="list-style-type: none"> April 2 through September 18. Seven days per week. All salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B). See gear restrictions and definitions (C.2, C.3). | | | |
| In 2012, season opens April 7 for all salmon except coho, two fish per day (C.1). Chinook minimum size limit of 24 inches total length (B); and the same gear restrictions as in 2011 (C.2, C.3). | | | |
| California State regulations require all salmon be made available to a CDFG representative for sampling immediately at port of landing. Any person in possession of a salmon with a missing adipose fin, upon request by an authorized agent or employee of the CDFG, shall immediately relinquish the head of the salmon to the state. (California Fish and Game Code §8226) | | | |

| B. MINIMUM SIZE (Inches) (See C.1) | | | |
|------------------------------------|---------|------|------|
| Area (when open) | Chinook | Coho | Pink |
| North of Cape Falcon | 24.0 | 16.0 | None |
| Cape Falcon to OR/CA Border | 24.0 | 16.0 | None |
| OR/CA Border to U.S./Mexico Border | 24.0 | - | 24.0 |

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS

C.1. Compliance with Minimum Size and Other Special Restrictions: All salmon on board a vessel must meet the minimum size or other special requirements for the area being fished and the area in which they are landed if that area is open. Salmon may be landed in an area that is closed only if they meet the minimum size or other special requirements for the area in which they were caught.

Ocean Boat Limits: Off the coast of Washington, Oregon, and California, each fisher aboard a vessel may continue to use angling gear until the combined daily limits of salmon for all licensed and juvenile anglers aboard has been attained (additional state restrictions may apply).

C.2. Gear Restrictions: Salmon may be taken only by hook and line using barbless hooks. All persons fishing for salmon, and all persons fishing from a boat with salmon on board, must meet the gear restrictions listed below for specific areas or seasons.

- U.S./Canada Border to Point Conception, California:* No more than one rod may be used per angler; and no more than two single point, single shank barbless hooks are required for all fishing gear. [Note: ODFW regulations in the state-water fishery off Tillamook Bay may allow the use of barbed hooks to be consistent with inside regulations.]
- Horse Mt., California, to Point Conception, California:* Single point, single shank, barbless circle hooks (see gear definitions below) are required when fishing with bait by any means other than trolling, and no more than two such hooks shall be used. When angling with two hooks, the distance between the hooks must not exceed five inches when measured from the top of the eye of the top hook to the inner base of the curve of the lower hook, and both hooks must be permanently tied in place (hard tied). Circle hooks are not required when artificial lures are used without bait.

C.3. Gear Definitions:

- Recreational fishing gear defined:* Angling tackle consisting of a line with no more than one artificial lure and/or natural bait attached. Off Oregon and Washington, the line must be attached to a rod and reel held by hand or closely attended; the rod and reel must be held by hand while playing a hooked fish. No person may use more than one rod and line while fishing off Oregon or Washington. Off California, the line must be attached to a rod and reel held by hand or closely attended; weights directly attached to a line may not exceed four pounds (1.8 kg). While fishing off California north of Point Conception, no person fishing for salmon, and no person fishing from a boat with salmon on board, may use more than one rod and line. Fishing includes any activity which can reasonably be expected to result in the catching, taking, or harvesting of fish.
- Trolling defined:* Angling from a boat or floating device that is making way by means of a source of power, other than drifting by means of the prevailing water current or weather conditions.
- Circle hook defined:* A hook with a generally circular shape and a point which turns inward, pointing directly to the shank at a 90° angle.

TABLE V-2. Recreational management measures adopted by the Council for non-Indian ocean salmon fisheries, 2011.
(Page 4 of 4)

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS

C.4. Control Zone Definitions:

- a. *The Bonilla-Tatoosh Line:* A line running from the western end of Cape Flattery to Tatoosh Island Lighthouse (48°23'30" N. lat., 124°44'12" W. long.) to the buoy adjacent to Duntze Rock (48°28'00" N. lat., 124°45'00" W. long.), then in a straight line to Bonilla Point (48°35'30" N. lat., 124°43'00" W. long.) on Vancouver Island, British Columbia.
- b. *Grays Harbor Control Zone* - The area defined by a line drawn from the Westport Lighthouse (46° 53'18" N. lat., 124° 07'01" W. long.) to Buoy #2 (46° 52'42" N. lat., 124°12'42" W. long.) to Buoy #3 (46° 55'00" N. lat., 124°14'48" W. long.) to the Grays Harbor north jetty (46° 36'00" N. lat., 124°10'51" W. long.).
- c. *Columbia Control Zone:* An area at the Columbia River mouth, bounded on the west by a line running northeast/southwest between the red lighted Buoy #4 (46°13'35" N. lat., 124°06'50" W. long.) and the green lighted Buoy #7 (46°15'09" N. lat., 124°06'16" W. long.); on the east, by the Buoy #10 line which bears north/south at 357° true from the south jetty at 46°14'00" N. lat., 124°03'07" W. long. to its intersection with the north jetty; on the north, by a line running northeast/southwest between the green lighted Buoy #7 to the tip of the north jetty (46°15'48" N. lat., 124°05'20" W. long. and then along the north jetty to the point of intersection with the Buoy #10 line; and on the south, by a line running northeast/southwest between the red lighted Buoy #4 and tip of the south jetty (46°14'03" N. lat., 124°04'05" W. long.), and then along the south jetty to the point of intersection with the Buoy #10 line.
- d. *Stonewall Bank Yelloweye Rockfish Conservation Area:* The area defined by the following coordinates in the order listed:
 44°37.46' N. lat.; 124°24.92' W. long.;
 44°37.46' N. lat.; 124°23.63' W. long.;
 44°28.71' N. lat.; 124°21.80' W. long.;
 44°28.71' N. lat.; 124°24.10' W. long.;
 44°31.42' N. lat.; 124°25.47' W. long.;
 and connecting back to 44°37.46' N. lat.; 124°24.92' W. long.
- e. *Klamath Control Zone:* The ocean area at the Klamath River mouth bounded on the north by 41°38'48" N. lat. (approximately six nautical miles north of the Klamath River mouth); on the west, by 124°23'00" W. long. (approximately 12 nautical miles off shore); and, on the south, by 41°26'48" N. lat. (approximately 6 nautical miles south of the Klamath River mouth).

C.5. Inseason Management: Regulatory modifications may become necessary inseason to meet preseason management objectives such as quotas, harvest guidelines, and season duration. In addition to standard inseason actions or modifications already noted under the season description, the following inseason guidance is provided to NMFS:

- a. Actions could include modifications to bag limits, or days open to fishing, and extensions or reductions in areas open to fishing.
- b. Coho may be transferred inseason among recreational subareas north of Cape Falcon on a fishery impact equivalent basis to help meet the recreational season duration objectives (for each subarea) after conferring with representatives of the affected ports and the Council's SAS recreational representatives north of Cape Falcon.
- c. Chinook and coho may be transferred between the recreational and commercial fisheries north of Cape Falcon on a fishery impact equivalent basis if there is agreement among the representatives of the Salmon Advisory Subpanel (SAS).
- d. If retention of unmarked coho is permitted in the area from the U.S./Canada border to Cape Falcon, Oregon, by inseason action, the allowable coho quota will be adjusted to ensure preseason projected mortality of critical stocks is not exceeded.
- e. Marked coho remaining from the June/July through August Cape Falcon to OR/CA border recreational coho quota may be transferred inseason to the September Cape Falcon to Humbug Mt. non-mark-selective recreational fishery on a fishery impact equivalent basis.

C.6. Additional Seasons in State Territorial Waters: Consistent with Council management objectives, the States of Washington, Oregon, and California may establish limited seasons in state waters. Check state regulations for details.

TABLE V-3. Treaty Indian ocean troll management measures adopted by the Council for ocean salmon fisheries, 2011. (Page 1 of 1)

| A. SEASON DESCRIPTIONS | | | | | |
|--|--|--|--|--|--|
| Supplemental Management Information | | | | | |
| 1. Overall Treaty-Indian TAC: 41,000 Chinook and 42,000 coho. | | | | | |
| <p>May 1 through the earlier of June 30 or 19,750 Chinook quota.</p> <p>All salmon except coho. If the Chinook quota for the May-June fishery is not fully utilized, the excess fish cannot be transferred into the later all-salmon season. If the Chinook quota is exceeded, the excess will be deducted from the later all-salmon season. See size limit (B) and other restrictions (C).</p> <ul style="list-style-type: none"> July 1 through the earlier of September 15, or 21,250 preseason Chinook quota, or 42,000 coho quota. <p>All salmon. See size limit (B) and other restrictions (C).</p> | | | | | |

| B. MINIMUM SIZE (Inches) | | | | | |
|--------------------------|----------------|----------------|----------------|----------------|------|
| Area (when open) | Chinook | | Coho | | Pink |
| | Total Length | Head-off | Total Length | Head-off | |
| North of Cape Falcon | 24.0 (61.0 cm) | 18.0 (45.7 cm) | 16.0 (40.6 cm) | 12.0 (30.5 cm) | None |

C. REQUIREMENTS, DEFINITIONS, RESTRICTIONS, OR EXCEPTIONS

C.1. Tribe and Area Boundaries. All boundaries may be changed to include such other areas as may hereafter be authorized by a Federal court for that tribe's treaty fishery.

S'KLALLAM - Washington State Statistical Area 4B (All).

MAKAH - Washington State Statistical Area 4B and that portion of the FMA north of 48°02'15" N. lat. (Norwegian Memorial) and east of 125°44'00" W. long.

QUILEUTE - That portion of the FMA between 48°07'36" N. lat. (Sand Pt.) and 47°31'42" N. lat. (Queets River) and east of 125°44'00" W. long.

HOH - That portion of the FMA between 47°54'18" N. lat. (Quillayute River) and 47°21'00" N. lat. (Quinault River) and east of 125°44'00" W. long.

QUINAULT - That portion of the FMA between 47°40'06" N. lat. (Destruction Island) and 46°53'18"N. lat. (Point Chehalis) and east of 125°44'00" W. long.

C.2. Gear restrictions

- Single point, single shank, barbless hooks are required in all fisheries.
- No more than eight fixed lines per boat.
- No more than four hand held lines per person in the Makah area fishery (Washington State Statistical Area 4B and that portion of the FMA north of 48°02'15" N. lat. (Norwegian Memorial) and east of 125°44'00" W. long.)

C.3. Quotas

- The quotas include troll catches by the S'Klallam and Makah tribes in Washington State Statistical Area 4B from May 1 through September 15.
- The Quileute Tribe will continue a ceremonial and subsistence fishery during the time frame of September 15 through October 15 in the same manner as in 2004-2010. Fish taken during this fishery are to be counted against treaty troll quotas established for the 2011 season (estimated harvest during the October ceremonial and subsistence fishery: 100 Chinook; 200 coho).

C.4. Area Closures

- The area within a six nautical mile radius of the mouths of the Queets River (47°31'42" N. lat.) and the Hoh River (47°45'12" N. lat.) will be closed to commercial fishing.
- A closure within two nautical miles of the mouth of the Quinault River (47°21'00" N. lat.) may be enacted by the Quinault Nation and/or the State of Washington and will not adversely affect the Secretary of Commerce's management regime.

TABLE V-4. Stock status relative to overfished and overfishing criteria. A stock is approaching an overfished condition if the 3-year geometric mean of the most recent two years and the forecast spawning escapement is less than the minimum stock size threshold (MSST); a stock would experience overfishing if the total annual exploitation rate exceeds the maximum fishing mortality threshold (MFMT). 2012 spawning escapement and exploitation rate estimates are based on preliminary 2012 preseason abundance forecasts and 2011 Council regulations.

| | Spawning Escapement | | | | | | | | Total Exploitation Rate | | | | | |
|---|---------------------|--------|---------|--------------------|-----------------------------|---------------|-----------|------------------|-------------------------|------|------|------|--------------------|-------|
| | 2008 | 2009 | 2010 | 2011 ^{a/} | Forecast 2012 ^{b/} | 3-yr Geo Mean | MSST | S _{MSY} | 2008 | 2009 | 2010 | 2011 | 2012 ^{b/} | MFMT |
| Chinook | | | | | | | | | | | | | | |
| Sacramento Fall | 65,364 | 40,873 | 124,270 | 114,741 | 469,000 | 188,402 | 91,500 | 122,000 | 0.06 | 0.01 | 0.17 | 0.42 | 0.43 | 0.78 |
| Klamath River Fall | 30,850 | 44,409 | 37,225 | 47,754 | 179,000 | 68,270 | 30,525 | 40,700 | 0.45 | 0.37 | 0.42 | 0.38 | 0.34 | 0.71 |
| Southern Oregon | 13 | 66 | 52 | 35 | NA | 49 | 30-45 | 150,000 to | NA | NA | NA | NA | NA | 0.78 |
| Central and Northern | 40 | 61 | 87 | 92 | NA | 79 | fish/mile | 200,000 | 0.52 | 0.53 | NA | NA | NA | 0.78 |
| Upper River Bright - Fall ^{c/} | 51,757 | 62,428 | 114,230 | 93,510 | 121,910 | 109,201 | 19,182 | 39,625 | 0.54 | 0.67 | NA | NA | NA | 0.86 |
| Upper River - Summer ^{c/} | 38,171 | 44,295 | 47,220 | 44,432 | 55,032 | 48,694 | 6,072 | 12,143 | 0.53 | 0.46 | NA | NA | NA | 0.75 |
| Willapa Bay - Fall ^{d/} | 1,900 | 2,847 | 3,395 | 3,690 | NA | 3,292 | 1,696 | 3,393 | 0.52 | 0.56 | NA | NA | NA | 0.78 |
| Grays Harbor Fall ^{d/} | 13,570 | 7,215 | 16,951 | NA | NA | 11,840 | 5,694 | 11,388 | 0.52 | 0.56 | NA | NA | NA | 0.78 |
| Grays Harbor Spring | 995 | 1,132 | 3,495 | 2,563 | NA | 2,164 | 546 | 1,092 | NA | NA | NA | NA | NA | 0.78 |
| Queets - Fall ^{c/} | 2,971 | 2,960 | 3,861 | 3,767 | NA | 3,505 | 1,250 | 2,500 | 0.52 | 0.56 | NA | NA | NA | 0.87 |
| Queets - Sp/Su | 305 | 495 | 259 | 373 | NA | 363 | 350 | 700 | NA | NA | NA | NA | NA | 0.78 |
| Hoh - Fall ^{d/} | 2,999 | 2,081 | 2,599 | 1,293 | NA | 1,912 | 600 | 1,200 | 0.52 | 0.56 | NA | NA | NA | 0.90 |
| Hoh Sp/Su | 671 | 880 | 828 | 827 | NA | 845 | 450 | 900 | NA | NA | NA | NA | NA | 0.78 |
| Quillayute - Fall ^{d/} | 3,612 | 3,130 | 4,635 | 3,993 | NA | 3,869 | 1,500 | 3,000 | 0.52 | 0.56 | NA | NA | NA | 0.87 |
| Quillayute - Sp/Su | 949 | 555 | 815 | 600 | NA | 647 | 600 | 1,200 | NA | NA | NA | NA | NA | 0.78 |
| Hoko -Su/Fa ^{c/} | 483 | 375 | 793 | 1,504 | NA | 764 | 425 | 850 | 0.63 | 0.25 | NA | NA | NA | 0.78 |
| Coho | | | | | | | | | | | | | | |
| Willapa Bay | 16,419 | 47,333 | 77,784 | 26,122 | 27,283 | 38,130 | Undef | Undef | NA | NA | NA | NA | 0.67 | Undef |
| Grays Harbor | 34,054 | 69,222 | 102,237 | 68,504 | 316,969 | 130,451 | 18,320 | 24,426 | 0.31 | 0.33 | NA | NA | 0.44 | 0.65 |
| Queets | 4,629 | 9,200 | 11,261 | NA | 28,593 | 17,944 | 4,350 | 5,800 | 0.37 | 0.43 | NA | NA | 0.49 | 0.65 |
| Hoh | 2,461 | 6,595 | 7,864 | 5,903 | 12,314 | 8,299 | 1,890 | 2,520 | 0.43 | 0.52 | NA | NA | 0.51 | 0.65 |
| Quillayute Fall | 6,252 | 7,863 | 9,837 | 9,512 | 31,535 | 14,343 | 4,725 | 6,300 | 0.37 | 0.50 | NA | NA | 0.49 | Undef |
| Juan de Fuca | 3,339 | 14,957 | 19,282 | 17,167 | 11,295 | 15,521 | 7,000 | 11,000 | 0.13 | 0.30 | NA | NA | 0.11 | 0.60 |
| Hood Canal | 11,516 | 26,961 | 4,197 | NA | 44,399 | 13,651 | 10,750 | 14,350 | 0.63 | 0.59 | NA | NA | 0.40 | 0.65 |
| Skagit | 24,093 | 60,798 | 31,090 | 45,220 | 30,091 | 34,844 | 14,875 | 25,000 | 0.32 | 0.31 | NA | NA | 0.38 | 0.60 |
| Stillaguamish | 12,938 | 22,179 | 15,172 | NA | 34,562 | 22,899 | 6,100 | 10,000 | 0.23 | 0.28 | NA | NA | 0.28 | 0.50 |
| Snohomish | 36,015 | 98,945 | 49,100 | NA | 78,321 | 62,013 | 31,000 | 50,000 | 0.28 | 0.26 | NA | NA | 0.26 | 0.60 |

a/ Preliminary.

b/ Preliminary approximations based on preseason abundance projections and last year's regulations or season structures.

c/ CWT based exploitation rates from annual catch and escapement distribution from PSC-CTC 2011 Exploitation Rate Analysis.

d/ Queets River fall Chinook coded-wire-tag (CWT) exploitation rates used as a proxy. Exploitation rates in the terminal fisheries will differ from those calculated for Queets fall CWTs.

TABLE V-5. Estimated ocean escapements and exploitation rates for critical natural and Columbia River hatchery coho stocks (thousands of fish) based on preliminary 2012 preseason abundance forecasts and 2011 Council management measures.^{a/}

| Abundance of fish based on preliminary 2012 preseason abundance forecasts and 2011 Council management measures. | | | | | |
|---|--|-------------------|----------------|-------------------|---|
| Stock | Ocean Escapement and ER Estimates Under 2011 Regulations ^{b/} | | | | 2012 FMP Conservation Objective ^{c/} |
| | 2012 Preseason | | 2011 Preseason | | |
| | Abundance | Exploitation Rate | Abundance | Exploitation Rate | |
| Natural Coho Stocks | | | | | |
| Skagit | 40.2 | 38.0% | 116.6 | 35.3% | Exploitation Rate ≤35.0% ^{d/} |
| Stillaguamish | 37.6 | 27.5% | 53.6 | 25.9% | Exploitation Rate ≤50.0% ^{d/} |
| Snohomish | 83.5 | 28.5% | 142.3 | 25.8% | Exploitation Rate ≤40.0% ^{d/} |
| Hood Canal | 58.7 | 39.8% | 60.8 | 40.0% | Exploitation Rate ≤65.0% ^{d/} |
| Strait of Juan de Fuca | 11.6 | 10.8% | 11.3 | 10.8% | Exploitation Rate ≤40.0% ^{d/} |
| Quillayute Fall | 31.5 | 49.0% | 26.6 | | 6.3 - 15.8 Spawners |
| Hoh | 12.3 | 51.0% | 10.0 | | 2.0 - 5.0 Spawners |
| Queets | 29.4 | 49.0% | 10.5 | | 5.8 - 14.5 Spawners |
| Grays Harbor | 137.0 | 44.0% | 81.4 | | 35.4 Spawners |
| LCN | 26.4 | 15.2% | 20.1 | 15.0% | Exploitation Rate ≤15.0% |
| OCN | 256.5 | 12.2% | 217.4 | 13.2% | Exploitation Rate ≤15.0% |
| R/K | NA | 6.9% | NA | 8.5% | Exploitation Rate ≤13.0% |
| Hatchery Coho Stocks | | | | | |
| Columbia Early | 173.2 | | 162.0 | | 18.6 Hatchery Escapement |
| Columbia Late | 55.0 | | 101.0 | | 11.9 Hatchery Escapement |

a/ Quota levels include harvest and hooking mortality estimates used in planning the Council's 2011 ocean fisheries and a coho catch for the Canadian troll fishery off the West Coast of Vancouver Island (WCVI).

b/ 2011 preseason regulations include the following coho quota fisheries: U.S. Canada Border to Cape Falcon: Treaty Indian troll - 42,000 non-selective; non-Indian troll - 12,800 selective; recreational - 67,200 selective; Cape Falcon to OR/CA border: recreational - 15,000 selective and 3,000 non-selective; troll - none. Ocean escapement is generally the estimated number of coho escaping ocean fisheries and entering freshwater. For Puget Sound stocks, ocean escapement is the estimated number of coho entering Puget Sound (Area 4B) which are available for U.S. net fisheries in Puget Sound and spawning escapement after impacts associated with the Canadian and Puget Sound troll and recreational fisheries have been deducted. For the OCN coho stock, this value represents the estimated spawner escapement in SRS accounting. For Columbia River hatchery and LCN stocks, ocean escapement represents the number of coho before the Buoy 10 fishery; the LCN exploitation rate shown is the Council fisheries exploitation rate, which had an ER forecast of 10.7% and an ESA limit of 15% including mainstem Columbia River fisheries.

c/ Goals represent Salmon FMP conservation objectives, ESA consultation standards, or hatchery escapement needs. Spawning escapement goals are not directly comparable to ocean escapement because the latter occur before inside

d/ Assumed exploitation rate based on preliminary abundance forecasts.

TABLE V-6. Comparison of Lower Columbia natural (LCN), Oregon coastal natural (OCN), and Rogue/Klamath (RK) coho projected harvest mortality and exploitation rates by fishery under Council-adopted 2011 management measures and preliminary 2012 preseason abundance estimates.

| Fishery | Projected Harvest Mortality and Exploitation Rate | | | | | |
|-------------------------------|---|---------|--------|---------|------------------|---------|
| | LCN | | OCN | | RK ^{a/} | |
| | Number | Percent | Number | Percent | Number | Percent |
| SOUTHEAST ALASKA | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| BRITISH COLUMBIA | 37 | 0.1% | 998 | 0.3% | 28 | 0.2% |
| PUGET SOUND/STRAITS | 42 | 0.1% | 238 | 0.1% | 0 | 0.0% |
| NORTH OF CAPE FALCON | | | | | | |
| Recreational | 1,448 | 4.8% | 2,643 | 0.9% | 6 | 0.0% |
| Treaty Indian Troll | 606 | 2.0% | 1,522 | 0.5% | 0 | 0.0% |
| Non-Indian Troll | 438 | 1.5% | 1,186 | 0.4% | 1 | 0.0% |
| SOUTH OF CAPE FALCON | | | | | | |
| Recreational: | | | | | | |
| Cape Falcon to Humbug Mt. | 400 | 1.3% | 7,383 | 2.5% | 27 | 0.2% |
| Humbug Mt. to Horse Mt. (KMZ) | 50 | 0.2% | 2,621 | 0.9% | 457 | 3.4% |
| Fort Bragg | 12 | 0.0% | 1,093 | 0.4% | 130 | 1.0% |
| South of Pt. Arena | 11 | 0.0% | 964 | 0.3% | 105 | 0.8% |
| Troll: | | | | | | |
| Cape Falcon to Humbug Mt. | 243 | 0.8% | 2,647 | 0.9% | 17 | 0.1% |
| Humbug Mt. to Horse Mt. (KMZ) | 3 | 0.0% | 257 | 0.1% | 34 | 0.3% |
| Fort Bragg | 2 | 0.0% | 518 | 0.2% | 57 | 0.4% |
| South of Pt. Arena | 12 | 0.0% | 1,007 | 0.3% | 35 | 0.3% |
| BUOY 10 | 224 | 0.7% | 139 | 0.0% | 0 | 0.0% |
| ESTUARY/FRESHWATER | NA | 3.7% | 12,399 | 4.2% | 32 | 0.2% |
| TOTAL | 3,528 | 15.2% | 35,615 | 12.2% | 929 | 6.9% |

a/ Unmarked hatchery production used as a surrogate for Rogue/Klamath natural stock coho.

TABLE V-7 Maximum allowable fishery impact rate for OCN coho under Amendment 13 matrix (Appendix A, Table A-2) and the OCN work group matrix (Appendix A, Table A-3) based on parent escapement levels by stock component and marine survival category.^{a/}

| Fishery Year (t) | Estimated OCN Coho Spawners by Stock Component | | | | | Hatchery Jack Survival Rate (t-1) | Amendment 13 Matrix | | | OCN Work Group Matrix ^{b/} | | |
|---------------------|--|----------|-------------------|-------------------|----------|--|--------------------------------|---------------------------------|---------------------------------|-------------------------------------|---------------------------------|---------------------------------|
| | Parent Spawner Year (t-3) | Northern | North- Central | South- Central | Southern | | Marine Survival Category | Parental Spawner Category | Maximum Allowable Impacts | Marine Survival Category | Parental Spawner Category | Maximum Allowable Impacts |
| 1998 | 1995 | 3,900 | 13,600 | 36,500 | 3,400 | 0.04% | Low | Very Low | ≤10-13% | Extremely Low | Very Low | ≤8% |
| 1999 | 1996 | 3,300 | 18,100 | 52,600 | 5,200 | 0.10% | Med | Very Low | ≤15% | Low | Critical | 0-8% |
| 2000 | 1997 | 2,100 | 2,800 | 18,400 | 8,200 | 0.12% | Med | Very Low | ≤15% | Low | Critical | 0-8% |
| 2001 | 1998 | 2,600 | 3,300 | 25,900 | 2,300 | 0.27% | Med | Very Low | ≤15% | Medium | Critical | 0-8% |
| 2002 | 1999 | 8,900 | 11,800 | 29,100 | 1,400 | 0.09% | Med | Low | ≤15% | Low | Low | ≤15% |
| 2003 | 2000 | 17,900 | 14,300 | 36,500 | 11,000 | 0.20% | Med | Low | ≤15% | Med | Low | ≤15% |
| 2004 | 2001 | 33,500 | 25,200 | 112,000 | 12,600 | 0.14% | Med | Low | ≤15% | Med | Low | ≤15% |
| 2005 | 2002 | 52,500 | 104,000 | 104,100 | 8,400 | 0.11% | Med | High | ≤20% | Low | High | ≤15% |
| 2006 | 2003 | 59,600 | 68,900 | 99,800 | 6,800 | 0.12% | Med | High | ≤20% | Low | High | ≤15% |
| 2007 | 2004 | 28,800 | 42,100 | 101,900 | 24,500 | 0.17% | Med | Med | ≤20% | Med | Med | ≤20% |
| 2008 | 2005 | 16,500 | 51,400 | 86,700 | 10,000 | 0.07% | Low | High | ≤15% | Extremely Low | High | ≤8% |
| 2009 | 2006 | 24,100 | 21,200 | 83,500 | 3,900 | 0.27% | Med | Low | ≤15% | Med | Low | ≤15% |
| 2010 | 2007 | 17,500 | 12,300 | 36,500 | 5,200 | 0.12% | Med | Low | ≤15% | Low | Low | ≤15% |
| 2011 | 2008 | 25,600 | 68,100 | 86,000 | 400 | 0.12% | Med | High | ≤20% | Low | High | ≤15% |
| 2012 | 2009 | 48,100 | 86,400 | 128,200 | 2,600 | 0.09% | Med | High | ≤20% | Low | High | ≤15% |
| 2013 | 2010 | 55,000 | 56,600 | 171,900 | 3,100 | - | - | High | - | - | High | - |
| 2014 | 2011 | 47,800 | 105,000 | 138,600 | 3,900 | - | - | High | - | - | High | - |

a/ Under the NMFS ESA consultation standards, the southern stock component is managed for a total allowable Marine Exploitation rate of 13%, as represented by Rogue/Klamath hatchery stocks, which is separate from these OCN coho impact rates.

b/ Developed by the OCN Coho Work Group as a result of the 2000 Review of Amendment 13.

CHAPTER VI: REFERENCES

- National Marine Fisheries Service (NMFS). 2003. Final Programmatic environmental impact statement for Pacific salmon fisheries management off the coasts of Southeast Alaska, Washington, Oregon, and California, and in the Columbia River basin. National Marine Fisheries Service Northwest Region, Seattle.
- NMFS. 2008. Endangered Species Act-section 7 formal consultation biological opinion: Effects of the 2008 Pacific Coast salmon plan fisheries on the southern resident killer whale distinct population segment (*Orcinus orca*) and their critical habitat. National Marine Fisheries Service Northwest Region, Seattle.
- Pacific Fishery Management Council (PFMC). 2006. Environmental assessment for the proposed 2006 management measures for the ocean salmon fishery managed under the Pacific Coast salmon plan. Pacific Fishery Management Council, Portland, Oregon.
- PFMC. 2011a. Preseason Report II: Proposed Alternatives and Environmental Assessment - Part 2 for 2011 Ocean Salmon Fishery Regulations. Pacific Fishery Management Council, Portland, Oregon.
- PFMC. 2011b. Preseason Report III: Analysis of Council Adopted Management Measures for 2011 Ocean Salmon Fisheries. Pacific Fishery Management Council, Portland, Oregon.
- PFMC. 2012. Review of 2011 Ocean Salmon Fisheries: Stock Assessment and Fishery Evaluation Document for the Pacific Coast Salmon Fishery Management Plan. Pacific Fishery Management Council, Portland, Oregon.

APPENDIX A

SUMMARY OF COUNCIL STOCK MANAGEMENT GOALS

LIST OF TABLES

| | <u>Page</u> |
|---|-------------|
| TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP | 95 |
| TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13..... | 101 |
| TABLE A-3. Fishery impact rate criteria for OCN coho stock components based on the harvest matrix resulting from the OCN work group 2000 review of Amendment 13..... | 102 |
| TABLE A-4. Council adopted management objectives for Puget Sound natural coho management units, expressed as exploitation rate ceilings for critical, low and normal abundance based status categories, with runsize breakpoints..... | 103 |
| TABLE A-5. Council recommended management objectives for Lower Columbia River natural tule Chinook, expressed as exploitation rate ceilings for abundance based status categories, with runsize forecast bins expressed as adult river mouth return forecasts of Lower Columbia River hatchery tule Chinook. | 103 |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 1 of 6)

| CHINOOK | | | | | |
|--|---|-----------------------------------|-----------------------------------|--------------------------|--|
| Stocks In The Fishery | Conservation Objective | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| Sacramento River Fall Indicator stock for the Central Valley fall (CVF) Chinook stock complex. | 122,000-180,000 natural and hatchery adult spawners (MSY proxy adopted 1984). This objective is intended to provide adequate escapement of natural and hatchery production for Sacramento and San Joaquin fall and late-fall stocks based on habitat conditions and average run-sizes as follows: Sacramento River 1953-1960; San Joaquin River 1972-1977 (ASETF 1979; PFMC 1984; SRFCRT 1994). The objective is less than the estimated basin capacity of 240,000 spawners (Hallock 1977), but greater than the 118,000 spawners for maximum production estimated on a basin by basin basis before Oroville and Nimbus Dams (Reisenbichler 1986). | 122,000 | 91,500 | 78% Proxy (SAC 2011) | Based on F _{ABC} and annual ocean abundance. F _{ABC} is F _{MSY} reduced by Tier 2 (10%) uncertainty |
| Sacramento River Spring ESA Threatened | NMFS ESA consultation standard/recovery plan: Conform to Sacramento River Winter Chinook ESA consultation standard (no defined objective for ocean management prior to listing). | Undefined | Undefined | Undefined | ESA consultation standard applies. |
| Sacramento River Winter ESA Endangered | NMFS ESA consultation standard/recovery plan: Recreational seasons: Point Arena to Pigeon Point between the first Saturday in April and the second Sunday in November; Pigeon Point to the U.S./Mexico Border between the first Saturday in April and the first Sunday in October. Minimum size limit ≥ 20 inches total length. Commercial seasons: Point Arena to the U.S./Mexico border between May 1 and September 30, except Point Reyes to Point San Pedro between October 1 and 15 (Monday through Friday). Minimum size limit ≥ 26 inches total length. Guidance from NMFS in 2010 and 2011 required implementation of additional closures and/or increased sized limits in the recreational fishery South of Point Arena. A new winter-run management framework and consultation standard is expected to be in place for the 2012 fishing season, or no later than March 1, 2012. (NMFS ESA Guidance for 2011). | Undefined | Undefined | Undefined | |
| California Coastal Chinook ESA Threatened | NMFS ESA consultation standard/recovery plan: Limit ocean fisheries to no more than a 16.0% age-4 ocean harvest rate on Klamath River fall Chinook. | Undefined | Undefined | Undefined | |
| Klamath River Fall Indicator stock for the Southern Oregon Northern California (SONC) Chinook stock complex. | At least 32% of potential adult natural spawners, but no fewer than 40,700 naturally spawning adults in any one year. Brood escapement rate must average at least 32% over the long-term, but an individual brood may vary from this range to achieve the required tribal/nontribal annual allocation. Natural area spawners to maximize catch estimated at 40,700 adults (STT 2005). | 40,700 | 30,525 | 71% (STT 2005) | Based on F _{ABC} and annual ocean abundance. F _{ABC} is F _{MSY} reduced by Tier 1 (5%) uncertainty |
| Klamath River - Spring | Undefined | Undefined | Undefined | Undefined | Component stock of SONC complex; ACL indicator stock is KRFC |
| Smith River | Undefined | Undefined | Undefined | Undefined | |
| Southern Oregon | Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982) measured by 60-90 fish per mile in index streams. ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council. | 60 fish per mile in index streams | 30 fish per mile in index streams | Undefined | |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 2 of 6)

| CHINOOK | | | | | | |
|--|--|---|-----------------------------------|-----------------------------------|-----------------------------|--|
| Stocks In The Fishery | Conservation Objective | | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| Central and Northern Oregon | Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982) measured by 60-90 fish per mile in index streams. ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council. | | 60 Fish per mile in index streams | 30 Fish per mile in index streams | Undefined | Component stock(s) of FNMC complex; international exception applies, ACLs are not applicable |
| Willapa Bay Fall | Undetermined in FMP. WDFW spawning escapement objective of 4,350. | | 3,393 | 1,697 | 78% Proxy (SAC 2011) | |
| Grays Harbor Fall Indicator stock for the Far North Migrating Coastal (FNMC) Chinook stock complex | 14,600 natural adult spawners--MSP based on full seeding of spawning and rearing habitat (WDF 1979). | Annual natural spawning escapement targets may vary from FMP conservation objectives if agreed to by WDFW and treaty tribes under the provisions of <i>Hoh v. Baldrige</i> and subsequent U.S. District Court orders. | 11,388 | 5,694 | 78% Proxy (SAC 2011) | FNMC complex; international exception applies, ACLs are not applicable. |
| Queets Fall Indicator stock for the FNMC Chinook stock complex | Manage terminal fisheries for 40% harvest rate, but no less than 2,500 natural adult spawners, the MSY level estimated by Cooney (1984). | | 2,500 | 1,250 | 87% (Cooney 1984) | |
| Hoh Fall Indicator stock for the FNMC Chinook stock complex | Manage terminal fisheries for 40% harvest rate, but no less than 1,200 natural adult spawners, the MSY level estimated by Cooney (1984). | | 1,200 | 600 | 90% (Cooney 1984) | |
| Quillayute Fall Indicator stock for the FNMC Chinook stock complex | Manage terminal fisheries for 40% harvest rate, but no less than 3,000 natural adult spawners, the MSY level estimated by Cooney (1984). | | 3,000 | 1,500 | 87% (Cooney 1984) | |
| Hoko Summer/Fall Indicator stock for the FNMC Chinook stock complex | 850 natural adult spawners, the MSP level estimated by Ames and Phinney (1977). May include adults used for supplementation program. | | 850 | 425 | 78% Proxy (SAC 2011) | |
| Grays Harbor Spring | 1,400 natural adult spawners. | | 1,400 | 700 | 78% Proxy (SAC 2011) | FNMC complex; international exception applies, ACLs are not applicable. |
| Queets Sp/Su | Manage terminal fisheries for 30% harvest rate, but no less than 700 natural adult spawners. | | 700 | 350 | 78% Proxy (SAC 2011) | |
| Hoh Spring/Summer | Manage terminal fisheries for 31% harvest rate, but no less than 900 natural adult spawners. | | 900 | 450 | 78% Proxy (SAC 2011) | |
| Quillayute Spring/Summer | 1,200 natural adult spawners for summer component (MSY). | | 1,200 | 600 | Undefined | |
| Willapa Bay Fall (hatchery) | 8,200 adult return to hatchery. WDFW spawning escapement objective of 9,800 hatchery spawners. | | Not applicable to hatchery stocks | | | |
| Quinalt Fall (hatchery) | Hatchery production. | | | | | |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 3 of 6)

| CHINOOK | | | | | |
|---|--|--------------------------------------|------------------------------------|--------------------------------------|---|
| Stocks In The Fishery | Conservation Objective | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| North Lewis River Fall | NMFS consultation standard/recovery plan. Mclsaac (1990) stock-recruit analysis supports MSY objective of 5,700 natural adult spawners. | 5,700 | ESA consultation standard applies. | 76% | ESA consultation standard applies. |
| Snake River Fall | NMFS consultation standard/recovery plan. No more than 70.0% of 1988-1993 base period AEQ exploitation rate for all ocean fisheries. | Undefined | | Undefined | |
| Upper Willamette Spring | NMFS consultation standard/recovery plan. Not applicable for ocean fisheries. | Undefined | | Undefined | |
| Columbia Upper River Spring | NMFS consultation standard/recovery plan. Not applicable for ocean fisheries. | Undefined | | Undefined | |
| Snake River - Spring/Summer | NMFS consultation standard/recovery plan. Not applicable for ocean fisheries. | Undefined | | Undefined | |
| Columbia Lower River Hatchery - Fall | 12,600 adults for hatchery egg-take. | Not applicable to hatchery stocks | | | |
| Columbia Lower River Hatchery Spring | 2,700 adults to meet Cowlitz, Kalama, and Lewis Rivers broodstock needs. | | | | |
| Columbia Mid-River Bright Hatchery Fall | 4,700 adults for Bonneville Hatchery and 2,000 for Little White Salmon Hatchery egg-take. | | | | |
| Columbia Spring Creek Hatchery Fall | 7,000 adults to meet hatchery egg-take goal. | | | | |
| Columbia Upper River Bright Fall | 40,000 natural bright adults above McNary Dam (MSY proxy adopted in 1984 based on CRFMP). The management goal has been increased to 60,000 by Columbia River managers in recent years. | 39,625 (Langness and Reidinger 2003) | 19,812 | 85.91% (Langness and Reidinger 2003) | FNMC complex; international exception applies, ACLs are not applicable. |
| Columbia Upper River Summer | Hold ocean fishery impacts at or below base period; recognize CRFMP objective - MSY proxy of 80,000 to 90,000 adults above Bonneville Dam, including both Columbia and Snake River stocks (state and tribal management entities considering separate objectives for these stocks). | 12,143 (CTC 1999) | 6,071 | 75% (CTC 1999) | |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 4 of 6)

| CHINOOK | | | | | | |
|--|---|---|------------------|-----------------------------------|--------------------------|------------------------------------|
| Stocks In The Fishery | Conservation Objective | | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| Eastern Strait of Juan de Fuca Summer/Fall | NMFS consultation standard/recovery plan. No more than 10.0% Southern U.S. (SUS) Rebuilding Exploitation Rate (RER) for the Elwha River and for the Dungeness River. 2011 comanagers Resource Management Plan (RMP) | Annual natural spawning escapement targets may vary from FMP conservation objectives if agreed to by WDFW and treaty tribes under the provisions of U.S. v. Washington and subsequent U.S. District Court orders. | Undefined | ESA consultation standard applies | Undefined | ESA Consultation standard applies. |
| Skokomish Summer/Fall | NMFS consultation standard/recovery plan. No more than 50.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Mid Hood Canal Summer/Fall | NMFS consultation standard/recovery plan. No more than 15.0% preterminal SUS CERC. 2011 comanagers RMP | | Undefined | | Undefined | |
| Nooksack Spring early | NMFS consultation standard/recovery plan. No more than 7.0% SUS CERC. 2011 comanagers RMP | | Undefined | | Undefined | |
| Skagit Summer/Fall | NMFS consultation standard/recovery plan. No more than 50.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Skagit Spring | NMFS consultation standard/recovery plan. No more than 38.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Stillaguamish Summer/Fall | NMFS consultation standard/recovery plan. No more than 25.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Snohomish Summer/Fall | NMFS consultation standard/recovery plan. No more than 15.0% SUS RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Cedar River Summer/Fall | NMFS consultation standard/recovery plan. No more than 20.0% SUS RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| White River Spring | NMFS consultation standard/recovery plan. No more than 20.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Green River Summer/Fall | NMFS consultation standard/recovery plan. No more than 15.0% preterminal SUS RER, at least 5,800 adult spawners. | | Undefined | | Undefined | |
| Nisqually River Summer/Fall | NMFS consultation standard/recovery plan. No more than 65.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |
| Puyallup Summer/Fall | NMFS consultation standard/recovery plan. No more than 50.0% total RER. 2011 comanagers RMP | | Undefined | | Undefined | |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 5 of 6)

| COHO | | | | | | |
|--|--|-----------|-----------------------------------|-----------------------------------|--------------------------|------------------------------------|
| Stocks In The Fishery | Conservation Objective | | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| Central California Coast ESA Threatened | NMFS ESA consultation standard/recovery plan: No retention of coho south of the OR/CA border. | | Undefined | ESA consultation standard applies | Undefined | ESA consultation standard applies. |
| Southern Oregon/Northern California Coast ESA Threatened | NMFS ESA consultation standard/recovery plan: No more than a 13.0% AEQ exploitation rate in ocean fisheries on Rogue/Klamath hatchery coho. | | Undefined | | Undefined | |
| Oregon Coastal Natural ESA Threatened | NMFS ESA consultation standard/recovery plan: Total AEQ exploitation rate limit based on parental seeding level and marine survival matrix in FMP Table 3-2. | | Undefined | | Undefined | |
| Lower Columbia Natural ESA Threatened | NMFS ESA consultation standard/recovery plan: AEQ exploitation rate limit on ocean and mainstem Columbia fisheries indentified in annual NMFS guidance. | | Undefined | | Undefined | |
| Oregon Coast Hatchery | Hatchery production. | | Not applicable to hatchery stocks | | | |
| Columbia River Late Hatchery | Hatchery rack return goal of 14,200 adults. | | | | | |
| Columbia River Early Hatchery | Hatchery rack return goal of 6,200 adults. | | | | | |
| Willapa Bay - Hatchery | Hatchery rack return goal of 6,100 adults. | | | | | |
| Quinalt - Hatchery | Hatchery production. | | | | | |
| Quillayute - Summer Hatchery | Hatchery production. | | | | | |
| South Puget Sound Hatchery | Hatchery rack return goal of 52,000 adults. | | | | | |
| Willapa Bay Natural | Undefined | Undefined | Undefined | Undefined | Undefined | Undefined |

TABLE A-1. Conservation objectives and reference points governing harvest control rules and status determination criteria for salmon stocks and stock complexes in the Pacific Coast Salmon FMP. (Page 6 of 6)

| COHO | | | | | | |
|---------------------------|---|---|---|-----------------------------------|--|---|
| Stocks In The Fishery | Conservation Objective | | S _{MSY} | MSST | MFMT (F _{MSY}) | ACL |
| Grays Harbor | 35,400 natural adult spawners (MSP based on WDF [1979]) | Annual natural spawning escapement targets may vary from FMP conservation objectives if agreed to by WDFW and treaty tribes under the provisions of Hoh v. Baldrige, U.S. v. Washington, or subsequent U.S. District Court orders | 24,426 S _{MSP} (FMP) *F _{SMY} (SAC 2010b) | 18,320 (Johnstone et al. 2011) | MFMT=65% (Johnstone et al. 2011) F _{MSY} =69% (SAC 2011) | International exception applies, ACLs are not applicable. |
| Queets | MSY range of 5,800 to 14,500 natural adult spawners (Lestelle et al 1984) | | 5,800 (Johnston et al. 2011) | 4,350 (Johnstone et al. 2011) | MFMT=65% (Johnstone et al. 2011) F _{MSY} =68% (SAC 2011) | |
| Hoh | MSY range of 2,000 to 5,000 natural adult spawners (Lestelle et al. 1984) | | 2,520 (SAC 2010b) | 1,890 S _{MSY} *0.75 | MFMT=65% (Johnstone et al. 2011) F _{MSY} =69% (SAC 2011) | |
| Quillayute - Fall | MSY range of 6,300 to 15,800 natural adult spawners (Lestelle et al. 1984) | | 6,300 (Johnston et al. 2011) | 4,725 (Johnstone et al. 2011) | MFMT Undefined; F _{MSY} =59% (SAC 2011) | |
| Strait of Juan de Fuca | Total allowable MSY exploitation rate of: 0.60 for ocean age-3 abundance > 27,445; 0.40 for ocean age-3 abundance >11,679 and ≤27,445; 0.20 for ocean age-3 abundance ≤11,679 | | 11,000 (Bowhay et al. 2009) | 7,000 (Bowhay et al. 2009) | 60% (Bowhay et al. 2009) | |
| Hood Canal | Total allowable MSY exploitation rate of: 0.65 for ocean age-3 abundance > 41,000; 0.45 for ocean age-3 abundance >19,545 and ≤41,000; 0.20 for ocean age-3 abundance ≤19,545 | | 14,350 (Bowhay et al. 2009) | 10,750 (Bowhay et al. 2009) | 65% (Bowhay et al. 2009) | |
| Skagit | Total allowable MSY exploitation rate of: 0.60 for ocean age-3 abundance > 62,500; 0.35 for ocean age-3 abundance >22,857 and ≤62,500; 0.20 for ocean age-3 abundance ≤22,857 | | 25,000 (Bowhay et al. 2009) | 14,857 (Bowhay et al. 2009) | 60% (Bowhay et al. 2009) | |
| Stillaguamish | Total allowable MSY exploitation rate of: 0.50 for ocean age-3 abundance > 20,000; 0.35 for ocean age-3 abundance >9,385 and ≤20,000; 0.20 for ocean age-3 abundance ≤9,385 | | 10,000 (Bowhay et al. 2009) | 6,100 (Bowhay et al. 2009) | 50% (Bowhay et al. 2009) | |
| Snohomish | Total allowable MSY exploitation rate of: 0.60 for ocean age-3 abundance > 125,000; 0.40 for ocean age-3 abundance >51,667 and ≤125,000; 0.20 for ocean age-3 abundance ≤51,667 | 50,000 (Bowhay et al. 2009) | 31,000 (Bowhay et al. 2009) | 60% (Bowhay et al. 2009) | | |
| PINK (odd-numbered years) | | | | | | |
| Puget Sound | 900,000 natural spawners or consistent with provisions of the Pacific Salmon Treaty (Fraser River Panel). | | 900,000 | 450,000 | Undefined | International exception applies, ACLs are not applicable. |

TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13.

| | | MARINE SURVIVAL INDEX (based on return of jacks per hatchery smolt) | | | |
|---|---|--|------------------------------|---------------------------|--------------|
| | | Low (<0.0009) | Medium (0.0009 to 0.0034) | High (>0.0034) | |
| | | Allowable Total Fishery Impact Rate | | | |
| PARENT SPAWNER STATUS | | | | | |
| High: | Parent spawners achieved Level #2 rebuilding criteria; grandparent spawners achieved Level #1 | $\leq 15\%$ | $\leq 30\%$ ^{a/} | $\leq 35\%$ ^{a/} | |
| Medium: | Parent spawners achieved Level #1 or greater rebuilding criteria | $\leq 15\%$ | $\leq 20\%$ ^{a/} | $\leq 25\%$ ^{a/} | |
| Low: | Parent spawners less than Level #1 rebuilding criteria | $\leq 15\%$ | $\leq 15\%$ | $\leq 15\%$ | |
| | | $\leq 10\text{-}13\%$ ^{b/} | | | |
| | | | | | |
| OCN Coho Spawners by Stock Component | | | | | |
| Rebuilding Criteria | Northern | North-Central | South-Central | Southern | Total |
| Full Seeding at Low Marine Survival: | 21,700 | 55,000 | 50,000 | 5,400 | 132,100 |
| Level #2 (75% of full seeding): | 16,400 | 41,300 | 37,500 | 4,100 | 99,300 |
| Level #1 (50% of full seeding): | 10,900 | 27,500 | 25,000 | 2,700 | 66,100 |
| 38% of Level #1 (19% of full seeding): | 4,100 | 10,500 | 9,500 | 1,000 | 25,100 |
| | | | | | |
| Stock Component (Boundaries) | Full Seeding of Major Basins at Low Marine Survival (Number of Adult Spawners) | | | | |
| Northern: (Necanicum River to Neskowin Creek) | Nehalem | Tillamook | Nestucca | Ocean Tribs. | |
| | 17,500 | 2,000 | 1,800 | 400 | |
| North-Central: (Salmon River to Siuslaw River) | Siletz | Yaquina | Alsea | Siuslaw | Ocean Tribs. |
| | 4,300 | 7,100 | 15,100 | 22,800 | 5,700 |
| South-Central: (Siltcoos River to Sixes River) | Umpqua | Coos | Coquille | Coastal Lakes | |
| | 29,400 | 7,200 | 5,400 | 8,000 | |
| Southern: (Elk River to Winchuck River) | Rogue | | | | |
| | 5,400 | | | | |

a/ When a stock component achieves a medium or high parent spawner status under a medium or high marine survival index, but a major basin within the stock component is less than 10% of full seeding, (1) the parent spawner status will be downgraded one level to establish the allowable fishery impact rate for that component, and (2) no coho-directed harvest impacts will be allowed within that particular basin.

b/ This exploitation rate criteria applies when (1) parent spawners are less than 38% of the Level #1 rebuilding criteria, or (2) marine survival conditions are projected to be at an extreme low as in 1994-1996 (<0.0006 jack per hatchery smolt). If parent spawners decline to lower levels than observed through 1998, rates of less than 10% would be considered, recognizing that there is a limit to further bycatch reduction opportunities.

TABLE A-3. Fishery impact rate criteria for OCN coho stock components based on the harvest matrix resulting from the OCN work group 2000 review of Amendment 13.

| Parent Spawner Status ^{a/} | Marine Survival Index (based on return of jacks per hatchery smolt) | | | | | | |
|---|--|---------------------------------|-------------------------------------|-----------------------|------------------------------|---------------------|---------------------|
| | Extremely Low (<0.0008) | Low (0.0008 to 0.0014) | Medium (>0.0014 to 0.0040) | High (>0.0040) | | | |
| High Parent Spawners $> 75\%$ of full seeding | E $\leq 8\%$ | J $\leq 15\%$ | O $\leq 30\%$ | T $\leq 45\%$ | | | |
| Medium Parent Spawners $> 50\%$ & $\leq 75\%$ of full seeding | D $\leq 8\%$ | I $\leq 15\%$ | N $\leq 20\%$ | S $\leq 38\%$ | | | |
| Low Parent Spawners $> 19\%$ & $\leq 50\%$ of full seeding | C $\leq 8\%$ | H $\leq 15\%$ | M $\leq 15\%$ | R $\leq 25\%$ | | | |
| Very Low Parent Spawners > 4 fish per mile & $\leq 19\%$ of full seeding | B $\leq 8\%$ | G $\leq 11\%$ | L $\leq 11\%$ | Q $\leq 11\%$ | | | |
| Critical ^{b/} Parental Spawners ≤ 4 fish per mile | A $0 - 8\%$ | F $0 - 8\%$ | K $0 - 8\%$ | P $0 - 8\%$ | | | |
| Sub-aggregate and Basin Specific Spawner Criteria Data | | | | | | | |
| Sub-aggregate | Miles of Available Spawning Habitat | 100% of Full Seeding | "Critical" | | Very Low, Low, Medium & High | | |
| | | | 4 Fish per Mile | 12% of Full Seeding | 19% of Full Seeding | 50% of Full Seeding | 75% of full Seeding |
| Northern | 899 | 21,700 | 3,596 | NA | 4,123 | 10,850 | 16,275 |
| North - Central | 1,163 | 55,000 | 4,652 | NA | 10,450 | 27,500 | 41,250 |
| South - Central | 1,685 | 50,000 | 6,740 | NA | 9,500 | 25,000 | 37,500 |
| Southern | 450 | 5,400 | NA | 648 | 1,026 | 2,700 | 4,050 |
| Coastwide Total | 4,197 | 132,100 | 15,636 | | 25,099 | 66,050 | 99,075 |

a/ Parental spawner abundance status for the OCN aggregate assumes the status of the weakest sub-aggregate.

b/ "Critical" parental spawner status is defined as 4 fish per mile for the Northern, North-Central, and South-Central subaggregates. Because the ratio of high quality spawning habitat to total spawning habitat in the Rogue River Basin differs significantly from the rest of the basins on the coast, the spawner density of 4 fish per mile does not represent "Critical" status for that basin. Instead, "Critical" status for the Rogue Basin (Southern Sub-aggregate) is estimated as 12% of full seeding of high quality

TABLE A-4. Council adopted management objectives for Puget Sound natural coho management units, expressed as exploitation rate ceilings for critical, low and normal abundance based status categories, with runsize breakpoints (abundances expressed as ocean-age 3).

| Status | Management Unit | | | | |
|---------------------------------|------------------------|------------|--------|---------------|-----------|
| | Strait of Juan de Fuca | Hood Canal | Skagit | Stillaguamish | Snohomish |
| Critical/Low runsize breakpoint | 11,679 | 19,545 | 22,857 | 9,385 | 51,667 |
| Critical exploitation rate | 0.20 | 0.20 | 0.20 | 0.20 | 0.20 |
| Low/normal runsize breakpoint | 27,445 | 41,000 | 62,500 | 20,000 | 125,000 |
| Low exploitation rate | 0.40 | 0.45 | 0.35 | 0.35 | 0.40 |
| Normal exploitation rate | 0.60 | 0.65 | 0.60 | 0.50 | 0.60 |

TABLE A-5. Council recommended management objectives for Lower Columbia River natural tule Chinook, expressed as exploitation rate ceilings for abundance based status categories, with runsize forecast bins expressed as adult river mouth return forecasts of Lower Columbia River hatchery tule Chinook.

| Runsize Forecast Bins | <30,000 | 30,000 to 40,000 | 40,000 to 85,000 | >85,000 |
|---------------------------|---------|------------------|------------------|---------|
| Maximum Exploitation Rate | 0.30 | 0.35 | 0.38 | 0.41 |

APPENDIX B

SALMON HARVEST ALLOCATION SCHEDULES

TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| HARVEST ALLOCATION - SECTION 5.3 OF THE PACIFIC COAST SALMON FISHERY | |
| MANAGEMENT PLAN..... | 107 |
| 5.3 ALLOCATION..... | 107 |
| 5.3.1 Commercial (Non-Tribal) and Recreational Fisheries North of Cape Falcon | 107 |
| 5.3.1.1 Goal, Objectives, and Priorities..... | 107 |
| 5.3.1.2 Allocation Schedule Between Gear Types | 108 |
| 5.3.1.3 Recreational Subarea Allocations | 109 |
| 5.3.2 Commercial and Recreational Fisheries South of Cape Falcon..... | 110 |
| 5.3.3 Tribal Indian Fisheries | 113 |
| 5.3.3.1 California..... | 113 |
| 5.3.3.2 Columbia River | 113 |
| 5.3.3.3 U.S. v. Washington Area..... | 113 |
| MEASURES TO MANAGE THE HARVEST - SECTION 5.3 OF THE PACIFIC COAST | |
| SALMON FISHERY MANAGEMENT PLAN | 112 |
| 6.5 SEASONS AND QUOTAS..... | 115 |
| 6.5.2 Procedures for Calculating Seasons..... | 115 |
| 6.5.3 Species-Specific and Other Selective Fisheries | 115 |
| 6.5.3.1 Guidelines..... | 115 |
| 6.5.3.2 Selective Fisheries Which May Change Allocation Percentages North of Cape Falcon | 115 |
| 6.5.4 Procedures for Calculating Quotas | 116 |

5.3 ALLOCATION

“A Conservation and management measures shall not discriminate between residents of different states. If it becomes necessary to allocate or assign fishing privileges among various United States fishermen, such allocation shall be (A) fair and equitable to all such fishermen; (B) reasonably calculated to promote conservation; and (C) carried out in such manner that no particular individual, corporation, or other entity acquires an excessive share of such privileges.”

Magnuson-Stevens Act, National Standard 4

Harvest allocation is required when the number of fish is not adequate to satisfy the perceived needs of the various fishing industry groups and communities, to divide the catch between non-Indian ocean and inside fisheries and among ocean fisheries, and to provide Federally recognized treaty Indian fishing opportunity. In allocating the resource between ocean and inside fisheries, the Council considers both in-river harvest and spawner escapement needs. The magnitude of in-river harvest is determined by the states in a variety of ways, depending upon the management area. Some levels of in-river harvests are designed to accommodate federally recognized in-river Indian fishing rights, while others are established to allow for non-Indian harvests of historical magnitudes. Several fora exist to assist this process on an annual basis. The North of Cape Falcon Forum, a state and tribal sponsored forum, convenes the pertinent parties during the Council's preseason process to determine allocation and conservation recommendations for fisheries north of Cape Falcon. The individual states also convene fishery industry meetings to coordinate their input to the Council.

5.3.1 Commercial (Non-Tribal) and Recreational Fisheries North of Cape Falcon

5.3.1.1 Goal, Objectives, and Priorities

Harvest allocations will be made from a total allowable ocean harvest, which is maximized to the largest extent possible but still consistent with PST and treaty-Indian obligations, state fishery needs, and spawning escapement requirements, including consultation standards for stocks listed under the ESA. The Council shall make every effort to establish seasons and gear requirements that provide troll and recreational fleets a reasonable opportunity to catch the available harvest. These may include single-species directed fisheries with landing restrictions for other species.

The goal of allocating ocean harvest north of Cape Falcon is to achieve, to the greatest degree possible, the objectives for the commercial and recreational fisheries as follows:

- Provide recreational opportunity by maximizing the duration of the fishing season while minimizing daily and area closures and restrictions on gear and daily limits.
- Maximize the value of the commercial harvest while providing fisheries of reasonable duration.

The priorities listed below will be used to help guide establishment of the final harvest allocation while meeting the overall commercial and recreational fishery objectives.

At total allowable harvest levels up to 300,000 coho and 100,000 Chinook:

- Provide coho to the recreational fishery for a late June through early September all-species season. Provide Chinook to allow (1) access to coho and, if possible, (2) a minimal Chinook-only fishery prior to the all-species season. Adjust days per week and/or institute area restrictions to stabilize season duration.

- Provide Chinook to the troll fishery for a May and early June Chinook season and provide coho to (1) meet coho hooking mortality in June where needed and (2) access a pink salmon fishery in odd years. Attempt to ensure that part of the Chinook season will occur after June 1.

At total allowable harvest levels above 300,000 coho and above 100,000 Chinook:

- Relax any restrictions in the recreational all-species fishery and/or extend the all-species season beyond Labor Day as coho quota allows. Provide Chinook to the recreational fishery for a Memorial Day through late June Chinook-only fishery. Adjust days per week to ensure continuity with the all-species season.
- Provide coho for an all-salmon troll season in late summer and/or access to a pink fishery. Leave adequate Chinook from the May through June season to allow access to coho.

5.3.1.2 Allocation Schedule Between Gear Types

Initial commercial and recreational allocation will be determined by the schedule of percentages of total allowable harvest as follows:

TABLE 5-1. Initial commercial/recreational harvest allocation schedule north of Cape Falcon.

| Coho | | | Chinook | | |
|-----------------------------------|--------------------------|--------------|-----------------------------------|--------------------------|--------------|
| Harvest (thousands of fish) | Percentage ^{a/} | | Harvest (thousands of fish) | Percentage ^{a/} | |
| | Troll | Recreational | | Troll | Recreational |
| 0-300 | 25 | 75 | 0-100 | 50 | 50 |
| >300 | 60 | 40 | >100-150 | 60 | 40 |
| | | | >150 | 70 | 30 |

a/ The allocation must be calculated in additive steps when the harvest level exceeds the initial tier.

This allocation schedule should, on average, allow for meeting the specific fishery allocation priorities described above. The initial allocation may be modified annually by preseason and inseason trades to better achieve (1) the commercial and recreational fishery objectives and (2) the specific fishery allocation priorities. The final preseason allocation adopted by the Council will be expressed in terms of quotas, which are neither guaranteed catches nor inflexible ceilings. Only the total ocean harvest quota is a maximum allowable catch.

To provide flexibility to meet the dynamic nature of the fisheries and to assure achievement of the allocation objectives and fishery priorities, deviations from the allocation schedule will be allowed as provided below and as described in Section 6.5.3.2 for certain selective fisheries.

1. Preseason species trades (Chinook and coho) that vary from the allocation schedule may be made by the Council based upon the recommendation of the pertinent recreational and commercial SAS representatives north of Cape Falcon. The Council will compare the socioeconomic impacts of any such recommendation to those of the standard allocation schedule before adopting the allocation that best meets FMP management objectives.
2. Inseason transfers, including species trades of Chinook and coho, may be permitted in either direction between recreational and commercial fishery allocations to allow for uncatchable fish in one fishery to be reallocated to the other. Fish will be deemed "uncatchable" by a respective commercial or recreational fishery only after considering all possible annual management actions to allow for their harvest which meet framework harvest management objectives, including single species or exclusive

registration fisheries. Implementation of inseason transfers will require (1) consultation with the pertinent recreational and commercial SAS members and the STT, and (2) a clear establishment of available fish and impacts from the transfer.

3. An exchange ratio of four coho to one Chinook shall be considered a desirable guideline for preseason trades. Deviations from this guideline should be clearly justified. Inseason trades and transfers may vary to meet overall fishery objectives. (The exchange ratio of four coho to one Chinook approximately equalizes the species trade in terms of average ex-vessel values of the two salmon species in the commercial fishery. It also represents an average species catch ratio in the recreational fishery.)
4. Any increase or decrease in the recreational or commercial total allowable catch (TAC), resulting from an inseason restructuring of a fishery or other inseason management action, does not require reallocation of the overall north of Cape Falcon non-Indian TAC.
5. The commercial TACs of Chinook and coho derived during the preseason allocation process may be varied by major subareas (i.e., north of Leadbetter Point and south of Leadbetter Point) if there is a need to do so to decrease impacts on weak stocks. Deviations in each major subarea will generally not exceed 50 percent of the TAC of each species that would have been established without a geographic deviation in the distribution of the TAC. Deviation of more than 50 percent will be based on a conservation need to protect weak stocks and will provide larger overall harvest for the entire fishery north of Cape Falcon than would have been possible without the deviation. In addition, the actual harvest of coho may deviate from the initial allocation as provided in Section 6.5.3.2 for certain selective fisheries.
6. The recreational TACs of Chinook and coho derived during the preseason allocation process will be distributed among four major recreational port areas as described for coho and Chinook distribution in Section 5.3.1.3. The Council may deviate from subarea quotas (1) to meet recreational season objectives based on agreement of representatives of the affected ports and/or (2) in accordance with Section 6.5.3.2 with regard to certain selective fisheries. Additionally, based on the recommendations of the SAS members representing the ocean sport fishery north of Cape Falcon, the Council will include criteria in its preseason salmon management recommendations to guide any inseason transfer of coho among the recreational subareas to meet recreational season duration objectives. Inseason redistributions of quotas within the recreational fishery or the distribution of allowable coho catch transfers from the commercial fishery may deviate from the preseason distribution.

5.3.1.3 Recreational Subarea Allocations

Coho

The north of Cape Falcon preseason recreational TAC of coho will be distributed to provide 50 percent to the area north of Leadbetter Point and 50 percent to the area south of Leadbetter Point. The distribution of the allocation north of Leadbetter point will vary, depending on the existence and magnitude of an inside fishery in Area 4B, which is served by Neah Bay.

In years with no Area 4B fishery, the distribution of coho north of Leadbetter Point (50 percent of the total recreational TAC) will be divided to provide 74 percent to the area between Leadbetter Point and the Queets River (Westport), 5.2 percent to the area between Queets River and Cape Flattery (La Push), and 20.8 percent to the area north of the Queets River (Neah Bay). In years when there is an Area 4B (Neah Bay) fishery under state management, the allocation percentages north of Leadbetter Point will be modified to maintain more equitable fishing opportunity among the ports by decreasing the ocean harvest share for Neah Bay. This will be accomplished by adding 25 percent of the numerical value of the Area

4B fishery to the recreational TAC north of Leadbetter Point prior to calculating the shares for Westport and La Push. The increase to Westport and La Push will be subtracted from the Neah Bay ocean share to maintain the same total harvest allocation north of Leadbetter Point. Table 5-2 displays the resulting percentage allocation of the total recreational coho catch north of Cape Falcon among the four recreational port areas (each port area allocation will be rounded to the nearest hundred fish, with the largest quotas rounded downward if necessary to sum to the TAC).

TABLE 5-2. Percentage allocation of total allowable coho harvest among the four recreational port areas north of Cape Falcon.^{a/}

| Port Area | Without Area 4B Add-on | With Area 4B Add-on | |
|----------------|------------------------|---------------------|-----------------------------------|
| Columbia River | 50.0% | 50.0% | |
| Westport | 37.0% | 37.0% | plus 17.3% of the Area 4B add-on |
| La Push | 2.6% | 2.6% | plus 1.2% of the Area 4B add-on |
| Neah Bay | 10.4% | 10.4% | minus 18.5% of the Area 4B add-on |

a/ The Council may deviate from these percentages as described under #6 in Section 5.3.1.2.

TABLE 5-3. Example distributions of the recreational coho TAC north of Leadbetter Point.

| Sport TAC North of Cape Falcon | Without Area 4B Add-On | | | | With Area 4B Add-On ^{a/} | | | | | |
|---|------------------------|----------|---------|-------------|-----------------------------------|----------|---------|--------|--------------------|--------|
| | Columbia River | Westport | La Push | Neah Bay | Columbia River | Westport | La Push | Ocean | Neah Bay Add-on | Total |
| 50,000 | 25,000 | 18,500 | 1,300 | 5,200 | 25,000 | 19,900 | 1,400 | 3,700 | 8,000 | 11,700 |
| 150,000 | 75,000 | 55,500 | 3,900 | 15,600 | 75,000 | 57,600 | 4,000 | 13,600 | 12,000 | 25,600 |
| 300,000 | 150,000 | 111,000 | 7,800 | 31,200 | 150,000 | 114,500 | 8,000 | 27,500 | 20,000 | 47,500 |

a/ The add-on levels are merely examples. The actual numbers in any year would depend on the particular mix of stock abundances and season determinations.

Chinook

Subarea distributions of Chinook will be managed as guidelines and shall be calculated by the STT with the primary objective of achieving all-species fisheries without imposing Chinook restrictions (i.e., area closures or bag limit reductions). Chinook in excess of all-species fisheries needs may be utilized by directed Chinook fisheries north of Cape Falcon or by negotiating a Chinook/coho trade with another fishery sector.

Inseason management actions may be taken by the NMFS NW Regional Administrator to assure that the primary objective of the Chinook harvest guidelines for each of the four recreational subareas north of Cape Falcon are met. Such actions might include: closure from 0 to 3, or 0 to 6, or 3 to 200, or 5 to 200 nautical miles from shore; closure from a point extending due west from Tatoosh Island for 5 miles, then south to a point due west of Umatilla Reef Buoy, then due east to shore; closure from North Head at the Columbia River mouth north to Leadbetter Point; change species that may be landed; or other actions as prescribed in the annual regulations.

5.3.2 Commercial and Recreational Fisheries South of Cape Falcon

The allocation of allowable ocean harvest of coho salmon south of Cape Falcon has been developed to provide a more stable recreational season and increased economic benefits of the ocean salmon fisheries at varying stock abundance levels. When coupled with various recreational harvest reduction measures or the timely transfer of unused recreational allocation to the commercial fishery, the allocation schedule is designed to help secure recreational seasons extending at least from Memorial Day through Labor Day

when possible, assist in maintaining commercial markets even at relatively low stock sizes, and fully utilize available harvest. Total ocean catch of coho south of Cape Falcon will be treated as a quota to be allocated between troll and recreational fisheries as provided in Table 5-4.

(Note: The allocation schedule provides guidance only when coho abundance permits a directed coho harvest, not when the allowable impacts are insufficient to allow coho retention south of Cape Falcon. At such low levels, allocation of the allowable impacts will be accomplished during the Council's preseason process.)

TABLE 5-4. Allocation of allowable ocean harvest of coho salmon (thousands of fish) south of Cape Falcon.^{a/}

| Total Allowable Ocean Harvest | Recreational Allocation | | Commercial Allocation | |
|-------------------------------|-------------------------|-------------------|-----------------------|------------------|
| | Number | Percentage | Number | Percentage |
| <100 | #100 ^{b/c/} | 100 ^{b/} | b/ ^{b/} | b/ ^{b/} |
| 200 | 167 ^{b/c/} | 84 ^{b/} | 33 ^{b/} | 17 ^{b/} |
| 300 | 200 | 67 | 100 | 33 |
| 350 | 217 | 62 | 133 | 38 |
| 400 | 224 | 56 | 176 | 44 |
| 500 | 238 | 48 | 262 | 52 |
| 600 | 252 | 42 | 348 | 58 |
| 700 | 266 | 38 | 434 | 62 |
| 800 | 280 | 35 | 520 | 65 |
| 900 | 290 | 32 | 610 | 68 |
| 1,000 | 300 | 30 | 700 | 70 |
| 1,100 | 310 | 28 | 790 | 72 |
| 1,200 | 320 | 27 | 880 | 73 |
| 1,300 | 330 | 25 | 970 | 75 |
| 1,400 | 340 | 24 | 1,060 | 76 |
| 1,500 | 350 | 23 | 1,150 | 77 |
| 1,600 | 360 | 23 | 1,240 | 78 |
| 1,700 | 370 | 22 | 1,330 | 78 |
| 1,800 | 380 | 21 | 1,420 | 79 |
| 1,900 | 390 | 21 | 1,510 | 79 |
| 2,000 | 400 | 20 | 1,600 | 80 |
| 2,500 | 450 | 18 | 2,050 | 82 |
| 3,000 | 500 | 17 | 2,500 | 83 |

a/ The allocation schedule is based on the following formula: first 150,000 coho to the recreational base (this amount may be reduced as provided in footnote b); over 150,000 to 350,000 fish, share at 2:1, 0.667 to troll and 0.333 to recreational; over 350,000 to 800,000 the recreational share is 217,000 plus 14% of the available fish over 350,000; above 800,000 the recreational share is 280,000 plus 10% of the available fish over 800,000.

Note: The allocation schedule provides guidance only when coho abundance permits a directed coho harvest, not when the allowable impacts are insufficient to allow general coho retention south of Cape Falcon. At such low levels, allocation of the allowable impacts will be determined in the Council's preseason process. Deviations from the allocation may also be allowed to meet consultation standards for ESA-listed stocks (e.g., the 1998 biological opinion for California coastal coho requires no retention of coho in fisheries off California).

b/ If the commercial allocation is insufficient to meet the projected hook-and-release mortality associated with the commercial all-salmon-except-coho season, the recreational allocation will be reduced by the number needed to eliminate the deficit.

c/ When the recreational allocation is 167,000 coho or less, special allocation provisions apply to the recreational harvest distribution by geographic area (unless superseded by requirements to meet a consultation standard for ESA-listed stocks); see text of FMP as modified by Amendment 11 allocation provisions.

The allocation schedule is designed to give sufficient coho to the recreational fishery to increase the probability of attaining no less than a Memorial Day to Labor Day season as stock sizes increase. This increased allocation means that, in many years, actual catch in the recreational fishery may fall short of its allowance. In such situations, managers will make an inseason reallocation of unneeded recreational coho to the south of Cape Falcon troll fishery. The reallocation should be structured and timed to allow the commercial fishery sufficient opportunity to harvest any available reallocation prior to September 1, while still assuring completion of the scheduled recreational season (usually near mid-September) and, in any event, the continuation of a recreational fishery through Labor Day. This reallocation process will occur no later than August 15 and will involve projecting the recreational fishery needs for the remainder of the summer season. The remaining projected recreational catch needed to extend the season to its scheduled closing date will be a harvest guideline rather than a quota. If the guideline is met prior to Labor Day, the season may be allowed to continue if further fishing is not expected to result in any significant danger of impacting the allocation of another fishery or of failing to meet an escapement goal.

The allocation schedule is also designed to assure there are sufficient coho allocated to the troll fishery at low stock levels to ensure a full Chinook troll fishery. This hooking mortality allowance will have first priority within the troll allocation. If the troll allocation is insufficient for this purpose, the remaining number of coho needed for the estimated incidental coho mortality will be deducted from the recreational share. At higher stock sizes, directed coho harvest will be allocated to the troll fishery after hooking mortality needs for Chinook troll fishing have been satisfied.

The allowable harvest south of Cape Falcon may be further partitioned into subareas to meet management objectives of the FMP. Allowable harvests for subareas south of Cape Falcon will be determined by an annual blend of management considerations including:

1. abundance of contributing stocks
2. allocation considerations of concern to the Council
3. relative abundance in the fishery between Chinook and coho
4. escapement goals
5. maximizing harvest potential

Troll coho quotas may be developed for subareas south of Cape Falcon consistent with the above criteria. California recreational catches of coho, including projections of the total catch to the end of the season, would be included in the recreational allocation south of Cape Falcon, but the area south of the Oregon-California border would not close when the allocation is met; except as provided below when the recreational allocation is at 167,000 or fewer fish.

When the south of Cape Falcon recreational allocation is equal to or less than 167,000 coho:

1. The recreational fisheries will be divided into two major subareas, as listed in #2 below, with independent quotas (i.e., if one quota is not achieved or is exceeded, the underage or overage will not be added to or deducted from the other quota; except as provided under #3 below).
2. The two major recreational subareas will be managed within the constraints of the following impact quotas, expressed as a percentage of the total recreational allocation (percentages based on avoiding large deviations from the historical harvest shares):
 - a. Central Oregon (Cape Falcon to Humbug Mountain) - 70%
 - b. South of Humbug Mountain - 30%

In addition,

- (1) Horse Mountain to Point Arena will be managed for an impact guideline of 3 percent of the south of Cape Falcon recreational allocation, and
 - (2) there will be no coho harvest constraints south of Point Arena. However, the projected harvest in this area (which averaged 1,800 coho from 1986-1990) will be included in the south of Humbug Mountain impact quota.
3. Coho quota transfers can occur on a one-for-one basis between subareas if Chinook constraints preclude access to coho.

5.3.3 Tribal Indian Fisheries

5.3.3.1 California

On October 4, 1993 the Solicitor, Department of Interior, issued a legal opinion in which he concluded that the Yurok and Hoopa Valley Indian tribes of the Klamath River Basin have a federally protected right to the fishery resource of their reservations sufficient to support a moderate standard of living or 50 percent of the total available harvest of Klamath-Trinity basin salmon, whichever is less. The Secretary of Commerce recognized the tribes' federally reserved fishing right as applicable law for the purposes of the MSA (58 FR 68063, December 23, 1993). The Ninth Circuit Court of Appeals upheld the conclusion that the Hoopa Valley and Yurok tribes have a federally reserved right to harvest fish in Parravano v. Babbitt and Brown, 70 F.3d 539 (1995) (Cert. denied in Parravano v. Babbitt and Brown 110, S.Ct 2546 [1996]). The Council must recognize the tribal allocation in setting its projected escapement level for the Klamath River.

5.3.3.2 Columbia River

Pursuant to a September 1, 1983 Order of the U.S. District Court, the allocation of harvest in the Columbia River was established under the "Columbia River Fish Management Plan" which was implemented in 1988 by the parties of U.S. v. Oregon. This plan replaced the original 1977 plan (pages 16-20 of the 1978 FMP). Since the Columbia River Fishery Management Plan expired on December 31, 1998, fall Chinook in Columbia River fisheries were managed through 2007 under the guidance of annual management agreements among the U.S. v. Oregon parties. In 2008, a new 10 year management agreement was negotiated through the U.S. v. Oregon process, which included revisions to some in-river objectives. This most recent plan is the "2008-2017 U.S. v Oregon Management Agreement". The plan provides a framework within which the relevant parties may exercise their sovereign powers in a coordinated and systematic manner in order to protect, rebuild, and enhance upper Columbia River fish runs while providing harvest for both treaty Indian and non-Indian fisheries. The parties to the agreement are the United States, the states of Oregon, Washington, and Idaho, and four Columbia River treaty Indian tribes-Warm Springs, Yakama, Nez Perce, and Umatilla.

5.3.3.3 U.S. v. Washington Area

Treaty Indian tribes have a legal entitlement to the opportunity to take up to 50 percent of the harvestable surplus of stocks which pass through their usual and accustomed fishing areas. The treaty Indian troll harvest which would occur if the tribes chose to take their total 50 percent share of the weakest stock in the ocean, is computed with the current version of the Fishery Regulation Assessment Model (FRAM), assuming this level of harvest did not create conservation or allocation problems on other stocks. A quota may be established in accordance with the objectives of the relevant treaty tribes concerning allocation of the treaty Indian share to ocean and inside fisheries. The total quota does not represent a guaranteed ocean harvest, but a maximum allowable catch.

The requirement for the opportunity to take up to 50 percent of the harvestable surplus determines the treaty shares available to the inside/outside Indian and all-citizen fisheries. Ocean coho harvest ceilings off the Washington coast for treaty Indians and all-citizen fisheries are independent within the constraints that (1) where feasible, conservation needs of all stocks must be met; (2) neither group precludes the other from the opportunity to harvest its share, and; (3) allocation schemes may be established to specify outside/inside sharing for various stocks.

6.5 SEASONS AND QUOTAS

6.5.2 Procedures for Calculating Seasons

Seasons will be calculated using the total allowable ocean harvest determined by procedures described in Chapter 5, and further allocated to the commercial and recreational fishery in accordance with the allocation plan presented in Section 5.3, and after consideration of the estimated amount of effort required to catch the available fish, based on past seasons.

Recreational seasons will be established with the goal of encompassing Memorial Day and/or Labor Day weekends in the season, if feasible. Opening dates will be adjusted to provide reasonable assurance that the recreational fishery is continuous, minimizing the possibility of an in-season closure.

Criteria used to establish commercial seasons, in addition to the estimated allowable ocean harvests, the allocation plan, and the expected effort during the season, will be: (1) bycatch mortality; (2) size, poundage, and value of fish caught; (3) effort shifts between fishing areas; (4) harvest of pink salmon in odd-numbered years; and (5) protection for weak stocks when they frequent the fishing areas at various times of the year.

6.5.3 Species-Specific and Other Selective Fisheries

6.5.3.1 Guidelines

In addition to the all-species and single or limited species seasons established for the commercial and recreational fisheries, other species-limited fisheries, such as "ratio" fisheries and fisheries selective for marked or hatchery fish, may be adopted by the Council during the preseason regulatory process. In adopting such fisheries, the Council will consider the following guidelines:

1. Harvestable fish of the target species are available.
2. Harvest impacts on incidental species will not exceed allowable levels determined in the management plan.
3. Proven, documented, selective gear exists (if not, only an experimental fishery should be considered).
4. Significant wastage of incidental species will not occur or a written economic analysis demonstrates the landed value of the target species exceeds the potential landed value of the wasted species.
5. The species specific or ratio fishery will occur in an acceptable time and area where wastage can be minimized and target stocks are maximally available.
6. Implementation of selective fisheries for marked or hatchery fish must be in accordance with U.S. v. Washington stipulation and order concerning co-management and mass marking (Case No. 9213, Subproceeding No. 96-3) and any subsequent stipulations or orders of the U.S. District Court, and consistent with international objectives under the PST (e.g., to ensure the integrity of the coded-wire tag program).

6.5.3.2 *Selective Fisheries Which May Change Allocation Percentages North of Cape Falcon*

As a tool to increase management flexibility to respond to changing harvest opportunities, the Council may implement deviations from the specified port area allocations and/or gear allocations to increase harvest opportunity through fisheries that are selective for marked salmon stocks (e.g., marked hatchery salmon). The benefits of any selective fishery will vary from year to year and fishery to fishery depending on stock abundance, the mix of marked and unmarked fish, projected hook-and-release

mortality rates, and public acceptance. These factors should be considered on an annual and case-by-case basis when utilizing selective fisheries. The deviations for selective fisheries are subordinate to the allocation priorities in Section 5.3.1.1 and may be allowed under the following management constraints:

1. Selective fisheries will first be considered during the months of August and/or September. However, the Council may consider selective fisheries at other times, depending on year to year circumstances identified in the preceding paragraph.
2. The total impacts within each port area or gear group on the critical natural stocks of management concern are not greater than those under the original allocation without the selective fisheries.
3. Other allocation objectives (i.e., treaty Indian, or ocean and inside allocations) are satisfied during negotiations in the North of Cape Falcon Forum.
4. The selective fishery is assessed against the guidelines in Section 6.5.3.1.
5. Selective fishery proposals need to be made in a timely manner in order to allow sufficient time for analysis and public comment on the proposal before the Council finalizes its fishery recommendations.

If the Council chooses to deviate from the specified port and/or gear allocations, the process for establishing a selective fishery would be as follows:

1. Allocate the TAC among the gear groups and port areas according to the basic FMP allocation process described in Section 5.3.1 without the selective fishery.
2. Each gear group or port area may utilize the critical natural stock impacts allocated to its portion of the TAC to access additional harvestable, marked fish, over and above the harvest share established in step one, within the limits of the management constraints listed in the preceding paragraph.

6.5.4 Procedures for Calculating Quotas

Quotas will be based on the total allowable ocean harvest and the allocation plan as determined by the procedures of Chapter 5.

To the extent adjustable quotas are used, they may be subject to some or all of the following inseason adjustments:

1. For coho, private hatchery contribution to the ocean fisheries in the OPI area.
2. Unanticipated loss of shakers (bycatch mortality of undersized fish or unauthorized fish of another species that have to be returned to the water) during the season. (Adjustment for coho hooking mortality during any all-salmon-except-coho season will be made when the quotas are established.)
3. Any catch that take place in fisheries within territorial waters that are inconsistent with federal regulations in the EEZ.
4. If the ability to update inseason stock abundance is developed in the future, adjustments to total allowable harvest could be made, where appropriate.
5. The ability to redistribute quotas between subareas depending on the performance toward achieving the overall quota in the area.

Changes in the quotas as a result of the inseason adjustment process will be avoided unless the changes are of such magnitude that they can be validated by the STT and Council, given the precision of the original estimates.

The basis for determining the private hatchery contribution in (1) above will be either coded-wire tag analysis or analysis of scale patterns, whichever is determined by the STT to be more accurate, or another more accurate method that may be developed in the future, as determined by the STT and Council.

In reference to (4) and (5) above, if reliable techniques become available for making inseason estimates of stock abundance, and provision is made in any season for its use, a determination of techniques to be applied will be made by the Council through the Salmon Methodology Review process and discussed during the preseason regulatory process.

APPENDIX C

OREGON PRODUCTION INDEX DATA

LIST OF TABLES

| | <u>Page</u> |
|---|-------------|
| TABLE C-1. Millions of coho smolts released annually into the OPI area by geographic area and rearing agency | 121 |
| TABLE C-2. Data set used in predicting Oregon production index hatchery (OPIH) adult coho. | 122 |
| TABLE C-3. Estimated coho salmon natural spawner abundance in Oregon coastal basins for each OCN coho management component | 124 |
| TABLE C-4. Data set used in predicting 2011 Oregon coastal natural river (OCNR) coho ocean recruits with random survey sampling and Mixed Stock Model (MSM) accounting..... | 125 |

TABLE C-1. Millions of coho smolts^{a/} released annually into the OPI area by geographic area and rearing agency.

| | Columbia River | | | | | | Oregon Coast | | | | |
|--------------------|----------------|-------|------|----------|---------|-------|--------------------|-----------|-------|------------|-----------|
| Year or | Washington | | | | | | Private | | | | |
| Average | Oregon | Early | Late | Combined | Federal | Total | ODFW ^{b/} | Yearlings | Total | California | Total OPI |
| 1960-1965 | 5.6 | - | - | 6.1 | 4.5 | 16.2 | 2.0 | - | 2.0 | 0.4 | 18.6 |
| 1966-1970 | 6.0 | 10.2 | 4.9 | 15.1 | 6.5 | 27.6 | 2.9 | 0.0 | 2.9 | 1.3 | 31.8 |
| 1971-1975 | 6.8 | 10.7 | 6.8 | 17.5 | 4.5 | 28.8 | 3.9 | 0.0 | 3.9 | 1.2 | 33.9 |
| 1976-1980 | 8.0 | 7.3 | 10.1 | 17.4 | 4.7 | 30.1 | 3.8 | 1.4 | 5.2 | 0.7 | 36.0 |
| 1981-1985 | 7.1 | 4.3 | 14.4 | 18.7 | 3.2 | 29.0 | 3.9 | 3.3 | 7.2 | 0.7 | 36.9 |
| 1986-1990 | 7.3 | 3.1 | 15.6 | 18.7 | 4.1 | 30.1 | 5.2 | 1.9 | 7.1 | 1.4 | 38.6 |
| 1991 | 10.4 | 3.7 | 15.3 | 19.0 | 5.9 | 35.2 | 5.3 | - | 5.3 | 1.5 | 42.0 |
| 1992 | 11.5 | 4.3 | 14.3 | 18.6 | 2.7 | 32.8 | 6.2 | - | 6.2 | 0.7 | 39.7 |
| 1993 | 11.1 | 4.3 | 14.8 | 19.1 | 4.1 | 34.3 | 4.3 | - | 4.3 | 0.8 | 39.4 |
| 1994 | 9.1 | 2.5 | 12.0 | 14.5 | 3.0 | 26.6 | 5.2 | - | 5.2 | 0.6 | 32.4 |
| 1995 | 7.1 | 3.4 | 12.9 | 16.3 | 1.7 | 25.1 | 3.7 | - | 3.7 | 0.7 | 29.5 |
| 1996 | 8.4 | 3.4 | 12.9 | 16.3 | 3.4 | 28.1 | 3.3 | - | 3.3 | 0.3 | 31.7 |
| 1997 | 6.1 | 3.2 | 7.8 | 11.0 | 3.9 | 21.0 | 2.9 | - | 2.9 | 0.7 | 24.6 |
| 1998 | 6.1 | 5.8 | 11.4 | 17.2 | 3.6 | 26.8 | 1.7 | - | 1.7 | 0.6 | 29.1 |
| 1999 | 7.6 | 4.0 | 11.5 | 15.5 | 4.8 | 27.9 | 1.0 | - | 1.0 | 0.7 | 29.6 |
| 2000 | 7.8 | 6.2 | 10.8 | 17.0 | 5.9 | 30.7 | 0.9 | - | 0.9 | 0.6 | 32.2 |
| 2001 | 7.6 | 4.2 | 9.7 | 13.9 | 3.7 | 25.2 | 0.9 | - | 0.9 | 0.6 | 26.7 |
| 2002 | 7.5 | 3.3 | 8.6 | 11.9 | 4.3 | 23.7 | 1.0 | - | 1.0 | 0.6 | 25.3 |
| 2003 | 8.2 | 3.3 | 8.7 | 12.0 | 3.1 | 23.3 | 0.8 | - | 0.8 | 0.5 | 24.6 |
| 2004 | 6.7 | 3.0 | 8.8 | 11.8 | 3.6 | 22.1 | 0.8 | - | 0.8 | 0.6 | 23.5 |
| 2005 | 6.1 | 2.5 | 9.1 | 11.6 | 2.8 | 20.6 | 0.8 | - | 0.8 | 0.6 | 22.0 |
| 2006 | 6.1 | 2.8 | 9.0 | 11.7 | 2.6 | 20.4 | 0.8 | - | 0.8 | 0.6 | 21.8 |
| 2007 | 6.2 | 3.1 | 9.0 | 12.1 | 3.1 | 21.4 | 0.7 | - | 0.7 | 0.6 | 22.6 |
| 2008 | 6.9 | 2.8 | 9.2 | 12.0 | 2.9 | 21.9 | 0.4 | - | 0.4 | 0.5 | 22.8 |
| 2009 | 6.9 | 2.5 | 8.3 | 10.8 | 3.2 | 20.9 | 0.4 | - | 0.4 | 0.6 | 21.8 |
| 2010 | 5.9 | 2.0 | 7.5 | 9.5 | 3.1 | 18.5 | 0.3 | - | 0.3 | 0.5 | 19.4 |
| 2011 ^{c/} | 5.8 | 1.8 | 8.4 | 10.2 | 3.0 | 19.0 | 0.4 | - | 0.4 | 0.5 | 19.8 |

a/ Defined here as 30 fish per pound or larger and released in February or later.

b/ Beginning in 1989, does not include minor releases from STEP projects.

c/ Preliminary.

TABLE C-2. Data set used in predicting Oregon production index hatchery (OPIH) adult coho. Adults and jacks shown in thousands of fish and smolts in millions of fish.

| Year (t) | Adults (t) | | Jacks (t-1) | | | Columbia River Smolts (t-1) | | |
|--------------------|--------------------|---------------------|-------------------------|------------------------------|-------------------------------|-----------------------------|-------------------------------|--|
| | OPIH ^{a/} | MSM ^{b/} | Total OPI ^{c/} | Columbia River ^{d/} | OR Coast/ CA ^{e/} | Delayed ^{f/} | Normal Timed ^{g/} | Adjustment Proportion ^{h/} |
| | | | | | | | | |
| 1970 | 2,765.1 | - | | | | | | |
| 1971 | 3,365.0 | - | 179.4 | 172.8 | 6.6 | 0.0 | 24.0 | 0.0000 |
| 1972 | 1,924.8 | - | 103.7 | 100.8 | 2.9 | 0.0 | 28.3 | 0.0000 |
| 1973 | 1,817.0 | - | 91.4 | 85.7 | 5.7 | 1.8 | 29.9 | 5.1592 |
| 1974 | 3,071.1 | - | 144.2 | 132.0 | 12.1 | 2.9 | 28.5 | 13.4316 |
| 1975 | 1,652.8 | - | 76.2 | 75.1 | 1.1 | 1.8 | 27.8 | 4.8626 |
| 1976 | 3,885.3 | - | 171.5 | 146.2 | 25.3 | 2.0 | 29.0 | 10.0828 |
| 1977 | 987.5 | - | 53.8 | 46.3 | 7.5 | 0.2 | 28.9 | 0.3204 |
| 1978 | 1,824.1 | - | 103.2 | 99.2 | 4.0 | 0.0 | 31.4 | 0.0000 |
| 1979 | 1,476.7 | - | 72.5 | 64.1 | 8.4 | 5.0 | 32.6 | 9.8313 |
| 1980 | 1,224.0 | - | 57.7 | 51.6 | 6.0 | 6.7 | 28.9 | 11.9626 |
| 1981 | 1,064.5 | - | 48.7 | 40.6 | 8.1 | 5.6 | 28.1 | 8.0911 |
| 1982 | 1,266.8 | - | 61.3 | 55.0 | 6.3 | 6.8 | 32.4 | 11.5432 |
| 1983 ^{i/} | 599.2 | - | 68.3 | 61.0 | 7.2 | 5.0 | 27.7 | 11.0108 |
| 1984 | 691.3 | - | 31.6 | 28.0 | 3.6 | 5.1 | 27.0 | 5.2889 |
| 1985 | 717.5 | - | 26.0 | 18.2 | 7.8 | 9.1 | 29.2 | 5.6719 |
| 1986 | 2,435.8 | 2,412.0 | 77.5 | 64.6 | 12.9 | 12.2 | 28.8 | 27.3653 |
| 1987 | 887.2 | 779.4 | 32.9 | 24.2 | 8.7 | 9.0 | 32.9 | 6.6201 |
| 1988 | 1,669.3 | 1,467.8 | 85.2 | 72.3 | 12.9 | 7.7 | 28.8 | 19.3302 |
| 1989 | 1,720.2 | 1,922.0 | 60.8 | 55.0 | 5.8 | 7.2 | 29.5 | 13.4237 |
| 1990 | 718.4 | 713.6 | 46.6 | 37.1 | 9.6 | 8.5 | 29.6 | 10.6537 |
| 1991 | 1,874.8 | 1,816.5 | 68.6 | 60.7 | 7.9 | 7.1 | 30.3 | 14.2234 |
| 1992 | 543.6 | 512.6 | 25.6 | 19.9 | 5.7 | 6.0 | 35.3 | 3.3824 |
| 1993 | 261.7 | 223.3 | 27.1 | 19.6 | 7.5 | 5.5 | 32.8 | 3.2866 |
| 1994 | 202.3 | 214.1 | 5.2 | 3.9 | 1.3 | 6.0 | 34.4 | 0.6802 |
| 1995 | 147.2 | 139.4 | 11.8 | 9.1 | 2.7 | 3.1 | 26.6 | 1.0605 |
| 1996 | 185.2 | 176.5 | 17.4 | 14.1 | 3.2 | 4.2 | 25.2 | 2.3500 |
| 1997 | 200.7 | 195.6 | 20.4 | 15.8 | 4.6 | 3.4 | 28.0 | 1.9186 |
| 1998 | 207.5 | 228.3 | 9.7 | 6.8 | 3.0 | 2.5 | 21.0 | 0.7976 |
| 1999 | 334.5 | 372.5 | 29.5 | 23.6 | 5.9 | 3.0 | 26.8 | 2.6418 |
| 2000 | 673.2 | 673.1 | 34.8 | 31.3 | 3.5 | 4.1 | 27.9 | 4.5996 |
| 2001 | 1,417.1 | 1,478.7 | 87.4 | 71.7 | 15.7 | 2.0 | 30.6 | 4.6863 |
| 2002 | 649.8 | 689.5 | 25.2 | 18.9 | 6.3 | 1.4 | 23.5 | 1.1260 |
| 2003 | 936.6 | 1,009.9 | 49.9 | 41.7 | 8.2 | 0.3 | 23.7 | 0.5278 |
| 2004 | 622.1 | 693.6 | 35.4 | 29.4 | 6.0 | 2.0 | 23.2 | 2.5345 |
| 2005 | 443.2 | 454.0 | 25.0 | 21.2 | 4.7 | 0.8 | 22.0 | 0.7709 |
| 2006 | 440.6 | 523.4 | 25.9 | 20.9 | 5.4 | 0.4 | 20.6 | 0.4058 |
| 2007 | 476.6 | 545.3 | 36.3 | 34.2 | 2.5 | 0.1 | 20.4 | 0.1676 |
| 2008 | 565.3 | 576.9 | 16.0 | 14.0 | 1.4 | 0.6 | 21.4 | 0.3925 |
| 2009 | . | 1,051.0 | 60.4 | 58.4 | 2.6 | 1.1 | 21.9 | 2.9333 |
| 2010 | . | 546.5 | 25.1 | 23.8 | 1.5 | 0.2 | 21.3 | 0.2235 |
| 2011 | . | 442.3 | 23.3 | 22.2 | 1.1 | 0.3 | 18.5 | 0.3600 |
| 2012 | . | 341.7 ^{j/} | 17.8 | 13.8 | 3.9 | 0.9 | 19.0 | 0.6537 |

TABLE C-2. Data sets used in predicting Oregon production index hatchery (OPIH) adult coho. Adults and jacks shown in thousands of fish and smolts in millions of fish. (Page 2 of 2)

- a/ Adult OPIH = Harvest impacts plus escapement for public hatchery stocks originating in the Columbia River, Oregon coastal rivers, and the Klamath River, California.
- b/ Adult MSM = Harvest impacts plus escapement for public hatchery stocks originating in the Columbia River, Oregon coastal rivers, and the Klamath River. Estimates derived from the MSM and used for prediction beginning in 2008.
- c/ Jack OPI = Total Jack CR and Jack OC.
- d/ Jack CR = Columbia River jack returns corrected for small adults.
- e/ Jack OC = Oregon coastal and California hatchery jack returns corrected for small adults.
- f/ Sm D = Columbia River delayed smolt releases from the previous year expected to return as adults in the year listed.
- g/ Sm CR = Columbia River smolt release from the previous year expected to return as adults in the year listed.
- h/ Correction term for delayed smolts released from Columbia River hatcheries (proportion).
- i/ Data not used in subsequent predictions due to El Niño impacts.
- j/ Preseason predicted adults.

TABLE C-3. Estimated coho salmon natural spawner abundance in Oregon coastal basins for each OCN coho management component.

| Component and Basin ^{a/} | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 1996-2011 Avg. |
|-----------------------------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|---------------|----------------|----------------|----------------|----------------|----------------|
| NORTHERN | | | | | | | | | | | | | | | | | |
| Necanicum | 768 | 253 | 946 | 728 | 474 | 5,247 | 2,896 | 3,068 | 2,198 | 1,218 | 750 | 431 | 1,055 | 3,827 | 4,445 | 1,842 | 1,884 |
| Nehalem | 1,057 | 1,173 | 1,190 | 3,713 | 14,285 | 22,310 | 20,903 | 33,059 | 18,736 | 10,451 | 11,614 | 14,033 | 17,205 | 21,753 | 32,215 | 14,270 | 14,873 |
| Tillamook | 661 | 388 | 271 | 2,175 | 1,983 | 1,883 | 15,715 | 14,584 | 2,532 | 1,995 | 8,774 | 2,295 | 4,828 | 16,251 | 14,890 | 21,512 | 6,921 |
| Nestucca | 519 | 271 | 169 | 2,201 | 1,171 | 3,940 | 13,003 | 8,929 | 4,695 | 686 | 1,876 | 394 | 1,844 | 4,252 | 1,947 | 8,754 | 3,416 |
| Ind. Tribs. | 275 | 61 | 0 | 47 | 0 | 71 | 16 | 0 | 661 | 2,116 | 1,121 | 376 | 639 | 2,052 | 1,473 | 1,409 | 645 |
| TOTAL | 3,280 | 2,146 | 2,576 | 8,864 | 17,913 | 33,451 | 52,533 | 59,640 | 28,822 | 16,466 | 24,135 | 17,529 | 25,571 | 48,135 | 54,970 | 47,787 | 27,739 |
| NORTH CENTRAL | | | | | | | | | | | | | | | | | |
| Salmon | 271 | 237 | 8 | 175 | 0 | 310 | 372 | 0 | 1,642 | 79 | 513 | 59 | 652 | 753 | 1,382 | 3,528 | 624 |
| Siletz | 763 | 336 | 394 | 706 | 3,553 | 1,437 | 2,252 | 9,736 | 8,179 | 14,567 | 5,205 | 2,197 | 20,634 | 24,070 | 6,283 | 26,964 | 7,955 |
| Yaquina | 5,127 | 384 | 365 | 2,588 | 647 | 3,039 | 23,981 | 13,254 | 5,539 | 3,441 | 4,247 | 3,158 | 10,913 | 11,182 | 8,589 | 19,065 | 7,220 |
| Beaver Ck. | 1,340 | 425 | 1,041 | 3,366 | 738 | 5,274 | 8,754 | 5,812 | 4,569 | 2,264 | 1,950 | 611 | 1,218 | 3,575 | 2,072 | 2,882 | 2,868 |
| Alsea | 1,637 | 680 | 213 | 2,050 | 2,465 | 3,339 | 6,170 | 8,957 | 5,233 | 13,907 | 1,972 | 2,146 | 13,320 | 14,638 | 9,688 | 22,393 | 6,801 |
| Siuslaw | 7,625 | 668 | 1,089 | 2,724 | 6,767 | 11,024 | 57,129 | 29,257 | 8,729 | 16,907 | 5,869 | 3,552 | 17,491 | 30,607 | 25,983 | 24,475 | 15,619 |
| Ind. Tribs. | 1,364 | 112 | 173 | 150 | 91 | 816 | 5,308 | 1,852 | 8,179 | 242 | 1,468 | 547 | 3,910 | 1,610 | 2,548 | 5,708 | 2,130 |
| TOTAL | 18,127 | 2,842 | 3,283 | 11,759 | 14,261 | 25,239 | 103,966 | 68,868 | 42,070 | 51,407 | 21,224 | 12,270 | 68,138 | 86,435 | 56,545 | 105,015 | 43,216 |
| SOUTH CENTRAL | | | | | | | | | | | | | | | | | |
| Umpqua | 10,824 | 2,960 | 9,153 | 7,685 | 12,233 | 35,702 | 37,591 | 29,607 | 29,920 | 42,532 | 18,092 | 11,783 | 37,868 | 57,984 | 70,019 | 68,736 | 30,168 |
| Coos | 12,128 | 1,127 | 3,167 | 4,945 | 5,386 | 43,301 | 35,688 | 29,559 | 23,337 | 17,048 | 11,266 | 1,329 | 14,881 | 26,979 | 27,658 | 9,205 | 16,688 |
| Coquille | 16,169 | 5,720 | 2,466 | 3,001 | 6,130 | 13,310 | 8,610 | 23,909 | 22,138 | 11,806 | 28,577 | 13,968 | 8,791 | 22,286 | 23,564 | 35,800 | 15,390 |
| Floras Ck. | - | - | 252 | 164 | 1,440 | 1,945 | 20 | 310 | 7,446 | 506 | 1,104 | 340 | 786 | 3,203 | 11,329 | 4,118 | 2,355 |
| Sixes R. | - | - | - | - | - | - | - | - | 403 | 105 | 294 | 97 | 43 | 176 | 100 | 247 | 183 |
| Coastal Lakes | 13,493 | 8,603 | 11,107 | 12,710 | 12,747 | 19,669 | 22,162 | 16,688 | 18,642 | 14,725 | 24,127 | 8,955 | 23,608 | 17,349 | 38,744 | 20,392 | 17,733 |
| Ind. Tribs. | - | - | - | - | - | - | - | - | - | - | - | - | 0 | 188 | 484 | 75 | 187 |
| TOTAL | 52,614 | 18,410 | 26,145 | 28,505 | 37,936 | 113,927 | 104,071 | 100,073 | 101,886 | 86,722 | 83,460 | 36,472 | 85,977 | 128,165 | 171,898 | 138,573 | 82,177 |
| SOUTH | | | | | | | | | | | | | | | | | |
| Rogue ^{b/} | 5,241 | 8,213 | 2,257 | 1,389 | 10,978 | 12,579 | 8,403 | 6,754 | 24,486 | 9,957 | 3,937 | 5,242 | 414 | 2,566 | 3,073 | 3,917 | 6,838 |
| COASTWIDE | 79,262 | 31,611 | 34,261 | 50,517 | 81,088 | 185,196 | 268,973 | 235,335 | 197,264 | 164,552 | 132,756 | 71,513 | 180,100 | 265,301 | 286,486 | 295,292 | 159,969 |

a/ The sum of the individual basins may not equal the aggregate totals due to the use of independent estimates at different geographic scales.

b/ Mark recapture estimate based on seining at Huntley Park in the lower Rogue River.

TABLE C-4. Data set used in predicting 2012 Oregon coastal natural river (OCNR) coho ocean recruits with random survey sampling and Mixed Stock Model (MSM) accounting. All environmental data in year of ocean entry (t-1) except SST-J, which is January of adult return year (t). Spawners is parent brood (t-3). Recruits shown in thousands of fish.
(Page 1 of 2)

| Year (t) | Recruits | | Environmental Index-Month(s) ^{a/} | | | | | | | |
|----------|----------|----------|--|---------|---------|---------|---------|-------|--------|---------|
| | Adults | Spawners | PDO-MJJ | UWI-JAS | UWI-SON | SSH-AMJ | SST-AMJ | SST-J | MEI-ON | SPR.TRN |
| 1970 | 183.1 | 204.7 | -0.37 | 51.67 | -16.67 | -144.37 | 10.91 | - | -1.10 | 78 |
| 1971 | 416.3 | 198.9 | -1.77 | 32.33 | -10.33 | -63.70 | 11.69 | 8.67 | -1.31 | 106 |
| 1972 | 185.5 | 129.2 | -1.42 | 42.33 | -3.67 | -57.13 | 11.85 | 8.44 | 1.72 | 107 |
| 1973 | 235.0 | 51.2 | -0.77 | 60.67 | -15.33 | -150.47 | 12.23 | 9.46 | -1.53 | 80 |
| 1974 | 196.4 | 65.6 | -0.22 | 41.33 | -8.00 | -71.40 | 10.96 | 9.30 | -1.26 | 102 |
| 1975 | 208.4 | 24.1 | -0.86 | 48.67 | -29.67 | -148.50 | 10.86 | 9.49 | -1.79 | 83 |
| 1976 | 451.7 | 37.8 | -0.25 | 18.00 | -5.67 | -110.63 | 10.72 | 9.07 | 0.48 | 103 |
| 1977 | 161.2 | 28.1 | 0.31 | 40.33 | -22.33 | -134.93 | 11.22 | 9.78 | 0.97 | 74 |
| 1978 | 111.6 | 34.8 | -0.06 | 33.33 | -1.33 | -86.07 | 11.58 | 11.24 | 0.20 | 97 |
| 1979 | 188.8 | 39.2 | 0.70 | 20.33 | -45.00 | -91.17 | 11.24 | 8.74 | 0.73 | 73 |
| 1980 | 108.3 | 13.7 | 0.40 | 69.33 | -43.67 | -63.87 | 12.05 | 10.50 | 0.24 | 78 |
| 1981 | 174.5 | 18.2 | 1.43 | 48.67 | -36.33 | -81.37 | 12.14 | 11.72 | -0.06 | 88 |
| 1982 | 185.7 | 38.4 | -0.26 | 33.67 | -26.67 | -68.67 | 11.01 | 9.86 | 2.45 | 109 |
| 1983 | 96.0 | 25.6 | 2.56 | 26.00 | -47.33 | -4.97 | 12.12 | 11.10 | -0.18 | 126 |
| 1984 | 94.7 | 30.1 | 0.43 | 53.67 | -52.00 | -63.27 | 11.44 | 10.65 | -0.35 | 112 |
| 1985 | 124.9 | 68.3 | 0.42 | 47.00 | 0.00 | -80.43 | 10.98 | 9.99 | -0.05 | 48 |
| 1986 | 114.3 | 19.4 | 1.14 | 53.33 | -4.33 | -82.03 | 11.52 | 10.04 | 0.87 | 89 |
| 1987 | 77.8 | 59.7 | 1.53 | 50.33 | -23.00 | -80.23 | 11.43 | 10.58 | 1.25 | 81 |
| 1988 | 152.5 | 66.3 | 0.86 | 51.33 | -25.00 | -62.70 | 11.49 | 9.89 | -1.47 | 68 |
| 1989 | 114.9 | 57.2 | 0.55 | 46.00 | 5.00 | -65.23 | 11.62 | 9.43 | -0.07 | 97 |
| 1990 | 63.3 | 25.3 | 0.38 | 54.00 | -3.00 | -63.93 | 12.00 | 9.97 | 0.37 | 81 |
| 1991 | 84.1 | 45.7 | -0.69 | 54.67 | 7.33 | -110.40 | 10.95 | 8.96 | 1.20 | 99 |
| 1992 | 107.6 | 40.7 | 1.57 | 53.33 | -11.00 | -30.20 | 12.69 | 10.11 | 0.60 | 123 |
| 1993 | 74.9 | 16.9 | 2.27 | 57.00 | 13.00 | 59.37 | 13.19 | 9.38 | 0.82 | 161 |
| 1994 | 41.0 | 30.4 | 0.58 | 57.33 | -6.00 | -64.10 | 11.45 | 11.04 | 1.28 | 87 |
| 1995 | 47.8 | 40.2 | 1.48 | 33.33 | -24.33 | -64.50 | 11.19 | 10.57 | -0.50 | 95 |
| 1996 | 64.5 | 45.2 | 1.35 | 83.67 | 4.67 | -47.30 | 11.44 | 11.66 | -0.16 | 120 |
| 1997 | 16.3 | 38.3 | 2.31 | 20.00 | -38.00 | -14.50 | 12.10 | 10.76 | 2.52 | 146 |
| 1998 | 22.4 | 42.8 | 0.35 | 73.67 | -37.33 | -41.17 | 11.37 | 12.26 | -1.17 | 105 |
| 1999 | 38.3 | 60.5 | -0.88 | 70.33 | -17.33 | -110.77 | 10.67 | 9.54 | -1.08 | 91 |
| 2000 | 58.7 | 14.8 | -0.38 | 45.00 | -11.00 | -54.67 | 11.35 | 10.00 | -0.76 | 72 |

TABLE C-4. Data set used in predicting 2012 Oregon coastal natural river (OCNR) coho ocean recruits with random survey sampling and Mixed Stock Model (MSM) accounting. All environmental data in year of ocean entry (t-1) except SST-J, which is January of adult return year (t). Spawners is parent brood (t-3). Recruits shown in thousands of fish. (Page 2 of 2)

| Year (t) | Recruits | | Environmental Index-Month(s) ^{a/} | | | | | | | |
|--------------------|----------|----------|--|---------|---------|---------|---------|-------|--------|---------|
| | Adults | Spawners | PDO-MJJ | UWI-JAS | UWI-SON | SSH-AMJ | SST-AMJ | SST-J | MEI-ON | SPR.TRN |
| 2001 | 156.5 | 20.9 | -0.69 | 60.67 | -29.67 | -124.50 | 10.68 | 10.17 | -0.18 | 61 |
| 2002 | 246.1 | 36.4 | -0.43 | 72.67 | -26.00 | -146.90 | 10.11 | 10.07 | 1.06 | 80 |
| 2003 | 227.3 | 57.4 | 0.84 | 65.33 | -7.33 | -61.67 | 11.15 | 11.01 | 0.52 | 112 |
| 2004 | 164.0 | 152.9 | 0.45 | 30.33 | 6.33 | -60.73 | 11.86 | 10.30 | 0.79 | 110 |
| 2005 | 146.3 | 238.4 | 1.23 | 73.33 | 6.00 | -23.67 | 12.54 | 10.21 | -0.41 | 145 |
| 2006 | 113.1 | 211.9 | 0.62 | 84.00 | -14.00 | -34.30 | 11.15 | 11.46 | 1.29 | 112 |
| 2007 | 64.8 | 156.7 | 0.26 | 23.67 | 5.00 | -121.53 | 10.62 | 9.85 | -1.18 | 74 |
| 2008 | 157.0 | 139.4 | -1.46 | 33.33 | -2.33 | -110.93 | 9.62 | 8.92 | -0.63 | 89 |
| 2009 | 262.9 | 104.5 | -0.57 | 36.33 | -39.67 | -93.63 | 10.45 | 9.37 | 1.06 | 82 |
| 2010 | 251.3 | - | -0.22 | 57.00 | -15.33 | -46.03 | 11.67 | 10.76 | -1.61 | 100 |
| 2011 | 289.8 | - | -0.97 | 41.67 | -12.67 | -49.93 | 10.69 | 10.12 | -0.98 | 100 |
| 2012 ^{b/} | 262.4 | - | - | - | - | - | - | 9.19 | - | - |

a/ Environmental Index descriptions:

PDO - Pacific Decadal Oscillation

UWI - Upwelling wind index (mean upwelling winds index in months of ocean migration year at 42° N 125° W)

SSH - Sea surface height (South Beach, OR at 44° 37.5' N, 124 ° 02.6' W)


SST - Sea surface temperature (mean sea surface temperature in January of return year at Charleston, OR)

MEI - Multi-variate ENSO index

SPR.TRN - Spring transition date (Julian)

b/ Forecast.

Marine Fisheries Management Zones



0 25 50 100 km

Projection: UTM Zone 10, NAD83

Andrew Weiss
Fish Program
Biological Data Systems
Feb. 2009