

**Preferred 2017 and 2018 harvest specifications (overfishing limits (OFLs in mt), acceptable biological catches (ABCs in mt), and annual catch limits (ACLs in mt)), for west coast groundfish stocks and stock complexes under preferred harvest control rules (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in stock complexes in italics).**

Stock or Stock Complex	2017			2018			Harvest Control Rules
	OFL	ABC	ACL	OFL	ABC	ACL	
<b>OVERFISHED STOCKS</b>							
BOCACCIO S. of 40°10'	2,139	2,044	790	2,013	1,924	741	ABC (P* = 0.45), ACL (SPR = 77.7%)
COWCOD S. of 40°10'	70	63	10	71	64	10	ABCs sum of Con. and Mont. area ABCs, ACLs projected from 2013 rebuilding analysis (SPR = 82.7% (F = 0.007)) + Mont. area ABC contrib., ACT = 4 mt
<i>COWCOD (Conception)</i>	58	53	NA	59	54	NA	ABC (P* = 0.45)
<i>COWCOD (Monterey)</i>	12	10	NA	12	10	NA	ACL = ABC (P* = 0.45)
DARKBLOTTED ROCKFISH	671	641	490	689	659	490	ABC (P* = 0.45), constant catch ACL = 490 mt
PACIFIC OCEAN PERCH	961	919	171	985	942	176	Updated projections from 2011 rebuilding analysis using actual catches from 2011-2014 and assumed ACL removals thereafter. ABC (P* = 0.45), ACL (SPR = 86.4%)
YELLOWEYE ROCKFISH	57	47	20	58	48	20	Catch-only update from the 2011 stock assessment assuming ACL removals; ABC (P* = 0.4), ACL (SPR = 76.0%)
<b>NON-OVERFISHED STOCKS</b>							
Arrowtooth Flounder	16,571	13,804	13,804	16,498	13,743	13,743	ACL = ABC (P* = 0.4)
Big skate	NA	NA	NA	NA	NA	NA	EC Species (i.e., no harvest control rules or specifications)
<b>Black Rockfish (CA)</b>	<b>349</b>	<b>334</b>	<b>334</b>	<b>347</b>	<b>332</b>	<b>332</b>	<b>ACL = ABC (P* = 0.45)</b>
<b>Black Rockfish (OR)</b>	<b>577</b>	<b>527</b>	<b>527</b>	<b>570</b>	<b>520</b>	<b>520</b>	<b>ACL = ABC (P* = 0.45)</b>
<b>Black Rockfish (WA)</b>	<b>319</b>	<b>305</b>	<b>305</b>	<b>315</b>	<b>301</b>	<b>301</b>	<b>ACL = ABC (P* = 0.45)</b>
Blackgill Rockfish S. of 40°10'	NA	NA	NA	146	133	123	ACL = ABC (P* = 0.45) w/ 40-10 adjustment. Managed in the Southern Slope Rockfish complex in 2017.
Cabezon (CA)	157	150	150	156	149	149	ACL = ABC (P* = 0.45) w/ 40-10 adjustment (ACL rounds to ABC)
Cabezon (OR)	49	47	47	49	47	47	ACL = ABC (P* = 0.45)
California scorpionfish	289	264	150	278	254	150	150 mt constant catch ACL; ACT = 111 mt
<b>Canary Rockfish</b>	<b>1,793</b>	<b>1,714</b>	<b>1,526</b>	<b>1,677</b>	<b>1,603</b>	<b>1,526</b>	<b>1,526 mt ACL in 2017-18; ACL = ABC (P* = 0.45) thereafter</b>
Chilipepper S. of 40°10'	2,727	2,607	2,607	2,623	2,507	2,507	ACL = ABC (P* = 0.45)
Dover Sole	89,702	85,755	50,000	90,282	86,310	50,000	ABC (P* = 0.45), ACL = 50,000 mt annually
English Sole	10,914	9,964	9,964	8,255	7,537	7,537	ACL = ABC (P* = 0.45)
Lingcod N. of 40°10'	3,549	3,333	3,333	3,310	3,110	3,110	ACL = ABC (P* = 0.45)
Lingcod S. of 40°10'	1,502	1,251	1,251	1,373	1,144	1,144	ACL = ABC (P* = 0.4)
Longnose skate	2,556	2,444	2,000	2,526	2,415	2,000	ABC (P* = 0.45), ACL = 2,000 mt annually
Longspine Thornyhead N. of 34°27'	4,571	3,808	2,894	4,339	3,614	2,747	ACL = 76% of coastwide ABC (P* = 0.4)
Longspine Thornyhead S. of 34°27'			914			867	ACL = 24% of coastwide ABC (P* = 0.4)
Pacific Cod	3,200	2,221	1,600	3,200	2,221	1,600	ABC (P* = 0.4), ACL = 50% of OFL

Stock or Stock Complex	2017			2018			Harvest Control Rules
	OFL	ABC	ACL	OFL	ABC	ACL	
Petrale Sole	3,280	3,136	3,136	3,152	3,013	3,013	ACL = ABC (P* = 0.45)
Sablefish N. of 36°	8,050	7,350	6,041	8,329	7,604	6,299	ACL: 40-10 rule applied to 84.9% of coastwide ABC (P* = 0.4)
Sablefish S. of 36°			1,075			1,120	ACL: 40-10 rule applied to 15.1% of coastwide ABC (P* = 0.4)
Shortbelly	6,950	5,789	500	6,950	5,789	500	ABC (P* = 0.4), ACL = 500 mt annually
Shortspine Thornyhead N. of 34°27'	3,144	2,619	1,713	3,116	2,596	1,698	ACL = 65.4% of coastwide ABC (P* = 0.4)
Shortspine Thornyhead S. of 34°27'			906			898	ACL = 34.6% of coastwide ABC (P* = 0.4)
Spiny dogfish	2,514	2,094	2,094	2,500	2,083	2,083	ACL = ABC (P* = 0.4)
Splitnose S. of 40°10'	1,841	1,760	1,760	1,842	1,761	1,761	ACL = ABC (P* = 0.45)
Starry flounder	1,847	1,282	1,282	1,847	1,282	1,282	ACL = ABC (P* = 0.4)
<b>Widow Rockfish</b>	<b>14,130</b>	<b>13,508</b>	<b>13,508</b>	<b>13,237</b>	<b>12,655</b>	<b>12,655</b>	<b>ACL = ABC (P* = 0.45)</b>
Yellowtail N. of 40°10'	6,786	6,196	6,196	6,574	6,002	6,002	ACL = ABC (P* = 0.45)
<b>STOCK COMPLEXES</b>							
Nearshore Rockfish North	118	105	105	119	105	105	Sum of component species specifications
<i>Black and yellow</i>	0.0	0.0	0.0	0.0	0.0	0.0	ACL = ABC (P* = 0.45)
<i>Blue (CA)</i>	34.1	31.1	30.9	34.8	31.8	31.8	ACL = ABC (P* = 0.45) w/40-10 adjustment. The stock is projected to be above target in 2018 under the Expected Catch scenario.
<i>Blue (OR &amp; WA)</i>	32.3	26.9	26.9	32.3	26.9	26.9	ACL = ABC (P* = 0.45)
<i>Brown</i>	2.0	1.8	1.8	2.0	1.9	1.9	ACL = ABC (P* = 0.45)
<i>Calico</i>	-	-	-	-	-	-	ACL = ABC (P* = 0.45)
<i>China</i>	30.2	27.5	27.5	29.3	26.8	26.8	ACL = ABC (P* = 0.45)
<i>Copper</i>	11.2	10.3	10.3	11.6	10.6	10.6	ACL = ABC (P* = 0.45)
<i>Gopher</i>	-	-	-	-	-	-	ACL = ABC (P* = 0.45)
<i>Grass</i>	0.7	0.5	0.5	0.7	0.5	0.5	ACL = ABC (P* = 0.45)
<i>Kelp</i>	0.0	0.0	0.0	0.0	0.0	0.0	ACL = ABC (P* = 0.45)
<i>Olive</i>	0.3	0.3	0.3	0.3	0.3	0.3	ACL = ABC (P* = 0.45)
<i>Quillback</i>	7.4	6.2	6.2	7.4	6.2	6.2	ACL = ABC (P* = 0.45)
<i>Treefish</i>	0.2	0.2	0.2	0.2	0.2	0.2	ACL = ABC (P* = 0.45)
Shelf Rockfish North	2,303	2,049	2,049	2,302	2,048	2,047	Sum of component species specifications
<i>Bronzespotted</i>	-	-	-	-	-	-	ACL = ABC (P* = 0.45)
<i>Bocaccio</i>	284.0	236.9	236.9	284.0	236.9	236.9	ACL = ABC (P* = 0.45)
<i>Chameleon</i>	-	-	-	-	-	-	ACL = ABC (P* = 0.45)
<i>Chilipepper</i>	205.2	196.2	196.2	197.4	188.7	188.7	ACL = ABC (P* = 0.45)
<i>Cowcod</i>	0.4	0.3	0.3	0.4	0.3	0.3	ACL = ABC (P* = 0.45)
<i>Flag</i>	0.1	0.1	0.1	0.1	0.1	0.1	ACL = ABC (P* = 0.45)
<i>Freckled</i>	-	-	-	-	-	-	ACL = ABC (P* = 0.45)
<i>Greenblotched</i>	1.3	1.1	1.1	1.3	1.1	1.1	ACL = ABC (P* = 0.45)
<i>Greenspotted 40°10' to 42° N. lat.</i>	9.4	8.5	8.2	9.3	8.5	8.2	ACL: 40-10 rule applied to 22.2% of northern model (CA N of 34°27' N latitude) ABC (P* = 0.45)
<i>Greenspotted N. of 42 N. lat. (OR &amp; WA)</i>	6.1	5.1	5.1	6.1	5.1	5.1	ACL = ABC (P* = 0.45)

Stock or Stock Complex	2017			2018			Harvest Control Rules
	OFL	ABC	ACL	OFL	ABC	ACL	
<i>Greenstriped</i>	1,299.6	1,186.5	1,186.5	1,306.4	1,192.7	1,192.7	ACL = ABC ( $P^* = 0.45$ )
<i>Halfbanded</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Harlequin</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Honeycomb</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Mexican</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Pink</i>	0.0	0.0	0.0	0.0	0.0	0.0	ACL = ABC ( $P^* = 0.45$ )
<i>Pinkrose</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Puget Sound</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Pygmy</i>	-	-	-	-	-	-	ACL = ABC ( $P^* = 0.45$ )
<i>Redstripe</i>	269.9	225.1	225.1	269.9	225.1	225.1	ACL = ABC ( $P^* = 0.45$ )
<i>Rosethorn</i>	12.9	10.8	10.8	12.9	10.8	10.8	ACL = ABC ( $P^* = 0.45$ )
<i>Rosy</i>	3.0	2.5	2.5	3.0	2.5	2.5	ACL = ABC ( $P^* = 0.45$ )
<i>Silvergray</i>	159.4	133.0	133.0	159.4	133.0	133.0	ACL = ABC ( $P^* = 0.45$ )
<i>Speckled</i>	0.2	0.1	0.1	0.2	0.1	0.1	ACL = ABC ( $P^* = 0.45$ )
<i>Squarespot</i>	0.2	0.1	0.1	0.2	0.1	0.1	ACL = ABC ( $P^* = 0.45$ )
<i>Starry</i>	0.0	0.0	0.0	0.0	0.0	0.0	ACL = ABC ( $P^* = 0.45$ )
<i>Stripetail</i>	40.4	33.7	33.7	40.4	33.7	33.7	ACL = ABC ( $P^* = 0.45$ )
<i>Swordspine</i>	0.0	0.0	0.0	0.0	0.0	0.0	ACL = ABC ( $P^* = 0.45$ )
<i>Tiger</i>	1.0	0.8	0.8	1.0	0.8	0.8	ACL = ABC ( $P^* = 0.45$ )
<i>Vermilion</i>	9.7	8.1	8.1	9.7	8.1	8.1	ACL = ABC ( $P^* = 0.45$ )
Slope Rockfish North	1,897	1,755	1,755	1,896	1,754	1,754	Sum of component species specifications
<i>Aurora (assuming sigma = 0.39)</i>	17.5	16.6	16.6	17.5	16.6	16.6	ACL = ABC ( $P^* = 0.45$ )
<i>Bank</i>	17.2	14.4	14.4	17.2	14.4	14.4	ACL = ABC ( $P^* = 0.45$ )
<i>Blackgill</i>	4.7	3.9	3.9	4.7	3.9	3.9	ACL = ABC ( $P^* = 0.45$ )
<i>Redbanded</i>	45.3	37.7	37.7	45.3	37.7	37.7	ACL = ABC ( $P^* = 0.45$ )
<i>Rougheye/Blackspotted</i>	210.7	192.4	192.4	214.6	195.9	195.9	ACL = ABC ( $P^* = 0.45$ )
<i>Sharpchin</i>	364.0	332.3	332.3	358.4	327.2	327.2	ACL = ABC ( $P^* = 0.45$ )
<i>Shortraker</i>	18.7	15.6	15.6	18.7	15.6	15.6	ACL = ABC ( $P^* = 0.45$ )
<i>Splinose</i>	1,026.7	981.6	981.6	1,027.1	981.9	981.9	ACL = ABC ( $P^* = 0.45$ )
<i>Yellowmouth</i>	192.4	160.5	160.5	192.4	160.5	160.5	ACL = ABC ( $P^* = 0.45$ )
Nearshore Rockfish South	1,329	1,166	1,163	1,344	1,180	1,179	Sum of component species specifications
<i>Shallow Nearshore Species</i>	NA						
<i>Black and yellow</i>	27.5	23.0	23.0	27.5	23.0	23.0	ACL = ABC ( $P^* = 0.45$ )
<i>China</i>	13.3	12.2	10.8	13.8	12.6	11.5	ACL = ABC ( $P^* = 0.45$ ) w/ 40-10 adjustment
<i>Gopher (N of Pt. Conception)</i>	144.0	120.1	120.1	144.0	120.1	120.1	ACL = ABC ( $P^* = 0.45$ )
<i>Gopher (S of Pt. Conception)</i>	25.6	21.4	21.4	25.6	21.4	21.4	ACL = ABC ( $P^* = 0.45$ )
<i>Grass</i>	59.6	49.7	49.7	59.6	49.7	49.7	ACL = ABC ( $P^* = 0.45$ )
<i>Kelp</i>	27.7	23.1	23.1	27.7	23.1	23.1	ACL = ABC ( $P^* = 0.45$ )
<i>Deeper Nearshore Species</i>	NA						

Stock or Stock Complex	2017			2018			Harvest Control Rules
	OFL	ABC	ACL	OFL	ABC	ACL	
<i>Blue (assessed area)</i>	234.5	214.1	212.8	239.4	218.6	218.6	<i>ACL = ABC (P* = 0.45) w/40-10 adjustment. The stock is projected to be above target in 2018 under the Expected Catch scenario.</i>
<i>Blue (S of 34°27' N. lat.)</i>	72.9	60.8	60.8	72.9	60.8	60.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Brown</i>	170.0	155.2	155.2	174.0	158.8	158.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Calico</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Copper</i>	310.9	283.8	283.8	316.7	289.2	289.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Olive</i>	224.6	187.4	187.4	224.6	187.4	187.4	<i>ACL = ABC (P* = 0.45)</i>
<i>Quillback</i>	5.4	4.5	4.5	5.4	4.5	4.5	<i>ACL = ABC (P* = 0.45)</i>
<i>Treefish</i>	13.2	11.0	11.0	13.2	11.0	11.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Shelf Rockfish South</i>	1,917	1,624	1,623	1,918	1,625	1,624	<i>Sum of component species specifications</i>
<i>Bronzespotted</i>	3.6	3.0	3.0	3.6	3.0	3.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Chameleon</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Flag</i>	23.4	19.5	19.5	23.4	19.5	19.5	<i>ACL = ABC (P* = 0.45)</i>
<i>Freckled</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Greenblotched</i>	23.1	19.3	19.3	23.1	19.3	19.3	<i>ACL = ABC (P* = 0.45)</i>
<i>Greenspotted</i>	78.9	72.0	70.9	78.5	71.7	70.7	<i>ACL: 40-10 rule applied to 77.8% of northern model (CA N of 34°27' N latitude) ABC plus the southern model ABC (P* = 0.45)</i>
<i>Greenstriped</i>	238.4	217.7	217.7	239.6	218.8	218.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Halfbanded</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Harlequin</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Honeycomb</i>	9.9	8.2	8.2	9.9	8.2	8.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Mexican</i>	5.1	4.2	4.2	5.1	4.2	4.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Pink</i>	2.5	2.1	2.1	2.5	2.1	2.1	<i>ACL = ABC (P* = 0.45)</i>
<i>Pinkrose</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Pygmy</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Redstripe</i>	0.5	0.4	0.4	0.5	0.4	0.4	<i>ACL = ABC (P* = 0.45)</i>
<i>Rosethorn</i>	2.1	1.8	1.8	2.1	1.8	1.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Rosy</i>	44.5	37.1	37.1	44.5	37.1	37.1	<i>ACL = ABC (P* = 0.45)</i>
<i>Silvergray</i>	0.5	0.4	0.4	0.5	0.4	0.4	<i>ACL = ABC (P* = 0.45)</i>
<i>Speckled</i>	39.4	32.8	32.8	39.4	32.8	32.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Squarespot</i>	11.1	9.2	9.2	11.1	9.2	9.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Starry</i>	62.6	52.2	52.2	62.6	52.2	52.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Stripetail</i>	23.6	19.7	19.7	23.6	19.7	19.7	<i>ACL = ABC (P* = 0.45)</i>
<i>Swordspine</i>	14.2	11.9	11.9	14.2	11.9	11.9	<i>ACL = ABC (P* = 0.45)</i>
<i>Tiger</i>	0.0	0.0	0.0	0.0	0.0	0.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Vermilion</i>	269.3	224.6	224.6	269.3	224.6	224.6	<i>ACL = ABC (P* = 0.45)</i>
<i>Yellowtail</i>	1,064.4	887.7	887.7	1,064.4	887.7	887.7	<i>ACL = ABC (P* = 0.45)</i>
<i>Slope Rockfish South</i>	827	718	707	683	586	586	<i>Sum of component species specifications</i>
<i>Aurora (assuming sigma = 0.39)</i>	74.4	70.9	70.9	74.5	71.0	71.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Bank</i>	503.2	419.7	419.7	503.2	419.7	419.7	<i>ACL = ABC (P* = 0.45)</i>
<i>Blackgill</i>	143.0	130.6	120.2	NA	NA	NA	<i>ACL = ABC (P*=0.45) w/ 40-10 adjustment. Managed with stock-specific specifications in 2018.</i>

Stock or Stock Complex	2017			2018			Harvest Control Rules
	OFL	ABC	ACL	OFL	ABC	ACL	
<i>Pacific ocean perch</i>	-	-	-	-	-	-	<i>ACL = ABC (P* = 0.45)</i>
<i>Redbanded</i>	10.4	8.7	8.7	10.4	8.7	8.7	<i>ACL = ABC (P* = 0.45)</i>
<i>Rougheye/Blackspotted</i>	4.3	3.9	3.9	4.4	4.0	4.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Sharpchin</i>	91.0	83.1	83.1	89.6	81.8	81.8	<i>ACL = ABC (P* = 0.45)</i>
<i>Shortraker</i>	0.1	0.1	0.1	0.1	0.1	0.1	<i>ACL = ABC (P* = 0.45)</i>
<i>Yellowmouth</i>	0.8	0.7	0.7	0.8	0.7	0.7	<i>ACL = ABC (P* = 0.45)</i>
Other Flatfish	11,165	8,510	8,510	9,690	7,281	7,281	Sum of component species specifications
<i>Butter sole</i>	4.6	3.2	3.2	4.6	3.2	3.2	<i>ACL = ABC (P* = 0.4)</i>
<i>Curlfin sole</i>	8.2	5.7	5.7	8.2	5.7	5.7	<i>ACL = ABC (P* = 0.4)</i>
<i>Flathead sole</i>	35.0	24.3	24.3	35.0	24.3	24.3	<i>ACL = ABC (P* = 0.4)</i>
<i>Pacific sanddab</i>	4,801.0	3,331.9	3,331.9	4,801.0	3,331.9	3,331.9	<i>ACL = ABC (P* = 0.4)</i>
<i>Rex sole</i>	5,476	4,562	4,562	4,001	3,333	3,333	<i>ACL = ABC (P* = 0.4)</i>
<i>Rock sole</i>	66.7	46.3	46.3	66.7	46.3	46.3	<i>ACL = ABC (P* = 0.4)</i>
<i>Sand sole</i>	773.2	536.6	536.6	773.2	536.6	536.6	<i>ACL = ABC (P* = 0.4)</i>
Other Fish	537	474	474	501	441	441	Sum of component species specifications
<i>Cabezon (WA)</i>	4.5	3.8	3.8	4.8	4.0	4.0	<i>ACL = ABC (P* = 0.45)</i>
<i>Kelp greenling (CA)</i>	118.9	99.2	99.2	118.9	99.2	99.2	<i>ACL = ABC (P* = 0.45)</i>
<i>Kelp greenling (OR) assuming sigma = 0.44</i>	239.1	226.2	226.2	203.2	192.2	192.2	<i>Preferred ACL = ABC (P* = 0.45)</i>
<i>Kelp greenling (WA)</i>	7.1	5.9	5.9	7.1	5.9	5.9	<i>Preferred ACL = ABC (P* = 0.45)</i>
<i>Leopard shark</i>	167.1	139.4	139.4	167.1	139.4	139.4	<i>ACL = ABC (P* = 0.45)</i>