

# **Humpback whale bycatch in 2014 - 2015 in the U.S. West Coast Groundfish Fisheries**

M. Bradley Hanson<sup>1</sup>, Thomas P. Good<sup>1</sup>, Jason E. Jannot<sup>2</sup>, and Jon McVeigh<sup>2</sup>

<sup>1</sup> Ecosystem Science Program  
Conservation Biology Division  
Northwest Fisheries Science Center  
National Marine Fisheries Service  
2725 Montlake Boulevard East  
Seattle, Washington 98112

<sup>2</sup> Fisheries Observation Science Program  
Fishery Resource Analysis and Monitoring Division  
Northwest Fisheries Science Center  
National Marine Fisheries Service  
2725 Montlake Blvd East  
Seattle, Washington 98112

10 March 2017

# Contents

Executive Summary.....	3
Acknowledgments.....	3
Introduction and Background .....	4
Humpback Whale Status, Life History and Ecology .....	4
West Coast Groundfish Fishery.....	8
Northwest Fisheries Science Center Groundfish Observer Programs .....	10
Humpback whale bycatch in West coast Groundfish fisheries.....	10
Amount and Extent of Humpback Whale Take.....	10
Methods.....	11
Data Sources .....	11
Observer Data .....	11
Fish Ticket Data .....	11
Designation of ‘take’ and ‘serious injury’ interactions .....	12
Results and Discussion .....	13
Estimating Humpback Whale Bycatch .....	13
Interactions with other pot/trap fisheries .....	15
Conservation Recommendations.....	16
Additional information relevant to the BiOp/RPMs and Conservation Measures .....	17
Species-Specific Measures .....	18
Species-specific Terms and Conditions .....	19
References .....	19

# Executive Summary

In accordance with the National Marine Fisheries Service (NMFS) Biological Opinion (BiOp) on Continuing Operation of the Pacific Coast Groundfish Fishery, this document provides a summary of observed bycatch of U.S. Endangered Species Act (ESA)-listed humpback whales (*Megaptera novaeangliae*) in sectors of the west coast groundfish fishery from 2014–2015.

There was one documented take of a humpback whale from 2014-2015 in the Pacific Coast groundfish fisheries--an observed entanglement of a humpback whale in October 2014. Although there were no observed interactions between humpbacks and Pacific coast groundfish fisheries during the previous reporting period (2010-2013), pot and trap fisheries generally represent the majority of documented fishery interactions with humpbacks along the U.S. west coast. We thus used Bayesian procedures to estimate bycatch in the West Coast groundfish pot fisheries for the periods 2006-2010 and 2011-2015. The model with a fixed bycatch rate indicated that the probability of exceeding five humpback whale entanglements in this 5-year period was just under 40%, while the model with uncertainty indicated that the probability of exceeding five entanglements approached 0%.

## Acknowledgments

The authors gratefully acknowledge the hard work and dedication of observers from the West Coast Groundfish Observer Program and the At-Sea Hake Observer Program, as well as contributions from observer program staff. This document was a joint effort of the Marine Mammal Bycatch Reporting Team, which included at various times and in various roles, Jason Jannot (NWFSC, WCGOP), Vanessa Tuttle (NWFSC, ASHOP), Brad Hanson (NWFSC, CBD), Eric Ward (NWFSC, CBD), Jim Caretta (SWFSC, MMTD), Karin Forney (SWFSC, MMTD), Robyn Angliss (NMML), Teresa Mongillo (WCRO, PRD), and Strandings Network Coordinators: Kristin Wilkinson, Brent Norberg, Justin Viezbicke, and Justin Greenman).

# Introduction and Background

In accordance with the National Marine Fisheries Service (NMFS) Biological Opinion (BiOp) Regarding the Effects of the Continued Operation of the Pacific Coast Groundfish Fishery (NMFS 2012a) as governed by Pacific Coast Groundfish Fishery Management Plan, this document provides a reporting of observed takes of U.S. Endangered Species Act-listed humpback whales (*Megaptera novaeangliae*) in U.S. west coast groundfish fishery sectors. This report updates the initial assessment submitted in accordance with the Biological Opinion requirement, which reported on bycatch in the fisheries for 2010-2013 (Hanson et al. 2015).

## Humpback Whale Status, Life History and Ecology

Humpback whales (*Megaptera novaeangliae*; Borowski, 1781) are baleen whales of the family Balaenopteridae. This section is adapted from the recent NOAA Fisheries Humpback Whale Status Review; for details, see Bettridge et al. (2015). Humpback whales are found in all oceans of the world with a broad geographical range from tropical to temperate waters in the northern hemisphere and tropical to arctic waters in the southern hemisphere. All populations migrate seasonally between winter calving and breeding grounds and summer feeding grounds within ocean basins. Despite this potential for dispersal or interbreeding of individuals from different major ocean basins is extremely rare, and whales from the major ocean basins are differentiated by reproductive seasonality, behavior, color patterns, and genetics.

Humpback whales were listed worldwide as endangered under the ESA in 1970, and a Recovery Plan was finalized for this species in 1991 (NMFS 1991). Under the MMPA, humpback whales are classified as a strategic stock and considered depleted. On August 12, 2009, NMFS initiated an ESA status review of humpback whales (74 Fed. Reg. 40568) and produced a status review (Bettridge et al. 2015) that identified distinct population segments (DPS) of humpback whale and evaluated their risk of extinction. In September 2016, NMFS issued a final rule revising the listing status of the species (81 FR 62259), in which 14 distinct population segments were identified. Of these, nine did not warrant listing under the ESA, four were listed as endangered, and one was listed as threatened. In the North Pacific, there are four discrete and significant DPS, identified by breeding location: *Hawaii*, *Central America*, *Mexico*, and *Western North Pacific*. Humpback whales of the Oregon, Washington, and California coast are from the Central America, Mexico and Hawaii DPS (Barlow et al. 2011). Only the Mexico DPS and Central America DPS are listed, as threatened and endangered, respectively.

Breeding locations in the North Pacific are more geographically separated than feeding areas and include regions offshore of Hawaii, Central America; the west coast of Mexico, and the Ogasawara and Okinawa Islands and the Philippines. Feeding areas in the North Pacific range from California, USA to Hokkaido, Japan, with most feeding occurring in coastal waters. Humpback whales in the North Pacific rarely move between these breeding regions. Strong fidelity to both feeding and breeding sites has been observed but movements are complex (Calambokidis et al. 2008; Barlow et al. 2011). In general, Asia and Mexico/Central America are the dominant breeding areas for humpback whales that migrate to feeding areas in lower latitudes and coastal California and Russia. The Revillagigedo Islands and Hawaiian Islands

wintering areas for humpback whales that feed in the more central and higher latitude areas (Calambokidis *et al.* 2008). Exceptions to this pattern exist, and complex population structure and strong site fidelity appear to coexist with lesser known, but potentially high, levels of plasticity in the movements of humpback whales (Salden *et al.* 1999).

The Hawaii DPS is composed of humpback whales that breed within the main Hawaiian Islands. Whales from this DPS use most known feeding grounds in the North Pacific; half migrate to feeding grounds in southeast Alaska and northern British Columbia, with many also using northern Gulf of Alaska and Bering Sea feeding grounds. The Central America DPS is composed of humpback whales that breed along the Pacific coast of countries in Central America. Whales from this DPS feed almost exclusively offshore of California and Oregon in the eastern Pacific, with a few individuals in the northern Washington-southern British Columbia feeding grounds. The Mexican DPS is composed of humpback whales that breed along the Pacific coast of mainland Mexico, Baja California, and the Revillagigedo Islands. Whales from this DPS feed across a broad geographic range from California to the Aleutian Islands, with concentrations in California-Oregon, northern Washington-southern British Columbia, northern and western Gulf of Alaska and Bering Sea feeding grounds. The Western North Pacific DPS is a combined DPS composed of humpback whales that breed/winter around Okinawa and the Philippines (Okinawa/Philippines DPS) and a second population that transit the Ogasawara area but breed in an unknown location (Second West Pacific DPS). Whales from the Okinawa/Philippines portion of the DPS migrate to feeding grounds in the northern Pacific, primarily off the Russian coast, while whales from the Second West Pacific DPS are linked to the Aleutian Islands feeding grounds.

A recent analysis of genetic variation in >2000 humpback whales found support for DPS designation in substantial level of genetic divergence among breeding areas at the mtDNA control region (Baker *et al.* 2013). For example, humpback whales in Central America have a unique mtDNA signature (Baker *et al.* 2008a; Baker *et al.* 2008b). The Hawaii population is separated from distant but neighboring populations in both frequencies of mtDNA haplotypes and nDNA (microsatellite) alleles (Baker *et al.* 2013). In Mexico, mtDNA haplotype frequencies in mainland and the Revillagigedo Islands humpback populations were not significantly different (Baker *et al.* 2013) and were thus considered a single population.

Recent humpback whale abundance estimates for the entire North Pacific basin have ranged from 18,302 (Calambokidis *et al.* 2008) to 21,808 individuals (Barlow *et al.* 2011); the latter estimate may still be an underestimate of actual humpback whale abundance. For the lower estimate, whale populations in breeding areas have been estimated at 10,000 individuals in Hawaii, 500 for Central America, 6,000-7,000 animals in Mexico, and 1,000 for the Western Pacific, for a total of 17,500-18,500. Barlow *et al.* (2011) did not apportion the 21,808 individuals to breeding areas, but the proportions are likely to be like those estimated by Calambokidis *et al.* (2008). Growth rates have been calculated on regional scales and include ~8%/year for the U.S. West Coast (1991-2008; Calambokidis 2009), 6.6%/year for the Alaskan Peninsula and Aleutian Islands (2001-2003; Zerbini *et al.* 2010), and 10.6%/year in southeast Alaska (1991-2007; Dahlheim *et al.* 2009), 5.5-6.0%/year for Hawaii and 6.7%/year in the western Pacific (1990-1993, NPAC and 2004-2006, SPLASH; Calambokidis *et al.* 2008).

Humpback whales face a variety of threats, depending on the region in which they occur. Threats listed in the Recovery Plan include entrapment and entanglement in fishing gear, collisions with ships, acoustic disturbance, habitat degradation, and competition for resources with humans (NMFS 1991). Climate change and ocean acidification are also global threats to marine ecosystems that could indirectly affect humpback whales via trophic dynamics and available prey. Globally, entrapment and entanglement in fishing gear and collisions with ships represent most of the reported and observed serious injuries and mortalities for the species (review in Carretta et al. 2014b). Entanglement data are available for most stocks of humpback whales worldwide. These entanglements result from humpback whale interactions with a variety of fisheries and gear types and generally result in some level of serious injury and mortality. The absolute number of humpback whale entanglements is likely under-represented by these data, in part because observer programs and stranding networks do not exist in many parts of the world.

### ***Threats from Fishing Gear Entanglements***

Humpback whales may break through, carry away, or become entangled in fishing gear. Whales carrying gear may later die, become debilitated or seriously injured, or have normal functions impaired, all without having been recorded. Of nations reporting to the IWC, 64.7% (n=11) reported humpback whale bycatch from 2003-2008 (Mattila and Rowles 2010). Some countries (e.g., U.S., Canada, Australia, South Africa) have well-developed reporting and response networks collecting information on entanglements. Still, <10% of humpback whale entanglements in the Gulf of Maine are reported, despite strong outreach and a response network (Robbins and Mattila 2004). For whales off the U.S. East Coast, 89% of removed gear was pots/traps or gillnet gear, although other gear types were observed (Johnson *et al.* 2005). A wide range of entangling gear has also been reported in the South Pacific (Neilson 2006; Lyman 2009), Newfoundland (Lien *et al.* 1992) and by the IWC (Mattila and Rowles 2010). In the North Pacific, entanglement is pervasive but highest among coastal populations (Robbins *et al.* 2007a; Robbins 2009).

Entanglement may result in only minor injury, or potentially may significantly affect individual health, reproduction or survival. Studies of the fate of entangled whales in the Gulf of Maine suggest that juveniles are less likely than adults to survive (Robbins *et al.* 2008), and observed entanglement deaths and serious injuries in that region are known to exceed what is considered sustainable for the population (Glass *et al.* 2009). Most deaths likely go unobserved and preliminary studies suggest that entanglement may be responsible for 3-4% of total mortality, especially among juveniles (Robbins *et al.* 2009).

Much more is known about fishing gear entanglement in the Northern Hemisphere than is in the Southern Hemisphere. Off Japan, an entangled whale is legally allowed to be killed and sold on the market (Lukoschek *et al.* 2009), so entanglement often leads to death for humpback whales in this region. While the number of reported bycaught animals is not large (3 to 5), the number of reports has been increasing and may underestimate the actual number caught. The Mexico population has one of highest scar rates from nets and lines in the North Pacific, indicating a high entanglement rate. Based on this information, the severity of the threat of fishing gear entanglements varies among regions and ranges from low to high.

### *Hawaii DPS*

Recent studies indicate that the Hawaii DPS experiences a high rate of interaction with fishing gear (20-71%), with the highest rates recorded in southeast Alaska and northern British Columbia (Neilson *et al.* 2009). Fatal entanglements of humpback whales in fishing gear have been reported in all areas, and observed fatalities are almost certainly under-reported. Recent studies in another humpback whale feeding ground, which has similar levels of scarring, estimate that the actual annual mortality rate may be as high as 3.7% (Angliss and Outlaw 2008). The level of threat from fishing is considered medium and is not expected to significantly diminish population growth.

### *Central America DPS*

Vessel collisions and entanglement in fishing gear pose the greatest threat to this population, especially off Panama, southern California, and San Francisco. Between 2004 and 2008, there were 18 reports of humpback whale entanglements in commercial fishing gear off California, Oregon, and Washington (Carretta *et al.* 2010), and the actual number of entanglements may be higher. Effective fisheries monitoring and stranding programs exist in California, but are lacking in Central America and much of Mexico. Levels of mortality from entanglement are unknown, but entanglement scarring rates indicate a significant interaction with fishing gear. The Central America DPS is therefore considered to be at moderate risk of extinction over the next three generations.

### *Mexico DPS*

Of the 17 records of stranded North Pacific humpback whales in the NMFS stranding database, three involved fishery interactions, two were attributed to vessel strikes, and in five cases the cause of death could not be determined (Carretta *et al.* 2010). Specifically, between 2004 and 2008, 14 humpback whales were reported seriously injured in commercial fisheries offshore of California and two were reported dead. What proportion of these represent the Mexican breeding population is unknown, but the fishing gear involved included gillnet, pot, and trap gear (Carretta *et al.* 2010). The Mexico DPS is considered to be “not at risk” of extinction, although some voted for “moderate risk” reflect the threat of entanglement among other threats.

### *Western North Pacific DPS*

Whales along the coast of Japan and Korea are at risk of entanglement related mortality in fisheries gear, although overall rates of net and rope scarring are similar to other regions of the North Pacific (Brownell *et al.* 2000). The threat of mortality from any such entanglement is high given the incentive for commercial sale allowed under Japanese and Korean legislation (Lukoschek *et al.* 2009). The reported number of humpback whale entanglements/deaths has increased for Japan since 2001 due to improved reporting, although the actual number of entanglements may be underrepresented in both Japan and Korea (Baker *et al.* 2006). The level of confidence in understanding the minimum magnitude of this threat is medium for the Okinawa/Philippines DPS and low for the Second West Pacific DPS, given the unknown wintering grounds and primary migratory corridors.

Fishing gear entanglements are considered likely to moderately reduce the population size or the growth rate of the Hawaii, Central America, and Mexico DPSs and are likely to seriously reduce the population size or the growth rate of the Western North Pacific [Okinawa/Philippines] DPS.

## West Coast Groundfish Fishery

The West Coast Groundfish Fishery (WCGF) is a multi-species fishery that utilizes a variety of gear types. The fishery harvests species designated in the Pacific Coast Groundfish Fishery Management Plan (PFMC 2011) and is managed by the Pacific Fishery Management Council (PFMC). Over 90 species are listed in the groundfish FMP, including a variety of rockfish, flatfish, roundfish, skates, and sharks. These species are found in both federal (> 5.6 km off-shore) and state waters (0-5.6 km). Groundfish are both targeted and caught incidentally by trawl nets, hook-&-line gear, and fish pots.

Under the FMP, the groundfish fishery consists of four management components:

The Limited Entry (LE) component encompasses all commercial fishers who hold a federal limited entry permit. The total number of limited entry permits available is restricted. Vessels with an LE permit are allocated a larger portion of the total allowable catch for commercially desirable species than vessels without an LE permit.

The Open Access (OA) component encompasses commercial fishers who do not hold a federal LE permit. Some states require fishers to carry a state-issued permit for certain OA sectors.

The Recreational component includes recreational anglers who target or incidentally catch groundfish species. Recreational fisheries are not covered by this report.

The Tribal component includes native tribal commercial fishers in Washington State that have treaty rights to fish groundfish. Tribal fisheries are not included in this report, with the exception of the observed tribal at-sea Pacific hake (*Merluccius productus*) (also known as whiting) sector.

These four components are further subdivided into sectors based on gear type, target species, permits and other regulatory factors. This report includes data from the following sectors:

### *Limited Entry (LE) sectors*

Beginning in 2011, an Individual Fishing Quota (IFQ) program for the LE bottom trawl fleet and the at-sea Pacific hake fleet was implemented, under the West Coast Groundfish Trawl Catch Share Program.

- IFQ fishery (formerly LE bottom trawl and at-sea Pacific hake, 2002-2010): This sector is subdivided into the following components due to differences in gear type and target strategy:
  - Bottom trawl: Bottom trawl nets are used to catch a variety of non-hake groundfish species. Catch is delivered to shore-based processors. Some bottom trawl vessels began carrying electronic monitoring (EM) equipment in 2015 for the purposes of compliance with catch quotas under the IFQ program.



- Midwater non-hake trawl/Midwater Rockfish: Midwater trawl nets are used to target midwater non-hake species, mainly mid-water rockfish such as yellowtail and widow rockfishes. Catch is delivered to shore-based processors. Some midwater trawl vessels began carrying electronic monitoring (EM) equipment in 2015 for the purposes of compliance with catch quotas under the IFQ program.
- Pot: Pot gear is used to target groundfish species, primarily sablefish (*Anoplopoma fimbria*). Catch is delivered to shore-based processors. Some pot vessels began carrying electronic monitoring (EM) equipment in 2015 for the purposes of compliance with catch quotas under the IFQ program.
- Hook-and-line: Longlines are primarily used to target groundfish species, mainly sablefish. Catch is delivered to shore-based processors.
- LE California halibut (*Paralichthys californicus*) trawl: Bottom trawl nets are used to target California halibut by fishers holding both a state California halibut permit and an LE federal trawl groundfish permit. Catch is delivered to shore-based processors.
- Shoreside Pacific hake trawl/Midwater hake: Midwater trawl nets are used to catch Pacific hake. Catch is delivered to shore-based processors.
- At-sea motherships and catcher-processors: Midwater trawl nets are used to catch Pacific hake. Catcher vessels deliver unsorted catch to a mothership. The catch is sorted and processed aboard the mothership. Catcher-processors catch and process at-sea. This component also includes the at-sea processing component of the tribal sector. The tribal sector must operate within defined boundaries in waters off northwest Washington. Tribal catch can be delivered to a contracted mothership by catcher vessels for processing or be caught and processed by a contracted catcher-processor.
- LE fixed gear (non-nearshore): This sector is subdivided into two components due to differences in permitting and management:
  - LE sablefish endorsed season: Longlines and pots are used to target sablefish. Catch is generally delivered to shore-based processors.
  - LE sablefish non-endorsed: Longlines and pots are used to target groundfish, primarily sablefish and thornyheads. Catch is delivered to shore-based processors or sold live at the dock.

#### Open Access (OA) Federal sectors

- OA fixed gear (non-nearshore): Fixed gear, including longlines, pots, fishing poles, stick gear, etc. is used to target non-nearshore groundfish. Catch is delivered to shore-based processors.

#### Open Access (OA) state sectors

- OA ocean shrimp (*Pandalus jordani*) trawl: Trawl nets are used to target ocean shrimp. Catch is delivered to shore-based processors.
- OA California halibut trawl: Trawl nets are used to target California halibut by fishers holding a state California halibut permit. Catch is delivered to shore-based processors.
- Nearshore fixed gear: A variety of gear, including longlines, pots, fishing poles, stick gear, etc. are used to target nearshore rockfish and other nearshore species managed by state permits in Oregon and California. Catch is delivered to shore-based processors or sold live.

## **Northwest Fisheries Science Center Groundfish Observer Programs**

The NWFSC Groundfish Observer Program's goal is to improve estimates of total catch and discard by observing commercial sectors of groundfish fisheries along the U.S. west coast that target or take groundfish as bycatch. The observer program has two units: the West Coast Groundfish Observer Program (WCGOP) and the At-Sea Hake Observer Program (A-SHOP). The WCGOP Program was established in May 2001 by NOAA Fisheries (a.k.a., National Marine Fishery Service, NMFS) in accordance with the Pacific Coast Groundfish Fishery Management Plan (50 CFR Part 660) (50 FR 20609). This regulation requires all vessels that catch groundfish in the US EEZ from 3-200 miles offshore carry an observer when notified to do so by NMFS or its designated agent. Subsequent state rule-making has extended NMFS's ability to require vessels fishing in the 0-3 mile state territorial zone to carry observers.

The WCGOP and A-SHOP observe distinct sectors of the groundfish fishery. The WCGOP observes the following sectors: IFQ shore-based delivery of groundfish and Pacific hake, LE and OA fixed gear, and state-permitted nearshore fixed gear sectors. The WCGOP also observes several state-managed fisheries that incidentally catch groundfish, including the California halibut trawl and ocean shrimp trawl fisheries. The A-SHOP observes the IFQ fishery that delivers Pacific hake at-sea including: catcher-processor, mothership, and tribal vessels. Details on how fisheries observers operate in both the IFQ (Catch Share) and Non-IFQ sectors can be found at: <http://www.nwfsc.noaa.gov/research/divisions/fram/observation/index.cfm>.

### **Humpback whale bycatch in West Coast Groundfish Fisheries**

The primary objective of this report is to provide estimates of bycatch of humpback whales in observed U.S. West Coast federally-permitted groundfish fisheries since the last report in 2015, which covered the years 2010–2013. Previous reports on marine mammal bycatch in West Coast groundfish fisheries (Jannot et al. 2011) as well as reports on the NWFSC Protected Species Reports webpage ([http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_products/protected\\_species.cfm](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_products/protected_species.cfm)) have provided data on bycatch of humpback whales in U.S. west coast commercial fisheries, which were derived from Observer Program data.

#### **Amount and Extent of Humpback Whale Take**

The Biological Opinion (BiOp) Regarding the Effects of the Continued Operation of the Pacific Coast Groundfish Fishery (PCGF) (NMFS 2012a) stated that:

We anticipate that take of humpback whales will occur as a result of the proposed continued operation of the PCGF. Incidental take of humpback whales occurs as a result of entanglement with fishing gear, as a consequence of fishing activity. This take is expected to occur in the sablefish pot/trap fishery. In the effects section, we estimated an average of 1 humpback whale per year entangled by proposed fishing, with a maximum of 3 humpback whales entangled in a single year. Therefore, the incidental take limit for humpback whales is a 5-year average of 1 humpback whale injury or mortality per year, and up to 3 humpback whale injuries or mortalities in any single year. Available data on takes will be reviewed

periodically by a Pacific Coast Groundfish and Endangered Species Workgroup as described under Reasonable and Prudent Measures and Terms and Conditions below. In addition to these take limits, we will evaluate total human-caused serious injury and mortality of humpback whales annually, and if PBR is exceeded, we will determine whether the MMPA 101(a)(5)(E) permit and humpback whale ITS are still valid. Consistent with the analysis in this biological opinion, a portion of unidentified whale and gear entanglements would be counted against these take limits and for this PBR evaluation in addition to known humpback whale entanglements in gear of the proposed fishery (pro-rating criteria and methods described in Section 2.3.3 or as adjusted by the Workgroup). Data used to pro-rate unidentified whale and gear entanglements will be updated each year. These criteria and methods are conservative in light of uncertainty about proposed fishery impacts on humpback whales, because of the opportunistic nature of entanglement observation and reporting, potential for unobserved injury or mortality because of entanglements, and difficulty identifying entangled whales to species and entangling gear to specific fisheries.

This first biennial report represents the fulfillment of the take estimate requirement and associated reporting requirements.

## **Methods**

### **Data Sources**

Data sources for this analysis include onboard observer data from the WCGOP and A-SHOP and landing receipt data, referred to as fish tickets, and obtained from the Pacific Fisheries Information Network (PacFIN).

#### **Observer Data**

A list of fisheries, coverage priorities and data collection methods employed by WCGOP in each observed fishery can be found in the Catch Shares (IFQ) and Non-Catch Shares (Non-IFQ) WCGOP manuals (NWFSC 2015a, b). A-SHOP information and documentation on data collection methods can be found in the A-SHOP observer manual (NWFSC 2014).

The sampling protocol employed by the WCGOP is primarily focused on the discarded portion of catch. To ensure that the recorded weights for the retained portion of the observed catch are accurate, haul-level retained catch weights recorded by observers are adjusted based on trip-level fish ticket records. This process is described in detail on the WCGOP Data Processing webpage ([http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_processing.cfm](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_processing.cfm)). Data processing was applied prior to the analyses presented in this report. For a complete list of groundfish species defined in the Pacific Coast Groundfish Fishery Management Plan see PFMC (2011).

#### **Fish Ticket Data**

For bycatch estimation, the landed amount of a particular fish species or species group is used as the effort metric. Thus, the retained landing information from sales receipts (known as fish tickets) is crucial for fleet-wide total bycatch estimation for all sectors of the commercial groundfish fishery on the U.S. west coast. Fish ticket landing receipts are completed by fish-buyers in each port for each delivery of fish by a vessel. Fish tickets are trip-aggregated sales receipts for market categories that may represent single or multiple species. Fish tickets are issued to fish-buyers by a state agency and must be returned to the issuing agency for processing. Fish tickets are designed by the individual states (Washington, Oregon, and California) with slightly different formats by state. In addition, each state conducts species-composition sampling at the ports for numerous market categories that are reported on fish tickets. Fish ticket and species-composition data are submitted by state agencies to the PacFIN regional database. Annual fish ticket landings data for 2010-2013, with state species composition sampling applied, were retrieved from the PacFIN database in 2014 and subsequently divided into various sectors of the groundfish fishery. Observer and fish ticket data processing steps are described in detail on the WCGOP website under Data Processing Appendix ([http://www.nwfsc.noaa.gov/research/divisions/fram/observer/data\\_processing.cfm/](http://www.nwfsc.noaa.gov/research/divisions/fram/observer/data_processing.cfm/)). All data processing steps specific to this report are described in the bycatch estimation methods section below.

### **Designation of ‘take’ and ‘serious injury’ interactions**

NMFS has established guidelines for distinguishing serious from non-serious injury of marine mammals pursuant to the Marine Mammal Protection Act through a policy directive (NMFS 2012b).

### **Bycatch Estimation**

#### **Statistical Model**

We applied statistical models to observer program data to characterize uncertainty in humpback whale bycatch estimation in the sablefish pot fishery (Table 1). Because only one humpback whale was documented as bycatch, we were restricted to using simple statistical models while estimating variances of total bycatch. The first approach we used was the Poisson process model, where the total number of entanglements or bycatch events were assumed to follow a Poisson distribution,

$$n_{bycatch} \sim (\lambda_1 \cdot N_y).$$

In this approach, the Poisson rate or intensity parameter ( $\lambda_1$ , where  $0 \leq \lambda_1 \leq 1$ ) was fixed at the annual bycatch point estimate (*e.g.*, 1 bycatch events out of 1000 sets would lead to  $\lambda_1 = 0.001$ ), and the effort for a particular year ( $N_y$ ) was used to estimate the total bycatch. A caveat of this first approach was that by fixing  $\lambda_1$ , we were ignoring the uncertainty in the bycatch rate, making the 95% CIs overly narrow. For example, two fishery sectors might have the same bycatch point estimate, but if one sector fished with 10x as much effort, that second estimate would be more precise. To incorporate this uncertainty due to variable sample sizes, our second approach was to treat the rate parameter as a random variable ( $\lambda_2$ , where  $0 \leq \lambda_2 \leq 1$ ). We did not use a

common approach to model uncertainty in the proportion  $p$  of a Binomial distribution using the Normal approximation,

$$p \sim (\hat{p}, \sqrt{\hat{p}(1-\hat{p})}),$$

where  $\hat{p}$  is the estimated proportion and  $n$  is the sample size, because the 95% CIs can include negative values due to the small estimated proportion. To keep this parameter ( $p$ ) positive, we instead simulated the number of bycatch events that would have occurred given a certain level of effort, and divided that result by effort. Using our previous numbers as an example,  $\lambda_2 \sim (p = 0.001, N = 1000)/1000$ . Both approaches require at least one bycatch event. For each model, we generated 100,000 random draws from the distributions of potential bycatch and calculated summary statistics (mean, median, and variance) as well as measures of uncertainty (95% CIs).

Table 1. Observer data for pot sablefish fisheries used in the estimation of bycatch for humpback whales.

Year	Gear	# vessels	# trips	# hauls	# hooks or pots	Observed Landings (mt)	Total Landings (mt)	Coverage rate	Takes
2002	Pot	6	23	249	5,496	82.5	352.2	23%	0
2003	Pot	6	35	370	9,257	148.3	604.2	25%	0
2004	Pot	3	13	140	5,408	82.7	619.6	13%	0
2005	Pot	7	39	495	13,899	281.1	615	46%	0
2006	Pot	7	39	291	10,805	200.5	581.8	34%	0
2007	Pot	4	31	158	6,046	90	428.4	21%	0
2008	Pot	6	24	330	13,679	244.9	433	57%	0
2009	Pot	3	27	67	3,883	66.5	489.1	14%	0
2010	Pot	7	43	316	11,364	139.4	503.5	28%	0
2011	Pot	3	22	228	9,070	137.4	371.9	37%	0
2012	Pot	5	19	355	14,337	101.1	286	35%	0
2013	Pot	3	15	49	1,938	40.5	283.1	14%	0
2014	Pot	4	16	195	7,574	104	337.3	31%	1
2015	Pot	9	35	299	11,329	218.8	358.1	61%	0

## Results and Discussion

### Estimating Humpback Whale Bycatch

Estimation of the fleet-wide bycatch is challenging because only one humpback whale has been observed entangled in U.S. west coast groundfish fisheries since the observer program began in 2002. To align with the incidental take statement in the 2012 Biological Opinion, we estimated the total bycatch in the most recent two 5-year periods (2006-2010, 2011-2015). No humpback whales were observed as bycatch in the first 5-year period (2006-2010), thus the 5-year sum of take was zero. For the time period 2011-2015, one humpback whale was observed taken in the U.S. west coast groundfish fisheries; it was in 2014 in the Limited Entry sablefish fishery on a vessel fishing pot gear. Using observer data from this sector, the bycatch rate calculated for the 2011-2015 period was 0.002 whales/year. Given the observer coverage over this time period (14

– 61%), this produced a fleet-wide estimate of 1.4 individuals caught by the LE sablefish pot fleet during the 5-year period (2011-2015; Table 2).

Table 2. The number of humpback whale entanglements, landings, bycatch ratio and estimated takes from the Limited Entry fixed gear pot fishery during the last two 5-year periods. Landings and observed take values are summed for each 5-year period; bycatch ratio is calculated as the number of humpback whales/observed landings for each period. The 2011-2015 values were used in the probability models.

Five-year period	# observed takes (5-year sum)	Observed mean sablefish (mt)	Observed sablefish landings (mt)	Bycatch ratio	Fleet-wide mean sablefish (mt/yr)	Fleet-wide 5-year sum sablefish (mt)	Fleet-wide humpback whales take estimate
2006-2010	0	148.24	741.18	0	322.31	1611.56	0
2011-2015	1	120.37	601.85	0.002	165.61	828.04	1.38

We also calculated probabilistic estimates of exceeding numbers of entanglements for the period 2011-2015 (Fig. 1) based on the two versions of the Poisson statistical model described above.




Figure 1. The probability of exceeding the numbers of entanglements/takes for a model with a fixed bycatch rate (solid line) and a model including uncertainty in the bycatch rate (dashed line). The dotted

gray vertical line represents the maximum allowable entanglements across 5 years under the Biological Opinion for humpback whales in the U.S. west coast groundfish fisheries.

The model with a fixed bycatch rate (solid line, Fig. 1) indicated that the probability of exceeding five humpback whale entanglements in this 5-year period (grey dotted vertical line intersects with solid black line) was just under 40%. Exceeding five humpback whales is equivalent to exceeding an average of one humpback whale per year for a 5-year period. If we model the bycatch rate with uncertainty, the probability of exceeding five entanglements approaches 0% (grey dotted vertical line intersects with dashed black line, Fig. 1). The probability of more than five entanglements decreased more rapidly when uncertainty was included in the model, compared to the model without uncertainty. The consequence of observing only one bycatch incident in the 5-year period was large variation in entanglement in both models (Table 3). We could only apply these models for the period 2011-2015, as that was the only period in which a humpback whale was entangled in gear in a U.S. west coast groundfish fishery. Extrapolating these models beyond the 2011-2015 time period or to other gear types or fishery sectors would not be appropriate.

Table 3. Summary statistics of the number of humpback whale entanglements for the 2011-2015 period based on 100,000 random draws from the fixed bycatch rate model (no uncertainty) and the model with uncertainty in bycatch rate. See text for model descriptions.

Statistic	Fixed Rate	Rate with Uncertainty
Minimum	~ 0	~ 0
Mean	0.27	0.27
Median	~ 0	~ 0
Maximum	5	8
Variance	0.27	0.35
Standard Deviation	0.52	0.59

The 2012 Biological Opinion for the continuing operation of West Coast groundfish fisheries set an incidental take limit for humpback whales at three humpback whale injuries or mortalities in any single year, and an average of one humpback whale injury or mortality per year over a five-year period. The actual entanglements/takes observed in the LE sablefish pot fishery are well below this threshold, as are the fleet-wide entanglements/take estimates modeled using landings and observer coverage.

### **Additional relevant information**

#### **Interactions with other pot/trap fisheries**

Pot and trap fisheries in general are the most commonly documented source of serious injury and mortality of humpback whales in U.S. west coast waters (Carretta et al. 2013, 2015, 2016a). From 2010 to 2014, there were 27 documented interactions associated with pot and trap fisheries (Carretta et al. 2016a, Jannot et al. 2016). Eighteen instances involved serious injuries (prorated and non-prorated) attributed to unidentified pot/trap fisheries (12 total serious injuries), Washington coastal dungeness crab pot (1), California dungeness crab pot (1), and California

spot prawn pot (0.75); this totaled 14.75 serious injuries/5 years, or 2.95 humpback whales annually. Five records (three in the California spot prawn pot fishery and two in unidentified pot/trap fisheries) involved non-serious injuries resulting from human intervention to remove gear, or cases where animals were able to free themselves. Four records involved dead whales, including one case where a pair of severed humpback flukes was found in southern California waters with two sets of California dungeness crab gear attached (Carretta et al. 2016a). Including the four deaths attributed to pot/traps, the minimum level of annual mortality and serious injury across all pot/trap fisheries is  $14.75+4=18.75/5$  years or 3.75 whales annually. In some cases the entangling gear could not be identified to fishery. In order to evaluate the effects of different fishing gear on humpback whales and other ESA listed whales, the workgroup would like to again recommend that buoys from groundfish pot gear and longlines be marked sufficiently for identification of fishery. Improved marking will allow more accurate accounting of which fisheries are interacting with whales.

### **Changes in Humpback Whale occurrence**

In recent years the distribution and duration of time humpback whales stay on the feeding grounds has changed. More humpback whales have been observed in Puget Sound, the mouth of the Columbia River, San Francisco Bay, and closer to shore in general than has been observed since the end of commercial whaling. Hydrophones and vessel surveys have also reported humpback whale detections later into the winter that has been observed in the past with some evidence that individuals may be over-wintering. To date, we have not seen the changes in humpback distributions affect interactions with groundfish fisheries but it will be important to continue monitoring because the changes in humpback whale distributions and duration on the feeding ground may increase the opportunity for the whales to have interactions with fisheries that have not been previously documented.

### **Conservation Recommendations**

Section 7(a)(1) of the ESA directs Federal agencies to use their authorities to further the purposes of the ESA by carrying out conservation programs for the benefit of the threatened and endangered species. Specifically, conservation recommendations are suggestions regarding discretionary measures to minimize or avoid adverse effects of a proposed action on listed species or critical habitat or regarding the development of information (50 CFR 402.02). The following conservation recommendations for humpback whales described in the BiOp provide general guidance for unique, visual marking of sablefish pot/trap gear as identifiable to a specific fishery, as well as guidance to report, track, and retrieve pot/trap gear that becomes lost, and guidance to minimize the loss of pot/trap gear. Implementing these recommendations would improve our knowledge of incidental take of humpback whales in the PCGF and minimize that take. Washington and Oregon commercial Dungeness crab fisheries are example models where regulations for unique, visual marking of gear and programs to report, track, and retrieve lost gear are established. Citations regarding these regulations and programs are provided below. These measures shall be further discussed and developed by the PCGW, who may recommend adoption as conservation measures.



- 1) NMFS and the PCGW should work with the PFMC to require or recommend visual marking that can be used to uniquely identify sablefish pot/trap gear (e.g., OAR 635-005-0480 and WAC 220-52-040 for Dungeness Crab Buoy Tag and Gear Marking Requirements). Visual marking can help identify gear entangled on a whale to a specific fishery, while absence of visual markings can also help rule out a fishery that uses unique, visual markers.
- 2) NMFS and the PCGW should work with the PFMC to create electronic monitoring and logbook reporting requirements for the sablefish pot/trap fishery that require or recommend fishers to document effort and lost gear (see Appendix C for example logbook regulations, instructions, and entry forms that include lost gear reporting).
- 3) NMFS and the PCGW should work with the PFMC to develop a database to track sablefish pot/trap fishing effort, locations, and lost fixed-gear (see Appendix D for an example database).
- 4) NMFS and the PCGW should work with the PFMC to summarize data on lost gear from the sablefish pot/trap fishery to evaluate the magnitude of gear loss and factors that may influence loss (specific areas, times of year, etc.). Also, summarize fixed-gear fishing effort and locations to support overlap analysis with humpback whale (or other large whale) migrations or aggregation. Data summary should follow the reporting cycle developed for the PCGW above.
- 5) NMFS and the PCGW should work with the PFMC to promote retrieval of lost gear (see Appendix E and Appendix F for information about example programs for gear recovery).
- 6) NMFS and the PCGW should work with the PFMC to assess available technology to minimize loss of sablefish pot/trap gear (i.e., Gearfinder technology) and promote use of appropriate technology.
- 7) NMFS and the PCGW should work with the PFMC to investigate the practice of storing sablefish pot/trap gear in the ocean to evaluate the potential for conservation issues and any need for additional regulation.

*The NWFSC Observer Program presently collects information on sablefish pot/trap fishing effort, locations, and lost fixed gear on observed vessels, and this information is in the program database. The Fishing Effort Report, submitted as part of the Biological Opinion process, summarizes the Sablefish fishing effort by gear, area, and depth. In addition, that report also summarizes information regarding lost gear. As the Sablefish fleet is currently observed at less than 100% coverage and there is no logbook associated with the fishery, the data available represent the observed portion of the fleet. NMFS has consulted the council's Coast Guard representative about this issue, and the representative felt that the legality of the practice depends on where and how the gear is stored; importantly, the practice likely takes place inside state waters, which complicates matters.*

### **Additional information relevant to the BiOp/RPMs and Conservation Measures**

In the incidental take statement in the BiOp, we included reasonable and prudent measures for management planning and take reporting that is applicable to all species considered in the BiOp. “Reasonable and prudent measures” are nondiscretionary measures to minimize the amount or

extent of incidental take (50 CFR 402.02). “Terms and conditions” implement the reasonable and prudent measures (50 CFR 402.14). These must be carried out for the exemption in section 7(o)(2) to apply. Here we provide the reasonable and prudent measures, including species specific measures described in the BiOp, followed by additional information on the status of each measure. Included were measures to minimize the amount or extent of incidental take associated with NMFS observer program sampling and handling of protected species where these effects are not otherwise authorized or exempted.

(1) NMFS shall develop a Pacific Coast Groundfish and Endangered Species Workgroup

*NMFS has convened a Pacific Coast Groundfish and Endangered Species Workgroup (PCGW), invited PFMC and other entities to provide points of contact, and helped develop terms of reference for the workgroup. The 1<sup>st</sup> meeting of the Pacific Coast Groundfish and Endangered Species Workgroup was in May 2015.*

(2) NMFS shall characterize changes in fishing effort.

*A report has been compiled summarizing fishing effort in the U. S. Pacific Coast Groundfish Fisheries (2002-2013) by NOAA Fisheries’ West Coast Groundfish Observer Program (Somers et al. 2015).*

(3) NMFS shall update reporting of take considered in this opinion.

*NMFS updates reporting of take on a biennial basis per the BiOp.*

(4) NMFS shall update the NWFSC risk assessment, as needed.

*If necessary, NMFS will update the BA's risk assessment for humpback whale.*

### **Species-Specific Measures**

NMFS included the following reasonable and prudent measure to improve our knowledge of incidental take of humpback whales in the PCGF.

(1) NMFS shall provide all west coast observers with the Fixed Gear Guide ([http://swr.nmfs.noaa.gov/psd/Fixed%20Gear%20Guide-FINAL\\_12.14.11.pdf](http://swr.nmfs.noaa.gov/psd/Fixed%20Gear%20Guide-FINAL_12.14.11.pdf)) and the entangled whale hotline (877-SOS-WHALE) during observer training. The guide will help observers that may opportunistically sight an entangled whale identify the entangling gear to a specific fishery. The hotline provides a resource for reporting and response.

*All observers are trained to identify Humpback Whales and are deployed with a marine mammal identification guide. Observers are also provided with the Fixed Gear Guide and the Whale Hotline number and trained to contact the hotline if they observe a whale entanglement. Furthermore, observers are provided with the Marine Mammal Reporting Form, which is given to and submitted by the fisher, when an incidental mortality or injury occurs during commercial fishing activity.*

### **Species-specific Terms and Conditions**

The terms and conditions described in the BiOp are non-discretionary, and NMFS must comply with them in order to implement the reasonable and prudent measures (50 CFR 402.14). NMFS has a continuing duty to monitor the impacts of incidental take and must report the progress of the action and its impact on the species as specified in the incidental take statement (50 CFR 402.14). If the terms and conditions in the BiOp are not complied with, the protective coverage of section 7(o)(2) will likely lapse. Terms and conditions specific to humpback whales are provided below.

1.a. Reporting shall be directed from observers through the observer program.

1.b. Reporting shall be similar to or modeled after the attached form (Appendix B of the BiOp).

*The observer program has a data collection form for interactions of marine mammals and other protected species with fishing vessels. This form can be found at:  
[http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_collection/manuals/cs\\_manual\\_2015/CS%202015%20Chapter%208.pdf](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_collection/manuals/cs_manual_2015/CS%202015%20Chapter%208.pdf).*

## **References**

- Baker CS, Medrano-Gonzalez L, Calambokidis J, Perry A, Pichler F, Rosenbaum H, Straley JM, Urban-Ramirez J, Yamaguchi M, Von Ziegeler O. 1998. Population structure of nuclear and mitochondrial DNA variation among humpback whales in the North Pacific. *Molecular Ecology* 7: 695-707.
- Baker C. S. and D. Steel. 2010. geneSPLASH: genetic differentiation of ‘ecostocks’ and ‘breeding stocks’ in North Pacific humpback whales. In: Symposium on the results of SPLASH humpback whale study. Final report and recommendations (Calambokidis J., ed.). Quebec City, Canada, pp. 58-59.
- Barlow, J., Calambokidis, J., Falcone, E. A., Baker, C. S., Burdin, A. M., Clapham, P. J., Ford, J. K. B., Gabriele, C. M., LeDuc, R., Mattila, D. K., Quinn, T. J., Rojas-Bracho, L., Straley, J. M., Taylor, B. L., Urbán R., J., Wade, P., Weller, D., Witteveen, B. H. and Yamaguchi, M. 2011. Humpback whale abundance in the North Pacific estimated by photographic capture-

- recapture with bias correction from simulation studies. *Marine Mammal Science* 27: 793–818.
- Barlow J., M. Ferguson, E. Becker, J. Redfern, K. A. Forney, I. Vilchis, P. Fiedler, T. Gerrodette, and L. Ballance. 2009. Predictive modeling of cetacean densities in the eastern Pacific Ocean. NOAA Technical Memorandum NMFS-SWFSC-TM-444, 206 p.
- Benson S. R., 2002. Ecosystem Studies of Marine Mammals and Seabirds in Monterey Bay, CA, 1996-1999. M.S. Thesis, San Jose State University.
- Benson S. R., Croll DA, Marinovic B, Chavez FP, Harvey JT. 2002. Changes in the cetacean assemblage of a coastal upwelling ecosystem during El Niño 1997-98 and the La Niña 1999. *Progress in Oceanography* 54: 279-291.
- Bettridge, S., C. S. Baker, J. Barlow, P. J. Clapham, M. Ford, D. Gouveia, D. K. Mattila, R. M. Pace, III, P. E. Rosel, G. K. Silber, and P. R. Wade. 2015. Status Review of the Humpback Whale (*Megaptera novaeangliae*) under the Endangered Species Act. U.S. Dept. Commer., NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-540, 240 p.
- Calambokidis J, E.A. Falcone, T. Quinn, A. Burdin, P. Clapham, J. K. B. Ford, C. Gabriele, R. LeDuc, D. Matillia, L. Rojas-Bracho, J. M. Straley, B. Taylor, J. Urban-Ramirez, D. Weller, B. Witteveen, M. Yamaguchi, A. Bendlin, D. Camacho, K. R. Flynn, A. Havron, J. Huggins, and N. Maloney. 2008. SPLASH: Structure of populations, levels of abundance and status of humpback whales in the North Pacific. Final report for contract AB133F-03-RP-00078. Olympia, Washington: Cascadia Research.
- Calambokidis, J., E. Falcone, A. Douglas, L. Schlender, and J. Huggins. 2009. Photographic identification of humpback and blue whales off the U.S. West Coast: results and updated abundance estimates from 2008 field season. Final Report for Contract AB133F08SE2786 from Southwest Fisheries Science Center. 18 p.
- Calambokidis J, G. Steiger, D. Ellifrit, B. Troutman, and C. Bowlby. 2004. Distribution and abundance of humpback whales and other marine mammals off the northern Washington coast. *Fisheries Bulletin* 102: 563-580.
- Carretta, J.V., K.A. Forney, E. Oleson, K. Martien, M.M. Muto, M.S. Lowry, J. Barlow, J. Baker, B. Hanson, D. Lynch, L. Carswell, R.L. Brownell Jr., J. Robbins, D.K. Mattila, K. Ralls, and M.C. Hill. 2012. U.S. Pacific Marine Mammal Stock Assessments: 2011. NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-488. 356 p.
- Carretta, J.V, E. Oleson, D.W. Weller, A.R. Lang, K.A. Forney, J. Baker, B. Hanson, K. Martien, M.M. Muto, A.J. Orr, H. Huber, M.S. Lowry, J. Barlow, D. Lynch, L. Carswell, R.L. Brownell Jr., and D.K. Mattila. 2014a. U.S. Pacific Marine Mammal Stock Assessments, 2013. NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-532. 406 p.

- Carretta, J.V., S.M. Wilkin, M.M. Muto, K. Wilkinson, and J. Rusin. 2014b. Sources of human-related injury and mortality for U.S. Pacific west coast marine mammal stock assessments, 2008-2012. NOAA Technical Memorandum, NOAA-TM-NMFS-SWFSC-533. 110 p.
- Dalla Rosa, L. 2010. Modeling the foraging habitat of humpback whales. PhD thesis, University of British Columbia, Vancouver. 185 p.
- Federal Register, Volume 74, p. 40568. August 12, 2009. Notice of initiation of a status review; request for information: Endangered and Threatened Species; Initiation of a Status Review for the Humpback Whale and Request for Information.
- Federal Register, Volume 80, p. 22303. April 21, 2015. Notice of proposed rule: Endangered and Threatened Species; Identification of 14 Distinct Population Segments of the Humpback Whale (*Megaptera novaeangliae*).
- Fleming, A. and J. Jackson. 2011. Global Review of Humpback Whales (*Megaptera novaeanglia*). NOAA Technical Memorandum NMFS, NOAA-TM-NMFS-SWFSC-474. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center. 206 p..
- Ford, M. J., M. A. Bellman, S. Copps, C. K. Emmons, J. Drake, T. P. Good, R. G. Gustafson, M. B. Hanson, G. Hanshew, E. E. Holmes. 2012. Risk assessment of U.S. West Coast groundfish fisheries to threatened and endangered marine species. Report of the National Marine Fisheries Service, Northwest Fisheries Science Center. Seattle, Washington.
- Forney, K. A. 2007. Preliminary estimates of cetacean abundance along the U.S. west coast and within four National Marine Sanctuaries during 2005. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-406. 27 p.
- Forney, K. A., and J. Barlow. 1998. Seasonal Patterns in the Abundance and Distribution of California Cetaceans, 1991-1992. Marine Mammal Science 14(3): 460-489.
- Hanson, M. B., T. P. Good, and J. E. Jannot. Humpback whale bycatch in the 2010 - 2013 U.S. West Coast Groundfish Fisheries. Report of the National Marine Fisheries Service to the Pacific Fisheries Management Council Endangered Species Workgroup. April 2015.
- Kieckhefer, T. 1992. Feeding ecology of humpback whales in continental shelf waters near Cordell Bank, California. Master's thesis. Moss Landing, CA: Moss Landing Marine Laboratories.
- Lagerquist B, Mate B, Ortega-Ortiz J, Winsor M, Urbán-Ramirez J. 2008. Migratory movements and surfacing rates of humpback whales (*Megaptera novaeangliae*) satellite tagged at Socorro Island, Mexico. Marine Mammal Science 24: 815-830.

- NMFS. 1991. Final Recovery Plan for the humpback whale (*Megaptera novaeangliae*). November 1991. U.S. Department of Commerce. National Oceanic and Atmospheric Administration. National Marine Fisheries Service. Office of Protected Resources. 105 p.
- NMFS (National Marine Fisheries Service). 2012a. Continuing Operation of the Pacific Coast Groundfish Fishery - Endangered Species Act (ESA) Section 7(a)(2) Biological Opinion and Section 7(a)(2) "Not Likely to Adversely Affect" Determination. PCTS Number: NWR-2012-876. 194 p.
- NMFS (National Marine Fisheries Service). 2012b. Guidelines for Distinguishing Serious from Non-Serious Injury of Marine Mammals Pursuant to the Marine Mammal Protection Act. National Marine Fisheries Service Instruction 02-038-01 from J. Lecky. 6 January, 2102. 42 p.
- NWFSC (Northwest Fisheries Science Center). 2009. Data report and summary analyses of the U.S. West Coast limited entry groundfish bottom trawl fishery. West Coast Groundfish Observer Program. NWFSC, 2725 Montlake Blvd E., Seattle, WA 98112. 70 p. Available at: [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_products/datareport/docs/trawl\\_report\\_2009\\_final.pdf](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_products/datareport/docs/trawl_report_2009_final.pdf) [accessed February 2015].
- NWFSC (Northwest Fisheries Science Center). 2010. Data report and summary analyses of the U.S. west coast limited entry groundfish bottom trawl fishery. West Coast Groundfish Observer Program. NWFSC, 2725 Montlake Blvd E., Seattle, WA 98112. 67 p. Available at: [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_products/datareport/docs/trawl\\_report\\_2010.pdf](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_products/datareport/docs/trawl_report_2010.pdf) [accessed February 2015].
- NWFSC (Northwest Fisheries Science Center). 2014. Observer Sampling Manual. Fishery Resource Analysis and Monitoring, At-Sea Hake Observer Program. NWFSC, 2725 Montlake Blvd. East, Seattle, Washington 98112. Online at: [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_collection/manuals/A-SHOP\\_Manual\\_2014.pdf](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_collection/manuals/A-SHOP_Manual_2014.pdf) [accessed February 2015].
- NWFSC (Northwest Fisheries Science Center). 2015a. West Coast Groundfish Observer Program 2015 Catch Shares Training Manual. West Coast Groundfish Observer Program. NWFSC, 2725 Montlake Blvd. East, Seattle, Washington, 98112. Online at: [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_collection/training.cfm](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_collection/training.cfm) [accessed February 2015].
- NWFSC (Northwest Fisheries Science Center). 2015b. West Coast Groundfish Observer Program 2015 Non-Catch Share Training Manual. West Coast Groundfish Observer Program. NWFSC, 2725 Montlake Blvd. East, Seattle, Washington, 98112. Online at: [http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data\\_collection/training.cfm](http://www.nwfsc.noaa.gov/research/divisions/fram/observation/data_collection/training.cfm) [accessed February 2015].

- Olsen, E., and coauthors. 2009. First satellite-tracked long-distance movement of a sei whale (*Balaenoptera borealis*) in the North Atlantic. *Aquatic Mammals* 35(3): 313-318.
- PFMC (Pacific Fishery Management Council). 2011. Pacific Groundfish Fishery Management Plan for the California, Oregon, and Washington Groundfish Fishery. Pacific Fishery Management Council, Portland, OR. Online at: [http://www.pcouncil.org/wp-content/uploads/GF\\_FMP\\_FINAL\\_Dec2011.pdf](http://www.pcouncil.org/wp-content/uploads/GF_FMP_FINAL_Dec2011.pdf).
- Rice, D. W. 1978. The humpback whale in the North Pacific: Distribution, exploitation, and numbers. Pages 29–44 in K. S. Norris and R. R. Reeves, eds. Report on a workshop on problems related to humpback whales (*Megaptera novaeangliae*) in Hawaii. Contract Report to the U.S. Marine Mammal Commission. NTIS PB-280–794. 90 p.
- Somers, K. A., J. E. Jannot, J. Hastie, Y. W. Lee, J. McVeigh, and C. E. Whitmire. 2015. Fishing Effort in the 2002-2013 U.S. Pacific Coast Groundfish Fisheries. West Coast Groundfish Observer Program. National Marine Fisheries Service, NWFS, 2725 Montlake Blvd E., Seattle, WA 98112.
- Stevick, P., J. Allen, P. Clapham, N. Friday, S. Katona, F. Larson, J. Lien, D. Mattila, P. Palsboll, J. Sigurjonsson, T. Smith, N. Oien, and P. S. Hammond. 2003. North Atlantic humpback whale abundance and rate of increase four decades after protection from whaling. *Marine Ecology Progress Series* 258: 263-273.
- Tynan C., D. Ainley, J. Barth, T. Cowles, S. Pierce, and L. Spear. 2005. Cetacean distributions relative to ocean processes in the northern California Current system. *Deep-Sea Res. II* 52: 145-167.
- West Coast Region. 2014. 50 CFR Part 660, Subparts C – G Federal Pacific Coast Groundfish Regulations for Commercial and Recreational Fishing 3-200 Nautical Miles off Washington, Oregon, and California. Prepared by: West Coast Region, National Marine Fisheries Service, 7600 Sand Point Way NE, Seattle, WA 98115. Online at: [http://www.westcoast.fisheries.noaa.gov/publications/fishery\\_management/groundfish/regulations.pdf](http://www.westcoast.fisheries.noaa.gov/publications/fishery_management/groundfish/regulations.pdf) [accessed February 2015].
- Wilson, S. M., G. D. Raby, N. J. Burnett, S. G. Hinch, and S. J. Cooke. 2014. Looking beyond the mortality of bycatch: sublethal effects of incidental capture on marine animals. *Biological Conservation* 171: 61–72.
- Zerbini A, Clapham P, Wade P, 2010. Assessing plausible rates of population growth in humpback whales from life-history data. *Marine Biology* 157: 1225-1236.