

Status of Pacific ocean perch (*Sebastes alutus*) along the US west coast in 2017



Chantel R. Wetzel¹
Lee Cronin-Fine²
Kelli F. Johnson^{1,2}

¹Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East, Seattle, Washington 98112

²University of Washington, School of Aquatic and Fishery Sciences

DRAFT SAFE

Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by NOAA Fisheries. It does not represent and should not be construed to represent any agency determination or policy.

Please cite as:

Wetzel, C.R., Cronin-Fine, L., and Johnson, K.F. 2017. Status of Pacific ocean perch (*Sebastes alutus*) along the US west coast in 2017. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Status of Pacific ocean perch (*Sebastes alutus*) along the US west coast in 2017

Contents

Executive Summary	i
Stock	i
Landings	i
Data and Assessment	iii
Stock Biomass	iv
Recruitment	vii
Exploitation Status	ix
Ecosystem Considerations	xii
Reference Points	xii
Management Performance	xiii
Unresolved Problems and Major Uncertainties	xiv
Decision Table	xv
Research and Data Needs	xviii
 1 Introduction	 1
1.1 Distribution and Stock Structure	1
1.2 Historical and Current Fishery	1
1.3 Summary of Management History and Performance	2
1.4 Fisheries off Canada and Alaska	2
 2 Data	 3
2.1 Fishery-Independent Data	3
2.1.1 Northwest Fisheries Science Center (NWFSC) Shelf-Slope Survey . .	4
2.1.2 Northwest Fisheries Science Center (NWFSC) Slope Survey	5
2.1.3 Alaska Fisheries Science Center (AFSC) Slope Survey	5
2.1.4 Pacific Ocean Perch Survey	6
2.1.5 Fishery Independent Data Not Included in the Base Model	6

2.1.5.1	Triennial Shelf Survey	6
2.1.5.2	Washington Research Lengths	7
2.2	Fishery-Dependent Data	8
2.2.1	Commercial Fishery Landings	8
2.2.2	Discards	9
2.2.3	Fishery Length and Age Data	10
2.2.4	Fishery Data Not Included in the Base Model	10
2.2.4.1	Historical Commercial Catch-Per-Unit Effort	10
2.2.4.2	Oregon Special Projects Length and Age Data	11
2.3	Biological Data	11
2.3.1	Natural Mortality	11
2.3.2	Sex Ratio, Maturation, and Fecundity	12
2.3.3	Length-Weight Relationship	12
2.3.4	Growth (Length-at-Age)	12
2.3.5	Ageing Precision and Bias	13
2.4	History of Modeling Approaches Used for This Stock	13
2.4.1	Previous Assessments	13
3	Assessment	14
3.1	General Model Specifications and Assumptions	14
3.1.1	Changes Between the 2011 Assessment Model and Current Model . .	14
3.1.2	Summary of Fleets and Areas	15
3.1.3	Other Specifications	16
3.1.4	Modeling Software and Model Bridging	17
3.1.5	Priors	17
3.1.6	Data Weighting	18
3.1.7	Estimated and Fixed Parameters	18
3.2	Model Selection and Evaluation	19
3.2.1	Key Assumptions and Structural Choices	19
3.2.2	Alternate Models Considered	19
3.2.3	Convergence	20
3.3	STAR Panel Review and Recommendations	21

3.4	Response to the 2011 STAR Panel Recommendations	21
3.5	Response to the 2017 STAR Panel Requests	22
3.6	Base Model Results	23
3.6.1	Parameter Estimates	23
3.6.2	Fits to the Data	24
3.6.3	Population Trajectory	26
3.6.4	Uncertainty and Sensitivity Analyses	27
3.6.5	Retrospective Analysis	29
3.6.6	Likelihood Profiles	29
3.6.7	Reference Points	29
4	Harvest Projections and Decision Tables	30
5	Regional Management Considerations	30
6	Research Needs	31
7	Acknowledgments	31
8	Tables	33
9	Figures	68
10	Appendix A. Detailed Fit to Length Composition Data	154
11	Appendix B. Detailed Fit to Age Composition Data	165
12	Appendix C. Description of CPUE and Triennial Data	172
13	Appendix D. List of Auxiliary Files Available	178
14	References	

Executive Summary

Stock

This assessment reports the status of the Pacific ocean perch rockfish (*Sebastes alutus*) off the US west coast from Northern California to the Canadian border using data through 2016. Pacific ocean perch are most abundant in the Gulf of Alaska and have been observed off of Japan, in the Bering Sea, and south to Baja California, though they are sparse south of Oregon and rare in southern California. Although neither catches nor other data from north of the US-Canada border were included in this assessment, the connectivity of these populations and the contribution to the biomass possibly through adult migration and/or larval dispersion is not certain. To date, no significant genetic differences have been found in the range covered by this assessment.

Landings

Harvest of Pacific ocean perch first exceeded 1 mt off the US west coast in 1918. Catches ramped up in the 1940s with large removals in Washington waters. During the 1950s the removals primary occurred in Oregon waters with catches from Washington declining following the 1940s. The largest removals, occurring between 1966-1968, were largely a result of harvest by foreign vessels. The fishery proceeded with more moderate removals ranging between 1165 to 2619 metric tons (mt) per year between 1969 and 1980. Removals generally declined from 1981 to 1994 to between 1031 and 1617 mt per year. Pacific ocean perch was declared overfished in 1999, resulting in large reductions in harvest in years since the declaration. Since 2000, annual landings of Pacific ocean perch have ranged between 54-270 mt, with landings in 2016 totaling 68 mt.

Pacific ocean perch are a desirable market species and discarding has historically been low. However, management restrictions (e.g. trip limits) resulted in increased discarding starting in the early 1990s. During the 2000s discarding increased for Pacific ocean perch due to harvest restrictions imposed to allow rebuilding, with estimated discard rates from the fishery peaking in 2009 and 2010 to approximately 50%, prior to implementation of catch shares in 2011. Since 2011, discarding of Pacific ocean perch has been estimated to be less than 3.5%.

Table a: Landings (mt) for the past 10 years for Pacific ocean perch by source.

Year	California	Oregon	Washington	At-sea hake	Survey	Total Landings
2007	0.15	83.65	45.12	4.05	0.58	133.55
2008	0.39	58.64	16.61	15.93	0.80	92.36
2009	0.92	58.74	33.22	1.56	2.72	97.17
2010	0.14	58.00	22.29	16.87	1.68	98.98
2011	0.12	30.26	19.66	9.17	1.94	61.14
2012	0.18	30.41	21.79	4.52	1.62	58.51
2013	0.08	34.86	14.83	5.41	1.71	56.89
2014	0.18	33.91	15.82	3.92	0.57	54.40
2015	0.12	38.05	11.41	8.71	1.59	59.88
2016	0.23	40.81	13.12	10.30	3.10	67.56




Figure a: Landings of Pacific ocean perch for California, Oregon, Washington, the foreign fishery (1966-1976), at-sea hake fishery, and fishery-independent surveys.

Data and Assessment

This a new full assessment for Pacific ocean perch, which was last assessed in 2011. In this assessment, aspects of the model including landings, data, and modelling assumptions were re-evaluated. The assessment was conducted using the length- and age-structured modeling software Stock Synthesis (version 3.30.03.05). The coastwide population was modeled allowing separate growth and mortality parameters for each sex (a two-sex model) from 1918 to 2017 and forecasted beyond 2017.

All of the data sources included in the base model for Pacific ocean perch have been re-evaluated for 2017. Changes of varying degrees have occurred in the data from those used in previous assessments. The landings history has been updated and extended back to 1918. Harvest was negligible prior to that year. Survey data from the Alaska and Northwest Fisheries Science Centers have been used to construct indices of abundance analyzed using a spatio-temporal delta-model. Length, marginal age, or conditional age-at-length compositions were also created for each fishery-dependent and -independent data source.

The definition of fishing fleets have changed from those in the 2011 assessment. Three fishing fleets were specified within the model: 1) a combined bottom trawl, mid-water trawl, and fixed gear fleet, where only a small fraction of Pacific ocean perch were captured by fixed gear (termed the fishery fleet), 2) the historical foreign fleet, and 3) the at-sea hake fishery. The fleet grouping were based on discarding practices. The fishery fleet estimated a retention curve based on discarding data and known management restrictions. However, very little if any discarding is assumed to have occurred by the foreign fleet and the catch reported by the at-sea hake fishery accounts for both discarded and landed fish and hence, no additional mortality was estimated for each of these fleets.

The assessment uses landings data and discard-fraction estimates; catch-per-unit-effort and survey indices; length- or age-composition data for each year and fishery or survey (with conditional age-at-length compositional data for the NWFSC shelf-slope survey); information on weight-at-length, maturity-at-length, and fecundity-at-length; information on natural mortality and the steepness of the Beverton-Holt stock-recruitment relationship; and estimates of ageing error. Recruitment at “equilibrium spawning output”, length-based selectivity of the fisheries and surveys, retention of the fishery, catchability of the surveys, growth, the time-series of spawning output, age and size structure, and current and projected future stock status are outputs of the model. Natural mortality (0.054 yr^{-1}) and steepness (0.72) were fixed in the final model. This was done due to relatively flat likelihood surfaces, such that fixing parameters and then varying them in sensitivity analyses was deemed the best way to characterize uncertainty.

Although this assessment using many types of data since the 1980s, there is little information about steepness and natural mortality. Estimates of steepness are uncertain partly because of highly variable recruitment. Uncertainty in natural mortality is common in many fish stock assessments even when length and age data are available.

A number of sources of uncertainty are explicitly included in this assessment. This assessment includes gender differences in growth, a non-linear relationship between individual spawner biomass and effective spawning output, and an updated relationship between length and maturity, based upon non-published information (Melissa Head, personal communication, NOAA, NWFSC). As is always the case, overall uncertainty is greater than that predicted by a single model specification. Among other sources of uncertainty that are not included in the current model are the degree of connectivity between the stocks of Pacific ocean perch off of Vancouver Island, British Columbia and those in US waters, and the effect of climatic variables on recruitment, growth and survival.

A base model was selected that best captures the central tendency for those sources of uncertainty considered in the model.

Stock Biomass

The predicted spawning output from the base model generally showed a slight decline prior to 1966 when fishing by the foreign fleet commenced. A short, but sharp decline occurred between 1966 and 1970, followed by a period of the spawning output stabilizing or with a minimal decline until the late 1990s. The stock showed increases in stock size following the year 2000 due to a combination of strong recruitment and low catches. The 2017 estimated spawning output relative to unfished equilibrium spawning output is above the target of 40% of unfished spawning output at 96.4% ($\sim 95\%$ asymptotic interval: $\pm 77.1\%$ -116%). Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in the estimated spawning output is high.

Table b: Recent trend in estimated spawning output (million eggs) and estimated relative spawning output (depletion).

Year	Spawning Output (million eggs)	$\sim 95\%$ confidence interval	Estimated depletion	$\sim 95\%$ confidence interval
2008	4955.00	2542 - 7368	0.69	0.538 - 0.833
2009	5143.00	2647 - 7639	0.71	0.559 - 0.864
2010	5267.00	2716 - 7818	0.73	0.574 - 0.884
2011	5345.00	2761 - 7930	0.74	0.583 - 0.896
2012	5394.00	2793 - 7996	0.75	0.590 - 0.903
2013	5427.00	2818 - 8037	0.75	0.594 - 0.907
2014	5566.00	2904 - 8228	0.77	0.612 - 0.928
2015	5979.00	3144 - 8814	0.83	0.661 - 0.994
2016	6515.00	3449 - 9582	0.90	0.722 - 1.081
2017	6966.00	3702 - 10230	0.96	0.771 - 1.156




Figure b: Time-series of spawning output trajectory (circles and line: median; light broken lines: 95% credibility intervals) for the base assessment model.




Figure c: Estimated relative spawning output (depletion) with approximate 95% asymptotic confidence intervals (dashed lines) for the base assessment model.

Recruitment

Recruitment deviations were estimated for the entire assessment period. There is little information regarding recruitment prior to 1965, and the uncertainty in these estimates is expressed in the model. Past assessments estimated large recruitments in 1999 and 2000. In recent years, a recruitment of unprecedented size is estimated to have occurred in 2008. Additionally, there is early evidence of a strong recruitment in 2013. The four lowest recruitments estimated within the model (in ascending order) occurred in 2012, 2003, 2005, and 2007.

Table c: Recent estimated trend in recruitment and estimated recruitment deviations determined from the base model

Year	Estimated Recruitment	~ 95% confidence interval	Estimated Recruitment Devs.	~ 95% confidence interval
2008	150412.00	90890 - 248913	2.69	2.385 - 2.997
2009	5928.00	2614 - 13441	-0.55	-1.311 - 0.216
2010	9600.00	4808 - 19171	-0.07	-0.666 - 0.529
2011	18888.00	9954 - 35841	0.61	0.076 - 1.137
2012	2661.00	1127 - 6280	-1.35	-2.178 - -0.532
2013	37397.00	18056 - 77455	1.20	0.549 - 1.854
2014	5672.00	2000 - 16087	-0.77	-1.836 - 0.289
2015	12354.00	3501 - 43585	-0.00	-1.372 - 1.365
2016	12515.00	3540 - 44248	0.00	-1.372 - 1.372
2017	12602.00	3565 - 44546	0.00	-1.372 - 1.372




Figure d: Time-series of estimated Pacific ocean perch recruitments for the base model with 95% confidence or credibility intervals.

Exploitation Status

The spawning output of Pacific ocean perch reached a low in 1975. Landings for Pacific ocean perch decreased significantly in 2000 compared to previous years. The estimated relative depletion was possibly below the target biomass level between the 1970s and 1990s, but has likely remained above the target otherwise, and currently is significantly greater than the 40% unfished spawning output target. Throughout the late 1960s and the early 1970s the exploitation rate and values of relative spawning potential $((1-SPR)/(1-SPR_{50\%}))$ were mostly above target levels. Recent exploitation rates on Pacific ocean perch were predicted to be significantly below target levels.

Table d: Recent trend in spawning potential ratio $(1-SPR)/(1-SPR_{50})$ and summary exploitation rate for Pacific ocean perch.

Year	$(1-SPR)/$ $(1-SPR_{50\%})$	~ 95% confidence interval	Exploitation rate	~ 95% confidence interval
2007	0.066	0.034 - 0.097	0.001	0.001 - 0.002
2008	0.054	0.027 - 0.082	0.001	0.001 - 0.002
2009	0.073	0.035 - 0.111	0.002	0.001 - 0.003
2010	0.069	0.034 - 0.105	0.002	0.001 - 0.002
2011	0.024	0.013 - 0.036	0.000	0.000 - 0.001
2012	0.023	0.012 - 0.034	0.000	0.000 - 0.001
2013	0.022	0.012 - 0.032	0.000	0.000 - 0.001
2014	0.019	0.010 - 0.029	0.000	0.000 - 0.001
2015	0.020	0.011 - 0.029	0.000	0.000 - 0.001
2016	0.020	0.011 - 0.030	0.000	0.000 - 0.001




Figure e: Estimated relative spawning potential ratio $(1-SPR)/(1-SPR_{50\%})$ for the base model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the SPR50% harvest rate. The last year in the time-series is 2016.




Figure f: Phase plot of estimated $(1-SPR)/(1-SPR_{50\%})$ vs. relative spawning output (depletion) for the base case model.

Ecosystem Considerations

Rockfish are an important component of the California Current ecosystem along the US west coast, with more than sixty five species filling various niches in both soft and hard bottom habitats from the nearshore to the continental slope, as well as near bottom and pelagic zones. Pacific ocean perch are generally considered to be semi-demersal, but there can, at times, be a significant pelagic component to their distribution.

Recruitment is one mechanism by which the ecosystem may directly impact the population dynamics of Pacific ocean perch. The 1999 cohort for many species of rockfish was large – sometimes significantly so. Long-term averages suggest that environmental conditions may influence the spawning success and survival of larvae and juvenile rockfish. Pacific ocean perch showed above average recruitment deviations in 1999 and 2000. The specific pathways through which environmental conditions exert influence on Pacific ocean perch dynamics are unclear; however, changes in water temperature and currents, distribution of prey and predators, and the amount and timing of upwelling are all possible linkages. Changes in the environment may also result in changes in length-at-maturity, fecundity, growth, and survival which can affect the status of the stock and its susceptibility to fishing. Unfortunately, there are few data available for Pacific ocean perch that provide insights into these effects.

Fishing has effects on both the age-structure of a population, as well as the habitat with which the target species is associated. Fishing often targets larger, older fish and years of fishing mortality results in a truncated age-structure when compared to unfished conditions. Rockfish are often associated with habitats containing living structure such as sponges and corals, and fishing may alter that habitat to a less desirable state. This assessment provides a look at the effects of fishing on age structure, and recent studies on essential fish habitat are beginning to characterize important locations for rockfish throughout their life history; however, there is little current information available to evaluate the specific effects of fishing on the ecosystem issues specific to Pacific ocean perch.

Reference Points

This stock assessment estimates that the spawning output of Pacific ocean perch is above the management target. Due to reduced landing and the large 2008 year-class, an increasing trend in spawning output was estimated in the base model. The estimated depletion in 2017 is 96.4% ($\sim 95\%$ asymptotic interval: $\pm 77.1\%$ -116%), corresponding to an unfished spawning output of 6,966 million eggs ($\sim 95\%$ asymptotic interval: 3,702-10,230 million eggs). Unfished age 3+ biomass was estimated to be 154,084 mt in the base model. The target spawning output based on the biomass target ($SB_{40\%}$) is 2,891.2 million eggs, with an equilibrium catch of 2,680.8 mt. Equilibrium yield at the proxy F_{MSY} harvest rate corresponding to $SPR_{50\%}$ is 2,555.8 mt. Estimated MSY catch is at a 2,860.7 spawning output of 1,942.8 million eggs (26.9% depletion)

Table e: Summary of reference points and management quantities for the base case.

Quantity	Estimate	~95% Confidence Interval
Unfished spawning output (million eggs)	7228	5126.5 - 9329.4
Unfished age 3+ biomass (mt)	154084	109381.3 - 198786.7
Unfished recruitment (R0, thousands)	12647.9	9483.8 - 16867.7
Spawning output(2017 million eggs)	6966	3701.9 - 10230.1
Relative spawning output (depletion) (2017)	0.964	0.771 - 1.2
Reference points based on $SB_{40\%}$		
Proxy spawning output ($B_{40\%}$)	2891.2	2050.6 - 3731.8
SPR resulting in $B_{40\%}$ ($SPR_{B40\%}$)	0.458	0.458 - 0.458
Exploitation rate resulting in $B_{40\%}$	0.038	0.038 - 0.039
Yield with $SPR_{B40\%}$ at $B_{40\%}$ (mt)	2680.8	1906.1 - 3455.6
Reference points based on SPR proxy for MSY		
Spawning output	3224.8	2287.2 - 4162.4
SPR_{proxy}	0.5	
Exploitation rate corresponding to SPR_{proxy}	0.033	0.033 - 0.034
Yield with SPR_{proxy} at SB_{SPR} (mt)	2555.8	1816.8 - 3294.7
Reference points based on estimated MSY values		
Spawning output at MSY (SB_{MSY})	1942.8	1374.4 - 2511.1
SPR_{MSY}	0.34	0.338 - 0.342
Exploitation rate at MSY	0.057	0.056 - 0.058
MSY (mt)	2860.7	2034.9 - 3686.5

Management Performance

Exploitation rates on Pacific ocean perch exceeded MSY proxy target harvest rates during the 1960s and 1970s, resulting in sharp declines in the spawning output. Exploitation rates subsequently declined to rates at or below the management target in the late 1970s. Management restrictions imposed in the 1990s further reduced exploitation rates. An overfished declaration for Pacific ocean perch resulted in very low exploitation rates since 2001 with Annual Catch Limits (ACLs) being set far below the Overfishing Limit (OFL) and Acceptable Biological Catch (ABC) values.

Table f: Recent trend in total catch and landings (mt) relative to the management guidelines. Estimated total catch reflect the landings plus the model estimated discarded biomass.

Year	OFL (mt; ABC prior to 2011)	ABC (mt)	ACL (mt; OY prior to 2011)	Total landings (mt)	Estimated total catch (mt)
2007	900		150	134	159
2008	911		150	92	134
2009	1,160		189	97	193
2010	1,173		200	99	182
2011	1,026	981	180	61	62
2012	1,007	962	183	59	60
2013	844	807	150	57	58
2014	838	801	153	54	56
2015	842	805	158	60	61
2016	850	813	164	68	68

Unresolved Problems and Major Uncertainties

1. Pacific ocean perch off the US west coast may be a fraction of a much large population extending into Canada or even Alaska. Modelling only a part of the total population might contribute to the lack of correspondence between the survey indices and other data sources, as seen in the $\ln(R0)$ profiles and age-structured production model diagnostics as well as some of the observation variability. While this comment is not intended to reflect badly on the STAT's capabilities, it is important to recognize that stock structure could potentially be a major source of uncertainty regarding the assessment results.
2. The indices of abundance used in the final base model provide almost no information on population scale, as demonstrated in the $\ln(R0)$ profiles examined during the review. The Triennial survey was the only index that provided signal with respect to population scale. However, this survey was removed in the final base model due to concerns about the quality of the survey and conflicts with other data. There are large amounts of composition data in the model, with both age- and length-compositions being included for some fleets. The compositional data and catch are providing the majority of the information on the estimated and derived quantities.
3. Use of conditional-age-at-length composition data provides information on parameters beyond those of the length-at-age relationship. The conditional-age-at-length data are robust to length-based processes (Piner et al. 2016), however they are also influenced by age-based processes (Lee et al. 2017). No age-based observation model processes were used in the assessment model as a link to the data, meaning that the conditional-age-at-length data were assumed to be unbiased with respect to the population. The conditional-age-at-length data were shown to be very influential on the estimated dynamics beyond growth estimates. More theoretical work in this area is needed to understand how to best the use this type of information and what potential systems or observation model processes could invalidate the assumption of randomness at length.

Decision Table

Model uncertainty has been described by the estimated uncertainty within the base model and by the sensitivities to different model structure. The estimated parameter that resulted in the most variability of predicted status and yield advice was natural mortality (M). The 12.5% and 87.5% quantiles based on spawning output uncertainty were used to determine the low and high values for M of 0.045 and 0.06 yr⁻¹. An alternative decision table was also provided which based the range of natural mortality values on the uncertainty in the parameter prior. This approach was suggestion post-STAR panel review by Owen Hamel as a method to explore a greater range of uncertainty. The low and high stated of nature values were based on the 12.5% and 87.5% quantiles when the uncertainty was divided between the data used to generate the prior and the uncertainty surrounding the estimated value.

This assessment synthesizes many sources of data and estimates recruitment variability, thus it is classified as a Category 1 stock assessment. Therefore, the sigma for P* to determine the catch reduction to account for scientific uncertainty is 0.36, since the estimated sigma in the assessment is less than this for current spawning biomass (0.24).

Table g: Projections of potential OFL (mt) and ACL (mt) and the estimated spawning output and relative depletion based on ACL removals. The ACL values for 2017 and 2018 are set at the harvest limits currently set by management.

Year	OFL	ACL	Spawning Output (million eggs)	Relative Depletion
2017	5861	281	6966	0.964
2018	6116	281	7299	1.010
2019	6251	5981	7559	1.046
2020	6091	5827	7539	1.043
2021	5894	5639	7485	1.036
2022	5685	5439	7382	1.021
2023	5475	5238	7246	1.002
2024	5270	5042	7089	0.981
2025	5077	4857	6921	0.958
2026	4899	4688	6748	0.934
2027	4738	4533	6572	0.909
2028	4590	4391	6398	0.885

Table h: Summary of 10-year projections beginning in 2019 for alternate states of nature based on an axis of uncertainty for the base model. The range of natural mortality values corresponded to the 12.5 and 87.5th quantile from the uncertainty around final spawning biomass. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. The SPR50 catch stream is based on the equilibrium yield applying the SPR50 harvest rate.

		States of nature					
		M = 0.045		M = 0.054		M = 0.060	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
ABC	2019	5981	5533	86	7559	105	9565
	2020	5827	5486	86	7541	104	9564
	2021	5639	5414	84	7488	104	9517
	2022	5439	5307	83	7388	102	9411
	2023	5238	5177	81	7253	100	9263
	2024	5042	5032	79	7098	98	9089
	2025	4857	4881	76	6932	96	8902
	2026	4688	4726	74	6760	94	8706
	2027	4533	4572	71	6586	91	8508
	2028	4391	4419	69	6413	89	8310
SPR50	2019	2556	5533	86	7559	105	9565
	2020	2556	5640	88	7694	106	9716
	2021	2556	5721	89	7792	108	9819
	2022	2556	5762	90	7835	108	9854
	2023	2556	5770	90	7834	108	9837
	2024	2556	5755	90	7802	108	9782
	2025	2556	5721	89	7747	107	9701
	2026	2556	5673	88	7673	106	9599
	2027	2556	5613	88	7585	105	9482
	2028	2556	5545	87	7487	104	9354

Table i: Alternative decision table. Summary of 10-year projections beginning in 2019 for alternate states of nature based on an axis of uncertainty for the base model. The range of natural mortality values are based on the 12.5 and 87.5th quantiles of the natural mortality prior. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. The SPR50 catch stream is based on the equilibrium yield applying the SPR50 harvest rate.

		States of nature					
		M = 0.038		M = 0.054		M = 0.077	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
ABC	2019	5981	4214	69	7559	105	41065
	2020	5827	4141	67	7541	104	41501
	2021	5639	4047	66	7488	104	41736
	2022	5439	3927	64	7388	102	41699
	2023	5238	3789	62	7253	100	41471
	2024	5042	3643	59	7098	98	41125
	2025	4857	3493	57	6932	96	40699
	2026	4688	3342	54	6760	94	40219
	2027	4533	3192	52	6586	91	39706
	2028	4391	3045	50	6413	89	39174
SPR50	2019	2556	4214	69	7559	105	41065
	2020	2556	4295	70	7694	106	41651
	2021	2556	4356	71	7792	108	42032
	2022	2556	4387	71	7835	108	42130
	2023	2556	4393	72	7834	108	42024
	2024	2556	4381	71	7802	108	41786
	2025	2556	4354	71	7747	107	41455
	2026	2556	4316	70	7673	106	41057
	2027	2556	4270	70	7585	105	40611
	2028	2556	4216	69	7487	104	40136

Research and Data Needs

There are many areas of research that could be improved to benefit the understanding and assessment of Pacific ocean perch. Below, are issues that are considered of importance.

1. **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for Pacific ocean perch. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life history of Pacific ocean perch may reduce that uncertainty.
2. **Steepness:** The amount of stock resilience, steepness, dictates the rate at which a stock can rebuild from low stock sizes. Improved understating regarding the steepness parameter for US west coast Pacific ocean perch will reduce our uncertainty regarding current stock status.
3. **Basin-wide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for Pacific ocean perch off of the west coast of the US and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and Alaska to determine if there are similarities with the US west coast observations would help to define the connectivity between Pacific ocean perch north and south of the US-Canada border.

Table j: Base model results summary.

Quantity	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
OFL (mt)	911	1,160	1,173	1,026	1,007	844	838	842	850	964
ACL (mt)	150	189	200	180	183	150	153	158	164	281
Landings (mt)	92	97	99	61	59	57	54	60	68	68
Total Est. Catch (mt)	134	193	182	62	60	58	56	61	68	68
(1-SPR)(1-SPR _{50%})	0.054	0.073	0.069	0.024	0.023	0.022	0.019	0.020	0.020	
Exploitation rate	0.001	0.002	0.002	0.000	0.000	0.000	0.000	0.000	0.000	
Age 3+ biomass (mt)	113935	114656	114685	129549	136687	143823	151230	156447	163853	168288
Spawning Output	4955	5143	5267	5345	5394	5427	5566	5979	6515	6966
95% CI	2542 - 7368	2647 - 7639	2716 - 7818	2761 - 7930	2793 - 7996	2818 - 8037	2904 - 8228	3144 - 8814	3449 - 9582	3702 - 10230
Relative Depletion	0.686	0.712	0.729	0.740	0.746	0.751	0.770	0.827	0.901	0.964
95% CI	0.538 - 0.833	0.559 - 0.864	0.574 - 0.884	0.583 - 0.896	0.590 - 0.903	0.594 - 0.907	0.612 - 0.928	0.661 - 0.994	0.722 - 1.081	0.771 - 1.156
Recruits	150412	5928	9600	18888	2661	37397	5672	12354	12515	12602
95% CI	90890 - 248913	2614 - 13441	4808 - 19171	9954 - 35841	1127 - 6280	18056 - 77455	2000 - 16087	3501 - 43585	3540 - 44248	3565 - 44546




Figure g: Equilibrium yield curve for the base case model. Values are based on the 2016 fishery selectivity and with steepness fixed at 0.72.

1 Introduction

1.1 Distribution and Stock Structure

Pacific ocean perch (*Sebastes alutus*) are most abundant in the Gulf of Alaska and have been observed off of Japan, in the Bering Sea, and south to Baja California, although they are sparse south of Oregon and rare in southern California. While genetic studies have found three populations of Pacific ocean perch off of British Columbia related to unique geography and oceanic conditions (Seeb and Gunderson 1988, Withler et al. 2001) with, notably, a separate stock off of Vancouver Island, no significant genetic differences have been found in the range covered by this assessment. Pacific ocean perch show dimorphic growth, with females reaching a slightly larger size than males. Males and females are equally abundant on rearing grounds at age 1.5.

The Pacific ocean perch population has been modeled as a single stock off of the US west coast (essentially northern California to the Canadian border, since Pacific ocean perch are seen extremely rarely in central and southern California). Good recruitments show up in size-composition data throughout all portions of this area, which supports the single stock hypothesis. This assessment includes landings and catch data for Pacific ocean perch from the states of Washington, Oregon and California, along with records from foreign fisheries, the at-sea hake fleet, and fishery-independent surveys.

1.2 Historical and Current Fishery

Prior to 1966, the Pacific ocean perch resource off of the northern portion of the US west coast was harvested almost entirely by Canadian and US vessels. Harvest was negligible prior to 1940, reached 1,367 mt in 1950, 3,243 mt in 1961 and 7,636 mt in 1965. Catches increased dramatically after 1965, with the introduction of large distant-water fishing fleets from the Soviet Union and Japan. Both nations employed large factory stern trawlers as their primary method for harvesting Pacific ocean perch. Peak removals are estimated at 18,883 mt in 1966 and 14,591 mt in 1967. These numbers are based upon a re-analysis of the foreign catch data (Rogers 2003), which focused on deriving a more realistic species composition for catches previously identified only as Pacific ocean perch. Catches declined rapidly following these peak years, and Pacific ocean perch stocks were considered to be severely depleted throughout the Oregon-Vancouver Island region by 1969 (Gunderson 1977, Gunderson et al. 1977). Landed harvest averaged 1,381 mt over the period 1977-94. Landings have continued to decline since 1994, primarily due to more restrictive management (Table 1 and Figure 1).

1.3 Summary of Management History and Performance

Prior to 1977, Pacific ocean perch in the northeast Pacific were managed by the Canadian Government in its waters and by the individual states in waters off of the US. With the implementation of the Magnuson Fishery Conservation and Management Act (MFCMA) in 1977, US territorial waters were extended to 200 nautical miles from shore and primary responsibility for management of the groundfish stocks off Washington, Oregon, and California shifted from the states to the Pacific Fishery Management Council (PFMC) and the National Marine Fisheries Service (NMFS). At that time, however, a Fishery Management Plan for the West Coast groundfish stocks had not yet been approved. In the interim, the state agencies worked with the PFMC to address conservation issues. In 1981, the PFMC adopted a management strategy to rebuild the depleted Pacific ocean perch stocks to levels that would produce Maximum Sustainable Yield (MSY) within 20 years. On the basis of cohort analysis (Gunderson 1978), the PFMC set Acceptable Biological Catch (ABC) levels at 600 mt for the US portion of the Vancouver INPFC area and 950 mt for the Columbia International North Pacific Fishery Commission (INPFC) area. To implement this strategy, the states of Oregon and Washington each established landing limits for Pacific ocean perch. Trawl trip limits of various forms remained in effect through 2016 (Table 2).

The landings of Pacific ocean perch have been historically governed by harvest guidelines and trip limits, while recently management has imposed total catch harvest limits in the form of overfishing limits (OFLs), acceptable biological catches (ABCs), and annual catch limits (ACLs). A trawl rationalization program, consisting of an individual fishing quota (IFQ) catch shares system was implemented in 2011 for the limited entry trawl fleet targeting non-whiting groundfish, including Pacific ocean perch and the trawl fleet targeting and delivering whiting to shore-based processors. The limited entry at-sea trawl sectors (motherships and catch-processors) that target whiting and process at-sea are managed in a system of harvest cooperatives.

Limits on Pacific ocean perch were first established in 1983 (Table 2). These were implemented as area closures, trip limits, and cumulative landing limits. In 1999, Pacific ocean perch was declared overfished with the assessment estimating the spawning output below the management limit (25% of virgin biomass or output). In reaction to the overfished declaration, harvest limits were reduced relative to previous years and a rebuilding plan was implemented in 2001 with recent ACLs being set well below the estimated OFLs (Table 3).

1.4 Fisheries off Canada and Alaska

Pacific ocean perch can be found in waters off the US west coast and northward through Alaskan waters. In contrast to the Pacific ocean perch stock off the US west coast, each assessed portion of the stock in Canadian and Alaskan waters have historically been estimated to be above management targets. The subset of the stock off the US west coast represents the tail of the species distribution with little to no Pacific ocean perch being encountered south

of northern California. The most recent updated assessments for the Bering Sea and the Gulf of Alaska stocks determined that neither stock is in an overfished state and recommended acceptable biological catches of 43,723 mt and 23,918 mt, respectively, for 2017.

In Canadian waters Pacific ocean perch has the largest single-species quota, accounting for approximately 25% of all rockfish landings by weight in the bottom trawl fleet. The Canadian Pacific ocean perch stock is broken into three separate areas that are individually assessed. The status of the stock within each area is above Canadian management targets.

2 Data

Data used in the Pacific ocean perch assessment are summarized in Figure 2. A description of each data source is provided below.

2.1 Fishery-Independent Data

Research surveys have been used to provide fishery-independent information about the abundance, distribution, and biological characteristics of Pacific ocean perch. A coast-wide survey was conducted in 1977 (Gunderson and Sample 1980) and repeated every three years through 2004 (referred to as the ‘Triennial shelf survey’). The NMFS coordinated a cooperative research survey of the Pacific ocean perch stocks off Washington and Oregon with the Washington Department of Fish and Wildlife (WDFW) and the Oregon Department of Fish and Wildlife (ODFW) in March-May 1979 (Wilkins and Golden 1983). This survey was repeated in 1985 (referred to as the Pacific ocean perch survey). Two slope surveys have been conducted off the West Coast in recent years, one using the research vessel Miller Freeman, which ended in 2001 (referred to as the ‘AFSC slope survey’), and another ongoing cooperative survey using commercial fishing vessels which began in 1998 as a DTS (Dover sole, thornyhead, and sablefish) survey and was expanded to other groundfish in 1999 (referred to as the ‘NWFSC slope survey’). In 2003, this survey was expanded spatially to include the shelf. This last survey, conducted by the NWFSC, continues to cover depths from 30-700 fathoms (55-1280 meters) on an annual basis (referred to as the ‘NWFSC shelf-slope survey’).

Age estimates for Pacific ocean perch prior to the 1980s were made via surface ageing of otoliths, which misses the very tight annuli at the edge of the otolith once the fish reaches near maximum size. Ages are highly biased by age 14, and maximum age was estimated to be in the 20s, which lead to an overestimate of the natural mortality rate and the productivity of the stock. Using break and burn methods, Pacific ocean perch have been aged to over 100 years. Otoliths from fishery-independent and -dependent sources that were only surface age reads were excluded from this assessment due to the bias associated with these age reads.

2.1.1 Northwest Fisheries Science Center (NWFSC) Shelf-Slope Survey

The NWFSC shelf-slope survey is based on a random-grid design; covering the coastal waters from a depth of 55 m to 1,280 m (Bradburn et al. 2011). This design uses four chartered industry vessels in most years, assigned to a roughly equal number of randomly selected grid cells. The survey, which has been conducted from late-May to early-October each year, is divided into two 2-vessel passes off the coast, which are executed from north to south. This design therefore incorporates both vessel-to-vessel differences in catchability as well as variance associated with selecting a relatively small number (approximately 700) of cells from a very large population of possible cells (greater than 11,000) distributed from the Mexican to the Canadian border.

The data from the NWFSC shelf-slope survey was analyzed using a spatio-temporal delta-model (Thorson et al. 2015), implemented as an R package, VAST (Thorson and Barnett 2017), which is publicly available online (<https://github.com/James-Thorson/VAST>). Spatial and spatio-temporal variation is specifically included in both encounter probability and positive catch rates, a logit-link for encounter probability and a log-link for positive catch rates. Vessel-year effects were included for each unique combination of vessel and year in the data to account for the random selection of commercial vessels used during sampling (Helser et al. 2004, Thorson and Ward 2014). Spatial variation was approximated using 1,000 knots, and the model used the bias-correction algorithm (Thorson and Kristensen 2016) in Template Model Builder (Kristensen et al. 2016). Further details regarding model structure are available in the user manual (https://github.com/James-Thorson/VAST/blob/master/examples/VAST_user_manual.pdf). The stratification and modeling configuration are provided in Table 4.

The smallest Pacific ocean perch tend to occur in the shallower depths (< 200 m) with only larger individuals occurring at depths deeper than 300 m. Data collected by the NWFSC shelf-slope survey between depths of 55 - 549 m and north of 42° and south of 49° were used to generate an index of abundance from 2003-2016. The estimated index of abundance is shown in Table 5. For contrast, the design based values are shown in Table 6. The lognormal distribution with random strata-year and vessel effects had the lowest AIC and was chosen as the final model. The Q-Q plot does not show any departures from the assumed distribution (Figure 4). The indices for the NWFSC shelf-slope survey show a tentative decline in the population between 2003 and 2009, with an increasing trend in biomass between the 2009 and 2016 median point estimates.

Length compositions were expanded based upon the stratification and the age data was used as conditional age-at-length data. The number of tows with length data ranged from 33 in 2006 to 69 in 2015 (Table 7), where ages were collected for Pacific ocean perch in nearly every tow (Table 8). The expanded length frequencies from this survey show an increase in small fish starting in 2010 (Figure 5). The age frequencies provide clear evidence of large year-classes moving through the population from the 1999, 2000, and 2008 recruitments; with early indications of a large 2013 recruitment (Figure 6).

The input sample sizes for length and marginal age-composition data for all fishery-independent surveys were calculated according to Stewart and Hamel (2014), which determined that the approximate realized sample size for shelf/slope rockfish species was $2.43 * N_{\text{tow}}$. The effective sample size of conditional-age-at-length data was set at the number of fish at each length by sex and by year.

2.1.2 Northwest Fisheries Science Center (NWFSC) Slope Survey

The NWFSC slope survey covered waters throughout the summer from 183 m to 1,280 m north of $34^{\circ}30'$ S, which is near Point Conception, from 1999 and 2002. Tows conducted between the depths of 183 and 549 m were used to create an index of abundance using a bayesian delta-GLMM and the VAST delta-GLMM models. The estimated index of abundance is shown in Table 5. The stratification and modeling configuration are provided in Table 4. Based on the diagnostics of the bayesian delta-GLMM, which does not account for spatial effects, a gamma distribution allowing for additional probability of extreme catch events with year-vessel random effects was selected as the final model. The Q-Q plot does show a minimal departure from the assumed distribution (Figure 7), but was determined to be acceptable based on the alternative model distributions. The trend of abundance across the four surveys years was generally flat with high estimated annual variance. Sensitivities (not shown) were done evaluating the excluding of this index within the base model or using the VAST estimated index and neither approach was found to be influential on the model estimates.

Length and age compositions were available for 2001 and 2002 and were expanded based upon the survey stratification (Tables 9 and 10). The expanded length frequencies from this survey shows that primarily only large fish were captured both years (Figure 8). The majority of fish observed by this survey were aged at greater than 10 years (Figure 9).

The input sample sizes for length and marginal age-composition data were calculated according to Stewart and Hamel (2014) described in Section 2.1.1.

2.1.3 Alaska Fisheries Science Center (AFSC) Slope Survey

The AFSC slope survey operated during autumn (October-November) aboard the R/V Miller Freeman. Partial survey coverage of the US west coast occurred during 1988-96 and complete coverage (north of $34^{\circ}30'$ S) during 1997, 1999, 2000, and 2001. Only the four years of consistent and complete surveys plus 1996, which surveyed north of 43° N latitude to the US-Canada border, were used in this assessment. These same data years were used in the last assessment. The number of tows with length data ranged from 19 in 2000 to 48 in 1996 (Table 11). Because a large number of positive tows occurred in 1996, it was decided to include that year, which surveyed from 43° N latitude to the US-Canada border. Therefore, only tows from 43° N latitude to the US-Canada border were used.

An index of abundance was estimated based on the data using the VAST delta-GLMM model. The estimated index of abundance is shown in Table 5. The stratification and modeling configuration are provided in Table 4. The lognormal distribution with random strata-year had the lowest AIC and was chosen as the final model. The Q-Q plot does not show any departures from the assumed distribution (Figure 10). The trend in the indices was generally flat over time.

Length compositions were available for each year the survey was conducted. No age data were available from this survey. The expanded length frequencies from this survey were generally of larger fish (> 30 cm), for 1997 where the highest frequency of fish were between 20 and 30 cm for both females and males (Figure 11).

The input sample sizes for length and marginal age composition data were calculated according to Stewart and Hamel (2014) described in Section 2.1.1.

2.1.4 Pacific Ocean Perch Survey

A survey designed to sample Pacific ocean perch was conducted in 1979 and again in 1985 (for a detailed description see Ianelli et al. (1992)). An index of abundance was estimated based on the data using the VAST delta-GLMM model. The estimated index of abundance is shown in Table 5. The stratification and modeling configuration are provided in Table 4. The lognormal distribution with random strata-year had the lowest AIC and was chosen as the final model. The Q-Q plot does not show any departures from the assumed distribution (Figure 12). The index shows a clear decline in abundance between the two survey years.

Length and age compositions were expanded based on the survey stratification. The survey had 125 and 126 Pacific ocean perch tows (Table 12) and ages were only available in 1985 due to surface reads for the 1979 data (Table 13). The length frequencies for both years are highest between the 30-45 cm range (Figure 13) with ages in 1985 having a large number of fish age 40 and greater (Figure 14).

The input sample sizes for length and marginal age-composition data were calculated according to Stewart and Hamel (2014) described in Section 2.1.1.

2.1.5 Fishery Independent Data Not Included in the Base Model

The follow datasets were evaluated but not included in the base model.

2.1.5.1 Triennial Shelf Survey

The Triennial shelf survey was first conducted by the AFSC in 1977 and spanned the time-frame from 1977-2004. The survey's design and sampling methods are most recently described in Weinberg et al. (2002). Its basic design was a series of equally-spaced transects from which searches for tows in a specific depth range were initiated. The survey design has changed slightly over the period of time. In general, all of the surveys were conducted in the mid-summer through early fall: the 1977 survey was conducted from early July through late September; the surveys from 1980 through 1989 ran from mid-July to late September; the 1992 survey spanned from mid-July through early October; the 1995 survey was conducted from early June to late August; the 1998 survey ran from early June through early August; and the 2001 and 2004 surveys were conducted in May-July.

Haul depths ranged from 91-457 m during the 1977 survey with no hauls shallower than 91 m. The surveys in 1980, 1983, and 1986 covered the West Coast south to 36.8° N latitude and a depth range of 55-366 m. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5° N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55-500 m and surveyed south to 34.5° N. In the final year of the Triennial series, 2004, the NWFSC's Fishery Resource and Monitoring division (FRAM) conducted the survey and followed very similar protocols as the AFSC.

The Triennial shelf survey was not used in the final base model for a number of reasons. First, there were concerns regarding the varying sampling and targeting of specific species by year across the time-series. Secondly, the Triennial shelf survey targeted the shelf of the West Coast and would not be expected to sample well slope species such as Pacific ocean perch. There were limited observations of Pacific ocean perch relative to other surveys (e.g. NWFSC shelf-slope survey) and the length and age distributions varied in such a manner that would indicate either poor sampling of Pacific ocean perch or inconsistent sampling of the population.

Information regarding the Triennial shelf survey index of abundance and the number of samples available and plots of the composition data are available in Appendix 12.

2.1.5.2 Washington Research Lengths

Research length and ages were provided by WDFW. However, the information regarding the nature of the research cruise and collection methods have been lost to time. The data set includes lengths and ages that were collected between 1967-1972 and in 1979. The distribution of lengths across years collected were consistent with primarily only larger Pacific ocean perch, 35-40 cm, being selected. All age data were based upon surface reads which unfortunately are highly biased at relatively young ages for Pacific ocean perch. Due to the lack of information regarding the collection of these data, they were not selected to be apart of the base model but a sensitivity was conducted which evaluated the impact of these data.

2.2 Fishery-Dependent Data

2.2.1 Commercial Fishery Landings

Washington

Historical commercial fishery landings of Pacific ocean perch in Washington for the years 1908-2016 were obtained from Theresa Tsou (WDFW) and Phillip Weyland (WDFW). This assessment is the first Pacific ocean perch assessment to include a historical catch reconstruction provided by Washington state and, hence, the historical catches for Washington differ from those used in the 2011 assessment. WDFW also provided catches for 1981-2016 period to include re-distribution of the “URCK” landings in the PacFIN database. These data are currently not available from PacFIN.

Oregon

Historical commercial fishery landings of Pacific ocean perch in Oregon for the years 1892-1986 were obtained from Alison Whitman (ODFW). A description of the methods can be found in Karnowski et al. (2014). Recent landings (1987-2016) were obtained from PacFIN (retrieval dated May 2, 2017, Pacific States Marine Fisheries Commission, Portland, Oregon; www.psmfc.org). The catch data from the POP and POP2 categories contained within PacFIN for Pacific ocean perch were used for this assessment. Additional catches from 1987-1999 for Pacific ocean perch under the URCK category not yet available in PacFIN were received directly from the state and combined with the landings data available for that period within PacFIN (Patrick Mirrick, personal communication, ODFW).

California

Historical commercial fishery landings of Pacific ocean perch were obtained directly from John Field at the SWFSC due to database issues for the historical period for the California Cooperative Groundfish Survey data system, also known as CALCOM Database (128.114.3.187) for the years 1916-1980. The catches received included revisions in the catch history from 1948-1960 based on fish that were landed in northern California that were not included in the original reconstruction. A description of the historical reconstruction methods can be found in Ralston et al. (2010). Recent landings (1981-2016) were obtained from PacFIN (retrieval dated May 2, 2017, Pacific States Marine Fisheries Commission, Portland, Oregon; www.psmfc.org).

At-Sea Hake Fishery

Catches of Pacific ocean perch are monitored aboard the vessel by observers in the at-sea hake Observer program (ASHOP) and were available for the years of 1975-2016. Observers use a spatial sample design, based on weight, to randomly choose a portion of the haul to sample for species composition. For the last decade, this is typically 30-50% of the total

weight. The total weight of the sample is determined by all catch passing over a flow scale. All species other than hake are removed and weighed by species on a motion compensated flatbed scale. Observers record the weights of all non-hake species. Non-hake species total weights are expanded in the database by using the proportion of the haul sampled to the total weight of the haul. The catches of non-hake species in unsampled hauls is determined using bycatch rates determined from sampled hauls. Since 2001, more than 97% of the hauls have been observed and sampled.

Foreign Catches

From the 1960s through the early 1970s, foreign trawling enterprises harvested considerable amounts of rockfish off Washington and Oregon, and along with the domestic trawling fleet, landed large quantities of Pacific ocean perch. Foreign catches of individual species were estimated by Rogers (2003) and attributed to INPFC areas for the years of 1966-1976 for Pacific ocean perch. The foreign catches were combined across areas for a coastwide removal total.

2.2.2 Discards

Data on discards of Pacific ocean perch are available from two different data sources. The earliest source is referred to as the Pikitch data and comes from a study organized by Ellen Pikitch that collected trawl discards from 1985-1987 (Pikitch et al. 1988). The northern and southern boundaries of the study were 48°42' N latitude and 42°60' N latitude respectively, which is primarily within the Columbia INPFC area (Pikitch et al. 1988, Rogers and Pikitch 1992). Participation in the study was voluntary and included vessels using bottom, midwater, and shrimp trawl gears. Observers of normal fishing operations on commercial vessels collected the data, estimated the total weight of the catch by tow, and recorded the weight of species retained and discarded in the sample. Results of the Pikitch data were obtained from John Wallace (personal communication, NWFSC, NOAA) in the form of ratios of discard weight to retained weight of Pacific ocean perch and sex-specific length frequencies. Discard estimates are shown in Table 14.

The second source is from the West Coast Groundfish Observer Program (WCGOP). This program is part of the NWFSC and has been recording discard observations since 2003. Table 14 shows the discard ratios (discarded/(discarded + retained)) of Pacific ocean perch from WCGOP. Since 2011, when the trawl rationalization program was implemented, observer coverage rates increased to nearly 100% for all the limited entry trawl vessels in the program and discard rates declined compared to pre-2011 rates. Discard rates were obtained for both the catch-share and the non-catch share sector for Pacific ocean perch. A single discard rate was calculated by weighting discard rates based on the commercial landings by each sector. Coefficient of variations were calculated for the non-catch shares sector and pre-catch share years by bootstrapping vessels within ports because the observer program randomly chooses vessels within ports to be observed. Post-ITQ, all catch-share vessels have 100% observer

coverage and discarding is assumed to be known. Discard length composition for the trawl fleet varied by year, with larger fish being discarded prior to 2011 (Figure 15).

2.2.3 Fishery Length and Age Data

Biological data from commercial fisheries that caught Pacific ocean perch were extracted from PacFIN on May 4, 2017. Lengths taken during port sampling in Oregon and Washington were used to calculate length and age compositions. There were no biological data from California for Pacific ocean perch available within PacFIN or CALCOM databases. The overwhelming majority of these data were collected from the mid-water and bottom trawl gear, but additional biological data were collected from non-trawl gear which was grouped together with trawl gear data. Tables 15 and 16 show the number of trips and fish sampled, along with the calculated sample sizes. Length and age data were acquired at the trip level and then aggregated to the state level. The input sample sizes were calculated via the Stewart method (Ian Stewart, personal communication, IPHC):

$$\text{Input effN} = N_{\text{trips}} + 0.138 * N_{\text{fish}} \text{ if } N_{\text{fish}}/N_{\text{trips}} \text{ is } < 44$$

$$\text{Input effN} = 7.06 * N_{\text{trips}} \text{ if } N_{\text{fish}}/N_{\text{trips}} \text{ is } \geq 44$$

The fishery fleet observed Pacific ocean perch that were generally greater than 30 cm across all years of available data (Figure 16). The fishery fleet age data has clear patterns showing a large cohort moving through the population (Figure 17). Lengths and ages were also available for the at-sea hake fishery and are shown in Figures 18 and 19.

2.2.4 Fishery Data Not Included in the Base Model

Several dataset available from the fishery were explored but not used in the final assessment.

2.2.4.1 Historical Commercial Catch-Per-Unit Effort

Data on catch-per-unit-effort (CPUE) in mt/hr from the domestic fishery were combined for the INPFC Vancouver and Columbia areas from Gunderson (1977). Although these data reflect catch rates for the US fleet, the highest catch rates coincided with the beginning of removals by the foreign fleet. This suggests that, barring unaccounted changes in fishing efficiency during this period, the level of abundance was high at that time. Unfortunately, the original data and the analysis methods used to create this CPUE series have been lost to time precluding a re-analysis of these data. Due to the inability to examine the assumptions made during the original analysis or the data used this time-series has been excluded from the base model. These data were included in the previous assessment but were deemed not influential in the model estimates. Information regarding the fishery CPUE are available in Appendix 12.

2.2.4.2 Oregon Special Projects Length and Age Data

Oregon special project data were provided by ODFW. These data represent samples made at either the dock or at processing plants from fishery landings. Length data were collected primarily from 1970-1986, with limited samples from more recent years. Age data were primarily available from 1981-1984. These data were collected for special projects and may not have been sampled randomly from the fishery landings. Due to these concerns, these data were not included in the base model but were included in a model sensitivity. This was the first time these data were explored for consideration in the assessment.

2.3 Biological Data

2.3.1 Natural Mortality

Historical Pacific ocean perch ages determined using scales and surface reading methods of otoliths resulted in estimates of natural mortality (M) between 0.10 and 0.20 yr^{-1} with a longevity less than 30 years (Gunderson 1977). Based on break-and-burn method of age determination using otoliths, the maximum age of Pacific ocean perch was revised to be 90 years (Chilton and Beamish 1982). The updated understanding concerning Pacific ocean perch longevity reduced the estimate of natural mortality based on Hoenig’s (1983) relationship to 0.059 yr^{-1} . The previous assessment applied a prior distribution on natural mortality based upon multiple life-history correlates (including Hoenig’s method, Gunderson (1997) gonadosomatic index, and McCoy and Gillooly’s (2008) theoretical relationship) developed separately for female and male Pacific ocean perch.

Hamel (2015) developed a method for combining meta-analytic approaches relating the M rate to other life-history parameters such as longevity, size, growth rate, and reproductive effort to provide a prior on M . In that same issue of *ICES Journal of Marine Science*, Then et al. (2015) provided an updated data set of estimates of M and related life history parameters across a large number of fish species from which to develop an M estimator for fish species in general. They concluded by recommending M estimates be based on maximum age alone, based on an updated Hoenig non-linear least squares estimator $M = 4.899A_{\max}^{-0.916}$. The approach of basing M priors on maximum age alone was one that was already being used for West Coast rockfish assessments. However, in fitting the alternative model forms relating M to A_{\max} , Then et al. (2015) did not consistently apply their transformation. In particular, in real space, one would expect substantial heteroscedasticity in both the observation and process error associated with the observed relationship of M to A_{\max} . Therefore, it would be reasonable to fit all models under a log transformation. This was not done. Re-evaluating the data used in Then et al. (2015) by fitting the one-parameter A_{\max} model under a log-log transformation (such that the slope is forced to be -1 in the transformed space (Hamel 2015)), the point estimate for M is:

$$M = \frac{5.4}{A_{\max}}$$

The above is also the median of the prior. The prior is defined as a lognormal distribution with mean $\ln(5.4/A_{\max})$ and $SE = 0.438$. Using a maximum age of 100, the point estimate and median of the prior is 0.054 yr^{-1} . The maximum age was selected based on available age data from all West Coast data sources. The oldest aged rockfish was 120 years, captured by the commercial fishery in 2007. However, age data are subject to ageing error which could impact this estimate of longevity. The selection of 100 years was based on the range of other ages available with multiple observations of fish between 90 and 102 years of age.

2.3.2 Sex Ratio, Maturation, and Fecundity

Examining all biological data sources, the sex ratio of young fish are within 5% of 1:1 by length until larger sizes which are dominated by females who reach a larger maximum size relative to males (Figure 20), with the sex ratio being approximately equal across ages (Figure 21), and hence this assessment assumed the sex ratio at birth was 1:1. This assessment assumed a logistic maturity-at-length curve based on analysis of 537 fish maturity samples collected from the NWFSC shelf-slope survey. This is revised from the previous assessment that assumed maturity-at-age based on the work of Hannah and Parker (2007). Additionally, the new maturity-at-length curve is based on the estimate of functional maturity, an approach that classifies rockfish maturity with developing oocytes as mature or immature based on the proportion of vitellogenin in the cytoplasm and the measured frequency of atretic cells (Melissa Head, personal communication, NWFSC, NOAA). The 50% size-at-maturity was estimated at 32.1 cm with maturity asymptoting to 1.0 for larger fish (Figure 22). Comparison between the maturity-at-age used in the previous assessment and the updated functional maturity-at-length is shown in Figure 23.

The fecundity-at-length has also been updated from the previous assessment based on new research. Dick et al. (2017) estimated new fecundity relationships for select West Coast stocks where fecundity for Pacific ocean perch was estimated equal to $8.66e-10L^{4.98}$ in millions of eggs where L is length in cm. Fecundity-at-length is shown in Figure 24.

2.3.3 Length-Weight Relationship

The length-weight relationship for Pacific ocean perch was estimated outside the model using all biological data available from fishery-dependent and -independent data sources, where the female weight-at-length in grams was estimated at $1.003e-05L^{3.1}$ and males at $9.881e-06L^{3.1}$ where L is length in cm (Figures 25 and 26).

2.3.4 Growth (Length-at-Age)

The length-at-age was estimated for male and female Pacific ocean perch using data collected from both fishery-dependent and -independent data sources that were collected from 1981-

2016. Figure 27 shows the lengths and ages for all years and all data as well as predicted von Bertalanffy fits to the data. Females grow larger than males and sex-specific growth parameters were estimated at the following values:

$$\text{Females } L_{\infty} = 42.32; k = 0.169; t_0 = -1.466$$

$$\text{Males } L_{\infty} = 39.03; k = 0.212; t_0 = -1.02$$

These values were used as starting parameter values within the base model prior to estimating each parameter for male and female Pacific ocean perch.

2.3.5 Ageing Precision and Bias

Uncertainty surrounding the age-reading error process for Pacific ocean perch was incorporated by estimating ageing error by age. Age-composition data used in the model were from break-and-burn otolith reads aged by the Cooperative Ageing Project (CAP) in Newport, Oregon. Break-and-burn double reads of more than 1500 otoliths were provided by the CAP lab. An ageing-error estimate was made based on these double reads using a computational tool specifically developed for estimating ageing error (Punt et al. 2008) and using release 1.0.0 of the R package `nwfscAgeingError` (Thorson et al. 2012) for input and output diagnostics, publicly available at: <https://github.com/nwfsc-assess/nwfscAgeingError>. A non-linear standard error was estimated by age, where there is more variability in the age of older fish (Table 19 and Figure 28).

2.4 History of Modeling Approaches Used for This Stock

2.4.1 Previous Assessments

The status of Pacific ocean perch off British Columbia, Washington, and Oregon have been periodically assessed since the intensive exploitation that occurred in the 1960s. Concerns regarding Pacific ocean perch status off the coast the US west coast were raised in the late 1970s (Gunderson 1978, 1981) and in 1981 the PFMC adopted a 20-year plan to rebuild the stock.

The 1992 assessment determined that Pacific ocean perch remained at low levels relative to the population size in 1960 (Ianelli et al. 1992) and recommended additional harvest restrictions to allow for stock rebuilding. The 1998 assessment (Ianelli and Zimmermann 1998) estimated that the stock was 13% of the unfished level, leading the National Marine Fishery Service (NMFS) to declare the stock overfished in 1999. A formal rebuilding plan was implemented in 2001. The rebuilding plan reduced the SPR harvest rate used to determine catches to 0.864 (in contrast to the default harvest rate of 0.50). The last full assessment of

Pacific ocean perch was conducted in 2011 (Hamel and Ono 2011), which concluded that the stock was still well below the target biomass of $40\%SB_0$ estimating the relative stock status at 19.1%.

3 Assessment

3.1 General Model Specifications and Assumptions

Stock Synthesis version 3.30.03.05 was used to estimate the parameters in the model. R4SS, version 1.27.0, along with R version 3.3.2 were used to investigate and plot model fits. A summary of the data sources used in the model (details discussed above) is shown in Figure 2.

3.1.1 Changes Between the 2011 Assessment Model and Current Model

The current model for Pacific ocean perch has many similar assumptions as the 2011 assessment but differs in some key ways. In this assessment, fleets were disaggregated into a trawl/other gear, at-sea hake, historical foreign fleet, and research fleets. The previous assessment implemented a single fleet where removals from all sources were aggregated together. The separating of fleets applied in this assessment allowed for differing assumptions regarding current and historical discarding practices. Although there are no compositional data available from the foreign fleet, it is assumed that very little to no discarding of fish occurred. Additionally, the at-sea hake fishery removals represent both discarded and retained fish and hence an additional discard rate would not be appropriate. Similar logic was applied in regard to survey removals.

The historical landings used in the model differ from those used in 2011. This assessment includes the first state provided historical reconstruction landings for Washington. The historical reconstruction has removals starting in 1908 and has larger removals in the 1940s relative to those used in the 2011 assessment (Figure 30). The starting year for modeling the stock was revised to 1918, the first year Pacific ocean perch landings exceeded 1 mt, rather than 1940 as modeled in the previous assessment, given the new information regarding historical removals prior to 1940. Explorations were conducted relative to the model starting year and no differences were found between the 1918 start year compared to starting the model in 1892, which is the first year there is record landings of Pacific ocean perch between California, Oregon, and Washington.

Selectivity in this model is assumed to be length-based and is modeled using double-normal selectivity for all fleets, except the Pacific ocean perch survey which retained the assumption used in previous assessment of logistic selectivity. The previous assessment mirrored selectivity

among the Pacific ocean perch and both slope surveys (AFSC and NWFSC). This assessment allows for survey-specific selectivity.

All fishery-independent indices have been re-evaluated for this assessment using a spatial-temporal delta generalized linear mixed model (VAST delta-GLMM) which is an updated approach from that used in 2011, which did not incorporate spatial effects. This assessment opted to not include the fishery CPUE and the Triennial shelf index and composition data based upon discussions during the STAR panel. The data used to create the CPUE index were not available for reanalysis and hence were excluded from this assessment due to questions regarding this index that could not be addressed. In regards to the Triennial survey, Pacific ocean perch is considered a slope species off the US west coast and this survey did not sample the prime habitat for Pacific ocean perch and had limited observations relative to the other surveys. It was concluded during the STAR panel that this data set was not a good source of information regarding this species and would not be included in the base model.

Maturity and fecundity were updated for this assessment based upon new research. Fecundity for Pacific ocean perch used in this assessment was based on a re-evaluation of the fecundity of West Coast rockfish by Dick et al. (2017), updating the previous fecundity estimates used in the 2011 assessment (Dick 2009) (Figure 24). Maturity in this assessment was based on examination of 537 fish samples which were used to estimate functional maturity, an approach that classifies rockfish maturity with developing oocytes as mature or immature based on the proportion of vitellogenin in the cytoplasm and the measured frequency of atretic cells (Melissa Head, personal communication, NWFSC, NOAA). The updated maturity curve was based on maturity-at-length where the previous estimates used in 2011 were based on maturity-at-age (Figure 23).

In this assessment, the beta prior for steepness developed from a meta-analysis of West Coast groundfish was updated to the 2017 value (James Thorson, personal communication, NWFSC, NOAA), with steepness fixed at the median of the prior in the final base model (0.72). Additionally, the prior for natural mortality was updated based on an analysis conducted by Owen Hamel (personal communication, NWFSC, NOAA), where female and male natural mortality were fixed at the median of the prior (0.054 yr^{-1}).

3.1.2 Summary of Fleets and Areas

Pacific ocean perch are most frequently observed in Oregon and Washington waters in survey and fishery observations. Multiple fisheries encounter Pacific ocean perch. Bottom trawl, mid-water trawl, fixed gear, and the at-sea (mid-water) hake fisheries account for the majority of the current Pacific ocean perch landings.

The majority of removals of Pacific ocean perch are attributable to trawl gears with fixed gear accounting for a small fraction of the catches available within PacFIN. Trawl and fixed gears were combined into a coast-wide fleet. For the period from 1918 to the early 1990s, prior

to the introduction of trip limits for rockfish, limited discarding of Pacific ocean perch was assumed. Observations of Pacific ocean perch in the Pikitch et al. (1988) data (1986-1987) allowed for a formal analysis of discard rates that were applied to the historical period of the fishery. Foreign trawl catches (1966-1976) were modeled as a single fleet. The at-sea hake fishery operates as a mid-water fishery targeting Pacific whiting but encounters Pacific ocean perch as a bycatch species. This fleet was also modeled as a single fleet.

3.1.3 Other Specifications

The specifications of the assessment are listed in Table 20. The model is a two-sex, age-structured model starting in 1918 with an accumulated age group at 60 years. Growth and natural mortality were assumed time invariant with a constant growth estimated and natural mortality fixed at the median of the prior. The lengths in the population were tracked by 1 cm intervals and the length data were binned into 1 cm intervals. A curvilinear ageing imprecision relationship was estimated and used to model ageing error. Fecundity-at-length was fixed at the values from Dick et al. (2017) for Pacific ocean perch and spawning output was defined in millions of eggs.

Age data for the commercial and at-sea hake fisheries, as well as the Pacific ocean perch, the NWFSC slope, and the NWFSC shelf-slope surveys were used in this assessment. The ages from the NWFSC shelf-slope survey were entered into the model as conditional age-at-length. The assessment used length-frequencies collected by the fishery fleet, the at-sea hake fishery, and Pacific ocean perch, AFSC slope, NWFSC slope, and the NWFSC shelf-slope surveys.

The specification of when to estimate recruitment deviations is an assumption that likely affects model uncertainty. Recruitment deviations were estimated from 1900-2014 to appropriately quantify uncertainty. The earliest length-composition data occur in 1966 and the earliest age data were in 1981. The most informed years for estimating recruitment deviations were from about the mid-1970s to 2013. The period from 1900-1974 was fit using an early series with little or no bias adjustment, the main period of recruitment deviates occurred from 1975-2014 with an upward and downward ramping of bias adjustment (Figure 29), and 2015 onward were fit using forecast recruitment deviates with no bias adjustment. Methot and Taylor (2011) summarize the reasoning behind varying levels of bias adjustment based on the information available to estimate the deviates. The standard deviation of recruitment variability was assumed to be 0.70 based on the estimated variation in recruitment from the base model.

The recommended selectivity in Stock Synthesis is the double-normal parameterization and it was used in this assessment for the all fleets, except the Pacific ocean perch survey, which was assumed logistic based on the length-composition data. Changes in retention curves were estimated for the fishery fleet.

Time blocks for the fishery fleet are provided in Table 20. Fishery selectivity and retention has changed over the modeled period due to management changes. The time block for fishery

selectivity was set from 1918-1999 and 2000-2017 based on changes in selectivity arising from the overfished deceleration. The time blocks on the retention curves for the fishery were set from 1918-1991, 1992-2001, 2002-2007, 2008, 2009-2010, 2011-2016 based on available discarding data and changes in trip limits that likely resulted in changes to discarding patterns of Pacific ocean perch. No discarding was assumed in the at-sea hake and the foreign fisheries. The length data are not available from the foreign fleet. The selectivity from this fleet was mirrored to the main fishery fleet.

The following distributions were assumed for data fitting: survey indices were lognormal, total discards were lognormal, and multinomial error structure for compositional data.

3.1.4 Modeling Software and Model Bridging

The STAT team used Stock Synthesis version 3.30.03.05 developed by Dr. Richard Methot at the NWFSC (Methot and Wetzel 2013). This most recent version was used because it included improvements and corrections to older versions. The previous assessment of Pacific ocean perch also used Stock Synthesis but an earlier version, 3.24; model bridging was performed between both versions of Stock Synthesis and are shown in Figure 31.

3.1.5 Priors

A prior distribution was developed for natural mortality (M) from an analysis based on an assumed maximum age of 100 years. The analysis was performed by Owen Hamel (personal communications, NWFSC, NOAA) and used data from Then et al. (2015) to provide a lognormal distribution for natural mortality. The lognormal prior has a median of 0.054 and a standard error of 0.438.

The prior for steepness (h) assumed a beta distribution with parameters based on an update of the Thorson-Dorn rockfish prior (commonly used in past West Coast rockfish assessments) conducted by James Thorson (personal communication, NWFSC, NOAA) which was reviewed and endorsed by the Scientific and Statistical Committee in 2017. The prior is a beta distribution with $\mu=0.72$ and $\sigma=0.15$. The previous assessment estimated and fixed steepness equal to 0.40. The current data do not contain information regarding steepness. This change in perception is likely due to the observation of large recruitment events in this assessment and varying model specifications between the 2011 and the current model.

The final base model had both of these parameters fixed at the median of the priors in the base model due to lack of information in the data concerning natural mortality and steepness.

3.1.6 Data Weighting

Length and age-at-length compositions from the NWFSC shelf-slope survey were fit along with length and marginal age compositions from the fishery and other survey fleets. Length data started with a sample size determined from the equation listed in Sections 2.1.1 (survey data) and 2.2.3 (fishery data). It was assumed for age-at-length data that each age was a random sample within the length bin and the model started with a sample size equal to the number of fish in that length bin.

One extra variability parameter was estimated and added to the input variance for the NWFSC shelf-slope survey index. Estimating additional variance for the other surveys was explored and determined to not be required. WCGOP data were bootstrapped to provide uncertainty of the total discards (Table 14).

The base assessment model was weighted using the “Francis method”, which was based on equation TA1.8 in Francis (2011). This formulation looks at the mean length or age and the variance of the mean to determine if across years, the variability is explained by the model. If the variability around the mean does not encompass the model predictions, then that data source should be down-weighted. This method accounts for correlation in the data (i.e., the multinomial distribution) as opposed to the McAllister and Ianelli (1997) method (Harmonic Mean weighting) of looking at the difference between individual observations and predictions. A sensitivity was performed examining the difference between the weighting approaches. The weights applied to each length and age data set for the base model are shown in Table 21.

3.1.7 Estimated and Fixed Parameters

There were 163 estimated parameters in the base model. These included one parameter for R_0 , 8 parameters for growth, 1 parameters for extra variability for the NWFSC shelf-slope survey index, 24 parameters for selectivity, retention, and time blocking of the fleets and the surveys, 117 recruitment deviations, and 12 forecast recruitment deviations (Table 22).

Fixed parameters in the model were as follows. Steepness was fixed at 0.72, the median of the prior. A sensitivity analysis and a likelihood profile were performed for steepness. Natural mortality was fixed at 0.054 yr^{-1} for females and males, which is the median of the prior. The standard deviation of recruitment deviates was fixed at 0.70. Maturity-at-length was fixed as described in Section 2.3.2. Length-weight parameters were fixed at estimates using all length-weight observations (Figure 26).

Dome-shaped selectivity was explored for all fleets within the model. Older Pacific ocean perch are often found in deeper waters and may move into areas that limit their availability to fishing gear, especially trawl gear. The final base model estimated dome-shaped selectivity for only the fishery. The selectivities for the the at-sea hake fishery and all surveys were estimated asymptotic.

3.2 Model Selection and Evaluation

The base assessment model for Pacific ocean perch was developed to balance parsimony and realism, and the goal was to estimate a spawning output trajectory for the population of Pacific ocean perch off the west coast of the US. The model contains many assumptions to achieve parsimony and uses many different sources of data to estimate reality. A series of investigative model runs were done to achieve the final base model.

3.2.1 Key Assumptions and Structural Choices

The key assumptions in the model were that the assessed population is a single stock with biological parameters characterizing the entire coast; natural mortality, maturity-at-length, length-at-age, and weight-at-length have remained constant over the period modeled; the standard deviation in recruitment deviation is 0.70; and steepness is 0.72. These are simplifying assumptions that unfortunately cannot be verified or disproved. Sensitivity analyses were conducted for most of these assumptions to determine their effect on the results.

Structurally, the model assumed that the landings from each fleet were representative of the coastwide population, instead of specific areas, and fishing mortality prior to 1918 was negligible. It also assumed that discards were low prior to 1992.

3.2.2 Alternate Models Considered

The exploration of models began by bridging from the 2011 assessment to Stock Synthesis version 3.30.03.05, which produced no discernible difference (Figure 31). The updated landings data and discard rates added to the 2011 assessment produced insignificant differences in the relative scale of the population although the updated historical removals resulted in an increase in the estimate of unfished spawning output (Figure 32). Updating the survey indices produced small differences in the relative scale of the population. Adding age and length data each resulted in less of a population decline from the 1970s to pre-2000, resulting in an increase in the estimated 2017 final stock status. However, the addition of new data resulted in an early pattern within recruitment, indicating that the assumptions within the previous model may not represent the best fit to the current data.

This assessment estimated discards in the model, so time was spent investigating time blocks for changes in selectivity and retention to match the discard data as best as possible. Using major changes in management and observed changes in landings, a set of blocks for retention were determined for the fishery fleet. In the spirit of parsimony, as few blocks as possible were used by only allowing blocks during time periods with data or when we felt they were justified by changes in management.

Natural mortality was also investigated and a new prior was developed assuming a maximum age of 100 years for females and males. The previous assessment estimated male natural mortality as an offset from a fixed female natural mortality. This assessment attempted to estimate natural mortality for both sexes using the 2017 updated prior, but based on the results of profiles over the parameter it was determined to be little to no information on natural mortality within the data and hence opted to fix the value for females within the base model. Upon additional exploration, the model estimated very little difference in male natural mortality relative to females (< 0.002) and in the interest of selecting the model that fit the data with the fewest parameters required, natural mortality for males was fixed equal to the female natural mortality in the base model.

Finally, multiple models were investigated where steepness was either estimated, fixed at the prior, or at an alternate value. The assessment in 2011 determined that there was sufficient information concerning steepness where the parameter was estimated and then fixed at the estimated value of 0.40. Based upon likelihood profiles performed on the current model, there was no longer support for a steepness value of 0.40. The likelihood profile was flat across various levels of steepness with a very small improvement in likelihood (< 0.50 log likelihood units) at the lowest steepness values. Estimating steepness starting at the median of the “type C” prior, the meta-analysis prior evaluated omitting information from Pacific ocean perch, of 0.76 resulted in very little if any movement from the median value due to the flat likelihood surface across values for this parameter with the final relative stock status for 2017 being estimated to $> 100\%$ of unfished spawning output. The base model fixed steepness at the median of the prior ($h = 0.72$). A model sensitivity is provided when steepness was fixed at the 2011 value of 0.40.

3.2.3 Convergence

Proper convergence was determined by starting the minimization process from dispersed values of the maximum likelihood estimates to determine if the model found a better minimum. Starting parameters were jittered by 10%. This was repeated 50 times and a better minimum was not found (Table 23). The model did not experience convergence issues when provided reasonable starting values. Through the jittering done as explained above and likelihood profiles, we are confident that the base model as presented represents the best fit to the data given the assumptions made. There were no difficulties in inverting the Hessian to obtain estimates of variability, although much of the early model investigation was done without attempting to estimate a Hessian.

3.3 STAR Panel Review and Recommendations

3.4 Response to the 2011 STAR Panel Recommendations

Recommendation: Considering trans-boundary stock effects should be pursued. In particular, the consequences of having spawning contributions from external stock components should be evaluated relative to the steepness estimates obtained in the present assessment.

STAT response: The STAT team agrees that this should be an ongoing area of research and collaboration between the US and Canada. This assessment presents a sensitivity where the inclusion of Canadian data are included within the model.

Recommendation: The benefits of adopting the complex model used this year should be evaluated relative to simpler assumptions and models. While the transition from the simpler old model to Stock Synthesis was shown to be similar for the historical period, the depletion estimates in the most recent years were different enough to warrant further investigation.

STAT response: This assessment was performed in Stock Synthesis, an integrated model, which can be modified to either simple or complex structural forms based upon the available data and the processes being modeled. There were not additional explorations of alternative modeling platforms.

Recommendation: Discard estimates from observer programs should be presented, reviewed (similar to the catch reconstructions), and be made available to the assessment process.

STAT response: This assessment uses discard rates and discard lengths collected by the WCGOP from 2003-2015.

Recommendation: The ability to allow different “plus groups” for specific data types should be evaluated (and implemented in Stock Synthesis). For example, this would provide the ability to use the biased surface-aged data in an appropriate way.

STAT response: The STAT team agrees that this should be explored, but additional research needs to be completed which evaluates the amount of bias and imprecision in surface-read ages. Evaluating available surface-read ages within the PacFIN database fish of lengths between 23-44 cm can be aged at 10 years old. This large range of lengths at the same age indicates considerable bias in ages for fish surface-read younger aged fish.

Recommendation: Historical catch reconstruction estimates should be formally reviewed prior to being used in assessments and should be coordinated so that interactions between stocks are appropriately treated. The relative reliability of the catch estimates over time could provide an axis of uncertainty in future assessments.

STAT response: California and Oregon have undergone extensive work to create historical catch reconstructions. This is the first assessment for Pacific ocean perch which includes

a Washington historical catch reconstruction. The data used in this assessment represent Washington state's current best estimate for historical catches. An historical catch reconstruction meeting was held in November of 2016 where states discussed methods and approaches to improve historical catch estimates. Additionally, both California and Washington are conducting research to estimate uncertainty surrounding historical catches which could be used to propagate uncertainty within the assessment.

3.5 Response to the 2017 STAR Panel Requests

The stock assessment review (STAR) panel for this assessment was held at the NWFSC in Seattle, WA from June 26-30, 2017. David Sampson was the chair, while Norman Hall, Kevin Piner, and Yiota Apostolaki were invited reviewers. It was a productive and busy review that thoroughly reviewed many facets of the assessments. As mentioned above, changes to the data used in this assessment were made during the panel.

Recommendation: Further investigation of Pacific ocean perch stock structure is recommended. One approach would be to look for correlations of US west coast recruitment deviations and survey biomass estimates with corresponding results from Pacific ocean perch assessments in Canada and Gulf of Alaska.

STAT response: We agree. A preliminary analysis using a subset of Canadian data was provided as a sensitivity, but further investigations should be conducted.

Recommendation: The next iteration of this assessment could be an update assessment.

STAT response: We agree.

Additionally, a number of general recommendations were made for all West Coast assessments:

Recommendation: Comprehensively evaluate the appropriateness of using the Triennial survey in assessments for other rockfish species and whether the survey should be split into early and late segments. The lingcod assessment reviewed during this STAR split the Triennial survey into separate early and late surveys, whereas the draft Pacific ocean perch assessment brought to the STAR had a single Triennial survey.

STAT response: We agree. As a whole this dataset should be evaluated to determine which West Coast species were well sampled. However, the treatment of keeping the data set as a single time-series or splitting into early or late periods may need to be considered on a species specific level. Changes in sampling range and timing may not impact or having differing levels of impact for West Coast species.

Recommendation: Explore the assumption that conditional age-at-length data are random samples of the age-composition.

STAT response: We agree. The conditional age-at-length data are highly influential in the model. Some explorations were conducted during the STAR panel examining the impact of these data and to determine if the underlying assumptions of the data were violated based on age-based processes. Further research should be conducted examining the assumptions of these data.

Recommendation: A standard approach for combining conditional age-at-length sample data into annual conditional-age-at-length compositions should be developed and reviewed. If age data are not selected in proportion to the available lengths, simple aggregation of the ages by length-bin may provide biased views of the overall age-composition and year-class strength.

STAT response: We agree.

Recommendation: Further explore the VAST approach for constructing relative abundance indices. The upcoming workshop at the Center for the Advancement of Population Assessment Methodology (CAPAM) will address this issue.

STAT response: The trend of the indices created using VAST and the Bayesian delta-glm which did not explicitly account for spatial dynamics were consistent with each other. However, assessments in general will benefit, from continued research regarding the best way to generate indices of abundance for fishery and non-fishery data.

3.6 Base Model Results

The base model parameter estimates along with approximate asymptotic standard errors are shown in Table 22 and the likelihood components are shown in Table 24. Estimates of derived reference points and approximate 95% asymptotic confidence intervals are shown in Table 25. Estimates of stock size over time are shown in Table 26.

3.6.1 Parameter Estimates

The estimates of maximum length and the von Bertalanffy growth coefficient, k , were less than the the external estimates for males and female but were well within the 95% confidence interval given the estimated uncertainty (Table 22, Figures 33 and 34). Female and male Pacific ocean perch grow quickly at younger ages, reaching near maximum length by age 20, with female Pacific ocean perch reaching larger maximum lengths.

Selectivity curves were estimated for the fishery and survey fleets. The estimated selectivities for all fleets within the model are shown in Figure 35. The fishery selectivity was estimated to be dome shaped, reaching maximum selectivity for fish between 35 and 40 cm. An shift in selectivity for the final asymptotic selectivity was estimated for the fishery for prior to the overfished decleration and post (1918-1999 and 2000-2016). The at-sea hake fishery was

estimated to have little selectivity for smaller Pacific ocean perch reaching full selectivity at the largest sizes. The foreign fleet for which only catch data are available was assumed to be identical to the main fishery, although a sensitivity was performed (not shown) that mirrored the foreign selectivity to that of the Pacific ocean perch survey selectivity resulting in a negligible difference in stock status. Survey selectivities were estimated to be asymptotic during model explorations with the final selectivity forced to be asymptotic in the final base model.

Retention curves were estimated for the fishery fleet only and were allowed to vary based upon discard data within the model over time (Figure 36). Historical retention was estimated to be high and declined over time due to management restriction on landings of Pacific ocean perch with the lowest retention occurring in 2009 and 2010 prior to the implementation of ITQs. Post-2011 retention was estimated to be nearly 100% for the fishery fleet.

Additional survey variability (process error added directly to each year's input variability) for the NWFSC shelf-slope survey was estimated within the model. The model estimated a small added variance for the NWFSC shelf-slope survey of 0.018. Preliminary models explored estimating added variance for each of the other indices, but resulted in no added variance being estimated and hence the added variance parameters were not estimated in the base model.

Estimates of recruitment suggest that the Pacific ocean perch population is characterized by variable recruitment with occasional strong recruitments and periods of low recruitment (Figures 37 and 38). There is little information regarding recruitment prior to 1970 and the uncertainty in those estimates is expressed in the model. The four lowest recruitments (in ascending order) occurred in 2012, 2003, 2005, and 2007. There are very large, but uncertain, estimates of recruitment in 2008, 2013, 2000, and 1999. The 2008 recruitment event is estimated to be larger by an order of magnitude compared to other recruitments estimated in the model. The uncertainty interval around the number of recruits is large based on the uncertainty surrounding the spawning output in that year. However, the uncertainty around the recruitment deviation estimated is low.

3.6.2 Fits to the Data

There are numerous types of data for which the fits are discussed: survey abundance indices, discard data (biomass and length compositions), length-composition data for the fisheries and surveys, marginal age compositions for the fisheries and surveys, and conditional age-at-length observations for the NWFSC shelf-slope survey.

The fits to the survey indices are shown in Figure 39. Extra standard error was estimated for the NWFSC shelf-slope survey. The Pacific ocean perch survey index were fit well by the model. Both the AFSC and NWFSC slope survey indices were generally flat and fit well by the model. The recent NWFSC shelf-slope survey showed a variable trend over the

time period with the 2016 data point being the highest estimate of the series and given the uncertainty around each data point (input and model estimated added variance) the model fit fell within the uncertainty interval for all years.

Fits to the total observed discards required time blocks (Figure 40). Fits to the trawl discards from the Pikitch data in 1985-1987 were quite good. The change in the discard rate modeled over 1992-2001 was based on management restrictions, which were assumed to have increased discarding practices in the fishery fleet. The next required time block was based on the WCGOP data from 2002-2007 and were fit well by the model. Discarding increased prior to the implementation of ITQs requiring blocks for 2008 and the 2009-2010 periods. The model fit the very low post-ITQ discard rates based on the WCGOP data well. The total estimated discard amount over time is shown in Figure 41.

Fits to the length data are shown based on the proportions of lengths observed by year and the Pearson residuals-at-length for all fleets. Detailed fits to the length data by year and fleet are provided in Appendix 10. Aggregate fits by fleet are shown in Figure 42. There are a few things that stand out when examining the aggregated length composition data. First, the sexed discard lengths appear to be poorly fit by the model but this is related to small sample sizes. The NWFSC slope survey lengths were under estimated by the model, but these data only represent two years.

Discard lengths from WCGOP were fit well by the model and show no obvious pattern in the residuals (Figure 43). The residuals to the fishery lengths clearly showed the growth differential between males and females where the majority of positive residuals at larger sizes were from female fish (Figure 44). The fishery showed large positive residuals for smaller fish for 2013-2016 which were attributed to the strong 2008 year class moving through the fishery. The at-sea hake fishery did not show an obvious pattern in residuals but clearly showed the selectivity of larger fish (Figure 45). The residuals for each of the surveys are shown in Figures 46, 47, 48, and 49. The Pearson residuals from the NWFSC shelf-slope survey clearly showed the strong year classes moving through the population.

Length data were weighted according to the Francis weights that adjust the weight given to a data set based on the fit to the mean lengths by year. The mean lengths from the fishery were consistent across the sampled period, showing only a decline in the mean length in 2013-2015 likely due to the large 2008 cohort (Figure 50). The at-sea hake fishery showed an increase in the mean length of fish observed to 2009 and then fluctuated at larger mean lengths thereafter (Figure 51). The mean lengths were consistent across the two sample years of the Pacific ocean perch survey (Figure 52). However, the model expected a decline in mean length over the period. The trend in the mean lengths observed by the AFSC slope survey was generally flat excluding the samples from 1997 which were smaller fish (Figure 53). The NWFSC slope length data from 2001 and 2002 were highly variable with differing mean lengths between the years which were not fit well by the model (Figure 54). The mean length for the NWFSC shelf-slope survey declined in 2012 and 2016 due to observations of young, small fish by the survey (Figure 55).

Age data were fitted to as marginal age compositions for the main fishery fleet, the at-sea

hake fishery, the Pacific ocean perch survey, and the NWFSC slope survey. The NWFSC shelf-slope ages were treated as conditional age-at-length data to facilitate the estimation of growth within the model. The aggregated fits to the marginal age data are shown in Figure 56. The aggregated age data were fit well for the fishery fleet which had the largest sample of ages. The at-sea hake fishery and the surveys had significantly lower sample sizes that resulted in spiky patterns in the aggregated data. However, the model generally captured the pattern of the data. Detailed fits to the age data by year and fleet are provided in Appendix 11.

The Pearson residuals for the main fishery fleet are shown in Figure 57. There are diagonal patterns in the residuals across years, which likely are cohorts moving through the fishery. The at-sea hake fishery only had age data for four non-consecutive years, combined with the tendency of this fleet to select older fish, preventing general conclusions regarding fits to the data and cohort strength over time (Figure 58). The Pacific ocean perch survey only had one year of age data (the 1979 were all surface reads), but both sexes had a larger observed number of older fish relative to the model estimates (Figure 59). The Pearson residuals for the two years of age data from NWFSC slope survey are shown in Figure 60. The residual pattern differs between the years and by sex with positive residuals of male fish across ages in the 2001 data.

The observed and expected conditional age-at-length fits are shown in Figures 61, 62, 63, 64, and 65 for the NWFSC shelf-slope survey observations. The fits generally match the observations. Some outliers are apparent with large residuals. The 2016 data varies from previous years, where larger fish across all ages have higher observations compared to the model expectations.

The age data were also weighted according to Francis weighting which adjusted the weight given to a data set based on the fit to the mean age by year. The mean ages from the fishery appear to have declined in recent years which could be due to incoming cohorts (Figure 66). The at-sea hake fishery mean ages are similar for 2006 and 2007, but both 2003 and 2014 have lower average ages in the samples (Figure 67). The NWFSC slope had a decline in the mean age between the two data years (Figure 68). The mean age for the NWFSC shelf-slope survey generally showed a declining trend over the time-series excluding 2013 and 2016 which sampled older fish relative to the other years (Figure 69).

3.6.3 Population Trajectory

The predicted spawning output (in millions of eggs) is given in Table 26 and plotted in Figure 70. The predicted spawning output from the base model generally showed a slight decline over the time-series until when the foreign fleet began. A short, but sharp decline occurred during the period of the foreign fishery in the late 1960s. The stock continued to decline minimally until 1975 (41.8%) when a combination of strong recruitment and low catches resulted in an increase in spawning output at the end of the time-series. The recent increase is even

faster for total biomass (Figure 71) because not all fish from the 2008 recruitment are mature (Figure 23) with the model estimating the final year total biomass higher than unfished conditions. The 2017 spawning output relative to unfished equilibrium spawning output is above the target of 40% of unfished spawning output (96.4%) (Figure 72). Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in the estimated spawning output is high, especially in the early years. The standard deviation of the log of the spawning output in 2017 is 0.24.

Recruitment deviations were estimated for the entire time-series that was modeled (Figure 37 and discussed in Section 3.6.1) and provide a realistic portrayal of uncertainty. Recruitment predictions from the mid-1970s and early 1980s were mostly below average, with the 1999, 2000, 2008, and 2013 cohorts being the strongest over the modeled period. Many other stock assessments of rockfish along the west coast of the US have estimated a large recruitment event in 1999 (e.g., greenstriped rockfish (Hicks et al. 2009), chilipepper rockfish (Field 2007), darkblotched rockfish (Gertseva et al. 2015)). The 2008 year class was estimated as the strongest year class measured to date for Pacific ocean perch. This year has been estimated to have very strong year classes for other West Coast stocks (e.g., darkblotched rockfish (Gertseva et al. 2015), widow rockfish (Hicks and Wetzel 2015)). It may be worthwhile to investigate the periods of strong and weak year classes further to see if it is an artifact of the data, a consistent autocorrelation, or a result of the environment.

The stock-recruit curve resulting from a fixed value of steepness is shown in Figure 73 with estimated recruitments also shown. The stock is predicted to have never fallen to low enough levels that the effects of steepness are obvious. However, the lowest levels of predicted spawning output showed some of the smallest recruitments and very few above average recruitments. Steepness was not estimated in this model, but a sensitivity to an alternative value of steepness is discussed below.

3.6.4 Uncertainty and Sensitivity Analyses

A number of sensitivity analyses were conducted. Each of the sensitivities conducted were single explorations from the base model assumptions and data and were not performed in a cumulative fashion.

1. Data weighting according to the harmonic mean (McAllister, and Ianelli, 1997). The data weights used in this sensitivity are showing in Table 29 and can be compared to the weights used in the base model shown in Table 21.
2. Fixed steepness at the value assumed in the 2011 assessment of 0.40.
3. Maturity relationship used in the previous assessment.
4. Fecundity relationship used in the previous assessment.

5. Remove the influence of the large 2008 year-class by setting the 2008 recruitment deviation to zero (estimated straight from the stock recruitment curve)
6. Include Triennial shelf survey (1980-2004) and composition data.
7. Remove all other surveys and associated length and age data, except for the Triennial shelf survey. Fishery length and ages were retained.
8. Include the historical commercial CPUE index.
9. Inclusion of available Canadian fishery and survey data (does not constitute all data used in Canadian assessments). This sensitivity includes Canadian fishery landings (1997-2016 with landings ranging from 260-400 mt by year) and survey removals (2004, 2006, 2008, 2010, 2012, 2014, 2016), no fishery or survey index of abundance, but length- and age-composition data from both the fishery and survey.
10. Inclusion of historical Washington research lengths.
11. Inclusion of Oregon special projects length and age data, which are sampled at the dockside or processing facilities.

Likelihood values and estimates of key parameters from each sensitivity are available in Tables 27 and 28. Plots of the estimated time-series of spawning output and relative depletion are shown in Figures 74, 75, 76, and 77.

The sensitivities that explored steepness and using only Triennial survey data exhibited the largest changes in estimated stock status relative to the base model. Fixing steepness at the 2011 value ($h = 0.40$) resulted in the final stock status being above the management target but having historically declining below the target. Using only the Triennial shelf survey data resulted in a reduction in stock size and status relative to the base model. The model estimated extended positive recruitments in the early period of the model in order to create an population age-structure that was consistent with the composition data resulting in an increase in biomass prior to the start of the foreign fishing fleet, indicating that potential model misspecification in the absence of the other survey data.

Weighting the data according to the harmonic means resulted in a largest decrease in the estimated stock status relative to the base model with the stock being estimated at 82% of unfished spawning output.

The sensitivity that removed the large 2008 year-class resulted in a large change in estimated stock status relative to the base model. Assuming a recruitment even straight from the stock recruitment curve resulted in an estimated stock status of 75%.

Including additional data from either Canada, Washington research lengths, and or Oregon special projects data resulted in minimal reduction in the stock status relative to the base model.

The sensitivities that explored the inclusion of the CPUE index, the 2011 maturity, or fecundity relationship had little impact relative to the base model estimated stock status.

3.6.5 Retrospective Analysis

A 5-year retrospective analysis was conducted by running the model using data only through 2011, 2012, 2013, 2014, and 2015, progressively (Figure 78 and 79). The initial scale of the spawning population was basically unchanged for all of these retrospectives. The estimation of the 2008 recruitment deviation decreased as more data were removed. Overall, no alarming patterns were present in the retrospective analysis.

3.6.6 Likelihood Profiles

Likelihood profiles were conducted for R_0 , steepness, and natural mortality values separately. These likelihood profiles were conducted by fixing the parameter at specific values and estimated the remaining parameters based on the fixed parameter value.

For steepness, the negative log-likelihood was essentially flat between values of 0.30 - 0.80 (Figure 80). Likelihood components by data source show that the fishery length and age data support a low steepness value, but the NWFSC shelf-slope age data support a higher value for steepness. The surveys generally do not provide information concerning steepness. The relative depletion for Pacific ocean perch has a wide range across different assumed values of steepness (Figure 81).

The negative log-likelihood was minimized at a natural mortality value of 0.06, but the 95% confidence interval extends over values ranging from 0.035 - 0.08. The age and length data likelihood contribution was minimized at natural mortality values ranging from 0.055-0.06 (Figure 82). The relative depletion for Pacific ocean perch widely varied across alternative values of natural mortality (Figure 83).

In regards to values of R_0 , the negative log-likelihood was minimized at approximately $\log(R_0)$ of 9.45 (Figure 84). The fishery and survey composition data were in opposition regarding values of R_0 where the fishery length and age data indicated lower values of R_0 while the survey ages from the Pacific ocean perch and the NWFSC shelf-slope surveys indicated a higher value. The survey indices were uninformative concerning R_0 , an issue that was explored and discussed in depth during the week of the STAR panel.

3.6.7 Reference Points

Reference points were calculated using the estimated selectivities and catch distributions among fleets in the most recent year of the model (2016). Sustainable total yields (landings

plus discards) were 2,555.8 mt when using an $SPR_{50\%}$ reference harvest rate and with a 95% confidence interval of 1,816.8 - 3,294.7 mt based on estimates of uncertainty. The spawning output equivalent to 40% of the unfished spawning output ($SB_{40\%}$) was 2,891.2 millions of eggs. The recent catches (landings plus discards) have been below the point estimate of potential long-term yields calculated using an $SPR_{50\%}$ reference point and the population has been increasing sharply over the last 15 years.

The predicted spawning output from the base model generally showed a sharp decline during the 1960s followed by less of a decline until 1975 (Figure 70). Since 2001, the spawning output has been rapidly increasing due to small catches, and recently, above average recruitment. The 2017 spawning output relative to unfished equilibrium spawning output is above the target of 40% of unfished spawning output (Figure 72). The fishing intensity, $(1 - SPR)/(1 - SPR_{50\%})$, exceeded the current estimates of the harvest rate limit ($SPR_{50\%}$) throughout the 1960s as seen in Figure 85. Recent exploitation rates on Pacific ocean perch were predicted to be much less than target levels. In recent years, the stock has experienced exploitation rates that have been below the target level while the spawning output level has remained above the target level.

Table 25 shows the full suite of estimated reference points for the base model and Figure 86 shows the equilibrium curve based on a steepness value fixed at 0.72.

4 Harvest Projections and Decision Tables

A ten year projection of the base model with catches equal to the estimated ACL for years 2019-2028 and a catch allocation equal to the percentages for each fleet over the period of 2014-2016 predicts an increase in the spawning output due to large 2008 cohort, with a slight downturn beginning in 2023 (Table 30). Additional projections with the current ACL or the SPR_{50} MSY using the low and high states of nature are provided in Table 31 and show the spawning output remaining well above the management target for either catch level.

5 Regional Management Considerations

The distribution of Pacific ocean perch occur primarily in the US west coast waters of Washington, Oregon, and northern California and is currently managed to a species level with harvest limits set for the stock north of the 40°10' latitude. The population within this area is treated as a single stock due to the lack of biological and genetic data indicating the presence of multiple stocks. Analysis conducted within this assessment did not find support for regional management within the area that Pacific ocean perch occur.

6 Research Needs

There are many areas of research that could be improved to benefit the understanding and assessment of Pacific ocean perch. Below, are issues that are considered of importance.

1. **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for Pacific ocean perch. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life history of Pacific ocean perch may reduce that uncertainty.
2. **Steepness:** The amount of stock resilience, steepness, dictates the rate at which a stock can rebuild from low stock sizes. Improved understanding regarding the steepness of US west coast Pacific ocean perch will reduce our uncertainty regarding current stock status.
3. **Basin-wide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for Pacific ocean perch off of the west coast of the US and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and Alaska to determine if there are similarities with the US west coast observations would help to define the connectivity between Pacific ocean perch north and south of the U.S.-Canada border.

7 Acknowledgments

Many people were instrumental in the successful completion of this assessment and their contribution is greatly appreciated. Jason Cope (NWFSC), Ian Taylor (NWFSC), and Owen Hamel (NWFSC), contributed greatly with discussions about data, modeling, and SS. Allan Hicks (IPHC) provided guidance on model approaches and data exploration. We are grateful to Theresa Tsou (WDFW) and Phillip Wyland (WDFW) who provided research data and the first historical reconstruction of catch for Washington state. Alison Whitman (ODFW), Patrick Mirrick (ODFW), and Ted Calavan (ODFW) provided Oregon composition data, historical catches, corrected PacFIN catches, and quickly uploaded age data that were critical to this assessment. We appreciate Vanessa Tuttle's patience and responsiveness to providing data. John Field (SWFSC) provided historical catch information and Don Pearson (SWFSC) compiled the extensive management changes for Pacific ocean perch which were critical in understanding and modeling fishery behavior. John Wallace provided multiple last minute PacFIN extractions and analyzed historical discard rates for use in the assessment.

We are very grateful to Patrick McDonald and the team of agers at CAP for their hard work reading numerous otoliths and availability to answer questions when needed. Beth

Horness was always eager to help, quick to supply survey extractions, and answered numerous questions we had. Jason Jannot and Kayleigh Sommers assisted with data from the WCGOP and entertained our many questions. We appreciate Melissa Head's effort to produce updated maturity estimates for the assessment. We would like to acknowledge our survey team and their dedication to improving the assessments we do. The assessment was greatly improved through the many discussions within the Population Ecology team in the FRAM division at the NWFSC.

The reviews of the STAR panel (Norman Hall, Kevin Piner, and Yiota Apostolaki) and STAR panel chair (David Sampson), the GMT representative (Lynn Mattes), and the GAP representative (Louie Zimm) are greatly appreciated for their patience, advice, focus, understanding, and comments during and after the STAR panel meeting. John Devore was an indispensable resource of the past, current, and future management as well as a great organizer and representative of the Pacific Fishery Management Council.

8 Tables

Table 1: Landings for each state (all gears combined), the at-sea hake fishery, the foreign fleet, and surveys for the modeled years.

Year	California	Oregon	Washington	At-Sea Hake	Foreign	Survey
1918	0.1	0.0	1.1	0.0	0	0.0
1919	0.0	0.0	0.4	0.0	0	0.0
1920	0.0	0.0	0.3	0.0	0	0.0
1921	0.0	0.0	0.3	0.0	0	0.0
1922	0.0	0.0	0.1	0.0	0	0.0
1923	0.0	0.0	0.2	0.0	0	0.0
1924	0.1	0.0	0.5	0.0	0	0.0
1925	0.1	0.0	0.6	0.0	0	0.0
1926	0.1	0.0	1.0	0.0	0	0.0
1927	0.1	0.0	1.4	0.0	0	0.0
1928	0.1	0.1	1.2	0.0	0	0.0
1929	0.3	0.1	0.7	0.0	0	0.0
1930	0.2	0.1	0.9	0.0	0	0.0
1931	0.4	0.1	0.4	0.0	0	0.0
1932	0.3	0.1	0.4	0.0	0	0.0
1933	0.6	0.1	0.5	0.0	0	0.0
1934	0.4	0.0	2.3	0.0	0	0.0
1935	0.4	0.1	7.7	0.0	0	0.0
1936	0.2	0.2	1.6	0.0	0	0.0
1937	0.5	0.4	2.0	0.0	0	0.0
1938	0.6	0.1	5.1	0.0	0	0.0
1939	0.9	0.4	8.7	0.0	0	0.0
1940	0.9	9.1	12.2	0.0	0	0.0
1941	1.3	14.0	13.6	0.0	0	0.0
1942	0.4	26.6	18.6	0.0	0	0.0
1943	1.0	94.3	453.6	0.0	0	0.0
1944	2.8	164.5	739.3	0.0	0	0.0
1945	6.7	247.1	1887.1	0.0	0	0.0
1946	7.3	193.2	845.9	0.0	0	0.0
1947	2.6	167.2	385.3	0.0	0	0.0
1948	4.2	177.8	491.1	0.0	0	0.0
1949	2.2	472.9	409.5	0.0	0	0.0
1950	1.5	690.1	675.7	0.0	0	0.0
1951	4.3	840.1	735.1	0.0	0	0.0
1952	3.1	2030.5	305.6	0.0	0	0.0
1953	146.4	1223.5	361.6	0.0	0	0.0
1954	123.6	1837.5	538.8	0.0	0	0.0
1955	50.6	1346.4	555.6	0.0	0	0.0
1956	4.1	2563.8	548.2	0.0	0	0.0

Year	California	Oregon	Washington	At-Sea Hake	Foreign	Survey
1957	1.7	2128.1	538.5	0.0	0	0.0
1958	3.1	1564.9	530.4	0.0	0	0.0
1959	1.6	892.6	337.0	0.0	0	0.0
1960	20.9	1358.8	928.1	0.0	0	0.0
1961	1.2	2061.9	1179.8	0.0	0	0.0
1962	0.6	2584.9	1725.2	0.0	0	0.0
1963	33.1	3693.9	2006.0	0.0	0	0.0
1964	47.1	4261.6	1770.7	0.0	0	0.0
1965	36.3	5627.8	1972.1	0.0	0	0.0
1966	5.3	1591.2	1725.5	0.0	15561	0.0
1967	18.1	354.7	1861.0	0.0	12357	0.0
1968	22.3	466.4	2501.2	0.0	6639	0.0
1969	8.4	422.3	1236.0	0.0	469	0.0
1970	8.7	507.4	1293.3	0.0	441	0.0
1971	12.2	290.4	673.6	0.0	902	0.0
1972	11.4	105.3	796.5	0.0	950	0.0
1973	11.9	121.2	713.1	0.0	1773	0.0
1974	15.7	136.7	641.8	0.0	1457	0.0
1975	11.4	181.3	413.9	62.3	496	0.0
1976	17.1	663.7	521.1	31.9	239	0.0
1977	16.7	457.1	752.0	3.8	0	11.9
1978	42.5	498.7	1391.5	15.4	0	0.0
1979	136.7	735.9	581.4	15.1	0	34.5
1980	19.2	948.6	666.2	47.0	0	4.6
1981	10.8	929.7	390.3	15.4	0	0.0
1982	145.9	584.0	273.0	28.3	0	0.0
1983	102.0	1032.7	437.7	10.9	0	4.4
1984	47.6	750.4	815.7	2.3	0	0.9
1985	70.9	789.5	503.2	11.4	0	13.6
1986	52.8	676.5	588.9	19.8	0	1.4
1987	120.9	550.0	399.4	5.4	0	0.0
1988	75.4	749.8	509.8	4.5	0	0.5
1989	29.5	927.8	466.2	4.3	0	4.2
1990	18.3	567.8	427.2	80.9	0	0.0
1991	8.4	853.2	530.1	46.1	0	0.0
1992	15.3	623.4	435.2	373.3	0	4.9
1993	11.0	797.8	464.7	0.9	0	0.2
1994	6.7	626.4	352.0	83.8	0	0.0
1995	9.2	515.0	289.8	46.6	0	2.8
1996	18.4	531.1	236.7	6.3	0	1.2
1997	15.8	439.1	184.9	6.4	0	0.1
1998	21.6	436.7	172.4	22.3	0	3.8
1999	19.8	326.8	145.8	16.5	0	1.4

Year	California	Oregon	Washington	At-Sea Hake	Foreign	Survey
2000	6.8	95.1	33.0	10.1	0	0.6
2001	0.5	193.4	51.8	21.0	0	2.8
2002	0.8	107.0	39.5	3.9	0	0.3
2003	0.2	94.6	30.2	6.3	0	3.6
2004	2.1	97.7	22.3	1.1	0	2.5
2005	0.1	51.2	10.4	1.7	0	1.8
2006	0.2	52.2	15.8	3.1	0	1.2
2007	0.2	83.7	45.1	4.0	0	0.6
2008	0.4	58.6	16.6	15.9	0	0.8
2009	0.9	58.7	33.2	1.6	0	2.7
2010	0.1	58.0	22.3	16.9	0	1.7
2011	0.1	30.3	19.7	9.2	0	1.9
2012	0.2	30.4	21.8	4.5	0	1.6
2013	0.1	34.9	14.8	5.4	0	1.7
2014	0.2	33.9	15.8	3.9	0	0.6
2015	0.1	38.1	11.4	8.7	0	1.6
2016	0.2	40.8	13.1	10.3	0	3.1

Table 2: West Coast history of regulations.

Date	Area	Regulation
11/10/1983	Columbia	Closed Columbia area to Pacific ocean perch fishing until the end of the year, as 950 mt OY for this species has been reached;
11/10/1983	Vancouver	retained 5,000-pound trip limit or 10% of total trip weight on landings of Pacific ocean perch in the Vancouver area.
1/1/1984	ALL	Continued 5,000-pound trip limit or 10% of total trip weight on Pacific ocean perch as specified in FMP. Fishery to close when area OYs are reached (see action effective November 10, 1983 above).
8/1/1984	Vancouver Columbia	Reduced trip limit for Pacific ocean perch in the Vancouver and Columbia areas to 20% by weight of all fish on board, not to exceed 5,000 pounds per vessel per trip.
8/16/1984	Columbia	Commercial fishing for Pacific ocean perch in the Columbia area closed for remainder of the year.
1/10/1985	Vancouver Columbia	Established Vancouver and Columbia areas Pacific ocean perch trip limit of 20% by weight of all fish on board (no 5,000-pound limit as specified in last half of 1984).
4/28/1985	Vancouver Columbia	Reduced the Vancouver and Columbia areas Pacific ocean perch trip limit to 5,000 pounds or 20% by weight of all fish on board, whichever is less.
4/28/1985	ALL	Landings of Pacific ocean perch less than 1,000 pounds will be unrestricted. The fishery for this species will close when the OY in each area is reached.
6/10/1985	ALL	Landings of Pacific ocean perch up to 1,000 pounds per trip will be unrestricted regardless of the percentage of these fish on board.
1/1/1986	Cape Blanco North	Established the Pacific ocean perch trip limit north of Cape Blanco (4250) at 20% (by weight) of all fish on board or 10,000 pounds whichever is less;
1/1/1986	ALL	landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board; Vancouver area OY = 600 mt; Columbia area OY = 950 mt.
12/1/1986	Vancouver	OY quota for Pacific ocean perch reached in the Vancouver area; fishery closed until January 1, 1987.
1/1/1987	ALL	Established coastwide Pacific ocean perch limit at 20% of all legal fish on board or 5,000 pounds whichever is less (in round weight); landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board; Vancouver area OY = 500 mt; Columbia area OY = 800 mt.
1/1/1988	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 5,000 pounds, whichever is less; landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board;
1/1/1989	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 5,000 pounds whichever is less;
1/1/1989	ALL	landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board (Vancouver area OY = 500 mt; Columbia area OY = 800 mt).
7/26/1989	ALL	Reduced the coastwide trip limit for Pacific ocean perch to 2,000 pounds or 20% of all fish on board, whichever is less, with no trip frequency restriction.
12/13/1989	Columbia	Closed the Pacific ocean perch fishery in the Columbia area because 1,040 mt OY reached.
1/1/1990	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board. (Vancouver area OY = 500 mt; Columbia area OY = 1,040 mt).
1/1/1991	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,000 mt).
1/1/1992	ALL	For Pacific ocean perch, established the coastwide trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,550 mt).

Date	Area	Regulation
1/1/1993	Cape Mendocino Coos Bay	For Pacific ocean perch, continued the coastwide trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,550 mt).
1/1/1994	ALL	Pacific Ocean Perch trip limit of 3,000 pounds or 20% of all fish on board, whichever is less, in landings of Pacific ocean perch above 1,000 pounds.
1/1/1995	ALL	For Pacific Ocean Perch, established a cumulative trip limit of 6,000 pounds per month
1/1/1996	ALL	Pacific Ocean Perch cumulative trip limit of 10,000 pounds per two-month period.
7/1/1996	4030 North	Reduced the cumulative 2-month limit for Pacific ocean perch to 8,000 pounds, and established the cumulative 2-month limit for Dover sole north of Cape Mendocino at 38,000 pounds
1/1/1997	ALL	Pacific Ocean Perch limited entry fishery cumulative trip limit of 8,000 pounds per two-month period
1/1/1998	ALL	Pacific Ocean Perch: limited entry fishery Cumulative trip limit of 8,000 pounds per two-month period.
7/1/1998	ALL	Open Access Rockfish: removed overall rockfish monthly limit and replaced it with limits for component rockfish species: for Sebastes complex, monthly cumulative limit is 33,000 pounds, for widow rockfish, monthly cumulative trip limit is 3,000 pounds, for Pacific Ocean Perch, monthly cumulative trip limit is 4,000 pounds.
1/1/1999	ALL	for the limited entry fishery A new three phase cumulative limit period system is introduced for 1999. Phase 1 is a single cumulative limit period that is 3months long, from January 1 - March 31. Phase 2 has 3 separate 2 month cumulative limit periods of April 1 - May 31, June 1 - July 31, and August 1 - September 30. Phase 3 has 3 separate 1 month cumulative limit periods of October 1-31, November 1-30, and December 1-31. For all species except Pacific ocean perch and Bocaccio, there will be no monthly limit within the cumulative landings limit periods. An option to apply cumulative trip limits lagged by 2 weeks (from the 16th to the 15th) was made available to limited entry trawl vessels when their permits were renewed for 1999. Vessels that are authorized to operate in this "B" platoon may take and retain, but may not land, groundfish during January 1-15, 1999.
1/1/1999	ALL	for the limited entry fishery Pacific Ocean Perch: cumulative limit, Phase 1: 4,000 pounds per month; Phase 2: 4,000 pounds per month; Phase 3: 4,000pounds per month.
1/1/1999	ALL	for open access gear: Pacific Ocean Perch: coastwide, 100 pounds per month.
1/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 500 lbs per month
1/1/2000	ALL	Pacific Ocean Perch, Open Access gear except exempted trawl, 100 lbs per month
1/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 500 lbs per month
5/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 2500 lbs per 2 months
5/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 2500 lbs per month
11/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 500 lbs per month
11/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 500 lbs per month
1/1/2001	3600 North	Pacific Ocean Perch, open access, 100 lbs per month
1/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 1500 lbs per month
1/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 1500 lbs per month
5/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 2500 lbs per month
5/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 2500 lbs per month
10/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 1500 lbs per month
11/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 1500 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, open access, 100 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 2000 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 2000 lbs per month
4/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 4000 lbs per month
5/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 4000 lbs per month
11/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 2000 lbs per month
11/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 2000 lbs per month
1/1/2003	3800 South	minor slope rockfish south including pacific ocean perch, open access gear, 10000 lbs per 2 months

Date	Area	Regulation
1/1/2003	3800 South	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 30000 lbs per 2 months
1/1/2003	3800 South	Minor slope rockfish south including Pacific ocean perch , limited entry trawl, 30000 lbs per 2 months
1/1/2003	3800 4010	minor slope rockfish south including pacific ocean perch, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch , limited entry trawl, 1800 lbs per 2 months
1/1/2003	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2003	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2003	4010 North	Pacific Ocean Perch, Limited entry trawl gear, 3000 lbs per 2 months
3/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, no more than 25% of the weight of sablefish landed per trip
11/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2004	3800 South	Minor slope rockfish south including Pacific ocean perch, open access gear, 10000 lbs per 2 months
1/1/2004	3800 South	minor slope rockfish south inclding pacific ocean perch, limited entry fixed gear, 40000 lbs per 2 months
1/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 40000 lbs per 2 months
1/1/2004	3800 4010	Minor slope rockfish south including Pacific ocean perch, open access gear, per trip no more than 25% of the weight of sablefish landed
1/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 7000 lbs per 2 months
1/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 7000 lbs per 2 months
1/1/2004	4010 North	pacific ocean perch, open access gear, 100 lbs per month
1/1/2004	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2004	4010 North	pacific ocean perch, limited entry trawl, 3000 lbs per 2 months
5/1/2004	3800 South	minor slope rockfish south inclding pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
5/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
5/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
5/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
11/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
11/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
11/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 10000 lbs per 2 months
11/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 10000 lbs per 2 months
1/1/2005	3800 South	minor slope rockfish south including darkblotched and pacific ocean perch, open access gear, 10000 lbs per 2 months
1/1/2005	3800 South	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, closed
1/1/2005	3800 4010	minor slope rockfish south including darkblotched and pacific ocean perch, open access gear, per trip no more than 25% of weight of sablefish onboard
1/1/2005	3800 4010	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, 4000 lbs per 2 months
1/1/2005	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2005	4010 North	pacific ocean perch, limited entry trawl gear, 3000 lbs per 2 months
1/1/2005	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2005	4010 South	minor slope rockfish south including darkblotched and pacific ocean perch, limited entry fixed gear, 40000 lbs per 2 months
5/1/2005	3800 4010	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, 8000 lbs per 2 months

Date	Area	Regulation
1/1/2008	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2008	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
1/1/2009	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2009	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2009	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2009	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2009	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 55000 lbs per 2 months
1/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2009	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
7/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 10000 lbs per 2 months
11/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2010	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2010	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2010	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2010	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2010	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2010	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 55000 lbs per 2 months
1/1/2010	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2010	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
1/1/2011	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2011	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2011	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2011	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2011	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2011	ALL	Pacific Ocean Perch managed in part by IFQ
1/1/2012	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2012	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2012	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2012	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2012	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2013	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2013	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2013	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months no more than 1375 lbs may be blackgill
1/1/2013	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months no more than 475 lbs of which may be blackgill rockfish
1/1/2014	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2014	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1375 lbs may be blackgill rockfish

Date	Area	Regulation
1/1/2014	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2014	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
1/1/2015	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2015	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1375 lbs may be blackgill rockfish
1/1/2015	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2015	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
7/1/2015	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1600 lbs may be blackgill rockfish
7/1/2015	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 550 lbs may be blackgill rockfish
1/1/2016	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2016	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2016	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
7/1/2016	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 550 lbs may be blackgill rockfish

Table 3: Recent trend in estimated total catch relative to management guidelines.

Year	OFL (mt; ABC prior to 2011)	ABC (mt)	ACL (mt; OY prior to 2011)	Total landings (mt)	Estimated total catch (mt)
2007	900		150	134	159
2008	911		150	92	134
2009	1,160		189	97	193
2010	1,173		200	99	182
2011	1,026	981	180	61	62
2012	1,007	962	183	59	60
2013	844	807	150	57	58
2014	838	801	153	54	56
2015	842	805	158	60	61
2016	850	813	164	68	68

Table 4: Description of the data used to create the indices, the modeling platform used to generate the estimates, and the model configuration.

	Pacific ocean perch	AFSC Slope	NWFSC Slope	NWFS Shelf-Slope
Depth	155-500	183-549	183-549	55-549
Latitude	44-48.5	42-49	42-49	42-49
Model	VAST	VAST	Bayesian Delta GLMM	VAST
Error Structure	Lognormal	Lognormal	Gamma	Lognormal
Knots	1000	1000	-	1000
Spatial	Y	Y	N	Y
Temporal	Y	Y	N	Y
Vessel-Year	N	N	Y	Y

Table 5: Summary of the fishery-independant biomass/abundance time-series used in the stock assessment. The standard error includes the input annual standard error and model estimated added variance.

Year	POP		AFSC Slope		NWFSC Slope		NWFSC Shelf-Slope	
	Obs	SE	Obs	SE	Obs	SE	Obs	SE
1979	56461	0.27	-	-	-	-	-	-
1985	34645	0.29	-	-	-	-	-	-
1996	-	-	7621	0.51	-	-	-	-
1997	-	-	3807	0.51	-	-	-	-
1999	-	-	4694	0.50	3643	0.63	-	-
2000	-	-	4243	0.53	4120	0.58	-	-
2001	-	-	4187	0.49	2325	0.59	-	-
2002	-	-	-	-	1903	0.60	-	-
2003	-	-	-	-	-	-	9646	0.36
2004	-	-	-	-	-	-	5284	0.39
2005	-	-	-	-	-	-	7528	0.39
2006	-	-	-	-	-	-	6010	0.41
2007	-	-	-	-	-	-	6268	0.36
2008	-	-	-	-	-	-	3867	0.39
2009	-	-	-	-	-	-	2745	0.36
2010	-	-	-	-	-	-	5404	0.34
2011	-	-	-	-	-	-	7533	0.34
2012	-	-	-	-	-	-	9289	0.34
2013	-	-	-	-	-	-	8093	0.34
2014	-	-	-	-	-	-	4914	0.34
2015	-	-	-	-	-	-	5752	0.31
2016	-	-	-	-	-	-	11770	0.36

Table 6: Summary of the design based estimates of fishery-independent biomass/abundance time-series.

Year	POP		AFSC Slope		NWFSC Slope		NWFSC Shelf-Slope	
	Obs	SE	Obs	SE	Obs	SE	Obs	SE
1979	34135	0.25	-	-	-	-	-	-
1985	16675	0.18	-	-	-	-	-	-
1996	-	-	6472	0.29	-	-	-	-
1997	-	-	2965	0.43	-	-	-	-
1999	-	-	19063	0.48	6472	0.45	-	-
2000	-	-	4438	0.50	2965	0.48	-	-
2001	-	-	14570	0.69	19063	0.40	-	-
2002	-	-	-	-	4438	0.45	-	-
2003	-	-	-	-	-	-	21055	0.36
2004	-	-	-	-	-	-	4623	0.55
2005	-	-	-	-	-	-	9674	0.60
2006	-	-	-	-	-	-	9609	0.53
2007	-	-	-	-	-	-	3769	0.57
2008	-	-	-	-	-	-	5723	0.59
2009	-	-	-	-	-	-	14790	0.78
2010	-	-	-	-	-	-	11133	0.47
2011	-	-	-	-	-	-	6186	0.46
2012	-	-	-	-	-	-	10208	0.46
2013	-	-	-	-	-	-	14306	0.58
2014	-	-	-	-	-	-	4040	0.29
2015	-	-	-	-	-	-	9766	0.56
2016	-	-	-	-	-	-	19859	0.52

Table 7: Summary of NWFSC shelf-slope survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2003	46	1426	111
2004	34	565	82
2005	38	526	92
2006	33	659	80
2007	50	628	121
2008	39	539	94
2009	46	471	111
2010	53	907	128
2011	53	921	128
2012	50	1175	121
2013	45	732	109
2014	52	991	126
2015	69	1165	167
2016	50	1150	121

Table 8: Summary of NWFSC shelf-slope survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2003	45	432	109
2004	34	219	82
2005	38	257	92
2006	33	254	80
2007	50	439	121
2008	39	328	94
2009	45	331	109
2010	53	579	128
2011	53	674	128
2012	49	699	119
2013	44	553	106
2014	52	626	126
2015	68	840	165
2016	44	703	106

Table 9: Summary of NWFSC slope survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2001	18	173	43
2002	24	368	58

Table 10: Summary of NWFSC slope survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2001	17	172	41
2002	24	359	58

Table 11: Summary of AFSC slope survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1996	48	1396	116
1997	21	347	51
1999	21	562	51
2000	19	353	46
2001	23	390	55

Table 12: Summary of Pacific ocean perch survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1979	125	2375	303
1985	126	2558	306

Table 13: Summary of Pacific ocean perch survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1985	29	1635	70

Table 14: Summary of discard rates used in the model by each data source.

Year	Source	Discard	Standard Error
1985	Pikitch	0.027	0.068
1986	Pikitch	0.024	0.063
1987	Pikitch	0.039	0.083
1992	Management Restrictions	0.100	0.300
2002	WCGOP	0.150	0.164
2003	WCGOP	0.183	0.268
2004	WCGOP	0.203	0.206
2005	WCGOP	0.175	0.346
2006	WCGOP	0.148	0.243
2007	WCGOP	0.171	0.261
2008	WCGOP	0.362	0.172
2009	WCGOP	0.504	0.153
2010	WCGOP	0.487	0.195
2011	WCGOP	0.015	0.053
2012	WCGOP	0.028	0.054
2013	WCGOP	0.027	0.054
2014	WCGOP	0.035	0.050
2015	WCGOP	0.010	0.053

Table 15: Summary of commercial fishery length samples used in the stock assessment (continued on next page).

Year	Trips	Fish	Sample Size
1966	1	238	7
1967	5	1020	35
1968	3	912	21
1969	4	1213	28
1970	13	1830	92
1971	22	4698	155
1972	23	4561	162
1973	17	4134	120
1974	20	4806	141
1975	19	3637	134
1976	21	3677	148
1977	32	4846	226
1978	52	7715	367
1979	34	3414	240
1980	55	5425	388
1981	40	3921	282
1982	48	4824	339
1983	39	3944	275
1984	31	3102	219
1985	45	4508	318
1986	40	4002	282
1987	43	3053	304
1988	9	601	64
1989	16	798	113
1990	12	599	85
1991	8	216	38
1994	43	2608	304
1995	49	3161	346
1996	64	3085	452
1997	76	3570	537
1998	56	3450	395
1999	58	2812	409
2000	49	2004	326
2001	59	1696	293
2002	50	1666	280

Year	Trips	Fish	Sample Size
2003	67	1661	296
2004	53	1202	219
2005	51	1277	227
2006	59	1486	264
2007	81	2248	391
2008	101	3058	523
2009	107	3207	550
2010	134	2872	530
2011	100	1943	368
2012	97	1873	355
2013	117	2167	416
2014	140	2850	533
2015	110	2504	456
2016	131	2158	429

Table 16: Summary of commercial fishery age samples used in the stock assessment.

Year	Trips	Fish	Sample Size
1981	20	1901	141
1982	40	2776	282
1983	33	3317	233
1984	27	2625	191
1985	21	2096	148
1986	17	1693	120
1987	24	1193	169
1988	4	199	28
1994	8	238	41
1999	18	863	127
2000	14	677	99
2001	40	1349	226
2002	38	1414	233
2003	40	1309	221
2004	30	854	148
2005	37	1018	177
2006	49	1258	223
2007	63	1825	315
2008	44	1129	200
2009	75	1548	289
2010	54	1264	228
2011	85	1230	255
2012	7	331	49
2013	10	265	47
2014	91	587	172
2015	78	513	149
2016	21	254	56

Table 17: Summary of at-sea hake fishery length samples used in the stock assessment.

Year	Trips	Fish	Sample Size
2003	153	805	263
2004	128	329	172
2005	221	734	321
2006	210	751	312
2007	319	1119	470
2008	26	2491	162
2009	12	366	63
2010	22	1794	155
2011	36	1748	226
2012	26	881	148
2013	26	834	140
2014	31	532	103
2015	23	925	150
2016	35	1947	240

Table 18: Summary of at-sea hake fishery age samples used in the stock assessment.

Year	Trips	Fish	Sample Size
2003	142	378	194
2006	198	410	255
2007	297	620	383
2014	22	101	36

Table 19: Estimated ageing error from the CAPS lab used in the assessment model

True Age (yr)	SD of Observed Age (yr)	True Age (yr)	SD of Observed Age (yr)
0.5	0.156	31.5	2.772
1.5	0.156	32.5	2.854
2.5	0.249	33.5	2.935
3.5	0.341	34.5	3.016
4.5	0.433	35.5	3.097
5.5	0.524	36.5	3.177
6.5	0.615	37.5	3.257
7.5	0.706	38.5	3.337
8.5	0.796	39.5	3.416
9.5	0.886	40.5	3.495
10.5	0.976	41.5	3.574
11.5	1.065	42.5	3.652
12.5	1.154	43.5	3.73
13.5	1.242	44.5	3.808
14.5	1.33	45.5	3.885
15.5	1.418	46.5	3.962
16.5	1.505	47.5	4.039
17.5	1.592	48.5	4.115
18.5	1.679	49.5	4.191
19.5	1.765	50.5	4.267
20.5	1.851	51.5	4.342
21.5	1.937	52.5	4.417
22.5	2.022	53.5	4.492
23.5	2.107	54.5	4.566
24.5	2.191	55.5	4.641
25.5	2.275	56.5	4.714
26.5	2.359	57.5	4.788
27.5	2.442	58.5	4.861
28.5	2.525	59.5	4.934
29.5	2.608	60.5	5.007
30.5	2.69		

Table 20: Specifications of the base model for Pacific ocean perch.

Model Specification	Base Model
Starting year	1918
<u>Population characteristics</u>	
Maximum age	60
Gender	2
Population lengths	5-50 cm by 1 cm bins
Summary biomass (mt)	Age 3+
<u>Data characteristics</u>	
Data lengths	11-47 cm by 1 cm bins
Data ages	1-40 ages
Minimum age for growth calculations	3
Maximum age for growth calculations	20
First mature age	0
Starting year of estimated recruitment	1940
<u>Fishery characteristics</u>	
Fishing mortality method	Continuous
Maximum F	3
Catchability	Analytical estimate
Fishery selectivity	Double Normal
At-Sea Hake selectivity	Double Normal
POP survey selectivity	Logistic
Triennial survey	Double Normal
AFSC slope survey	Double Normal
NWFSC slope survey	Double Normal
NWFSC shelf/slope survey	Double Normal
<u>Fishery time blocks</u>	
Fishery selectivity	None
Fishery retention	1918-1991, 1992-2001, 2002-2007, 2008, 2009-2010, 2011-2016

Table 21: Data weights applied when using Francis data weighting in the base model.

Fleet	Lengths	Ages
Fishery	0.096	0.217
At-sea hake	0.104	0.031
Pacific ocean perch survey	1.000	1
AFSC slope survey	0.077	-
NWFSC slope survey	0.565	0.304
NWFSC shelf-slope survey	0.031	0.363

Table 22: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
NatM_p_1_Fem_GP_1	0.054	-5	(0.02, 0.1)			Log_Norm (-2.92, 0.44)
L_at_Amin_Fem_GP_1	20.7718	3	(15, 25)	OK	0.14	None
L_at_Amax_Fem_GP_1	41.6366	2	(35, 45)	OK	0.15	None
VonBert_K_Fem_GP_1	0.165916	3	(0.1, 0.4)	OK	0.00	None
SD_young_Fem_GP_1	1.34174	5	(0.03, 5)	OK	0.06	None
SD_old_Fem_GP_1	2.56502	5	(0.03, 5)	OK	0.12	None
Wtlen_1_Fem	1.003e-05	-99	(0, 3)			None
Wtlen_2_Fem	3.1026	-99	(2, 4)			None
Mat50%_Fem	32.1	-99	(20, 40)			None
Mat_slope_Fem	-1	-99	(-2, 4)			None
Eggs_scalar_Fem	8.66e-10	-99	(0, 6)			None
Eggs_exp_len_Fem	4.9767	-99	(-3, 5)			None
NatM_p_1_Mal_GP_1	0.054	-5	(0, 0.3)			Normal (0.05, 0.1)
L_at_Amin_Mal_GP_1	20.7718	-2	(6, 68)			None
L_at_Amax_Mal_GP_1	38.929	2	(13, 122)	OK	0.00	None
VonBert_K_Mal_GP_1	0.197	3	(0.04, 1.09)	OK	0.04	None
SD_young_Mal_GP_1	1.34174	-5	(0, 742.07)			None
SD_old_Mal_GP_1	2.33	5	(0, 742.07)	OK	0.06	None
Wtlen_1_Mal	9.881e-06	-99	(0, 3)			None
Wtlen_2_Mal	3.1039	-99	(2, 4)			None
CohortGrowDev	1	-99	(0, 2)			None
FracFemale_GP_1	0.5	-99	(0.01, 0.99)			None
SR_LN(R0)	9.44525	1	(5, 20)	OK	0.15	None
SR_BH_steep	0.72	-2	(0.2, 1)			Full_Beta (0.72, 0.15)
SR_sigmaR	0.7	-6	(0.5, 1.2)			None
SR_regime	0	-99	(-5, 5)			None

Continued on next page

Table 22: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
SR_autocorr	0	-99	(0, 2)			None
Early_InitAge_18	0.00152404	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_17	0.00160538	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_16	0.00169047	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_15	0.00177936	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_14	0.00187211	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_13	0.00196872	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_12	0.00206916	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_11	0.00217339	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_10	0.00228098	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_9	0.00239223	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_8	0.00250635	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_7	0.00262303	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_6	0.00274268	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_5	0.00286644	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_4	0.00299538	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_3	0.00312957	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_2	0.00326871	3	(-6, 6)	act	0.70	dev (NA, NA)
Early_InitAge_1	0.00341298	3	(-6, 6)	act	0.70	dev (NA, NA)
LnQ_base_POP(4)	-0.409011	-1	(-15, 15)			None
LnQ_base_AFSCSlope(6)	-2.9448	-1	(-15, 15)			None
LnQ_base_NWFSCSlope(7)	-3.32004	-1	(-15, 15)			None
LnQ_base_NWFSCcombo(8)	-3.00892	-1	(-15, 15)			None
Q_extraSD_NWFSCcombo(8)	0.0180677	2	(0, 0.5)	OK	0.07	None
SizeSelP1_Fishery(1)	37.9605	1	(20, 45)	OK	0.15	None
SizeSelP2_Fishery(1)	-5	-2	(-6, 4)			None

Continued on next page

Table 22: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
SizeSel_P3_Fishery(1)	3.75994	3	(-1, 9)	OK	0.13	None
SizeSel_P4_Fishery(1)	-1.65	-3	(-9, 9)			None
SizeSel_P5_Fishery(1)	-3.3326	4	(-5, 9)	OK	0.24	None
SizeSel_P6_Fishery(1)	-0.143674	2	(-5, 9)	OK	0.28	None
Retain_P1_Fishery(1)	28.2406	1	(15, 45)	OK	0.36	None
Retain_P2_Fishery(1)	1.06649	1	(0.1, 10)	OK	0.13	None
Retain_P3_Fishery(1)	7.00664	1	(-10, 10)	OK	1.63	None
Retain_P4_Fishery(1)	0	-3	(0, 0)			None
SizeSel_P1_ASHOP(2)	49.4953	1	(20, 49.5)	HI	0.15	None
SizeSel_P2_ASHOP(2)	-5	-2	(-6, 4)			None
SizeSel_P3_ASHOP(2)	5.16735	3	(-1, 9)	OK	0.18	None
SizeSel_P4_ASHOP(2)	1	-3	(-1, 9)			None
SizeSel_P5_ASHOP(2)	-4.35	-4	(-9, 9)			None
SizeSel_P6_ASHOP(2)	999	-2	(-5, 999)			None
SizeSel_P1_POP(4)	25.2002	1	(20, 70)	OK	2.31	None
SizeSel_P2_POP(4)	11.6319	3	(0.001, 50)	OK	4.10	None
SizeSel_P1_AFSCSlope(6)	21.5277	1	(20, 45)	OK	6.01	None
SizeSel_P2_AFSCSlope(6)	-5	-2	(-6, 4)			None
SizeSel_P3_AFSCSlope(6)	1.15414	3	(-1, 9)	OK	6.49	None
SizeSel_P4_AFSCSlope(6)	1	-3	(-1, 9)			None
SizeSel_P5_AFSCSlope(6)	-9	-4	(-9, 9)			None
SizeSel_P6_AFSCSlope(6)	999	-2	(-5, 999)			None
SizeSel_P1_NWFSCSlope(7)	35.9362	1	(20, 45)	OK	2.35	None
SizeSel_P2_NWFSCSlope(7)	-5	-2	(-6, 4)			None
SizeSel_P3_NWFSCSlope(7)	1.84194	3	(-1, 9)	OK	1.93	None
SizeSel_P4_NWFSCSlope(7)	1	-3	(-1, 9)			None

Continued on next page

Table 22: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
SizeSel_P5_NWFSCSlope(7)	-9	-4	(-9, 9)			None
SizeSel_P6_NWFSCSlope(7)	999	-2	(-5, 999)			None
SizeSel_P1_NWFSCcombo(8)	21.1415	1	(18, 49.5)	OK	4.10	None
SizeSel_P2_NWFSCcombo(8)	-5	-2	(-6, 4)			None
SizeSel_P3_NWFSCcombo(8)	3.04272	3	(-1, 9)	OK	2.22	None
SizeSel_P4_NWFSCcombo(8)	1	-3	(-1, 9)			None
SizeSel_P5_NWFSCcombo(8)	-9	-4	(-9, 9)			None
SizeSel_P6_NWFSCcombo(8)	999	-2	(-5, 999)			None
SizeSel_P6_Fishery(1)_BLK4repl_1918	1.31282	4	(-5, 9)	OK	0.63	None
Retain_P2_Fishery(1)_BLK2add_1918	1.23704	4	(0.1, 10)	OK	0.14	None
Retain_P3_Fishery(1)_BLK1repl_1918	9.54767	4	(-10, 10)	OK	11.85	None
Retain_P3_Fishery(1)_BLK1repl_1992	2.5765	4	(-10, 10)	OK	0.47	None
Retain_P3_Fishery(1)_BLK1repl_2002	1.90938	4	(-10, 10)	OK	0.14	None
Retain_P3_Fishery(1)_BLK1repl_2008	0.686957	4	(-10, 10)	OK	0.29	None
Retain_P3_Fishery(1)_BLK1repl_2009	0.0318378	4	(-10, 10)	OK	0.24	None

Table 23: Results from 50 jitters from the base model.

Status	Base.Model
Returned to base case	45
Found local minimum	5
Found better solution	0
Total	50

Table 24: Likelihood components from the base model

Likelihood Component	Value
Total	1638.71
Survey	0
Discard	-13.39
Length-frequency data	-33.83
Age-frequency data	142.62
Recruitment	1531.69
Forecast Recruitment	11.47
Parameter Priors	0

Table 25: Summary of reference points and management quantities for the base case.

Quantity	Estimate	95% Confidence Interval
Unfished spawning output (million eggs)	7228	5126.5 - 9329.4
Unfished age 3+ biomass (mt)	154084	109381.3 - 198786.7
Unfished recruitment (R0, thousands)	12647.9	9483.8 - 16867.7
Spawning output(2017 million eggs)	6966	3701.9 - 10230.1
Depletion (2017)	0.964	0.771 - 1.2
Reference points based on $SB_{40\%}$		
Proxy spawning output ($B_{40\%}$)	2891.2	2050.6 - 3731.8
SPR resulting in $B_{40\%}$ ($SPR_{B40\%}$)	0.458	0.458 - 0.458
Exploitation rate resulting in $B_{40\%}$	0.038	0.038 - 0.039
Yield with $SPR_{B40\%}$ at $B_{40\%}$ (mt)	2680.8	1906.1 - 3455.6
Reference points based on SPR proxy for MSY		
Spawning output	3224.8	2287.2 - 4162.4
SPR_{proxy}	0.5	
Exploitation rate corresponding to SPR_{proxy}	0.033	0.033 - 0.034
Yield with SPR_{proxy} at SB_{SPR} (mt)	2555.8	1816.8 - 3294.7
Reference points based on estimated MSY values		
Spawning output at MSY (SB_{MSY})	1942.8	1374.4 - 2511.1
SPR_{MSY}	0.34	0.338 - 0.342
Exploitation rate at MSY	0.057	0.056 - 0.058
MSY (mt)	2860.7	2034.9 - 3686.5

Table 26: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1918	154,227	7,232	153,446	1.00	12,694	0	0	0
1919	154,239	7,233	153,458	1.00	12,696	1	0	0
1920	154,253	7,234	153,471	1.00	12,698	0	0	0
1921	154,267	7,234	153,485	1.00	12,700	0	0	0
1922	154,281	7,235	153,499	1.00	12,702	0	0	0
1923	154,296	7,235	153,514	1.00	12,705	0	0	0
1924	154,312	7,236	153,530	1.00	12,707	0	0	0
1925	154,327	7,237	153,545	1.00	12,709	1	0	0
1926	154,343	7,237	153,561	1.00	12,711	1	0	0
1927	154,359	7,238	153,577	1.00	12,714	1	0	0
1928	154,375	7,239	153,593	1.00	12,716	1	0	0
1929	154,392	7,239	153,609	1.00	12,718	1	0	0
1930	154,410	7,240	153,627	1.00	12,720	1	0	0
1931	154,427	7,241	153,644	1.00	12,723	1	0	0
1932	154,446	7,242	153,663	1.00	12,726	1	0	0
1933	154,465	7,242	153,682	1.00	12,730	1	0	0
1934	154,484	7,243	153,700	1.00	12,735	1	0	0
1935	154,502	7,244	153,718	1.00	12,744	3	0	0
1936	154,516	7,245	153,731	1.00	12,757	8	0	0

Table 26: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1937	154,536	7,245	153,752	1.00	12,778	2	0	0
1938	154,558	7,246	153,772	1.00	12,808	3	0	0
1939	154,580	7,247	153,793	1.00	12,851	6	0	0
1940	154,602	7,247	153,812	1.00	12,951	10	0.005	0
1941	154,618	7,247	153,825	1.00	13,021	23	0.005	0
1942	154,638	7,247	153,840	1.00	13,104	30	0.005	0
1943	154,657	7,246	153,854	1.00	13,198	47	0.085	0
1944	154,188	7,220	153,379	1.00	13,296	561	0.135	0.004
1945	153,395	7,177	152,581	0.99	13,401	926	0.275	0.006
1946	151,409	7,073	150,589	0.98	13,509	2188	0.155	0.015
1947	150,623	7,026	149,796	0.97	13,651	1070	0.085	0.007
1948	150,404	7,006	149,569	0.97	13,829	567	0.105	0.004
1949	150,126	6,983	149,282	0.97	14,032	688	0.135	0.005
1950	149,703	6,952	148,848	0.96	14,235	905	0.195	0.006
1951	148,872	6,899	148,005	0.95	14,372	1398	0.225	0.009
1952	147,923	6,839	147,044	0.95	14,350	1616	0.305	0.011
1953	146,309	6,745	145,426	0.93	14,073	2393	0.245	0.016
1954	145,427	6,686	144,550	0.92	13,546	1772	0.325	0.012
1955	143,849	6,593	142,993	0.91	12,844	2559	0.27	0.018
1956	142,890	6,533	142,070	0.90	12,029	2000	0.385	0.014
1957	140,760	6,420	139,985	0.89	11,330	3192	0.35	0.023
1958	139,071	6,336	138,344	0.88	10,851	2734	0.295	0.02
1959	137,894	6,286	137,205	0.87	10,696	2150	0.195	0.016
1960	137,488	6,282	136,821	0.87	11,020	1262	0.32	0.009
1961	135,856	6,225	135,191	0.86	11,969	2364	0.405	0.017
1962	133,194	6,119	132,498	0.85	13,024	3321	0.49	0.025
1963	129,443	5,955	128,692	0.82	12,375	4413	0.58	0.034
1964	124,334	5,714	123,548	0.79	10,480	5868	0.61	0.047
1965	119,008	5,450	118,282	0.75	9,205	6223	0.69	0.053
1966	112,202	5,106	111,580	0.71	8,581	7819	0.89	0.07
1967	94,457	4,222	93,904	0.58	8,215	18967	0.88	0.202
1968	81,293	3,562	80,772	0.49	8,810	14652	0.835	0.181
1969	73,235	3,160	72,713	0.44	12,530	9718	0.475	0.134
1970	72,806	3,134	72,187	0.43	19,502	2187	0.49	0.03
1971	72,430	3,110	71,571	0.43	9,504	2307	0.435	0.032
1972	72,762	3,104	71,731	0.43	6,856	1908	0.43	0.027
1973	73,366	3,096	72,824	0.43	6,891	1891	0.525	0.026
1974	73,097	3,048	72,675	0.42	7,039	2646	0.485	0.036
1975	73,059	3,021	72,628	0.42	8,944	2277	0.315	0.031
1976	73,951	3,063	73,492	0.42	6,986	1185	0.37	0.016
1977	74,348	3,116	73,825	0.43	8,867	1514	0.325	0.021
1978	74,829	3,186	74,375	0.44	6,323	1281	0.435	0.017
1979	74,410	3,207	73,903	0.44	7,081	2007	0.365	0.027
1980	74,318	3,228	73,916	0.45	7,375	1546	0.395	0.021
1981	73,872	3,227	73,428	0.45	9,193	1731	0.34	0.024
1982	73,697	3,235	73,202	0.45	14,078	1382	0.28	0.019
1983	73,866	3,254	73,221	0.45	14,290	1058	0.38	0.014
1984	73,648	3,242	72,788	0.45	10,278	1629	0.385	0.022
1985	73,702	3,221	72,889	0.45	9,745	1659	0.35	0.023
1986	74,198	3,205	73,573	0.44	10,153	1425	0.345	0.019
1987	74,825	3,191	74,220	0.44	9,788	1376	0.295	0.019
1988	75,796	3,198	75,171	0.44	12,226	1107	0.345	0.015

Table 26: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1989	76,576	3,212	75,913	0.44	20,778	1382	0.355	0.018
1990	77,426	3,243	76,537	0.45	20,190	1478	0.29	0.019
1991	78,986	3,297	77,744	0.46	9,058	1127	0.35	0.014
1992	80,642	3,333	79,587	0.46	5,847	1483	0.365	0.019
1993	82,343	3,359	81,836	0.46	6,266	1571	0.33	0.019
1994	83,931	3,395	83,549	0.47	12,684	1417	0.285	0.017
1995	85,492	3,456	85,002	0.48	13,082	1181	0.24	0.014
1996	87,090	3,559	86,318	0.49	6,755	956	0.215	0.011
1997	88,674	3,691	87,972	0.51	6,183	883	0.175	0.01
1998	90,180	3,824	89,778	0.53	4,590	717	0.175	0.008
1999	91,320	3,924	90,907	0.54	28,557	725	0.135	0.008
2000	92,515	4,004	91,811	0.55	42,825	563	0.04	0.006
2001	94,507	4,097	92,592	0.57	12,949	161	0.07	0.002
2002	97,480	4,185	95,341	0.58	7,130	297	0.04	0.003
2003	101,307	4,266	100,612	0.59	3,538	179	0.04	0.002
2004	104,853	4,326	104,458	0.60	9,047	158	0.035	0.002
2005	108,025	4,386	107,729	0.61	4,334	149	0.02	0.001
2006	110,732	4,510	110,250	0.62	4,753	78	0.02	0.001
2007	112,881	4,723	112,608	0.65	4,461	86	0.035	0.001
2008	114,580	4,955	113,935	0.69	150,412	159	0.025	0.001
2009	116,934	5,143	114,656	0.71	5,928	134	0.035	0.001
2010	121,621	5,267	114,685	0.73	9,600	193	0.035	0.002
2011	129,996	5,345	129,549	0.74	18,888	182	0.01	0.001
2012	137,389	5,394	136,687	0.75	2,661	62	0.01	0
2013	144,808	5,427	143,823	0.75	37,397	60	0.01	0
2014	151,877	5,566	151,230	0.77	5,672	58	0.01	0
2015	158,254	5,979	156,447	0.83	12,354	56	0.01	0
2016	164,311	6,515	163,853	0.90	12,515	61	0.01	0
2017	169,051	6,966	168,288	0.96	12,602	68	0.04	0
2018	172,729	7,299	171,957	1.01	12,660	-	-	-
2019	175,536	7,559	174,759	1.05	12,702	-	-	-
2020	171,852	7,539	171,073	1.04	12,699	-	-	-
2021	167,707	7,485	166,926	1.03	12,690	-	-	-
2022	163,300	7,382	162,519	1.02	12,674	-	-	-
2023	158,782	7,246	158,002	1.00	12,651	-	-	-
2024	154,266	7,089	153,487	0.98	12,624	-	-	-
2025	149,834	6,921	149,057	0.96	12,594	-	-	-
2026	145,544	6,748	144,768	0.93	12,561	-	-	-
2027	141,428	6,572	140,655	0.91	12,526	-	-	-
2028	137,510	6,398	136,738	0.88	12,490	-	-	-

Table 27: Sensitivity of the base model

Label	Base	Harmonic weights	Steepness = 0.40	Old Maturity	Old Fecundity	2008 Recruitment
Total Likelihood	1638.710	2470.480	1640.330	1638.710	1638.710	1885.200
Survey Likelihood	-13.392	-13.700	-13.653	-13.392	-13.391	-12.734
Discard Likelihood	-33.831	-16.899	-33.491	-33.833	-33.833	57.169
Length Likelihood	142.621	776.472	141.795	142.620	142.627	193.643
Age Likelihood	1531.690	1705.140	1532.410	1531.700	1531.690	1642.890
Recruitment Likelihood	11.469	19.316	11.383	11.472	11.470	4.080
Forecast Recruitment Likelihood	0.000	0.000	0.000	0.000	0.000	0.000
Parameter Priors Likelihood	0.125	0.125	1.870	0.125	0.125	0.125
Parameter Deviation Likelihood	0.000	0.000	0.000	0.000	0.000	0.000
log(R0)	9.445	9.320	9.338	9.445	9.446	9.450
SB Virgin	7227.970	6467.270	6482.640	7092.700	8603.800	7322.850
SB 2017	6966.030	5320.090	3552.640	6969.750	8476.350	5474.610
Depletion 2017	0.964	0.823	0.548	0.983	0.985	0.748
Total Yield - SPR 50	2555.780	2279.200	1029.410	2551.430	2584.350	2585.200
Steepness	0.720	0.720	0.400	0.720	0.720	0.720
Natural Mortality - Female	0.054	0.054	0.054	0.054	0.054	0.054
Length at Amin - Female	20.772	20.641	20.769	20.772	20.772	20.370
Length at Amax - Female	41.637	41.718	41.622	41.637	41.637	41.730
Von Bert. k - Female	0.166	0.169	0.166	0.166	0.166	0.175
SD young - Female	1.342	1.336	1.343	1.342	1.342	1.392
SD old - Female	2.565	2.777	2.569	2.565	2.565	2.513
Natural Mortality - Male	0.054	0.054	0.054	0.054	0.054	0.054
Length at Amin - Male	20.772	20.641	20.769	20.772	20.772	20.370
Length at Amax - Male	38.929	38.916	38.909	38.929	38.929	39.096
Von Bert. k - Male	0.197	0.202	0.197	0.197	0.197	0.204
SD young - Male	1.342	1.336	1.343	1.342	1.342	1.392
SD old - Male	2.330	2.600	2.330	2.330	2.330	2.201

Table 28: Sensitivity of the base model

Label	Base	Include Triennial	Only Triennial	Include CPUE	Canadian Data	WA Research Lengths	OR Special Projects
Total Likelihood	1638.71	1683.18	406.16	1638.71	1731.96	1660.91	1704.90
Survey Likelihood	-13.39	-12.34	12.14	-13.39	-13.58	-13.39	-13.42
Discard Likelihood	-33.83	-33.74	-45.95	-33.83	-33.50	-33.81	-33.06
Length Likelihood	142.62	168.29	260.85	142.62	182.17	163.78	170.39
Age Likelihood	1531.69	1548.48	154.63	1531.69	1584.48	1532.64	1567.61
Recruitment Likelihood	11.47	12.34	24.25	11.47	12.25	11.55	13.24
Forecast Recruitment Likelihood	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Parameter Priors Likelihood	0.13	0.13	0.23	0.13	0.13	0.13	0.13
Parameter Deviation Likelihood	0.00	0.00	0.00	0.00	0.00	0.00	0.00
log(R0)	9.45	9.41	8.91	9.45	9.44	9.42	9.37
SB Virgin	7227.97	6999.21	4664.85	7228.02	7222.61	7017.92	6741.97
SB 2017	6966.03	6657.76	3204.74	6966.14	6661.04	6660.94	6293.13
Depletion 2017	0.96	0.95	0.69	0.96	0.92	0.95	0.93
Total Yield - SPR 50	2555.78	2470.58	1579.24	2555.80	2580.38	2501.20	2375.86
Steepness	0.72	0.72	0.72	0.72	0.72	0.72	0.72
Natural Mortality - Female	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Length at Amin - Female	20.77	20.80	22.31	20.77	20.76	20.75	20.79
Length at Amax - Female	41.64	41.65	42.29	41.64	41.70	41.57	41.67
Von Bert. k - Female	0.17	0.17	0.15	0.17	0.16	0.17	0.16
SD young - Female	1.34	1.35	1.12	1.34	1.35	1.34	1.33
SD old - Female	2.57	2.58	2.60	2.57	2.55	2.56	2.58
Natural Mortality - Male	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Length at Amin - Male	20.77	20.80	22.31	20.77	20.76	20.75	20.79
Length at Amax - Male	38.93	38.93	39.38	38.93	38.96	38.90	38.98
Von Bert. k - Male	0.20	0.20	0.19	0.20	0.20	0.20	0.19
SD young - Male	1.34	1.35	1.12	1.34	1.35	1.34	1.33
SD old - Male	2.33	2.34	2.66	2.33	2.33	2.34	2.39

Table 29: Data weights applied when using harmonic data weighting.

Fleet	Lengths	Ages
Fishery	0.401	0.775
At-sea hake	0.633	0.141
Pacific ocean perch survey	1.000	1
AFSC slope survey	0.696	1
NWFSC slope survey	0.457	-
NWFSC shelf-slope survey	0.544	0.346

Table 30: Projection of potential OFL, spawning biomass, and depletion for the base case model.

Year	OFL (mt)	ACL (mt)	Age 3+ biomass (mt)	Spawning Output	Depletion
2017	5861	281	168288	6966	0.96
2018	6116	281	171957	7299	1.01
2019	6251	5981	174759	7559	1.05
2020	6091	5827	171073	7539	1.04
2021	5894	5639	166926	7485	1.04
2022	5685	5439	162519	7382	1.02
2023	5475	5238	158002	7246	1.00
2024	5270	5042	153487	7089	0.98
2025	5077	4857	149057	6921	0.96
2026	4899	4688	144768	6748	0.93
2027	4738	4533	140655	6572	0.91
2028	4590	4391	136738	6398	0.89

Table 31: Summary of 10-year projections beginning in 2019 for alternate states of nature based on an axis of uncertainty for the base model. Columns range over low, mid, and high states of nature over natural mortality, and rows range over different assumptions of catch levels. An entry of "—" indicates that the stock is driven to very low abundance under the particular scenario.

		States of nature							
		M = 0.045		M = 0.054		M = 0.06			
	Year	Catch	Spawning Output	Depletion (%)	Spawning Output	Depletion (%)	Spawning Output	Depletion (%)	
ABC	2019	5981	5533	86	7559	105	9565	114	
	2020	5827	5486	86	7541	104	9564	114	
	2021	5639	5414	84	7488	104	9517	114	
	2022	5439	5307	83	7388	102	9411	112	
	2023	5238	5177	81	7253	100	9263	111	
	2024	5042	5032	79	7098	98	9089	109	
	2025	4857	4881	76	6932	96	8902	106	
	2026	4688	4726	74	6760	94	8706	104	
	2027	4533	4572	71	6586	91	8508	102	
	2028	4391	4419	69	6413	89	8310	99	
SPR50	2019	2556	5533	86	7559	105	9565	114	
	2020	2556	5640	88	7694	106	9716	116	
	2021	2556	5721	89	7792	108	9819	117	
	2022	2556	5762	90	7835	108	9854	118	
	2023	2556	5770	90	7834	108	9837	117	
	2024	2556	5755	90	7802	108	9782	117	
	2025	2556	5721	89	7747	107	9701	116	
	2026	2556	5673	88	7673	106	9599	115	
	2027	2556	5613	88	7585	105	9482	113	
	2028	2556	5545	87	7487	104	9354	112	

Table 32: Alternative decision table. Summary of 10-year projections beginning in 2019 for alternate states of nature based on an axis of uncertainty for the base model. The range of natural mortality values are based on the 12.5 and 87.5th quantiles of the natural mortality prior. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. The SPR50 catch stream is based on the equilibrium yield applying the SPR50 harvest rate.

		States of nature					
		M = 0.038		M = 0.054		M = 0.077	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
ABC	2019	5981	4214	69	7559	105	41065
	2020	5827	4141	67	7541	104	41501
	2021	5639	4047	66	7488	104	41736
	2022	5439	3927	64	7388	102	41699
	2023	5238	3789	62	7253	100	41471
	2024	5042	3643	59	7098	98	41125
	2025	4857	3493	57	6932	96	40699
	2026	4688	3342	54	6760	94	40219
	2027	4533	3192	52	6586	91	39706
	2028	4391	3045	50	6413	89	39174
SPR50	2019	2556	4214	69	7559	105	41065
	2020	2556	4295	70	7694	106	41651
	2021	2556	4356	71	7792	108	42032
	2022	2556	4387	71	7835	108	42130
	2023	2556	4393	72	7834	108	42024
	2024	2556	4381	71	7802	108	41786
	2025	2556	4354	71	7747	107	41455
	2026	2556	4316	70	7673	106	41057
	2027	2556	4270	70	7585	105	40611
	2028	2556	4216	69	7487	104	40136

9 Figures




Figure 1: Total catches Pacific ocean perch through 2016.




Figure 2: Summary of data sources used in the base model.




Figure 3: Fishery-dependent and fishery-independent indices for Pacific ocean perch.




Figure 4: Q-Q plots for the VAST lognormal distribution for the NWFSC shelf-slope survey.




Figure 5: NWFSC shelf-slope survey length frequency distributions for Pacific ocean perch.




Figure 6: NWFSC shelf-slope survey age frequency distributions for Pacific ocean perch.




Figure 7: Q-Q plots for the VAST lognormal distribution for the NWFSC slope survey.




Figure 8: NWFSC slope survey length frequency distributions for Pacific ocean perch.




Figure 9: NWFSC slope survey age frequency distributions for Pacific ocean perch.




Figure 10: Q-Q plots for the VAST lognormal distribution for the AFSC slope survey.




Figure 11: AFSC slope survey length frequency distributions for Pacific ocean perch.




Figure 12: Q-Q plots for the VAST lognormal distribution for the Pacific ocean perch survey.




Figure 13: Pacific ocean perch survey length frequency distributions for Pacific ocean perch.




Figure 14: Pacific ocean perch survey age frequency distributions for Pacific ocean perch.




Figure 15: Discard length frequency distributions from WCGOP for Pacific ocean perch.




Figure 16: Commercial fishery length frequency distributions for Pacific ocean perch.




Figure 17: Commercial fishery age frequency distributions for Pacific ocean perch.




Figure 18: At-sea hake fishery length frequency distributions for Pacific ocean perch.




Figure 19: At-sea hake fishery age frequency distributions for Pacific ocean perch.




Figure 20: The estimated sex ratio of Pacific ocean perch at length from all biological data sources.




Figure 21: The estimated sex ratio of Pacific ocean perch at age from all biological data sources.




Figure 22: The estimated functional maturity of Pacific ocean perch at length.




Figure 23: Comparison between estimated maturity-at-length used in this assessment and maturity-at-age applied in the 2011 assessment of Pacific ocean perch.




Figure 24: Fecundity at length of Pacific ocean perch in the base model and a comparison of the fecundity in the 2011 assessment.




Figure 25: Weight-at-length for Pacific ocean perch from all data sources.




Figure 26: Estimated weight-at-length for Pacific ocean perch from all data sources.




Figure 27: Estimated length-at-age for Pacific ocean perch from all data sources.




Figure 28: The estimated ageing error used in this assessment compared to the ageing error assumed in the previous assessment for Pacific ocean perch.




Figure 29: Recruitment bias ramp applied in the base model.




Figure 30: Comparison of the catches assumed by this assessment and the previous assessment for Pacific ocean perch.





Figure 31: Comparison of model bridging estimates from Stock Synthesis version 3.30 and 3.24 for Pacific ocean perch for the 2011 assessment.




Figure 32: Each of the data sets used in the current assessment was added to the 2011 model without updating model assumptions. Each data source was included in an additive fashion where the final model “+ Age” is the 2011 model with all data sources updated.




Figure 33: Estimated length-at-age for male and female for Pacific ocean perch with estimated CV.




Figure 34: Comparison between the estimated length-at-age for male and female (solid lines) for Pacific ocean perch with estimated CV to the external estimates based on the data (dashed lines).




Figure 35: Estimated selectivity by length by each fishery and survey for Pacific ocean perch.




Figure 36: Estimated retention by length by the fishery fleet for Pacific ocean perch.




Figure 37: Estimated time-series of recruitment for Pacific ocean perch.




Figure 38: Estimated time-series of recruitment deviations for Pacific ocean perch.




Figure 39: Estimated fits to the CPUE and survey indices for Pacific ocean perch.




Figure 40: Estimated fits to the discard rates for Pacific ocean perch.




Figure 41: Estimated total discards for Pacific ocean perch. Estimated discard contributes less than 3.5 percent of the total mortality across all years from the fishery.




Figure 42: Length compositions aggregated across time by fleet. Labels ‘retained’ and ‘discard’ indicate retained or discarded samples for each fleet. Panels without this designation represent the whole catch.




Figure 43: Pearson residuals, discard, Fishery (max=3.88)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 44: Pearson residuals, retained, Fishery (max=3.21) (plot 4 of 4)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 45: Pearson residuals, whole catch, At-sea hake (max=2.4)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 46: Pearson residuals, whole catch, Pacific ocean perch survey (max=1.81)
Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 47: Pearson residuals, whole catch, AFSC slope survey (max=2.87)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 48: Pearson residuals, whole catch, NWFSC slope survey (max=3.41)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 49: Pearson residuals, whole catch, NWFSC shelf_slope survey (max=2.89)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 50: Mean length for Fishery with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from Fishery: 0.9999 (0.6659_1.7973) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 51: Mean length for At-sea hake with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from At-sea hake: 1.0081 (0.5106_5.2646) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 52: Mean length for Pacific ocean perch survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from Pacific ocean perch survey: 14.0397 (14.0397_Inf) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 53: Mean length for AFSC slope survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from AFSC slope survey: 1.0032 (0.5781_15.7101) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 54: Mean length for NWFSC slope survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from NWFSC slope survey: 0.9996 (0.9996_Inf) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 55: Mean length for NWFSC shelf_slope survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for len data from NWFSC shelf_slope survey: 1.0029 (0.6139_3.8241) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 56: Age compositions aggregated across time by fleet.




Figure 57: Pearson residuals, retained, Fishery (max=5.52) (plot 2 of 2)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 58: Pearson residuals, whole catch, At-sea hake (max=3.87)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 59: Pearson residuals, whole catch, Pacific ocean perch survey (max=2.66)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 60: Pearson residuals, whole catch, NWFSC slope survey (max=2.28)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).

Conditional AAL plot, whole catch, NWFSC shelf-slope survey




Figure 61: Conditional AAL plot, whole catch, NWFSC shelf_slope survey (plot 1 of 5) These plots show mean age and std. dev. in conditional AAL. Left plots are mean AAL by size_class (obs. and pred.) with 90% CIs based on adding 1.64 SE of mean to the data. Right plots in each pair are SE of mean AAL (obs. and pred.) with 90% CIs based on the chi-square distribution.




Figure 62: Conditional AAL plot, whole catch, NWFSC shelf_slope survey (plot 2 of 5)




Figure 63: Conditional AAL plot, whole catch, NWFSC shelf_slope survey (plot 3 of 5)




Figure 64: Conditional AAL plot, whole catch, NWFSC shelf_slope survey (plot 4 of 5)




Figure 65: Conditional AAL plot, whole catch, NWFSC shelf_slope survey (plot 5 of 5)




Figure 66: Mean age for Fishery with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for age data from Fishery: 0.9994 (0.647_2.0865) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 67: Mean age for At-sea hake with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for age data from At-sea hake: 1.0044 (0.6702_2559.3148) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 68: Mean age for NWFSC slope survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for age data from NWFSC slope survey: 0.9993 (0.9993_Inf) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 69: Mean age from conditional data (aggregated across length bins) for NWFSC shelf_slope survey with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for conditional age_at_length data from NWFSC shelf_slope survey: 1.0058 (0.5567_3.4459) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138.




Figure 70: Estimated time-series of spawning output for Pacific ocean perch.




Figure 71: Estimated time-series of total biomass for Pacific ocean perch.




Figure 72: Estimated time-series of relative biomass for Pacific ocean perch.




Figure 73: Estimated recruitment (red circles) and the assumed stock-recruit relationship (black line). The green line shows the effect of the bias correction for the lognormal distribution




Figure 74: Time-series of spawning output for model sensitivities for Pacific ocean perch.




Figure 75: Time-series of relative biomass for model sensitivities for Pacific ocean perch.




Figure 76: Time-series of spawning output for model sensitivities for Pacific ocean perch.




Figure 77: Time-series of relative biomass for model sensitivities for Pacific ocean perch.




Figure 78: Retrospective pattern for spawning output.




Figure 79: Retrospective pattern for estimated recruitment deviations.




Figure 80: Likelihood profile across steepness values.




Figure 81: Trajectories of relative biomass across values of steepness.




Figure 82: Likelihood profile across natural mortality values.




Figure 83: Trajectories of relative biomass across values of natural mortality.




Figure 84: Likelihood profile across R_0 values.




Figure 85: Estimated spawning potential ratio $(1-SPR)/(1-SPR_{50\%})$ for the base-case model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the SPR50% harvest rate. The last year in the time series is 2016.




Figure 86: Equilibrium yield curve for the base case model. Values are based on the 2016 fishery selectivity and with steepness fixed at 0.72.

10 Appendix A. Detailed Fit to Length Composition Data




Figure 87: Length comps, discard, Fishery




Figure 88: Length comps, retained, Fishery (plot 1 of 4)




Figure 89: Length comps, retained, Fishery (plot 1 of 4) (plot 2 of 4)




Figure 90: Length comps, retained, Fishery (plot 1 of 4) (plot 2 of 4) (plot 3 of 4)

Length comps, retained, Fishery




Figure 91: Length comps, retained, Fishery (plot 1 of 4) (plot 2 of 4) (plot 3 of 4) (plot 4 of 4)





Figure 92: Length comps, whole catch, At-sea hake

Length comps, whole catch, Pacific ocean perch survey



Length (cm)

Figure 93: Length comps, whole catch, Pacific ocean perch survey





Figure 94: Length comps, whole catch, AFSC slope survey

Length comps, whole catch, NWFSC slope survey



Length (cm)

Figure 95: Length comps, whole catch, NWFSC slope survey




Figure 96: Length comps, whole catch, NWFSC shelf_slope survey

11 Appendix B. Detailed Fit to Age Composition Data




Figure 97: Age comps, retained, Fishery (plot 1 of 2)




Figure 98: Age comps, retained, Fishery (plot 1 of 2) (plot 2 of 2)

Age comps, whole catch, At-sea hake




Figure 99: Age comps, whole catch, At_sea hake

Age comps, whole catch, Pacific ocean perch survey





Figure 100: Age comps, whole catch, Pacific ocean perch survey

Age comps, whole catch, NWFSC slope survey



Age (yr)

Figure 101: Age comps, whole catch, NWFSC slope survey




Figure 102: Ghost age comps, whole catch, NWFSC shelf_slope survey

12 Appendix C. Description of CPUE and Triennial Data

Data on catch-per-unit-effort (CPUE) in mt/hr from the domestic fishery were combined for the INPFC Vancouver and Columbia areas Gunderson (1977)). Although these data reflect catch rates for the US fleet, the highest catch rates coincided with the beginning of removals by the foreign fleet. This suggests that, barring unaccounted changes in fishing efficiency during this period, the level of abundance was high at that time. The estimated index of abundance is shown in Table 33 and Figure 103.

An index of abundance was estimated based on the data using the VAST delta-GLMM model. The estimated index of abundance is shown in Table 33 and Figure 103. The lognormal distribution with random strata-year had the lowest AIC and was chosen as the final model. The index shows a decline in abundance in the early years of the time-series and abundance remaining flat for the latter years.

Length and age compositions were expanded based upon the survey stratification. The number of tows with length data ranged from 17 in 1986 to 81 in 1998 (Table 34). Ages were read using surface reading methods until 1989 when the break-and-burn method replaced surface reads as the best method to age Pacific ocean perch. Unfortunately, surface reading of Pacific ocean perch otoliths results in significant underestimates of age. Due to this, these otoliths were excluded from analysis. The available ages from the Triennial shelf survey and the number of tows where otoliths were collected are shown in Table 35. The expanded length and age frequencies from this survey are shown in Figures 104 and 105, respectively.

Table 33: Summary of the fishery CPUE and the Triennial shelf survey indices not used in the stock assessment.

Year	Fishery CPUE		Triennial	
	Obs	SE	Obs	SE
1956	0.40	0.40	-	-
1957	0.30	0.40	-	-
1958	0.32	0.40	-	-
1959	0.29	0.40	-	-
1960	0.28	0.40	-	-
1961	0.31	0.40	-	-
1962	0.29	0.40	-	-
1963	0.34	0.40	-	-
1964	0.35	0.40	-	-
1965	0.55	0.40	-	-
1966	0.47	0.40	-	-
1967	0.30	0.40	-	-
1968	0.17	0.40	-	-
1969	0.18	0.40	-	-
1970	0.17	0.40	-	-
1971	0.20	0.40	-	-
1972	0.20	0.40	-	-
1973	0.11	0.40	-	-
1980	-	-	10384	0.65
1983	-	-	8974	0.60
1986	-	-	2977	0.66
1989	-	-	4873	0.66
1992	-	-	3207	0.65
1995	-	-	2724	0.63
1998	-	-	4163	0.64
2001	-	-	1494	0.64
2004	-	-	2922	0.68

Table 34: Summary of Triennial shelf survey length samples.

Year	Tows	Fish	Sample Size
1980	18	1315	43
1983	40	2820	97
1986	17	877	41
1989	42	1851	102
1992	33	1182	80
1995	71	1136	172
1998	81	1482	196
2001	74	669	179
2004	63	1240	153

Table 35: Summary of Triennial shelf survey age samples.

Year	Tows	Fish	Sample Size
1989	15	577	36
1992	10	373	24
1995	12	275	29
1998	28	352	68
2001	43	342	104
2004	57	416	138




Figure 103: Fishery CPUE and Triennial shelf survey indices of abundance for Pacific ocean perch. The fishery CPUE was based on Gunderson 1977 and the Triennial shelf survey index was estimated using VAST.




Figure 104: Triennial shelf survey length frequency distributions for Pacific ocean perch.




Figure 105: Triennial shelf survey age frequency distributions for Pacific ocean perch.

13 Appendix D. List of Auxiliary Files Available

The listed files are also available as auxiliary files to accompany the assessment document:

1. Numbers at age for female and male Pacific ocean perch (POPnatagef.csv and POPnatagem.csv)
2. The Pacific ocean perch Stock Synthesis 3.30 model files
 - (a) 2017pop.dat
 - (b) 2017pop.ctl
 - (c) forecast.ss
 - (d) starter.ss

14 References

Bradburn, M., Keller, A., and Horness, B. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.

Chilton, D.E., and Beamish, R.J. 1982. Age determination methods for fishes studied by the Groundfish Program at the Pacific Biological Station. [Ottawa:] Minister of Supply; Services Canada.

Dick, E., Beyer, S., Mangel, M., and Ralston, S. 2017. A meta-analysis of fecundity in rockfishes (genus *Sebastes*). Fisheries Research **187**: 73–85. doi: [10.1016/j.fishres.2016.11.009](https://doi.org/10.1016/j.fishres.2016.11.009).

Dick, E.J. 2009. Modeling the Reproductive Potential of Rockfishes (*Sebastes* Spp.). ProQuest. Available from [http://books.google.com/books?hl=en&lr=&id=0d6-3rhfynkC&oi=fnd&pg=PR7&dq=%22Synthesis+of+findings+regarding+the+reproductive%22+%22C:+Linear+interpolation+algorithms%22+%22for+yellowtail+rockfish+\(S.+flavidus\)%22+%22greater+than+zero,+based+on+the+2-level+relative+fecundity%22+%22A:Methods+for+data+recovery+from+published%22+%22&ots=NR0UylgymD&sig=58IaN_a3pJeYTPYVmJ1NYMABmvE](http://books.google.com/books?hl=en&lr=&id=0d6-3rhfynkC&oi=fnd&pg=PR7&dq=%22Synthesis+of+findings+regarding+the+reproductive%22+%22C:+Linear+interpolation+algorithms%22+%22for+yellowtail+rockfish+(S.+flavidus)%22+%22greater+than+zero,+based+on+the+2-level+relative+fecundity%22+%22A:Methods+for+data+recovery+from+published%22+%22&ots=NR0UylgymD&sig=58IaN_a3pJeYTPYVmJ1NYMABmvE) [accessed 27 February 2017].

Field, J.C. 2007. Status of the Chilipepper rockfish, *Sebastes goodei*, in 2007. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Francis, R.C., and Hilborn, R. 2011. Data weighting in statistical fisheries stock assessment models. Canadian Journal of Fisheries and Aquatic Sciences **68**(6): 1124–1138. doi: [10.1139/f2011-025](https://doi.org/10.1139/f2011-025).

Gertseva, V., Matson, S., and Council, E. 2015. Status of the darkblotched rockfish resource off the continental U.S. Pacific Coast in 2015. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Gunderson, D.R. 1977. Population biology of Pacific ocean perch, *Sebastes alutus*, stocks in the WashingtonQueen Charlotte Sound region and their response to fishing. Fishery Bulletin **75**: 369–403. Available from <http://fishbull.noaa.gov/75-2/gunderson.pdf> [accessed 27 February 2017].

Gunderson, D.R. 1978. Results of cohort analysis for Pacific ocean perch stocks off British Columbia, Washington, and Oregon and an evaluation of alternative rebuilding strategies for

these stocks. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Gunderson, D.R. 1981. An updated cohort analysis for Pacific ocean perch stocks off Washington and Oregon. Unpublished report, Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Gunderson, D.R. 1997. Trade-off between reproductive effort and adult survival in oviparous and viviparous fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **54**(5): 990–998. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f97-019> [accessed 27 February 2017].

Gunderson, D.R., and Sample, T.M. 1980. Distribution and abundance of rockfish off Washington, Oregon and California during 1977. Northwest; Alaska Fisheries Center, National Marine Fisheries Service. Available from <http://spo.nmfs.noaa.gov/mfr423-4/mfr423-42.pdf> [accessed 28 February 2017].

Gunderson, D.R., Westrheim, S., Demory, R., and Fraidenburg, M. 1977. The status of Pacific ocean perch (*Sebastes alutus*) stocks off British Columbia, Washington, and Oregon in 1974.

Hamel, O.S. 2015. A method for calculating a meta-analytical prior for the natural mortality rate using multiple life history correlates. *ICES Journal of Marine Science: Journal du Conseil* **72**(1): 62–69. doi: [10.1093/icesjms/fsu131](https://doi.org/10.1093/icesjms/fsu131).

Hamel, O.S., and Ono, K. 2011. Stock Assessment of Pacific Ocean Perch in Waters off of the U.S. West Coast in 2011. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Hannah, R., and Parker, S. 2007. Age-modulated variation in reproductive development of female Pacific Ocean perch (*Sebastes alutus*) in waters off Oregon. Alaska Sea Grant, University of Alaska Fairbanks. pp. 1–20. doi: [10.4027/bamnpr.2007.01](https://doi.org/10.4027/bamnpr.2007.01).

Helser, T., Punt, A.E., and Methot, R.D. 2004. A generalized linear mixed model analysis of a multi-vessel fishery resource survey. **70**: 251–264.

Hicks, A.C., and Wetzel, C.R. 2015. The status of Widow Rockfish (*Sebastes entomelas*) along the U.S. west coast in 2015. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Hicks, A.C., Haltuch, M.A., and Wetzel, C.R. 2009. Status of greenstriped rockfish (*Sebastes elongatus*) along the outer coast of California, Oregon, and Washington. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. *Fishery*

Bulletin **82**: 898–903. Available from <http://fishbull.noaa.gov/81-4/hoenig.pdf> [accessed 28 February 2017].

Ianelli, J.N., and Zimmermann, M. 1998. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 1998. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Ianelli, J.N., Ito, D.H., and Wilkins, M. 1992. Status and future prospects for the Pacific ocean perch resource in waters off Washington and Oregon as assessed in 1992. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Karnowski, M., Gertseva, V., and Stephens, A. 2014. Historical Reconstruction of Oregon's Commercial Fisheries Landings. Oregon Department of Fish; Wildlife, Salem, OR.

Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H.J., and Bell, B. 2016. TMB: Automatic Differentiation and Laplace Approximation. *Journal of Statistical Software* **70**: 1–21.

McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling - importance resampling algorithm. *Canadian Journal of Fisheries and Aquatic Sciences* **54**: 284–300. Available from <http://www.nrcresearchpress.com/doi/pdf/10.1139/f96-285> [accessed 10 March 2017].

McCoy, M.W., and Gillooly, J.F. 2008. Predicting natural mortality rates of plants and animals. *Ecology Letters* **11**(7): 710–716. doi: [10.1111/j.1461-0248.2008.01190.x](https://doi.org/10.1111/j.1461-0248.2008.01190.x).

Methot, R.D., and Wetzel, C.R. 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* **142**: 86–99. doi: [10.1016/j.fishres.2012.10.012](https://doi.org/10.1016/j.fishres.2012.10.012).

Methot, R.D., Taylor, I.G., and Chen, Y. 2011. Adjusting for bias due to variability of estimated recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(10): 1744–1760. doi: [10.1139/f2011-092](https://doi.org/10.1139/f2011-092).

Pikitch, E.K., Erickson, D.L., and Wallace, J.R. 1988. An evaluation of the effectiveness of trip limits as a management tool. Northwest; Alaska Fisheries Center, National Marine Fisheries Service NWAFC Processed Report. Available from <https://www.afsc.noaa.gov/Publications/ProcRpt/PR1988-27.pdf> [accessed 28 February 2017].

Punt, A.E., Smith, D.C., KrusicGolub, K., and Robertson, S. 2008. Quantifying age-reading error for use in fisheries stock assessments, with application to species in Australia's southern and eastern scalefish and shark fishery. *Canadian Journal of Fisheries and Aquatic Sciences* **65**(9): 1991–2005. doi: [10.1139/F08-111](https://doi.org/10.1139/F08-111).

Ralston, S., Pearson, D.E., Field, J.C., and Key, M. 2010. Documentation of the California catch reconstruction project. US Department of Commerce, National Oceanic; Atmospheric

Administration, National Marine.

Rogers, J. 2003. Species allocation of *Sebastes* and *Sebastolobus* species caught by foreign countries off Washington, Oregon, and California, U.S.A. in 1965-1976. Unpublished document.

Rogers, J.B., and Pikitch, E.K. 1992. Numerical definition of groundfish assemblages caught off the coasts of Oregon and Washington using commercial fishing strategies. Canadian Journal of Fisheries and Aquatic Sciences **49**(12): 2648–2656. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f92-293> [accessed 9 March 2017].

Seeb, L.W., and Gunderson, D.R. 1988. Genetic variation and population structure of Pacific ocean perch (*Sebastes alutus*). Canadian Journal of Fisheries and Aquatic Sciences **45**(1): 78–88. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f88-010> [accessed 28 February 2017].

Stewart, I.J., and Hamel, O.S. 2014. Bootstrapping of sample sizes for length- or age-composition data used in stock assessments. Canadian Journal of Fisheries and Aquatic Sciences **71**(4): 581–588. doi: [10.1139/cjfas-2013-0289](https://doi.org/10.1139/cjfas-2013-0289).

Then, A.Y., Hoenig, J.M., Hall, N.G., and Hewitt, D.A. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. ICES Journal of Marine Science **72**(1): 82–92. doi: [10.1093/icesjms/fsu136](https://doi.org/10.1093/icesjms/fsu136).

Thorson, J.T., and Barnett, L.A.K. 2017. Comparing estimates of abundance trends and distribution shifts using single- and multispecies models of fishes and biogenic habitat. ICES Journal of Marine Science: Journal du Conseil: fsw193. doi: [10.1093/icesjms/fsw193](https://doi.org/10.1093/icesjms/fsw193).

Thorson, J.T., and Kristensen, K. 2016. Implementing a generic method for bias correction in statistical models using random effects, with spatial and population dynamics examples. Fisheries Research **175**: 66–74. doi: [10.1016/j.fishres.2015.11.016](https://doi.org/10.1016/j.fishres.2015.11.016).

Thorson, J.T., and Ward, E.J. 2014. Accounting for vessel effects when standardizing catch rates from cooperative surveys. Fisheries Research **155**: 168–176. doi: [10.1016/j.fishres.2014.02.036](https://doi.org/10.1016/j.fishres.2014.02.036).

Thorson, J.T., Shelton, A.O., Ward, E.J., and Skaug, H.J. 2015. Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. ICES Journal of Marine Science **72**(5): 1297–1310. doi: [10.1093/icesjms/fsu243](https://doi.org/10.1093/icesjms/fsu243).

Thorson, J.T., Stewart, I.J., and Punt, A.E. 2012. nwfsAgeingError: A user interface in R for the Punt et al. (2008) method for calculating ageing error and imprecision. Available from: <http://github.com/nwfs-assess/nwfsAgeingError/>.

Weinberg, J.R., Rago, P.J., Wakefield, W.W., and Keith, C. 2002. Estimation of tow distance

and spatial heterogeneity using data from inclinometer sensors: An example using a clam survey dredge. *Fisheries Research* **55**(1–3): 49–61. doi: [10.1016/S0165-7836\(01\)00292-2](https://doi.org/10.1016/S0165-7836(01)00292-2).

Wilkins, M., and Golden, J. 1983. Condition of the Pacific ocean perch resource off Washington and Oregon during 1979: Results of a cooperative trawl survey. *North American Journal of Fisheries Management* **3**: 103–122.

Withler, R., Beacham, T., Schulze, A., Richards, L., and Miller, K. 2001. Co-existing populations of Pacific ocean perch, *Sebastes alutus* , in Queen Charlotte Sound, British Columbia. *Marine Biology* **139**(1): 1–12. doi: [10.1007/s002270100560](https://doi.org/10.1007/s002270100560).