



NOAA FISHERIES SERVICE

Pacific Halibut Bycatch in US West Coast Fisheries (2002-2017)



Jason Jannot
Kayleigh Somers
Neil Riley
Vanessa Tuttle

Publication date: August 2018

This document should be cited as follows:


Jannot, J.E., K. A. Somers, N. B. Riley, V. Tuttle, J. McVeigh. 2018. Pacific halibut bycatch in the U.S. west coast fisheries (2002-2017). NOAA Fisheries, NWFSC Observer Program, 2725 Montlake Blvd E., Seattle, WA 98112.



NWFSC Observer Program
National Marine Fisheries Service
Northwest Fisheries Science Center
Fishery Resource Analysis and Monitoring Division
2725 Montlake Blvd. E.
Seattle, WA 98112




PACIFIC HALIBUT BYCATCH 2002-2017




2017 Catch Shares
Bycatch Allocation: 35.9 mt




Trawl Gears




△ = 107.3 mt
2002-2017 *LE Sablefish




Fixed Gears



Pacific Halibut Bycatch Observed 2002-2017



Fixed Gears



CONTACT INFO: Jason Jannot
NWFSC, Seattle, WA
jason.jannot@noaa.gov



WEBSITE: <http://tinyurl.com/PhalibutReports>

Pacific Halibut Bycatch in U.S. West Coast Groundfish Fisheries (2002-2017)

Jason E. Jannot*, Kate Richerson, Kayleigh Somers,
Neil B. Riley, Vanessa Tuttle, Jon McVeigh

NWFSC Observer Program
Fishery Resource Analysis and Monitoring Division
Northwest Fisheries Science Center
National Marine Fisheries Service
NOAA
Seattle, WA 98112

August 9, 2018

*jason.jannot@noaa.gov

Contents

1	EXECUTIVE SUMMARY	4
2	INTRODUCTION	13
2.1	Observed West Coast Groundfish Fisheries	13
2.2	NW Fisheries Science Center (NWFSC) Groundfish Observer Program	15
2.3	Pacific Halibut Management and Fishery Interaction	16
3	METHODS	17
3.1	Data Sources	17
3.2	Shore-based IFQ Fishery	17
3.2.1	Pacific Halibut Data Collection in the Shore-based IFQ Fishery	18
3.2.2	Shore-based IFQ fishery Bycatch Estimation	19
3.2.3	Viability Analysis	21
3.2.4	Length Frequencies	22
3.3	Non-nearshore Fixed Gear Fishery	23
3.3.1	Discard Estimation	24
3.3.2	Discard Mortality Rates	26
3.4	IPHC Pacific halibut Derby Fishery	28
3.5	Observed State Fisheries	28
3.6	Exempted Fishing Permits	29
3.7	Non-groundfish Fisheries Not Observed by NWFSC	29
4	RESULTS	30
4.1	IFQ Fishery	30
4.2	IFQ Electronic Monitoring EFP	31
4.3	Non-Nearshore Fixed Gear Fishery	31
4.4	IPHC Pacific halibut Derby	31
4.5	Observed State Fisheries, EFPs and Non-Groundfish Fisheries	32
5	SUMMARY & CONCLUSIONS	32
5.1	IFQ Fishery	32
5.2	Non-IFQ Fisheries	32
6	ACKNOWLEDGEMENTS	33
7	REFERENCES	33
8	TABLES	36
8.1	Tables: IFQ Fishery	36
8.2	Tables: Non-Nearshore Fixed Gear Fisheries	62
8.3	Tables: Legal-Sublegal P. halibut Lengths	78
8.4	Tables: IPHC P. halibut Derby Fishery	79
8.5	Tables: Observed State Fisheries	81
8.6	Tables: At-Sea Hake Fisheries	86
8.7	Tables: IFQ Electronic Monitoring	90

8.8	Tables: Other EFP fleet and PHLB catch summaries	92
9	FIGURES	94
A	Appendicies	101
A.1	IFQ Electronic Monitoring DMR comparison	101
A.2	Catch Shares Weighted Length Frequencies	105
A.3	Pacific Halibut IBQ Expansions for In-Season Management, Special Cases	122
A.3.1	In season reporting to the Vessel Account System	122
A.3.2	In season IBQ Weight Calculations for Bottom Trawl Gear	122
A.3.3	In season IBQ Weight Calculations for Pot Gear	123
A.3.4	In season IBQ Weight Calculations for Hook-&-Line Gear	123
A.3.5	In season IBQ Weight Alternative Calculation Scenarios	124
A.4	IPHC Length-Weight Table	130
A.5	Data flow	131

1 EXECUTIVE SUMMARY

Pacific halibut mortality estimates are provided for the years 2002 through 2017 from all fishery sectors observed by the Northwest Fishery Science Center Groundfish Observer Program (Table 1).

Table 1: Pacific halibut mortality estimates for 2017 and the years of observation, for all fishery sectors observed by the Northwest Fishery Science Center Groundfish Observer Program. Estimates include both individuals discarded at the dock and with mortality rates applied, where appropriate.

Sector	Years Observed	2017 Pacific halibut Discard Mortality (mt)
Individual Fishing Quota (IFQ) fisheries ¹	2011-2017	31.41
IFQ Electronic Monitoring (EM) EFP ²	2015-2017	5.47
At-sea Pacific hake	2002-2017	0.55
Non-nearshore fixed gear targeting groundfish	2002-2017	41.71
Nearshore fixed gear	2003-2017	1.55
Pink shrimp trawl	2004-2017	0.00
California halibut trawl	2002-2017	0.00
P. halibut Derby	2017	2.26
Ridgeback Prawn	2017	0.00
Sea Cucumber	2017	0.00

¹ Does not include estimates from IFQ vessels with Electronic Monitoring

² EFP = Exempted Fishing Permit, includes all gears

In addition, we provide historical estimates of P. halibut bycatch in the Limited Entry (LE) bottom trawl fishery for the 2002-2010 period and P. halibut bycatch estimates for observed, non-IFQ vessels with an exempted fishing permit (EFP) targeting groundfish (2002-2017). For completeness, we also include the P. halibut landed catch from PacFIN fish tickets reported by non-groundfish fisheries that are not observed by the NWFSC Observer Program for the period 2002-2017. Estimates of P. halibut bycatch from the International Pacific Halibut Commission P. halibut derby, the California sea cucumber, and the California ridgeback prawn fisheries are included in this report for the first time.

Final estimates of observed fishery sectors including the IFQ EM EFP are shown in Tables 1, 2 & 65 in the report. We include in these tables (and elsewhere in the report), the small amount of P. halibut landed and subsequently discarded at the dock by IFQ bottom and midwater trawl vessels. These landed and then discarded at the dock amounts are listed by strata in Tables 8 and 9 of the report. IFQ EM EFP P. halibut catch is included in the summaries found in Tables 1, 2, 62 and 65. A comparison of observed discard mortality rates (DMR) on IFQ vessels with and without EM is provided in Appendix A.1 and includes a comparison of the observer viability method to the PFMC's Groundfish Management Team Time-on-Deck model for mortality of P. halibut on IFQ EM vessels.

In 2017 the non-nearshore fixed gear sector had the largest estimated *P. halibut* discard mortality of any sector (41.71 mt, Tables 1, 2 & 39). Nearly all of that bycatch (37.95 mt, or 91%) occurred on the LE Sablefish Endorsed vessels. These vessels fish federally permitted sablefish tier quota during the primary season (April-October). Almost all of the LE Sablefish Endorsed bycatch occurred fishing longline gear north of Point Chehalis, WA (33.21 mt or 88%, Table 39). A smaller amount of *P. halibut* mortality also occurred on LE sablefish endorsed vessels fishing longline gear south of Pt. Chehalis (4.58 mt). Open access (OA) vessels targeting non-nearshore groundfish species with hook-&-line gear caught substantially less than the LE sector (3.57 mt).

In 2017, the IFQ sector accounted for the second largest source of discard mortality of *P. halibut* among the sectors analyzed (31.41 mt, Table 1 & 2), with the majority of this bycatch caught on bottom trawl vessels between Pt. Chehalis, WA and 40°10' N. lat., fishing deeper than 60 fathoms (18.99 mt, Table 18). IFQ bottom trawl and LE Sablefish Endorsed longline vessels together comprised approximately 88% of the 2017 *P. halibut* discard mortality in observed U.S. West Coast groundfish fisheries.

The 2017 IFQ fishery estimate of *P. halibut* discard mortality, coast-wide, was 31.41 mt, with an additional 5.47 mt caught by IFQ EM EFP vessels (Tables 1 & 62) which is included in the IFQ estimate in Tables 2 and 65. The IFQ total (IFQ + IFQ EM EFP: 36.88 mt) is 1.73 mt greater than the 2016 estimate (35.15 mt, see Table 2) but, as in past years, well below the IBQ¹ allocation (79.33 mt). As in prior years, bottom trawl gear produced the largest component of IFQ discard mortality, followed in decreasing magnitude by pot, hook-&-line, and midwater trawl gear.

In Appendix A.1, we present a comparison of alternative methods for calculating discard mortality rates (DMRs) in the IFQ EM fishery. Electronic monitoring does not yet allow for accurate estimates of Pacific halibut viability. Currently in the IFQ EM bottom trawl fishery a 0.90 mortality rate is applied to all *P. halibut* bycatch (Table 62). As an alternative to the 0.90 rate, we also present mortality estimates based on observer assessed viabilities and the PFMC Groundfish Management Team's Time-on-Deck model (see Appendix A.1). Small sample sizes preclude definitive conclusions from this analysis. The NWFSC Observer Program might revisit this analysis in future reports.

In 2017, the NWFSC Observer Program began observing the International Pacific Halibut Commission's (IPHC) Pacific halibut derby fishery, the California sea cucumber trawl fishery, and the California ridgeback prawn trawl fishery. For the first time in this report, we estimate *P. halibut* discard mortality in these fisheries. There was zero (0) observed catch of *P. halibut* in the sea cucumber and ridgeback prawn fisheries (Table 57 & 58). The Pacific halibut discard mortality estimate for the 2017 IPHC Pacific halibut derby fishery was 2.26 mt (Tables 1, 2). Observer coverage, discard ratios, fleet-wide estimates of gross discards, discard mortality, and retained *P. halibut* are presented in Tables 47, 48, & 50. Discard mortality estimates were calculated using the same methods as for the non-nearshore hook-and-line fishery, which uses observed estimates of *P. halibut* viability. Viabilities of observed *P. halibut* bycatch in the *P. halibut* derby fishery are given in Table 49. Observed lengths of discarded *P. halibut* in the derby fishery are given in Tables 51 & 52.

¹IBQ = Individual Bycatch Quota, which is used for *P. halibut* North of 40°10' N. lat.

Pacific halibut discard in the nearshore fixed gear, pink shrimp trawl, California halibut trawl, and at-sea Pacific hake fisheries combined represents a very small component of total P. halibut mortality (Table 1; Figure 1).

The NWFSC Observer Program data used in this report has been updated to include the most recent data available (2002-2017). Pacific Fisheries Information Network (PacFIN) data used in this report were accessed April 2017. The estimates for all sectors and years (except LE Trawl 2002-2010) have been recalculated based on these base data. In all other respects, this report uses the same methods as reported in last year's report (Jannot et al. 2016).

Table 2: Pacific halibut discard mortality estimates (mt, including a small amount discarded at the dock in IFQ Bottom Trawl, Midwater Rockfish, and Midwater Hake fisheries) for all sectors observed by the NWFSC Groundfish Observer Program. Mortality rates of less than 100% were applied in the bottom trawl fisheries (LE and IFQ), IFQ hook and line, IFQ pot, and non-IFQ, non-nearshore fixed gear sectors, for which some information regarding gear specific survivorship was available. For all other sectors, a 100% mortality rate was applied because gear specific survivorship information is not available. Rounding of values might mask very small weights in some categories and are presented here as zero (0). All weights are estimated based on whole fish (a.k.a. 'round weight', not head-&-gut). Ridgeback Prawn and Sea Cucumber fisheries had zero (0) observed P. halibut catch. * = confidential data, less than 3 vessels observed; - = no observer coverage.

Year	LE bottom trawl 2002-10	IFQ Fishery ⁹						Non-Nearshore fixed gear			Nearshore Fixed Gear ³	Pink Shrimp ³	CA Halibut _{3,6}	P. halibut Derby	At-sea Hake ³	Totals		
		Bottom Trawl _{1,2,9}	LE CA Halibut _{1,3}	Hook & Line	Pot ⁹	Midwater Rockfish _{3,4,9}	Midwater Hake ^{2,3,5,9}	LE Endorsed	LE Non- Endsd	OA						All sec- tors	Sectors w/ <100% mortal- ity rate ₇	Sectors w/100% mortal- ity rate ₈
Total Discard Mortality (mt)																		
2002	344.82							22.76	0.00	-	-	-	0.00		1.14	368.72	367.58	1.14
2003	124.43							31.54	0.03	-	0.00	-	0.00		2.65	158.65	156.00	2.65
2004	133.12							38.82	0.00	-	1.00	0.00	0.70		1.13	174.77	172.64	2.13
2005	286.52							38.10	0.00	-	2.20	0.04	0.03		1.97	328.86	324.65	4.21
2006	242.47							107.30	0.00	-	0.53	-	0.02		0.83	351.15	349.79	1.36
2007	208.81							21.24	0.28	3.48	0.09	0.21	0.03		1.18	235.32	233.84	1.48
2008	207.81							41.65	0.48	6.42	0.35	0.00	0.31		3.98	261.00	256.67	4.33
2009	251.1							51.47	0.04	5.65	1.28	0.00	0.00		0.33	309.87	308.26	1.61
2010	180.97							22.12	0.06	5.22	0.08	0.00	0.00		1.57	210.02	208.37	1.65
2011		31.30	0	0.97	0.89	*	0.35	12.07	3.20	2.09	3.07	0.19	0.00		0.61	54.74	50.52	4.22
2012		36.13	*	2.34	0.51	0.0	0.62	24.94	0.73	1.61	2.25	0.00	0.00		0.64	69.77	66.26	3.51
2013		32.41	see ¹	0.48	0.21	0.0	1.34	2.94	0.00	0.07	1.36	0.00	0.00		1.06	39.87	36.11	3.76
2014		26.28	see ¹	0.61	0.08	0.0	1.36	30.16	0.00	0.35	0.95	0.00	0.00		0.37	60.16	57.48	2.68
2015		33.36	see ¹	1.52	0.38	0.0	0.70	10.37	0.02	0.46	1.44	0.01	0.00		0.06	48.32	46.11	2.21
2016		33.28	see ¹	1.02	0.18	0.0	0.68	16.62	0.91	2.56	3.01	0.00	0.00		0.15	58.41	54.57	3.84
2017		35.11	see ¹	0.66	0.78	0.0	0.51	37.95	0.03	3.73	1.55	0.00	0.00	2.26	0.55	83.13	80.52	2.61

¹ Starting in 2013, LE CA Halibut estimates are combined with IFQ Bottom Trawl estimates.

² Includes a small amount landed and discarded at the dock.

³ 100% mortality rate

⁴ from 2011-14, 'Midwater Trawl'

⁵ from 2011-14, 'Shoreside Hake'

⁶ Starting in 2011, this sector only includes OA CA halibut

⁷ LE Bottom Trawl, IFQ Bottom Trawl, IFQ hook and line, IFQ pot, LE and OA CA Halibut, Non-Nearshore Fixed Gear

⁸ IFQ Midwater Rockfish, Midwater Hake, Nearshore fixed gear, Pink Shrimp, At-sea Hake

⁹ Includes P. halibut catch from IFQ electronic monitoring EFP

Note: Ridgeback Prawn and Sea Cucumber fisheries had zero (0) observed P. halibut catch

Table 3: Percent of legal-sized P. halibut bycatch, by weight (mt) in the IFQ Bottom Trawl fishery north of 40 °10' N. lat. (mortality rate applied).

Year	% legal-sized P. halibut in IFQ bottom trawl north of 40 °10' N. lat.
2011	67%
2012	67%
2013	64%
2014	60%
2015	68%
2016	67%
2017	76%




Figure 1: Total estimated *P. halibut* discard mortality (mt \pm 1 SE, with mortality rates applied if applicable) from all sectors observed by the NWFSC Groundfish Observer Program. estimates are not included for sectors and years where there were insufficient observer data. Values are reported in Table 2

¹ Individual Bycatch Quota (IBQ) allocated north of 40°10' N. latitude.

² IBQ catch observations includes all sectors and gears except At-sea Hake which is shown separately.

³ Other fisheries includes OR and CA Nearshore, WA, OR, and CA pink shrimp, California halibut, sea cucumber, ridgeback prawn, and IPHC *P. halibut* derby fisheries.

List of Tables

1	P. halibut mortality summary for 2017	4
2	Pacific halibut discard mortality estimates by sector	7
3	Percent of legal-sized P. halibut bycatch for the IFQ Bottom Trawl fishery north of 40°10' N. lat.	8
4	Data collected from P. halibut caught on IFQ vessels using different types of gear.	18
5	Mortality rates for IFQ bottom trawl vessels	19
6	Mortality rates for IFQ pot vessels	19
7	Mortality rates used for each of the condition categories for non-nearshore longline and hook & line vessels	27
8	IFQ bottom trawl coverage and P. halibut catch	37
9	IFQ midwater trawl coverage and P. halibut catch	38
10	IFQ fixed gear coverage and P. halibut catch	39
11	IFQ discard ratios for unsampled catch on bottom trawl vessels	40
12	IFQ discard ratios for unsampled catch on midwater trawl vessels	41
13	IFQ discard ratios for unsampled catch on fixed gear vessels	42
14	IFQ P. halibut bottom trawl viabilities by area, depth, and year	43
15	IFQ P. halibut midwater trawl viabilities by area and year	46
16	IFQ P. halibut Pot viabilities by area and year	47
17	IFQ P. halibut caught on hook and line vessels by area and year	48
18	IFQ bottom trawl gross at-sea discard and at-sea mortality by area, depth, and year	49
19	IFQ midwater trawl gross at-sea discard and at-sea mortality by area and year	51
20	IFQ pot gross at-sea discard and at-sea mortality by area and year	52
21	IFQ hook and line gross at-sea discard and at-sea mortality by area and year	53
22	IFQ legal-sized (82 cm) mortality from bottom trawl vessels	54
23	IFQ legal-sized (82 cm) mortality from midwater trawl vessels	56
24	IFQ legal-sized (82 cm) mortality from pot vessels	57
25	IFQ legal-sized (82 cm) mortality from hook and line vessels	58
26	IFQ bottom trawl P. halibut monthly bycatch	59
27	IFQ bottom trawl vessel actual length frequencies.	60
28	IFQ pot vessels actual length frequencies	61
29	Visual estimates of P. halibut lengths (cm) from IFQ vessels using bottom trawl, pot, and hook and line gear (2011-2017)	62
30	Number of observed vessels, trips, and sets by year and gear type in the LE Sablefish Endorsed fishery.	63
31	Number of observed vessels, trips, and sets by year and gear type in the LE Sablefish NonEndorsed fishery. The number of observed pot vessels in this fishery is too small to meet confidentiality and thus not reported.	64
32	Number of observed vessels, trips, and sets by year and gear type in the OA Fixed Gear fishery. OA Fixed Gear fishery was not observed until 2003.	64
33	Expansion factors and discard rates for Non-Nearshore Fixed Gear fishery	65
34	Expansion factors used in the non-nearshore fishery	66
35	Percent of observed trips that caught Pacific halibut in non-nearshore fishery.	67
36	Viabilities caught by longline vessels in the LE Sablefish Endorsed fishery	68

37	Viabilities caught by longline vessels in the LE Sablefish Non-Endorsed fishery . . .	68
38	Viabilities caught by longline vessels in the OA Fixed Gear fishery	69
39	Estimated gross discard (mt) and discard mortality (mt) in the non-nearshore fishery	70
40	Estimated P. halibut discard mortality (mt, mortality rate applied, see text for de- scription) from each sector of the non-nearshore fixed gear fishery by year.	71
41	LE Sablefish Endorsed hook and line vessels actual length frequencies	72
42	LE Sablefish Endorsed pot vessels actual length frequencies	73
43	LE Sablefish Non-Endorsed hook and line vessels actual length frequencies	74
44	OA Fixed Gear hook and line vessels actual length frequencies	75
45	Visual estimates of P. halibut lengths (cm) from Non-Nearshore fixed gear vessels (2002-2017)	77
46	Legal (82 cm) versus sub-legal actual and visual length frequencies	78
47	P. halibut Derby fishery observer coverage	79
48	Discard ratio and gross discard estimate of P. halibut for the P. halibut Derby fishery	79
49	Viability of P. halibut discards in the P. halibut Derby Fishery	79
50	P. halibut discard mortality in the P. halibut Derby Fishery	80
51	Physical length frequencies of P. halibut discarded in the P. halibut Derby Fishery . .	80
52	Visual length estimates of P. halibut discarded in the P. halibut Derby Fishery	81
53	Nearshore fishery coverage and P. halibut bycatch	82
54	WA and OR Pink Shrimp fishery coverage and P. halibut bycatch	83
55	CA Pink Shrimp fishery coverage and P. halibut bycatch	84
56	California halibut fishery coverage and P. halibut bycatch	85
57	Sea Cucumber fishery observer coverage and P. halibut bycatch	86
58	Ridgeback Prawn fishery observer coverage and P. halibut bycatch	86
59	At-sea Hake fishery coverage and P. halibut bycatch	87
60	Tribal At-sea Hake fishery coverage and P. halibut bycatch	88
61	At-sea Hake fishery actual P. halibut length frequencies	89
62	IFQ coverage and P. halibut bycatch on electronic monitoring EFP vessels	91
63	P. halibut catch on EFP vessels	92
64	P. halibut catch from non-groundfish fisheries not observed by the NWFSC Observer Program.	92
65	Discard estimates for all fishery sectors	93
66	IFQ EM bottom trawl vessels that caught P. halibut	102
67	P. halibut mortality estimates comparing the observer viability method to the Time- on-Deck model on EM and non-EM bottom trawl vessels	103
68	IFQ EM pot vessels that caught P. halibut	104
69	P. halibut mortality estimated using the observer viability method on EM pot vessels, compared with non-EM vessels.	104
70	Weighted length frequency distributions for bottom trawl vessels in the IFQ fishery .	105
71	Percentage of weighted length measurements in each viability category, for IFQ bottom trawl vessels	108
72	Table 71 continued for IFQ bottom trawl vessels	109
73	Weighted length frequency distributions for pot vessels in the IFQ fishery	109
74	Percentage of weighted length measurements in each viability category, for IFQ pot vessels	113

75	Table 75 continued for IFQ pot vessels	114
76	Weighted length frequency distributions for Pacific halibut in the limited entry bottom trawl fishery, 2002-10	115
77	Percentage of weighted length measurements in each condition category for the limited entry bottom trawl fishery, 2002-10.	116
78	Continuation of Table 77	117
79	Number of dead <i>P. halibut</i> in each length bin, summed across viability categories, for IFQ bottom trawl vessels	119
80	Number of dead <i>P. halibut</i> in each length bin, summed across viability categories, for IFQ pot vessels	120
81	Number of dead <i>P. halibut</i> in each length bin for Shoreside Hake vessels 2011-14.	121
82	Number of dead <i>P. halibut</i> in each length bin for Midwater Rockfish vessels by year	122
83	VAS calculations for unsampled <i>P. halibut</i>	128

List of Figures

1	Total estimated <i>P. halibut</i> discard mortality by sector	9
2	Number of vessels by month for IFQ bottom trawl vessels in 2017	94
3	Number of tows by month for IFQ bottom trawl vessels in 2017	95
4	Tow hours by month for IFQ bottom trawl vessels in 2017	96
5	Non-Nearshore Fixed Gear <i>P. halibut</i> mortality by sector and year	97
6	Length frequency distribution of Pacific halibut in the Non-Nearshore Fixed Gear fishery	98
7	Spatial distribution of Pacific halibut bycatch	99
8	Number of sets, trips, and vessels by opening day for the <i>P. halibut</i> Derby fishery	100
9	IPHC length-weight conversion table for Pacific halibut.	130
10	IFQ groundfish fishery data flow	131

2 INTRODUCTION

Pacific halibut (*Hippoglossus stenolepis*) is found in coastal waters throughout the North Pacific. Off the U.S. West Coast of the United States, it inhabits continental shelf areas (<150 fm) from Washington to central California (Clark and Hare 1998). Pacific halibut has long supported a directed commercial fishery in the U.S. and Canada, but it is also caught as bycatch in other fisheries that target demersal species inhabiting similar depths and seafloor habitat types (Chastain 2012). The objective of this report is to provide estimates of P. halibut bycatch in the U.S. West Coast groundfish fisheries from 2002-2017.

2.1 Observed West Coast Groundfish Fisheries

The U.S. West Coast groundfish fishery is a multi-species fishery that utilizes a variety of gear types. The fishery harvests species designated in the Pacific Coast Groundfish Fishery Management Plan (FMP; PFMC 2011) and is managed by the Pacific Fishery Management Council (PFMC). Over 90 species are listed in the groundfish FMP, including a variety of rockfish, flatfish, roundfish, skates, and sharks. These species are found in both state (0-4.8 km) and federal (>4.8 km off-shore to the EEZ) waters. Groundfish are both targeted and caught incidentally by trawl nets, hook-&-line gears, and fish pots. Under the FMP, the groundfish fishery consists of four management components:

- The Limited Entry (LE) component encompasses all commercial fisheries who hold a federal limited entry permit. The total number of limited entry permits available is restricted. Vessels with an LE permit are allocated a larger portion of the total allowable catch for commercially desirable species than vessels without an LE permit.
- The Open Access (OA) component encompasses commercial fishers who do not hold a federal LE permit. Some states require fishers to carry a state issued permit for certain OA sectors.
- The Recreational component includes recreational anglers who target or incidentally catch groundfish species. Estimate of P. halibut bycatch in recreational fisheries are compiled by the IPHC and are not covered by this report.
- The Tribal component includes native tribal commercial fishers in Washington state that have treaty rights to fish groundfish. Estimates of P. halibut bycatch from tribal fisheries are compiled by the IPHC and are not included in this report, with the exception of the observed tribal at-sea Pacific hake (a.k.a. Pacific whiting, henceforth referred to as hake) sector which are included as part of the “At-sea hake” values included in Tables 2 and 65.

These four components can be further subdivided into sectors based on gear type, target species, permits and other regulatory factors. This report includes data from the following sectors:

- IFQ fishery: This sector is subdivided into the following components due to differences in gear type and target strategy:
 - Bottom trawl (formerly LE bottom trawl 2002-2010): Bottom trawl nets are used to catch a variety of non-hake groundfish species. Catch is delivered to shore-based processors.

- Midwater rockfish trawl: Midwater trawl nets are used to target mid-water non-hake species, typically rockfish, and landings of Pacific hake are less than 50% (by weight) of total trip landings. Catch is delivered to shore-based processors. From 2011-14, reported as IFQ non-hake Midwater Trawl.
 - Pot: Pot gear is used to target groundfish species, primarily sablefish. Catch is delivered to shore-based processors.
 - Hook-and-Line: Longlines are primarily used to target groundfish species, mainly sablefish. Catch is delivered to shore-based processors.
 - LE California halibut trawl: Bottom trawl nets are used to target California halibut by fishers holding a state California halibut permit and an LE federal trawl groundfish permit. Catch is delivered to shore-based processors.
 - Midwater hake trawl: Midwater trawl nets are used to catch Pacific hake and more than 50% (by weight) of the total trip landings is P. hake. Catch is delivered to shore-based processors. From 2011-14, reported as Shoreside Hake.
 - At-sea motherships: Midwater trawl nets are used to catch Pacific hake. Catcher vessels deliver unsorted catch to a mothership. The catch is sorted and processed aboard the mothership.
 - At-sea catcher-processors: Midwater trawl nets are used to catch and process Pacific hake at sea.
 - At-sea tribal: Midwater trawl nets are used to catch and process Pacific hake at sea by Native American tribes. The tribes must operate within defined boundaries in waters off northwest Washington.
- OA pink shrimp trawl: Trawl nets are used to target pink shrimp on vessels carrying a state pink shrimp permit. Catch is delivered to shore-based processors.
 - OA ridgeback prawn trawl: Trawl nets are used to target ridgeback prawn. Catch is delivered to shore-based processors.
 - OA California halibut trawl: Trawl nets are used to target California halibut by fishers holding a state California halibut permit. Catch is delivered to shore-based processors.
 - OA sea cucumber trawl: Trawl nets are used to target sea cucumbers. Catch is delivered to shore-based processors.
 - LE fixed gear (non-nearshore): This sector is subdivided into two components based on differences in permitting and management:
 - LE sablefish endorsed: Longlines and pots are used to target sablefish. Catch is generally delivered to shore-based processors.
 - LE sablefish non-endorsed: Longlines and pots are used to target groundfish, primarily sablefish and thornyheads, by LE sablefish vessels that have caught their sablefish quota limit or are fishing outside the normal LE sablefish season. Catch is delivered to shore-based processors or sold alive.

- OA fixed gear (non-nearshore): Fixed gear, including longlines, pots, fishing poles, stick gear, etc. is used to target non-nearshore groundfish. Catch is delivered to shore-based processors.
- Pacific halibut fixed gear derby : Longlines and hook-and-lines are used to target Pacific halibut during one or more 24-hour fishing derbies. This fishery is managed by the International Pacific Halibut Commission. Catch is delivered to shore-based processors.
- Nearshore fixed gear: A variety of fixed gear, including longline, pots, fishing poles, stick gear, etc. are used to target nearshore rockfish and other nearshore species managed by state permits in Oregon and California. Catch is delivered to shore-based processors or sold live.

2.2 NW Fisheries Science Center (NWFSC) Groundfish Observer Program

The NWFSC Groundfish Observer Program observes commercial sectors that target or take groundfish as bycatch. The observer program has two units: the West Coast Groundfish Observer Program (WCGOP) and the At-Sea Hake Observer Program (A-SHOP).

The WCGOP was established in May 2001 by NOAA Fisheries (a.k.a., National Marine Fisheries Service, NMFS) in accordance with the Pacific Coast Groundfish Fishery Management Plan (50 CFR Part 660) (50 FR 20609). This regulation requires all vessels that catch groundfish in the U.S. EEZ from 4.8-322 km offshore carry an observer when notified to do so by NMFS or its designated agent. Subsequent state rule-making has extended NMFS's ability to require vessels fishing in the 0-4.8 km state territorial zone to carry observers.

The A-SHOP has conducted observations of the U.S. West Coast at-sea hake fishery since 2001. Prior to 2001, observer coverage of the U.S. West Coast at-sea hake fishery was conducted by the North Pacific Groundfish Observer Program. Current A-SHOP program information and documentation on data collection methods can be found in the A-SHOP observer manual (NWFSC 2018b). The at-sea hake fishery has mandatory observer coverage, with each vessel over 38 meters carrying two observers. Beginning in 2011, under IFQ/Co-op Program management, all catcher vessels that deliver catch to motherships are required to carry WCGOP observers or use electronic monitoring equipment.

The NWFSC Groundfish Observer Program's goal is to improve estimates of total catch and discard by observing groundfish fisheries along the U.S. West Coast. The WCGOP and A-SHOP observe distinct sectors of the groundfish fishery. The WCGOP observes multiple sectors of the groundfish fishery, including: IFQ shoreside delivery of groundfish and Pacific hake, at-sea mothership catcher-vessels fishing for Pacific hake, LE and OA fixed gear, and state-permitted nearshore fixed gear sectors. The WCGOP also observes several fisheries that incidentally catch groundfish, including the California halibut trawl and pink shrimp trawl fisheries. The A-SHOP observes the fishery that catches and delivers Pacific hake at-sea including non-tribal catcher-processor and mothership vessels.

2.3 Pacific Halibut Management and Fishery Interaction

The International Pacific Halibut Commission, a body founded through treaty agreement between the U.S. and Canada, sets the *P. halibut* annual total allowable catch (TAC) for IPHC Area 2A, the collective U.S. waters off the states of Washington, Oregon and California. The TAC is based on bycatch mortality, which takes into account potential survival after being discarded. Regulations for IPHC Area 2A are set by NOAA Fisheries West Coast Regional Office. Pacific halibut catch in Area 2A is divided between tribal and non-tribal fisheries, between commercial and recreational fisheries, and between recreational fisheries in different states (Washington, Oregon and California). The Pacific Fishery Management Council describes this *P. halibut* catch division each year in a catch-sharing plan. In 2017, the LE fixed gear sablefish endorsed sector was allowed to retain and land *P. halibut* north of Pt. Chehalis, WA. The IFQ midwater Pacific hake fishery is a maximized-retention fishery. Under this fishery, small amounts of incidental *P. halibut* take are allowed to be landed and subsequently donated to food banks or destroyed. In all other West Coast commercial groundfish fishery sectors, *P. halibut* must be discarded at-sea. However, small amounts of *P. halibut* are, on rare occasions, mixed with target species and accidentally landed. These individuals are subsequently donated or destroyed as in the IFQ Midwater hake fishery.

In 2011, the LE bottom trawl sector of the U.S. West Coast groundfish fishery began fishing under an IFQ management program. An IFQ is defined as a federal permit under a limited access system to harvest a quantity of fish, representing a portion of the total allowable catch of a fishery that can be received or held for exclusive use by a person (MSA 16 UIC 1802(23)). The implementation of the IFQ management program in 2011 resulted in changes to the method used for estimating fishing mortality, including the mandate that vessels must carry NMFS observers on all IFQ fishing trips. A full list of changes to the fishery can be found in Jannot et al. 2012.

Under the IFQ program, *P. halibut* is managed at the permit level, through Individual Bycatch Quota (IBQ) pounds. An IBQ accounts for bycatch mortality including any potential survivorship after capture. Currently, this is the only species managed under IBQ for the U.S. West Coast groundfish IFQ fishery. Each federal groundfish permit with a trawl endorsement is allocated IBQ pounds for *P. halibut* caught north of 40°10' N. latitude. Pacific halibut caught south of 40°10' N. latitude are not managed by an IBQ quota but are reported here under the IFQ fishery.

Data collection and reporting for this fishery is described in section 3.2.1 by gear type. The shore-based IFQ fishery includes all IFQ fishery components with the exception of at-sea motherships and catcher-processors. Motherships and catcher-processors have a bycatch quota for *P. halibut*, but it is not accounted for at the permit level.

With the exception of the IFQ fishery, *P. halibut* bycatch mortality is accounted for at the fishery sector level only. *P. halibut* is regularly caught as bycatch in the LE sablefish endorsed fixed gear, LE sablefish non-endorsed fixed gear, and OA fixed gear sectors.

3 METHODS

3.1 Data Sources

Data sources for this analysis include on-board observer data (from the WCGOP and A-SHOP), landing receipt data (referred to as fish tickets, obtained from PacFIN) and data generated from vessels carrying electronic monitoring (a.k.a. EM) equipment. Currently only vessels in the IFQ sector fishing on an exempted fishing permit (EFP) carry EM equipment. EM data are obtained from Pacific States Marine Fisheries Commission. To date, observer data is the sole source for discard estimation in the IFQ sectors, except for vessels using EM under an EFP, as stated above. All other sectors use a combination of observer and PacFIN data to estimate discard mortality. A list of fisheries, coverage priorities and data collection methods employed by WCGOP in each observed fishery can be found in the WCGOP manuals (NWFSC 2018b). A-SHOP program information, documentation and data collection methods can be found in the A-SHOP observer manual (NWFSC 2018b).

The sampling protocol employed by the WCGOP is primarily focused on the discarded portion of catch. To ensure that the recorded weights for the retained portion of the observed catch are accurate, haul-level retained catch weights recorded by observers are adjusted based on trip-level fish ticket records. This process is described in further detail on the WCGOP Data Processing webpage (NWFSC 2018a) and was conducted prior to the analyses presented in this report. All weights of *P. halibut* presented in this report are round weights, that is, whole fish. IPHC converts these weights to dressed weight (i.e., head and organs removed).

For data processing purposes, species and species groups were defined based on management (NWFSC 2018c). A complete listing of groundfish species is defined in the Pacific Coast Groundfish Fishery Management Plan (PFMC 2011).

Fish ticket landing receipts are completed by fish-buyers in each port for each delivery of fish by a vessel. Fish tickets are trip-aggregate sales receipts for market categories that may represent single or multiple species. Fish tickets are issued to fish-buyers by a state agency and must be returned to the agency for processing. Fish ticket and species-composition data are submitted by state agencies to the PacFIN regional database. Annual fish ticket landings data were retrieved from the PacFIN database (April 2017) and subsequently divided into various sectors of the groundfish fishery as indicated in Figure 1 of our online [PacFIN processing document](#) (NWFSC 2018c).

3.2 Shore-based IFQ Fishery

The methods used to report in-season IBQ estimates via the Vessel Account System (VAS) are separate from those methods used to estimate final fleet-wide *P. halibut* mortality. Methods for in-season IBQ estimation are discussed in Appendix A.3. Results obtained by methods described here resulted in fleet-wide estimates of *P. halibut* mortality that are very close to those reported by the VAS (Data not shown due to confidentiality).

3.2.1 Pacific Halibut Data Collection in the Shore-based IFQ Fishery

The WCGOP discard sampling methodologies ensure that P. halibut mortality can be estimated, regardless of the limitations imposed by the vessel, catch composition, or catch quantity. Three pieces of information are necessary to estimate P. halibut mortality (also see Table 4):

1. A count of individual P. halibut in the haul or sample
2. Actual or visual length measurements (cm)
3. A viability obtained by physical assessment of individual P. halibut using IPHC designed dichotomous keys that relate the physical condition of the fish to a viability code (NWFSC 2018b). A unique key is used for each gear type (trawl, longline, pot).

Observers could sample all or a subset of P. halibut caught in a haul/set. The proportion of P. halibut sampled is based on the number of P. halibut caught in the haul/set, the level of assistance provided by the crew, as well as other variables (e.g., physical space, weather). Sampling and assessment of P. halibut is dependent on crew assistance and cooperation. Regulations prohibit vessel crew from discarding any P. halibut without first notifying the observer. The vessel crew must comply with requests by the observer to ensure proper P. halibut sampling, including but not limited to: modifying P. halibut sorting procedure, assisting the observer by delivering the P. halibut to the observer, and modifying operations to ensure P. halibut sampling is completed. Table 4 describes the P. halibut data obtained on IFQ-permitted vessels fishing different gear types.

On vessels fishing fixed gear (pot or hook-&-line), observers must sample at least 50% of the gear per set. Actual length measurements are obtained on bottom trawl, midwater trawl, and pot vessels, but only visual length estimates are made on vessels fishing hook-&-line gear in the IFQ fishery. Visual estimates are in 10 cm increments (55-64 cm, 65-74 cm, etc.).

The crew's cooperation is vital to the observer's sampling success during hook-&-line fishing. When an observer samples for P. halibut, the crew are not permitted to shake loose or discard any P. halibut before the observer can estimate the fish length, nor can they restrict the observer's view of the line as it comes out of the water. If requested by the observer, the crew is required to physically hand individual fish to the observer or slow the gear retrieval.

Table 4: Data collected from P. halibut caught on IFQ vessels using different types of gear.

Gear	Count	Length Measurement	Viability
Bottom trawl	all in the haul	actual, all or subset	yes
Midwater trawl ¹	all in the sample	actual, all or subset	yes
Pot	all in sampled portion	actual, all or subset	yes
Hook & Line	all in sampled portion	visual, all or subset	no

¹Applies only to Catcher Processors and Mothership Catcher Vessels. Shoreside midwater trawl hauls are dumped directly into the hold and any P. halibut are delivered to the dock for discard or donation.

Viability is assessed at the point of fish release when returned to sea. On vessels using “re-suscitation boxes” or other techniques to increase the likelihood of survival, condition sampling is performed prior to the fish being returned to sea. Observations of several condition characteristics are used to assign each fish to one of three viability categories for trawl and pot gear: Excellent, Poor, or Dead (NWFSC 2016; Williams and Chen 2004). Observer field estimates of viability for P. halibut discarded in the IFQ fishery by vessels fishing bottom trawl or pot gear are used to compute the total estimated mortality of discarded P. halibut. IBQ weight (or simply IBQ) refers to the estimated mortality of discarded P. halibut, with the appropriate mortality rate applied based on viability (Tables 2 & 3).

Viability categories are used to assign mortality rates to P. halibut. Mortality rates for vessels fishing bottom trawl gear are based on mortality data collected by Hoag (1975), who found some survivorship among fish in the dead condition category. Mortality rates for vessels fishing pot gear are based on conservative assumptions of likely survival from pot-induced injuries (Williams and Wilderbuer 1995). Because of the difficulties of collecting P. halibut viability on hook & line vessels, we used a discard mortality rate (DMR) of 0.16, which represents an average of DMRs over all years for the Bering Sea/Aleutian region longline fishery (Williams 2008). Discard mortality was assumed to be 100% for all midwater trawl bycatch estimates.

Table 5: Mortality rates used for each of the condition categories (m_c) for IFQ bottom trawl vessels (Clark et al. 1992).

m_c	Rate
m_{exc}	0.20
m_{poor}	0.55
m_{dead}	0.90

Table 6: Mortality rates used for each of the condition categories (m_c) for IFQ pot gear vessels (IPHC, 2011).

m_c	Rate
m_{exc}	0.00
m_{poor}	1.00
m_{dead}	1.00

3.2.2 Shore-based IFQ fishery Bycatch Estimation

We stratified IFQ P. halibut bycatch data based on sector (shoreside non-hake groundfish, shore-side Pacific hake, at-sea Pacific hake, and LE California halibut) and gear (bottom trawl, midwater trawl, pot, hook-&-line). LE California halibut tows were separated from IFQ bottom trawl tows in

2011-12, but have been combined with IFQ bottom trawl since 2013 to maintain confidentiality. Within the shoreside non-hake groundfish sector, we further stratified using area and depth within each gear type. We maintained area and depth strata that were applied to bottom trawl, hook-&-line, and pot gear in previous reports (see Table 4 of this report for specific strata; Heery et al. 2010, Jannot et al. 2011, 2012, 2013) because prior work demonstrated that these variables were correlated with *P. halibut* bycatch (Heery et al. 2010). Observations from IFQ vessels fishing mid-water trawl gear targeting Pacific hake or other midwater target species were not post-stratified. In addition to the strata described above, we also provide bycatch estimates north and south of the groundfish management line (40°10' N. lat.) for each sector and gear type.

Despite the 100% observer coverage mandate since 2011, there were some rare occasions (e.g., observer illness, trawl net ripped) when tows or sets were either only partially sampled, not sampled or data failed quality control. In these cases, we used ratio estimators to apportion unsampled weight to *P. halibut*, within each stratum. To obtain the estimated weight of *P. halibut* (\hat{W}) when the entire haul or set was unsampled(or data failed), the unsampled discard weight, summed across unsampled hauls within the stratum, was multiplied by the ratio of the weight of *P. halibut* discard (summed across fully sampled hauls within a stratum) divided by the total discard weight of all species in all fully sampled hauls within a stratum:

$$\hat{W}_{u,s} = \sum_u x_{u,s} \times \frac{\sum w_{f,s}}{\sum x_{f,s}} \quad (1)$$

where, for each stratum:

s = stratum, which includes sector and year and could include, area, depth, gear

u = unsampled haul

f = fully sampled haul

x = weight of discarded catch

\hat{W} = estimated weight of unsampled *P. halibut* in the stratum

w = sampled weight of *P. halibut*

The unsampled weight of partially sampled hauls or sets was categorized into weight of non-IFQ species (NIFQ) or IFQ species. Unsampled IFQ species weight was further categorized into IFQ flatfish (IFQFF), IFQ rockfish (IFQRF), IFQ roundfish (IFQRD) and IFQ mixed species (IFQM). Unsampled *P. halibut* would only occur in NIFQ (south of 40°10' N. lat. only), IFQM, or IFQFF unsampled categories. Thus, those are the only categories for which *P. halibut* is estimated. IFQM included all 2017 IFQ managed species (see [76 FR 27508](#) for a listing of IFQ species). NIFQ included all species encountered that were not designated as an IFQ managed species. IFQFF included all IFQ flatfish species managed as a complex under the groundfish FMP. North of the 40°10' north latitude groundfish management line, *P. halibut* would be included in unsampled IFQFF or IFQM categories. South of the groundfish management line, *P. halibut* would only be included in the unsampled NIFQ category.

To obtain the estimated weight of *P. halibut* (\hat{W}) in partially sampled hauls or sets, the unsampled discard weight, summed across partially sampled hauls within the stratum, was multiplied by the

ratio of the weight of P. halibut (summed across fully sampled hauls within a stratum) divided by the total discard weight of all species occurring within a category (NIFQ, IFQFF, IFQM) in all fully sampled hauls within a stratum. Estimated P. halibut weight was summed across unsampled categories.

$$\hat{W}_{p,s} = \sum_y \left(\sum_p x_{p,y,s} \times \frac{\sum w_{f,s}}{\sum x_{f,y,s}} \right) \quad (2)$$

where, for each stratum:

s = stratum, which includes year and sector, and could include, area, depth, gear

y = unsampled category (either NIFQ, IFQFF, or IFQM)

p = partially sampled haul

f = fully sampled haul

x = weight of discarded catch

\hat{W} = estimated weight of unsampled P. halibut in the stratum

w = sampled weight of P. halibut

Expanded weights of P. halibut obtained using the equations above for unsampled or partially sampled hauls were then added to the sampled weight of P. halibut within each stratum to obtain the total P. halibut weight per stratum.

3.2.3 Viability Analysis

We used observer field estimates of viability for P. halibut discarded in the IFQ fishery by vessels fishing bottom or pot gear to compute the total estimated mortality of discarded P. halibut by IFQ gear/sector and stratum.

To account for the impact of fish size on survivorship, we computed a weighted mortality rate for each condition category. Length measurements associated with each viability record were converted to weight based on the IPHC length-weight table provided in Appendix A.4.

A discard mortality rate for each condition category was then computed as the proportion of P. halibut sampled weight in a viability category multiplied by the viability category-specific mortality rate (see Tables 5 and 6 above):

$$DMR_{csj} = m_c \times P_{csj} \quad (3)$$

where:

s = stratum, which could include, area, depth, gear, and sector

c = viability condition (Excellent, Poor, Dead)

j = year

m = mortality rate

P = proportion of sampled P. halibut weight (w)

DMR = discard mortality rate

Discard mortality rates for each condition category c and stratum s were then multiplied by gross

discard estimates to compute total estimated discard mortality for each gear type separately :

$$\hat{F}_{sj} = \sum_c (B_{sj} \times DMR_{sj}) \quad (4)$$

where:

s = stratum, which could include, area, depth, gear, and sector

c = viability condition (Excellent, Poor, Dead)

j = year

F = total estimated discard mortality

B = gross estimated discard weight

DMR = discard mortality rate

Viability data are collected from only a sub-sample of the P. halibut that observers encounter. Based on previous evaluations by Wallace and Hastie (2009), we expect that survivorship of P. halibut in bottom trawl tows are most directly affected by the length of the tow and the amount of catch that fills the net. These variables are not part of the bycatch ratio stratification process (above), and their use in stratifying viability data would make it difficult to then apply discard mortality rates to initial gross estimates of bycatch. We found that tow duration was directly related to depth, one of the variables used to stratify discard ratios and initial gross discard estimates for bottom trawl gear. Because depth and tow duration appeared to co-vary, we used depth and area to stratify IFQ viability data collected from bottom trawl gear. For IFQ viability data collected from pot gear, only area is used to stratify the data. For longline gear, we used a discard mortality rate of 16%, which represents an average of DMRs over all years for the Bering Sea/Aleutian region longline fishery (Williams 2008).

Final estimates of P. halibut bycatch and discard mortality are also presented in the context of the estimated mortality of legal-sized halibut. This was computed by applying the proportion of sampled P. halibut weighed in each depth stratum that was from legal-sized fish (82 cm or larger) to initial estimates. Viabilities were then applied to gross legal-sized discard estimates in the same manner as described above.

3.2.4 Length Frequencies

The length frequency distribution for P. halibut in the 2011-2017 IFQ fishery is provided in Tables 27 & 28. Pacific halibut pose unique challenges for observer sampling. Observers typically measure the length of P. halibut and then convert the measurement to weight using the IPHC length-weight conversion table (Table 9 in A.4). Occasionally, observers weigh individual fish. Sometimes crew members presort the catch by removing P. halibut and immediately return them to sea. Vessel crews presort P. halibut to increase the likelihood of survival of the discarded fish. Presorting is prevalent on vessels fishing with hook-&-line gear. Fishers have raised concerns regarding crew safety when landing large P. halibut. In addition, hook-&-line fishers are concerned that P. halibut individuals would be injured during landing because of their interaction with the vessel 'crucifier' (gear used to strip the bait and any catch off of the hook and ganglion line). Therefore, shake-offs prior to the crucifier (a form of pre-sorting) is almost universal on IFQ hook-&-line vessels. Another case of pre-sorting can occur when halibut are too heavy and/or

awkward to weigh in observer baskets. In all cases of pre-sorting, random samples are not available. Therefore, observers visually estimate the length of the halibut in ten-centimeter units (40cm, 50cm, 60cm, etc.), which are later converted to weight using the IPHC length-weight conversion table (Table 9 in Appendix A.4).

Tables 70 & 73 (Appendix A.2) provide the actual observed length frequency distributions of discarded *P. halibut* for vessels fishing IFQ using bottom trawl or pot gear. These length frequencies have been weighted based on the ratio of total estimated *P. halibut* discard weight to the weight of *P. halibut* that was measured in each stratum (see Appendix A.2 for further details). We have summarized the proportion of length measurements in each condition category (Excellent, Poor, and Dead) in Tables 71 and 74 (Appendix A.2) to inform size-specific modeling of mortality. The frequency of sampled fish within each condition category was weighed in the same manner as length frequency distributions and then summarized for each 2 cm length bin. In addition, we also provide a count of the number of dead individuals in each 2 cm length bin (Appendix A.2, Tables 79, 80 & 81). These values were obtained by multiplying the number of individuals in a length bin within a viability category, by the condition specific mortality rate (Tables 5 & 6; or 1.0 in the case of midwater trawl) and summed these values across viabilities to obtain the number of dead per length bin. This method assumes there is no size-specific mortality.

3.3 Non-nearshore Fixed Gear Fishery

The WCGOP samples each non-nearshore fixed gear sector through separate random selection processes, with the limited entry (LE) sablefish endorsed season permits receiving the highest level of coverage, then LE sablefish non-endorsed permits, and open access (OA) fixed gear the lowest. LE sablefish endorsed vessels that fish outside of the primary season or that have reached their tier quota in the primary season are not randomly chosen for observation. Given this sampling structure and anticipated differences in variance from one sector to the next, we chose to maintain sector as a stratification variable in our analysis. Testing of alternative stratification schemes (Heery et al. 2010) indicated that latitude and gear type were the most important variables with respect to *P. halibut* bycatch in the non-nearshore fixed gear groundfish fishery. Bycatch estimates were produced separately for each sector and gear combination. Two latitudinal strata were applied to the LE sablefish endorsed longline sector (north and south of Pt. Chehalis, WA = 46°53.30' N. lat.) because previous modeling demonstrated that these strata significantly improved the fit of predicted bycatch amounts to the amounts observed (Heery et al. 2010). Pt. Chehalis, WA was used in previous estimates of *P. halibut* bycatch in the LE sablefish endorsed season longline sector because of its relevance to groundfish management and its apparent ability to split out higher bycatch rates off the northern coast of Washington (Heery and Bellman 2009). Evaluations of latitudinal strata for the other fixed gear sectors did not improve the fit of models to an extent that justified their use. Thus, we maintained previous stratifications for the other groundfish fixed gear sectors (Heery and Bellman 2009, Heery et al. 2010, Jannot et al. 2011, 2012, 2013).

3.3.1 Discard Estimation

A deterministic approach was used to estimate *P. halibut* discard for all sectors of the non-nearshore groundfish fixed gear fishery. Discard ratios were computed from observer data as the discarded weight of *P. halibut* divided by the retained weight (Table 34). Retained weight varies by sector in this fishery and can be either sablefish or all FMP groundfish (except Pacific hake, see Table 33 for type of retained used; for list of FMP groundfish species, see: NWFSC 2018c). Ratio denominators were identified for each sector of the non-nearshore fixed gear fishery based on the targeting behavior of that sector. Discard ratios were then multiplied by the total sector landed weight of either sablefish or FMP groundfish (except Pacific hake), corresponding to the denominator used to compute the observed discard ratio for each sector. This provided an expanded gross estimate of *P. halibut* discard for each sector. A discard mortality rate (discussed below) was then applied to compute estimated discard mortality.

Total landed weights for each sector are obtained from fish ticket landing receipts. Fish tickets for fixed gear that included recorded weights for sablefish were included in the non-nearshore fixed gear sector. Commercial fixed gear fish tickets with recorded nearshore species weight were not used in this portion of the fixed gear analysis, regardless of whether they included recorded weights for sablefish (Figure 1 in [PacFIN Processing Document](#)). In addition, fixed gear fish tickets without recorded sablefish or nearshore species were included in the non-nearshore fixed gear sectors only if groundfish landings were greater than non-groundfish landings based on a unique vessel and landing date.

Fish tickets from the non-nearshore fixed gear sector were partitioned into the three commercial fixed-gear sectors (LE sablefish endorsed season, LE sablefish non-endorsed, and OA fixed gear) through the following process. Commercial fixed-gear fish tickets were first divided out by whether the vessel had a federal groundfish permit (limited entry) or no federal groundfish permit (open access). OA fish tickets were placed in the OA fixed gear groundfish sector. Next, LE fish tickets were separated based on whether the vessel's federal groundfish permit(s) had a sablefish endorsement with tier quota for the primary season or if it was not endorsed (also referred to as 'zero' tier). Fish tickets for all LE sablefish vessels with tier endorsements that were operating within this period and within their allotted tier quota were placed in the LE sablefish endorsed sector. If LE sablefish endorsed vessels fished outside of the primary season (November through March) or made trips within the season after they had reached their tier quota, the fish tickets were placed in the LE sablefish non-endorsed sector. In addition, fish tickets from non-endorsed LE vessels were also placed in the LE sablefish non-endorsed sector.

Further processing of fish tickets identified and removed the directed commercial *P. halibut* fishery landings from the non-nearshore fixed gear analysis. The directed *P. halibut* fishery occurs for only a few days each year, during 10-hour openings that are designated by the IPHC. LE and OA fixed gear vessels that typically target groundfish can participate in the directed fishery. For most fixed gear vessels, (other than LE sablefish endorsed vessels north of Pt. Chehalis) this is the only time during which they are allowed to land *P. halibut*. Fish tickets that included *P. halibut* landings on or within the 2 days after a directed fishery opening were considered to be part of the directed fishery and not part of the non-nearshore fixed gear fishery targeting federal FMP groundfish. These fish tickets were removed prior to our analysis. This

approach may have resulted in the removal of some non-directed fishery landings north of Pt. Chehalis, but any bias introduced by this step is considered to be extremely small given the short time period across which fish tickets were removed.

WCGOP observer data were stratified according to sector and gear type (longline and pot/trap). As previously described, one additional latitudinal stratum at Pt. Chehalis, WA (46°53.30' N. lat.) was used for the LE sablefish endorsed longline sector. Some retention of *P. halibut* was allowed in the LE sablefish endorsed season in the area north of Pt. Chehalis. The Pt. Chehalis line was the only latitudinal stratification incorporated into this portion of the analysis and was only applied to the LE sablefish endorsed sector. Discard amounts provided for the other two gear sectors represent coast-wide estimates.

The number of observed trips, sets, and vessels are summarized for each sector, gear type, and area (where applicable) (Tables 30, 31 & 32). The landed weight of sablefish and FMP groundfish (excluding Pacific hake) is used as a measure for expanding discard from observed trips to the entire fleet (Tables 33 & 34). Observed discard ratios were calculated by sector, gear type and area based on the following equation:

$$\hat{D}_s = \frac{\sum_t d_s}{\sum_t r_s} \times F_s \quad (5)$$

s = stratum, including gear, sector, gear type, and area

t = observed sets

d = observed discard (mt) of *P. halibut*

r = observed retained weight (mt) of sablefish or all FMP groundfish except Pacific hake

F = weight (mt) of retained sablefish or all FMP groundfish excluding Pacific hake recorded on fish tickets in strata s

\hat{D}_s = discard estimate for stratum s

For all strata except the LE sablefish non-endorsed longline and the OA sectors, discard ratios were calculated by dividing the stratum discard weight of *P. halibut* by the retained catch weight of sablefish. Retained groundfish was used as the ratio denominator for the LE sablefish non-endorsed longline and the OA sectors because these sectors target a wider range of groundfish species. A broader denominator was therefore necessary to effectively capture the level of fishing effort in these sectors.

Where FMP groundfish (excluding Pacific hake) was used to compute discard ratios, retained weights recorded by the observer not appearing on fish tickets were excluded from the denominator. This prevents double-counting associated with differences in the species codes used by observers and processors. For instance, while observers may record rockfish catch at the species level, various species of rockfish are often grouped, weighed, and recorded together on the fish ticket by the processor under a grouped market category, e.g., northern unspecified scope rockfish. In some cases, this difference in species coding prevents observer and fish ticket weights from being matched and adjusted properly. Species coding on fish tickets varies considerably between processors and over time, and it is not possible to make assumptions regarding which individual observer-recorded species likely coincide with species grouping codes

on fish tickets. By using only the retained groundfish weight from fish tickets in discard ratio denominators, we prevent double-counting of retained weights. This is not a factor when using a single species in the denominator, such as sablefish, as any retained weights in observer and fish ticket data that share the same species code will match and adjust properly.

The expansion factors for each fishery sector and gear type can be found in Table 34. The discard rate multiplied by the expansion factor yielded an expanded gross P. halibut discard estimate for each stratum (Table 39). If landings were made by a fixed gear sector for which there were zero or very few WCGOP observations, the most appropriate observed discard ratio was selected and applied to those landings based on similarities in the fishery management structure, fishing and discard behavior, and the gear fished. The LE sablefish endorsed vessels fishing outside of the primary season with pot gear often land a small amount of groundfish; however, this portion of the fleet is not observed by the WCGOP. Given similarities in gear type and catch composition, OA fixed gear pot observations were selected as the most appropriate source of information for an observed discard rate (Table 33).

3.3.2 Discard Mortality Rates

Once an initial gross P. halibut discard weight was estimated, this value was multiplied by a discard mortality rate (Table 39) to generate final discard mortality estimates (Tables 39 & 40, Figure 5). Discard mortality is approximated based on viabilities in a manner similar to the approach used for IFQ bottom trawl. Observers have systematically collected viability data on hook & line vessels in the Non-Nearshore Fixed Gear sector since 2011. Current methods require observers to collect a length and viability on the first 5 P. halibut observed in each set on these vessels and to ignore any injuries incurred during landing when assessing viability. For the period 2002-2010, we used a single mortality rate for all bycatch (16%) on longline and hook & line vessels, which represents an average of DMRs over all years for the Bering Sea/Aleutian region longline fishery (Williams 2008). For the period 2011-2017, we used observer field estimates of discarded P. halibut viability on Non-Nearshore Fixed Gear vessels fishing longline or hook & line gear to estimate mortality of discarded P. halibut. (Note: Observers currently do not take viability of P. halibut caught on IFQ hook & line vessels).

Methods used to calculate discard mortality based on viability condition are almost identical to those methods currently accepted for use with IFQ bottom trawl vessels (see subsection 3.2.3). To account for the impact of fish size on survivorship, we computed an annual weighted mortality rate for P. halibut in each condition category in the LE Sablefish Endorsed fishery (Table 36). For the LE Sablefish Non-Endorsed and OA Fixed Gear sectors, sample sizes were too small to calculate an annual rate. Therefore, we calculated a five year running average of weighted mortality rate for each condition category in these two sectors (Tables 37 & 38). Length measurements associated with each viability record were converted to weight based on the IPHC length-weight table provided in Appendix A.4.

The proportion of P. halibut sampled weight in a viability category multiplied by the viability category-specific mortality rate (Table 7 above):

$$DMR_{csj} = m_c \times P_{csj} \quad (6)$$

Table 7: Mortality rates used for each of the condition categories (m_c) for Non-Nearshore hook & line vessels: minor, mod = moderate, severe, dead (Trumble et al. 2000).

m_c	Rate
m_{minor}	0.035
m_{mod}	0.363
m_{severe}	0.662
m_{dead}	1.00

where:

s = stratum, which could include, area and sector
 c = viability condition (Minor, Moderate, Severe, Dead)
 j = year
 m = mortality rate
 P = proportion of sampled P. halibut weight (w)
 DMR = discard mortality rate

Discard mortality rates for each condition category c and stratum s were then multiplied by gross discard estimates to compute total estimated discard mortality for each sub-sector separately :

$$\hat{F}_{sj} = \sum_c (B_{sj} \times DMR_{sj}) \quad (7)$$

where:

s = stratum, which could include, area and sector
 c = viability condition ((Minor, Moderate, Severe, Dead)
 j = year
 F = total estimated discard mortality
 B = gross estimated discard weight
 DMR = discard mortality rate

Viabilities from pot gear would be appropriate to use in estimating discard mortality, however bycatch of P. halibut in pot gear is infrequent and the sample size is too small to utilize in this analysis. Consistent with past reports, we relied on discard mortality rates (DMR) computed for Alaska groundfish fisheries (Williams 2008). An 18% DMR was applied to estimates for pot gear, coinciding with the DMR used for the sablefish pot fishery in Alaska.

For additional context, we present the length frequency distribution of P. halibut from visual length estimates and physically measured lengths in non-nearshore fixed gear sectors (Tables 41, 42, 43, 44, & 45) and the proportion of sampled P. halibut discard of legal (>82 cm) and sub-legal (<82 cm) sizes in non-nearshore fixed gear sectors (Table 46). The majority of P. halibut lengths recorded in these fisheries were visual estimates of length, rounded to the nearest 10 cm. In

other words, specimens that are 76 cm and 82 cm are both visually estimated to be 80 cm. With this level of resolution, it was not possible to compute the exact proportion of sub-legal versus legal *P. halibut* from visually estimated lengths. Visual estimates were instead summarized in the manner in which they are recorded; with sub-legal and legal sized halibut falling within the 75-84 cm length bin.

3.4 IPHC Pacific halibut Derby Fishery

For the first time in 2017, the WCGOP observed the Pacific halibut derby fishery as a pilot study, and fleet-wide discard estimates were derived from WCGOP and fish ticket data. This fishery was defined based on using fixed gear and landing Pacific halibut within two days of the halibut derby openings (Somers et al. 2018). Prior to 2017, landings in this fishery were included in the non-groundfish fisheries not observed by the NWFSC and no estimates of discards were calculated. Effort in this fishery occurs primarily in Washington and Oregon and uses only hook-and-line gear. Gross discard and mortality estimates for *P. halibut* were computed based on the same methods as described above for the non-nearshore hook-and-line fisheries (Section 3.3). However, for the *P. halibut* derby fishery, we used Pacific halibut as the retained weight for both discard rates and expansion factors. We estimated landings, discard, and total mortality in the Pacific halibut derby fishery (Tables 47, 48, & 50). Because the gear and effort in this fishery is similar to the non-nearshore hook-and-line fisheries, the same mortality rates based on viability (Table 7) were applied to discarded *P. halibut* in the Derby fishery (Table 49). We also present the number of observed vessels, trips, and sets for each opening of the fishery (Figure 8) and the observed physical and visual length frequencies of discarded *P. halibut* (Tables 51 & 52).

3.5 Observed State Fisheries

Pacific halibut bycatch was also observed in the following state managed fisheries:

- Oregon and California nearshore groundfish fixed gear sectors (Table 53)
- Washington, Oregon, and California pink shrimp trawl fisheries (Tables 54 & 55)
- OA California halibut trawl fishery (Table 56)
- California sea cucumber trawl fishery (Table 57)
- California ridgeback prawn trawl fishery (Table 58)

Note that the LE California halibut fishery is covered under the IFQ fishery. Bycatch estimates for these fishery sectors were computed within each fishery based on the following equation:

$$\hat{B} = \frac{\sum_t b}{\sum_t r} \times F \quad (8)$$

b = observed discard (mt) of *P. halibut* on set/haul t

t = observed sets

r = observed retained weight (mt) of target species on set/haul t

F = weight (mt) of retained target species

\hat{B} = Discard estimate of P. halibut (mt)

The nearshore fixed gear fishery targets a variety of groundfish and state managed nearshore species that inhabit areas less than 50 fathoms deep. All species included in the nearshore target group, as listed in the WCGOP data processing appendix (NWFSC 2018c), were included in the denominator when calculating bycatch ratios for the nearshore fixed gear sector. Pink shrimp and California halibut were considered the target species in their respective fisheries. Discard mortality rates are not available for California halibut and pink shrimp fisheries due to a lack of information regarding survivorship. To maintain confidentiality, the Nearshore fisheries cannot be split out by gear type (hook & line vs. pot). For these reasons, we assumed 100% mortality in the Nearshore, Pink Shrimp, and CA halibut fisheries.

For the first time in 2017, the California sea cucumber trawl and the California ridgeback prawn trawl fisheries were observed by WCGOP as a pilot study. Prior to 2017, landings in these fisheries were included in non-groundfish fisheries not observed by the NWFSC and no estimates of discards were calculated. Effort in these fisheries occurs only in California, uses shrimp and bottom trawl gears, and targets sea cucumbers or ridgeback prawns. Discard estimates for each species was computed based on the same equation as described above for the OA California halibut fishery, but utilizing sea cucumber or ridgeback prawn as the retained weight for both discard rates and expansion factors. No mortality rates were applied. In the 2017, there was no observed catch of P. halibut in either the CA sea cucumber trawl fishery (Table 57) or the CA ridgeback prawn trawl fishery (Table 58).

3.6 Exempted Fishing Permits

EFPs are federal permits issued by NMFS authorizing vessels to engage in fishing operations that otherwise would be prohibited by regulation ([PFMC Council Operating Procedure 19](#)) EFPs directed toward groundfish species have been required to carry WCGOP observers on 100% of trips. Thus to obtain the catch from EFPs, we sum the at-sea discards and landed P. halibut catch.

Since 2015, vessels in the IFQ fishery could elect to participate in an electronic monitoring (EM) EFP. To obtain the catch from the IFQ EM EFP, we sum the P. halibut catch from the electronic monitoring data supplied to NWFSC Observer Program by the Pacific States Marine Fisheries Commission. Unlike the normal IFQ program, IFQ vessels fishing under an EM EFP are not required to carry an observer on every fishing trip because EM is used to ensure compliance with the IFQ program. The NWFSC Observer Program targets 30% of randomly selected IFQ EM trips for observer coverage for the purposes of scientific observation (e.g., biological sampling). A comparison of the discard mortality rates between the EM and non-EM IFQ vessels and between observer viability method versus the Time-on-deck model are presented in Appendix A.1.

3.7 Non-groundfish Fisheries Not Observed by NWFSC

Pacific halibut bycatch from non-groundfish fisheries that are not observed by the NWFSC Observer Program is recorded on fish tickets. Data from these fisheries are only available to the

the NWFSC Observer Program from PacFIN fish ticket records. We provide a summary of landed *P. halibut* from these fisheries by year.

4 RESULTS

4.1 IFQ Fishery

All participating vessels carry an observer on all fishing trips under IFQ management (100% trips observed,) except those participating in the EM EFP (see below for EM EFP results). For all 2017 strata, 99% or more of the observed IFQ tows or sets were sampled (Tables 8, 9, & 10). IFQ flatfish, IFQ mixed species, and unsorted catch all contributed to unsampled catch (Table 11; see NWFSC 2018b for IFQ sampling protocols). The total estimated weight of *P. halibut* from unsampled tows or sets in 2017 represents a small fraction (0.51 mt, or 0.7%) of the total 2017 IFQ gross discard weight of *P. halibut* (Tables 11, 12 & 13).

Gross bycatch estimates and total discard mortality estimates were largest for vessels fishing bottom trawl gear, north of the 40°10' N. latitude management line in depths greater than 60 fathoms (Table 18). This gear-area-depth stratum accounts for 89% of the 2017 *P. halibut* discard mortality in the IFQ fishery. The next largest fraction (6%) of total IFQ discard mortality was found in the same gear-area combination in shallow waters (<60 fm). Together, bottom trawl gear fishing north of the 40°10' N. latitude management line accounts for 95% of the 2017 *P. halibut* discard mortality in the IFQ fishery (Table 18).

In terms of viability, the majority of *P. halibut* on IFQ vessels were classified as either excellent or dead, depending on the stratum (Tables 14, 15, 16, & 17). In 2017, the majority of individuals caught with bottom trawl were in excellent condition in the areas north of Pt. Chehalis and north of 40°10' N. latitude, irrespective of depth, although deeper depths had more dead individuals than shallow depths in these areas (Table 14).

Estimated *P. halibut* discard mortality from all sectors and gears of the 2017 IFQ fishery is 2.2 mt greater than the average for the previous 5 years (2012-16 mean = 34.70 mt, 2017 = 36.88 mt, including IFQ EM EFP).

The 2017 IFQ estimated *P. halibut* discard mortality for all gears is 80% less than the estimated discard mortality from the 2010 LE bottom trawl fishery (Tables 2 & 65) and 83% less than the average mortality in the LE bottom trawl fishery over the years 2002-2010 (220 mt). The management change to Catch Shares in 2011 could explain this decrease in *P. halibut* catch. IBQs for *P. halibut* might have increased fisher incentives to avoid *P. halibut* bycatch and thereby changed fisher behavior (i.e., changing fishing grounds, gear, operations, or *P. halibut* handling).

Estimated bycatch weight of *P. halibut* from the At-sea hake component of the 2017 IFQ fishery increased from 2016 (2016 = 0.15 mt, 2017 = 0.55; Tables 60 & 65). There was no fishing in the Tribal sector. At-sea hake *P. halibut* length frequencies are given in Table 61.

4.2 IFQ Electronic Monitoring EFP

Estimated P. halibut discard mortality from the 2017 IFQ Electronic Monitoring Exempted Fishing Permit, including fish discarded at the dock, was 4.88 mt from bottom trawl vessels, 0.13 mt from pot vessels, and 0.46 mt from midwater trawl vessels (Table 62).

Both IFQ EM bottom trawl and IFQ EM pot vessels had slightly higher discard mortality rates (DMR) than non-EM IFQ vessels when using the observer viability method (Tables 67 & 69). However, the observer viability method on IFQ EM bottom trawl vessels appears to give a lower DMR than the Time-on-Deck model (Table 67). Caution must be used in interpreting the DMRs reported in Appendix A.1 because sample sizes were very small. The number of EM vessels catching P. halibut was a small subset of the overall EM fleet and those vessels that did catch P. halibut typically caught very few P. halibut during observer sampling (Tables 66 & 68).

4.3 Non-Nearshore Fixed Gear Fishery

The 2017 estimated discard mortality of P. halibut in the longline portion of the LE sablefish endorsed sector increased by 59% from 2016 (2016 = 15.59 mt, 2017 = 37.78 mt; Table 39) but is still well within the historical range for this fishery (2.94 - 104.45 mt; Table 39). Compared to 2016, the 2017 observed discard ratio increased north of Pt. Chehalis (Table 34). Estimated discard of P. halibut from the pot portion of the LE sablefish endorsed sector decreased by 84% compared to 2016 (2016 = 1.03 mt, 2017 = 0.16 mt; Table 39).

Discard of P. halibut among the LE sablefish non-endorsed longline vessels dropped during 2017 relative to 2016 (2016 = 0.91 mt, 2017 = 0.02 mt); pot vessels in this sector had a slight increase in P. halibut bycatch (2016 < 0.01 mt, 2017 = 0.01 mt) but remain a tiny fraction of total P. halibut discard (Table 39). P. halibut bycatch in both OA hook-and-line (2017 = 3.57 mt) and pot vessels (2017 = 0.16 mt) ticked up during 2017 but still account for only a small portion of total fixed gear bycatch.

Landings of target species increased for both LE and OA longline and hook-and-line vessels in all non-nearshore sectors in 2017 (Table 34), but observed P. halibut encounters were less than in 2016 (Table 35) even though observer coverage increased, or remained similar to, 2016 levels for these vessels (Table 31).

Physical measurements of P. halibut length frequency from the non-nearshore fixed gear sectors can be found in Tables 41, 42, 43, & 44. Visual estimates of length frequencies in the non-nearshore fixed gear sectors can be found in Table 45.

4.4 IPHC Pacific halibut Derby

The NWFSC Observer Program attained a 7% coverage rate (Table 47) in the first year of covering the IPHC P. halibut derby fishery. Observer coverage fairly evenly distributed among the three openings of the fishery in 2017 (Figure 8). The P. halibut discard ratio in this fishery was 0.19 leading to a gross discard weight of 25.7 mt (Table 48). The majority of discarded fish had only minor or moderate injuries (Table 49). Thus, despite the high discard ratio, the total discard

mortality after accounting for viability was 2.3 mt, or a realized discard mortality rate of 9%. The majority of observed P. halibut discards were less than legal-size (82 cm) although a few were above that size (Tables 51 & 52).

4.5 Observed State Fisheries, EFPs and Non-Groundfish Fisheries

Very small amounts of P. halibut bycatch were recorded in state managed observed fisheries. Even assuming 100% mortality, bycatch estimates for the nearshore groundfish fixed gear sector, pink shrimp trawl fishery, and the OA sector of the California halibut trawl fishery made up a minor portion of the 2017 total mortality estimate for P. halibut (Tables 53, 54, 55 & 56). Zero (0) catch of P. halibut was observed in the sea cucumber and ridgeback prawn fisheries of California (Tables 57 & 58).

Pacific halibut bycatch by year, from non-EM EFP vessels has been zero since 2011 (Table 63). Pacific halibut landings from non-groundfish fisheries not observed by NWFSC Observer Program were 20.05 mt in 2017. (Table 64).

5 SUMMARY & CONCLUSIONS

5.1 IFQ Fishery

- Estimated P. halibut discard mortality from the 2017 IFQ non-EM vessels was 31.41 mt and from IFQ EM vessels was 5.47 mt.
- EM vessels had higher discard mortality rates (DMR) than non-EM IFQ vessels. DMR on EM bottom trawl vessels was lower when using observer viabilities compared to the Time-on-Deck model. However samples sizes were very small, complicating interpretation.
- P. halibut discard from the at-sea Pacific hake fishery in 2017 (0.55 mt) increased relative to 2016 (0.15 mt), but remains below the historical average (2002-16: 1.18 mt).

5.2 Non-IFQ Fisheries

- The 2017 estimates of P. halibut discard mortality in the LE sablefish endorsed sector (37.95 mt) increased relative to 2016 (16.62 mt) possibly due to increases in the discard ratio and in effort, but it is not completely clear from available data. The 2017 Pacific halibut mortality estimates on LE sablefish non-endorsed vessels decreased for longline gear (0.02 mt) but increased for pot vessels (0.01 mt), relative to last year. P. halibut mortality increased relative to 2016 on OA fixed gear hook & line (3.57 mt) and pot vessels (0.16 mt). These increases were possibly due to increases in effort relative to 2016.
- In the first year of observer coverage in the IPHC P. halibut derby fishery, observer coverage was 7% and evenly distributed across the three openings. The total P. halibut discard mortality after accounting for viability was 2.3 mt.
- In the first year of observer coverage in the California sea cucumber and ridgeback prawn fisheries, zero (0) P. halibut catch was observed.

- Estimated P. halibut mortality in all other non-IFQ observed fisheries remained low relative to the IFQ and non-nearshore sectors, and were within the range observed in previous years.

6 ACKNOWLEDGEMENTS

The authors gratefully acknowledge the hard work and dedication of observers from the Northwest Fisheries Science Center Fisheries Observation Science Program, Ryan Shama for answering questions regarding WCGOP sampling strategies, Rebecca Hoch for designing the cover and all the NWFSC Observer Program staff for their hard work and dedication. We thank Su Kim of the NWFSC Scientific Communications Office for producing the infographic in this report. Comments and suggestions from our partners at the International Pacific Halibut Commission including: Ian Stewart, Claude Dykstra and Lara Erikson significantly improved this report. We thank our partners at Pacific States Marine Fisheries Commission who provide us with data from the IFQ Electronic Monitoring EFP and landings data from PacFIN and specifically Aileen Smith for providing us with the data to compare DMRs in the EM fishery.

7 REFERENCES

Clark, W. G., and S. R. Hare. 1998. Accounting for bycatch in management of the Pacific halibut fishery. *Journal of Fisheries Management* 18:809-821.

Clark, W. G., S. H. Hoag, R. J. Trumble, and G. H. Williams. 1992. Re-estimation of survival for trawl caught halibut released in different condition factors. International Pacific Halibut Commission, Report of Assessment and Research Activities 1992:197-206.

Heery, E., and M. Bellman. 2009. [Observed and Estimated Total Discard of Pacific Halibut in the 2002-2008 U.S. West Coast Groundfish Non-Nearshore Fixed Gear Fishery.](#)

Heery, E., M. Bellman, and J. Majewski. 2010. [Pacific halibut bycatch in the U.S. West Coast groundfish fishery from 2002 through 2009. West Coast Groundfish Observer Program.](#)

Hoag, S. H. 1975. Survival of Halibut Released after Capture By Trawls. International Pacific Halibut Commission, Scientific Report No. 57. 18 p.

IPHC Staff and Chastain, E. 2012. [International Pacific Halibut Commission Annual Report 2011.](#) International Pacific Halibut Commission, Seattle, WA, USA.

Jannot, J. E., A. W. Al-Humaidhi, M. A. Bellman, N. B. Riley, and J. Majewski. 2012. [Pacific halibut bycatch in the U.S. West Coast IFQ groundfish fishery \(2011\) and the non-IFQ groundfish fisheries \(2002-2011\).](#) NWFSC Observer Program.

Jannot, J. E., M. Bellman, and J. Majewski. 2011. [Pacific Halibut Bycatch in the U.S. West Coast Groundfish Fishery, 2002-2010](#). West Coast Groundfish Observer Program. NWFSC, 2725 Montlake Blvd E., Seattle, WA 98112.

Jannot, J. E., M. A. Bellman, M. Mandrup, N. Riley, and J. McVeigh. 2013. [Pacific halibut bycatch in the US West Coast groundfish fisheries \(2002-2012\)](#). NOAA Fisheries, NWFSC Observer Program, 2725 Montlake Blvd. E., Seattle, WA 98112.

Jannot, J. E., K. Somers, N. Riley, and J. McVeigh. 2014. [Pacific halibut bycatch in the US West Coast groundfish fisheries \(2002-2013\)](#). NOAA Fisheries, NWFSC Observer Program, 2725 Montlake Blvd. E., Seattle, WA 98112.

Jannot, J. E., K. Somers, N. Riley, V. Tuttle, and J. McVeigh. 2016. [Pacific halibut bycatch in the US West Coast groundfish fisheries \(2002-2016\)](#). NOAA Fisheries, NWFSC Observer Program, 2725 Montlake Blvd. E., Seattle, WA 98112.

Kaimmer, S. M., and R. J. Trumble. 1998. Injury, condition, and mortality of Pacific halibut bycatch following careful release of Pacific cod and sablefish longline fisheries. *Fisheries Research* 38:131-144.

NWFSC. 2018a. [FOS Observer Program Data Processing](#).

NWFSC. 2018b. [FOS Observer Program Manuals](#).

NWFSC. 2018c. [WCGOP Data Processing Appendix](#).

PFMC. 2011. [Pacific Groundfish Fishery Management plan for the California, Oregon, and Washington Groundfish Fishery](#).

Somers, K. A., J. E. Jannot, N. Riley, V. Tuttle, and J. McVeigh. 2016. [Estimated catch and discard of groundfish species in the 2016 U.S. west coast fisheries](#). NOAA Fisheries, NWFSC Observer Program, 2725 Montlake Blvd. E., Seattle, WA 98112.

Trumble, R. J., S. M. Krimmer, and G. H. Williams. 2000. Estimation of discard mortality rates for Pacific halibut bycatch in groundfish longline fisheries. *North American Journal of Fisheries Management* 20:931-939.

Wallace, J., and J. Hastie. 2009. Pacific halibut bycatch in IPHC Area 2A in the 2008 groundfish bottom trawl fishery. NWFSC, 2725 Montlake Blvd E., Seattle, WA 98112.

Williams, G. H. 2008. Pacific halibut discard mortality rates in the 2006 open access and CDQ groundfish fisheries, and recommendations for discard mortality rates needed for monitoring halibut bycatch in 2008 CDQ fisheries. International Pacific Halibut Commission, Seattle, WA.

Williams, G. H., and D. Chen. 2004. Pacific halibut discard mortality rates in the 1990-2002 Alaskan groundfish fisheries, with recommendations for monitoring in 2004 International Pacific Halibut

Commission. Report of Assessment and Research Activities 2003:227-244.

Williams, G. H., and T. K. Wilderbuer. 1995. Discard mortality rates of Pacific halibut bycatch: Fishery differences and trends during 1990-1993. *in* Proceedings of the International Symposium on North Pacific flatfish. Alaska Sea Grant College Program Report. University of Alaska Fairbanks.

8 TABLES

8.1 Tables: IFQ Fishery

Table 8: Number of vessels, trips, and tows observed and metric tons of sampled Pacific halibut discarded at-sea and the P. halibut landed and discarded at the dock (from PacFIN fish tickets) on IFQ bottom trawl vessels. All participating vessels carry an observer on all fishing trips under IFQ management (100% observed). For clarity, the number of hauls with unsampled catch categories is provided. Some tows are completely unsampled. See Table 62 for bottom trawl vessels fishing under the Electronic Monitoring EFP. (*) confidential data, (-) not applicable.

Area		Bottom Trawl								Unsampled categories			Sampled Rate	
Depth (fm)	Year	No. of vessels	No. of trips	No. sampled tows	No. unsampled tows	Sampled tow hours	Unsampled tow hours	Discarded at sea (mt)	Discarded at dock (mt)	IFQFF	IFQM	Non-IFQ	%tows sampled	% tow hrs. sampled
N. of Pt. Chehalis														
0-60														
	2011	13	46	296	3	804.25	11.25	7.28	0.00	2	5	10	99.00%	98.62%
	2012	14	66	312	5	662.80	6.80	4.35	0.00	0	1	10	98.42%	98.98%
	2013	11	94	448	1	1124.53	3.05	5.35	0.00	1	0	19	99.78%	99.73%
	2014	10	32	184	1	387.28	3.00	1.96	0.00	0	3	4	99.46%	99.23%
	2015	8	56	278	0	577.36	0.00	3.89	0.00	0	0	0	100.00%	100.00%
	2016	11	71	269	4	629.14	14.58	2.62	0.00	0	0	4	98.53%	97.74%
	2017	10	45	157	0	314.49	0.00	0.68	0.00	11	0	11	100.00%	100.00%
>60														
	2011	22	145	973	5	3870.62	27.73	18.07	0.01	3	8	138	99.49%	99.29%
	2012	19	167	1292	10	4933.33	39.21	28.60	0.03	0	20	58	99.23%	99.21%
	2013	17	200	1657	4	6013.21	15.70	28.90	0.14	2	3	54	99.76%	99.74%
	2014	13	147	1195	10	4834.45	32.22	24.45	0.08	0	3	19	99.17%	99.34%
	2015	16	147	1006	1	3467.43	4.50	23.76	0.01	0	4	54	99.90%	99.87%
	2016	17	137	958	1	2988.10	5.33	14.12	0.03	0	0	13	99.90%	99.82%
	2017	19	151	1033	1	3252.60	3.75	17.32	0.07	1	0	21	99.90%	99.88%
40°10' to Pt. Chehalis														
0-60														
	2011	21	139	1059	19	2004.60	36.72	9.71	0.00	12	2	65	98.24%	98.20%
	2012	21	152	947	8	1864.09	18.51	7.33	0.00	3	6	29	99.16%	99.02%
	2013	20	204	933	2	2167.95	5.25	8.31	0.00	0	8	23	99.79%	99.76%
	2014	19	198	1059	9	2391.97	30.43	9.92	0.00	0	17	29	99.16%	98.74%
	2015	15	190	1034	5	2241.72	17.33	10.00	0.00	0	8	30	99.52%	99.23%
	2016	18	180	1029	9	2603.97	32.67	7.08	0.01	0	1	25	99.13%	98.76%
	2017	25	155	619	1	1302.86	1.67	4.62	0.00	4	0	13	99.84%	99.87%
>60														
	2011	56	751	4984	28	25758.16	143.25	20.16	0.01	5	14	178	99.44%	99.45%
	2012	54	703	4450	26	23012.24	99.87	19.37	0.04	2	27	137	99.42%	99.57%
	2013	54	743	4883	15	24709.66	72.51	19.88	0.02	1	19	165	99.69%	99.71%
	2014	50	623	3783	10	19466.22	31.34	16.85	0.01	0	8	88	99.74%	99.84%
	2015	49	591	3685	4	17621.28	18.34	29.32	0.04	0	11	72	99.89%	99.90%
	2016	43	584	3523	2	16161.49	9.58	31.07	0.08	0	0	61	99.94%	99.94%
	2017	46	666	4024	5	18028.18	14.41	35.31	0.09	6	3	108	99.88%	99.92%
S. of 40°10' N. lat.														
0-60														
	2011	3	21	63	0	157.17	0.00	0.17	0.00	3	0	1	100.00%	100.00%
	2012	*	*	*	*	*	*	*	*	*	*	*	*	*
	2013†	4	56	171	0	453.42	0.00	0.03	0.00	0	0	0	100.00%	100.00%
	2014†	5	16	39	1	76.54	2.08	0.00	0.00	0	0	1	97.50%	97.35%
	2015†	5	29	75	0	143.22	0.00	0.00	0.00	0	0	0	100.00%	100.00%
	2016†	*	*	*	*	*	*	*	*	*	*	*	*	*
	2017†	*	*	*	*	*	*	*	*	*	*	*	*	*
>60														
	2011	15	240	1357	3	5838.74	12.07	0.16	0.00	3	0	34	99.78%	99.79%
	2012	13	255	1587	3	5881.45	4.08	0.75	0.00	1	1	69	99.81%	99.93%
	2013†	14	277	1727	2	6423.88	2.75	0.88	0.00	0	2	69	99.88%	99.96%
	2014†	14	277	1877	12	6318.95	50.11	0.56	0.00	1	0	35	99.36%	99.21%
	2015†	11	186	1231	3	4198.51	5.80	0.33	0.00	0	0	14	99.76%	99.86%
	2016†	7	91	616	0	1931.13	0.00	0.09	0.00	0	0	3	100.00%	100.00%
	2017†	7	63	335	1	1421.27	3.75	0.07	0.00	0	0	2	99.70%	99.74%
LE CA Halibut S. of 40°10' N. lat.														
All depths														
	2011	3	63	155	0	507.17	0.00	0.00	0.00	0	0	2	100.00%	100.00%
	2012	*	*	*	*	*	*	*	*	*	*	*	*	*

†Combined IFQ and LE CA Halibut bottom trawl.

Table 9: Number of vessels, trips, and tows observed and metric tons of sampled Pacific halibut discarded at-sea and the P. halibut landed and discarded at the dock (from PacFIN fish tickets) on IFQ midwater trawl vessels. All participating vessels carry an observer on all fishing trips under IFQ management (100% observed). For clarity the number of hauls with unsampled catch categories is provided. Some tows are completely unsampled. Note that starting in 2015, sector names have changed such that trips with P. hake comprising less than 50% of the total landings are renamed Midwater Rockfish whereas trips with P. hake greater than 50% of landings are renamed Midwater Hake. (*) confidential data, (-) not applicable.

Midwater Trawl Gear										Unsampled categories			Sampled Rate	
Sector-Area	Year	No. of vessels	No. of trips	No. sampled tows	No. unsampled tows	Sampled tow hours	Unsampled tow hours	Discarded at sea (mt)	Discarded at dock (mt)	IFQFF	IFQM	Non-IFQ	%tows sampled	% tow hrs. sampled
Non-hake Shoreside North of 40°10'														
	2011	*	*	*	*	*	*	*	*	*	*	*	*	*
	2012	4	9	35	0	72.96	0.00	0.00	0.00	0	0	0	100.00%	100.00%
	2013	6	22	77	0	137.49	0.00	0.00	0.00	0	0	1	100.00%	100.00%
	2014	9	34	133	0	268.46	0.00	0.00	0.00	0	0	0	100.00%	100.00%
Midwater Rockfish North of 40°10'														
	2015	7	43	146	0	243.97	0.00	0.00	0.00	0	0	1	100.00%	100.00%
	2016	3	13	42	0	84.07	0.00	0.00	0.00	0	0	0	100.00%	100.00%
	2017	11	133	279	0	547.16	0.00	0.03	0.01	0	0	0	100.00%	100.00%
Shoreside Hake North of 40°10'														
	2011	27	914	1715	0	3971.49	0.00	0.03	0.33	0	0	2	100.00%	100.00%
	2012	24	721	1598	0	5948.46	0.00	0.00	0.62	0	0	3	100.00%	100.00%
	2013	24	942	1732	0	4621.83	0.00	0.05	1.28	0	0	2	100.00%	100.00%
	2014	25	957	1718	1	4716.14	1.25	0.11	1.25	0	0	7	99.94%	99.97%
Midwater Hake North of 40°10'														
	2015	5	126	286	0	1159.49	0.00	0.00	0.14	0	0	3	100.00%	100.00%
	2016	4	97	207	0	652.59	0.00	0.00	0.03	0	0	0	100.00%	100.00%
	2017	4	136	242	0	612.59	0.00	0.00	0.04	0	0	0	100.00%	100.00%

Table 10: Number of vessels, trips, and sets observed and metric tons of sampled Pacific halibut discarded at-sea and the P. halibut landed and discarded at the dock (from PacFIN fish tickets) on IFQ fixed gear vessels. All participating vessels carry an observer on all fishing trips under IFQ management (100% observed). For clarity the number of hauls with unsampled catch categories is provided. Some sets are completely unsampled. Note in 2015, IFQ vessels using pot gear that fished north of Point Chehalis were all part of the Electronic Monitoring EFP (see Table 62 for summary of these vessels). (*) confidential data, (-) not applicable.

Gear								Unsampled categories			Coverage Rate
Area	Year	No. of vessels	No. of trips	No. sampled sets	No. unsampled sets	discarded at sea (mt)	discarded at dock (mt)	IFQFF	IFQM	Non-IFQ	%sets sampled
Hook and Line											
North of 40°10' N. lat.	2011	6	21	408	1	6.03	0.00	0	0	0	99.76%
South of 40°10' N. lat.	2011	6	71	212	0	0.00	0.00	0	0	1	100.00%
Coastwide	2012	8	32	506	0	14.66	0.00	0	0	0	100.00%
	2013	8	29	215	0	3.00	0.00	0	0	0	100.00%
	2014	8	31	227	32	3.43	0.00	0	0	0	87.64%
	2015	5	16	185	0	9.49	0.00	0	0	0	100.00%
	2016	5	30	351	0	6.39	0.00	0	0	0	100.00%
	2017	4	13	148	4	4.12	0.00	0	0	0	97.37%
	Pot										
North of Pt. Chehalis	2011	3	12	75	0	1.03	0.00	0	0	0	100.00%
	2012	5	45	418	0	1.27	0.00	0	0	7	100.00%
	2013	3	12	167	0	0.22	0.00	0	0	1	100.00%
Pt. Chehalis to 40°10' N. lat.	2011	8	76	719	18	2.30	0.00	0	0	1	97.56%
	2012	9	60	470	0	0.62	0.00	0	0	0	100.00%
	2013	5	40	504	0	0.76	0.00	0	0	2	100.00%
	2015	6	39	363	0	1.31	0.01	0	0	0	100.00%
South of 40°10' N. lat.	2011	11	148	737	0	0.00	0.00	0	0	2	100.00%
	2012	13	167	812	0	0.00	0.00	0	0	1	100.00%
	2013	6	41	409	0	0.00	0.00	0	0	2	100.00%
	2015	3	18	220	0	0.00	0.00	0	0	0	100.00%
Coastwide	2014	14	113	1278	0	0.32	0.00	0	0	9	100.00%
	2016	8	61	584	0	1.70	0.00	0	0	0	100.00%
	2017	6	43	573	0	1.09	0.00	0	0	0	100.00%

Table 11: Values used to calculate the expanded weight of Pacific halibut (PHLB) from each unsampled category on U.S. West Coast groundfish IFQ bottom trawl vessels by year. Unsampled catch weight could be assigned to one of four categories: IFQ flatfish species, IFQ mixed species, non-IFQ species, or unsorted (a mix of both IFQ and non-IFQ species). The sampled weight, discard ratio, unsampled weight and estimated P. halibut gross at-sea discard are presented within each category, as a function of sector, management area, depth, and area north or south of Pt. Chehalis, WA. The sum of expanded weight is the sum of the estimated gross P. halibut discard across categories. The sampled discarded PHLB weight is the sum of sampled PHLB. The total discard (gross) is the sum of the PHLB in unsampled hauls plus the sampled PHLB. All weights are metric tons (mt). (*) confidential data. Note that adding values across columns might give slightly different results because values are rounded to two decimals for reporting.

Bottom Trawl																					
Area	Depth (fm)	Year	IFQ Flatfish				Mixed IFQ species				Non-IFQ Species				Unsorted				Sum of Exp. Discard Weight	Samp. Discarded PHLB	Total Discard
			Samp. Weight	Discarc Ratio	Unsamp Weight	Est. Dis-card	Samp. Weight	Discarc Ratio	Unsamp Weight	Est. Dis-card	Samp. Weight	Discarc Ratio	Unsamp Weight	Est. Dis-card	Samp. Weight	Discarc Ratio	Unsamp Weight	Est. Dis-card			
N. of Pt. Chehalis																					
0-60																					
	2011	60.53	0.12	0.16	0.02	80.81	0.09	5.22	0.48	55.65	0.00	2.66	0.00	136.46	0.05	2.29	0.12	0.62	7.36	7.98	
	2012	50.77	0.09	0.00	0.00	56.29	0.08	0.05	0.00	45.51	0.00	1.02	0.00	101.80	0.05	0.56	0.03	0.03	4.77	4.80	
	2013	104.68	0.05	0.07	0.00	114.61	0.05	0.00	0.00	92.99	0.00	2.00	0.00	207.60	0.03	0.91	0.02	0.03	5.43	5.46	
	2014	26.44	0.07	0.00	0.00	32.70	0.06	1.62	0.10	27.58	0.00	0.85	0.00	60.28	0.03	0.02	0.00	0.10	1.97	2.07	
	2015	32.67	0.12	0.00	0.00	38.76	0.10	0.00	0.00	32.66	0.00	0.00	0.00	71.42	0.05	0.00	0.00	0.00	3.89	3.89	
	2016	43.01	0.06	0.00	0.00	57.97	0.05	0.00	0.00	64.58	0.00	0.78	0.00	122.55	0.02	2.56	0.06	0.06	2.71	2.76	
	2017	9.87	0.07	2.32	0.16	16.37	0.04	0.00	0.00	21.23	0.00	0.29	0.00	37.60	0.02	0.00	0.00	0.16	0.68	0.84	
>60																					
	2011	114.16	0.19	1.03	0.20	142.47	0.15	1.01	0.15	207.64	0.00	15.03	0.00	350.11	0.06	4.79	0.30	0.64	21.65	22.29	
	2012	84.84	0.36	0.00	0.00	122.87	0.25	2.42	0.59	268.93	0.00	6.84	0.00	391.80	0.08	24.85	1.90	2.49	30.18	32.67	
	2013	185.79	0.16	0.20	0.03	227.34	0.13	1.07	0.14	241.41	0.00	5.38	0.00	468.75	0.06	1.39	0.08	0.25	29.66	29.91	
	2014	192.81	0.13	0.00	0.00	233.86	0.11	0.87	0.09	293.94	0.00	1.81	0.00	527.80	0.05	29.12	0.61	0.70	24.88	25.58	
	2015	108.65	0.22	0.00	0.00	134.93	0.18	2.84	0.51	129.24	0.00	3.76	0.00	264.18	0.09	0.16	0.01	0.53	24.34	24.86	
	2016	114.43	0.12	0.00	0.00	157.47	0.09	0.00	0.00	204.62	0.00	0.94	0.00	362.08	0.04	0.45	0.00	0.00	14.14	14.14	
	2017	140.48	0.12	0.02	0.00	193.42	0.09	0.00	0.00	219.66	0.00	2.74	0.00	413.08	0.04	0.08	0.00	0.01	17.41	17.41	
40°10' to Pt. Chehalis																					
0-60																					
	2011	96.63	0.11	0.97	0.11	117.73	0.09	2.40	0.21	188.16	0.00	6.76	0.00	305.90	0.03	5.71	0.20	0.52	10.48	11.00	
	2012	72.35	0.11	0.45	0.05	86.10	0.09	2.35	0.21	142.99	0.00	2.56	0.00	229.09	0.03	1.95	0.07	0.33	7.73	8.06	
	2013	109.66	0.08	0.00	0.00	120.95	0.07	0.86	0.06	138.52	0.00	1.84	0.00	259.47	0.03	0.41	0.01	0.07	8.47	8.55	
	2014	176.72	0.06	0.00	0.00	194.49	0.05	6.19	0.32	204.19	0.00	4.48	0.00	398.67	0.03	13.96	0.11	0.42	10.05	10.48	
	2015	158.17	0.06	0.00	0.00	192.63	0.05	0.35	0.02	193.08	0.00	2.01	0.00	385.71	0.03	1.71	0.05	0.06	10.16	10.22	
	2016	203.22	0.04	0.00	0.00	258.27	0.03	0.05	0.00	217.24	0.00	3.05	0.00	475.52	0.02	6.73	0.10	0.10	7.22	7.32	
	2017	75.56	0.06	1.21	0.07	108.64	0.04	0.00	0.00	120.00	0.00	0.97	0.00	228.64	0.02	0.91	0.00	0.07	4.63	4.71	
>60																					
	2011	190.48	0.12	0.78	0.09	352.51	0.06	4.00	0.25	753.78	0.00	18.25	0.00	1106.30	0.02	7.54	0.15	0.49	22.02	22.51	
	2012	180.33	0.11	0.06	0.01	369.70	0.05	6.92	0.37	641.16	0.00	12.38	0.00	1010.86	0.02	7.26	0.14	0.52	19.94	20.46	
	2013	229.39	0.09	0.07	0.01	401.78	0.05	9.72	0.49	709.89	0.00	11.56	0.00	1111.67	0.02	9.68	0.14	0.63	20.44	21.08	
	2014	335.57	0.05	0.00	0.00	501.04	0.03	3.02	0.10	506.94	0.00	4.08	0.00	1007.97	0.02	10.38	0.06	0.15	16.96	17.11	
	2015	323.15	0.09	0.00	0.00	466.22	0.06	0.93	0.06	548.36	0.00	4.48	0.00	1014.58	0.03	2.95	0.02	0.08	29.67	29.75	
	2016	289.10	0.11	0.00	0.00	531.95	0.06	0.00	0.00	471.32	0.00	8.21	0.00	1003.27	0.03	4.54	0.09	0.09	31.39	31.48	
	2017	373.00	0.10	4.35	0.42	706.32	0.05	0.73	0.04	503.13	0.00	6.54	0.00	1209.45	0.03	4.06	0.05	0.49	35.68	36.17	
S. of 40°10' N. lat.																					
0-60																					
	2011	4.60	0.00	0.04	0.00	5.04	0.00	0.00	0.00	11.75	0.01	0.01	0.00	16.79	0.01	0.00	0.00	0.00	0.17	0.17	
	2012	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
	2013	4.55	0.00	0.00	0.00	6.65	0.00	0.00	0.00	66.93	0.00	0.00	0.00	73.58	0.00	0.00	0.00	0.00	0.03	0.03	
	2014	0.86	0.00	0.00	0.00	2.38	0.00	0.00	0.00	4.45	0.00	0.45	0.00	6.84	0.00	0.02	0.00	0.00	0.00	0.00	
	2015	6.11	0.00	0.00	0.00	17.97	0.00	0.00	0.00	7.47	0.00	0.00	0.00	25.44	0.00	0.00	0.00	0.00	0.00	0.00	
	2016	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
	2017	3.25	0.00	0.00	0.00	3.37	0.00	0.00	0.00	2.49	0.00	0.00	0.00	5.86	0.00	0.00	0.00	0.00	0.00	0.00	
>60																					
	2011	155.01	0.00	0.10	0.00	275.06	0.00	0.00	0.00	223.70	0.00	2.86	0.00	498.76	0.00	1.36	0.00	0.00	0.16	0.16	
	2012	80.42	0.00	0.01	0.00	266.50	0.00	0.03	0.00	222.92	0.00	7.14	0.03	489.41	0.00	1.93	0.00	0.03	0.81	0.84	
	2013	119.64	0.00	0.00	0.00	364.86	0.00	0.07	0.00	296.89	0.00	7.47	0.02	661.75	0.00	0.23	0.00	0.02	0.88	0.90	
	2014	169.03	0.00	0.03	0.00	363.29	0.00	0.00	0.00	341.56	0.00	1.22	0.00	704.84	0.00	5.64	0.00	0.01	0.56	0.57	
	2015	93.62	0.00	0.00	0.00	233.85	0.00	0.00	0.00	173.60	0.00	0.44	0.00	407.45	0.00	12.71	0.00	0.00	0.33	0.33	
	2016	48.17	0.00	0.00	0.00	99.67	0.00	0.00	0.00	88.29	0.00	0.11	0.00	187.96	0.00	0.00	0.00	0.00	0.10	0.10	
	2017	20.81	0.00	0.00	0.00	46.21	0.00	0.00	0.00	49.36	0.00	0.03	0.00	95.57	0.00	0.05	0.00	0.00	0.07	0.07	
LE CA Halibut S. of 40°10' N. lat.																					
All depths																					
	2011	0.73	0.00	0.00	0.00	0.74	0.00	0.00	0.00	75.42	0.00	0.01	0.00	76.16	0.00	0.00	0.00	0.00	0.00	0.00	
	2012	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*

‡Combined IFQ and LE CA Halibut bottom trawl.

Table 12: Values used to calculate the expanded weight of Pacific halibut (PHLB) from each unsampled category on U.S. West Coast groundfish IFQ midwater trawl vessels by year. Unsampled catch weight could be assigned to one of four categories: IFQ flatfish species, IFQ mixed species, non-IFQ species, or unsorted (a mix of both IFQ and non-IFQ species). The sampled weight, discard ratio, unsampled weight and estimated P. halibut gross at-sea discard are presented within each category, as a function of sector. All midwater trawling occurs north of 40°10' and all depths are included in the summaries. The sum of expanded weight is the sum of the estimated gross P. halibut discard across categories. The sampled discarded PHLB weight is the sum of sampled PHLB. The total discard (gross) is the sum of the PHLB in unsampled hauls plus the sampled PHLB. All weights are metric tons (mt). (*) confidential data.

Area		Midwater Trawl																		
Depth (fm)	Year	IFQ Flatfish				Mixed IFQ species				Non-IFQ Species				Unsorted				Sum of Exp. Discard Weight	Samp. Discarded PHLB	Total Discard
		Samp. Weig	Discard Ratio	Unsamp Weight	Est. Discard	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Discard	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Discard	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Discard			
Non-hake Shoreside																				
	2011	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
	2012	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00
	2013	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.05	0.00	0.02	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00
	2014	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.73	0.00	0.00	0.00	0.75	0.00	0.00	0.00	0.00	0.00	0.00
Midwater Rockfish																				
	2015	0.00	0.00	0.00	0.00	3.79	0.00	0.00	0.00	16.37	0.00	0.05	0.00	20.15	0.00	0.00	0.00	0.00	0.00	0.00
	2016	0.09	0.00	0.00	0.00	2.36	0.00	0.00	0.00	0.01	0.00	0.00	0.00	2.37	0.00	0.00	0.00	0.00	0.00	0.00
	2017	0.03	0.98	0.00	0.00	17.00	0.00	0.00	0.00	2.55	0.00	0.00	0.00	19.56	0.00	0.00	0.00	0.00	0.03	0.03
Shoreside Hake																				
	2011	0.03	0.99	0.00	0.00	521.49	0.00	0.00	0.00	3.82	0.00	1.37	0.00	525.31	0.00	0.00	0.00	0.00	0.03	0.03
	2012	0.00	0.00	0.00	0.00	128.31	0.00	0.00	0.00	8.19	0.00	0.36	0.00	136.50	0.00	0.00	0.00	0.00	0.00	0.00
	2013	0.05	1.00	0.00	0.00	460.78	0.00	0.00	0.00	7.24	0.00	0.27	0.00	468.03	0.00	0.00	0.00	0.00	0.05	0.05
	2014	0.16	0.71	0.00	0.00	498.24	0.00	0.00	0.00	13.04	0.00	0.23	0.00	511.28	0.00	0.05	0.00	0.00	0.11	0.11
Midwater Hake																				
	2015	0.00	0.00	0.00	0.00	43.76	0.00	0.00	0.00	4.47	0.00	0.12	0.00	48.23	0.00	0.00	0.00	0.00	0.00	0.00
	2016	0.00	0.00	0.00	0.00	59.29	0.00	0.00	0.00	1.18	0.00	0.00	0.00	60.47	0.00	0.00	0.00	0.00	0.00	0.00
	2017	0.00	0.00	0.00	0.00	110.53	0.00	0.00	0.00	1.50	0.00	0.00	0.00	112.03	0.00	0.00	0.00	0.00	0.00	0.00

Table 13: Values used to calculate the expanded weight of Pacific halibut (PHLB) from each unsampled category on U.S. West Coast groundfish IFQ fixed gear vessels by year. Unsampled catch weight could be assigned to one of four categories: IFQ flatfish species, IFQ mixed species, non-IFQ species, or unsorted (a mix of both IFQ and non-IFQ species). The sampled weight, discard ratio, unsampled weight and estimated P. halibut gross at-sea discard are presented within each category, as a function of gear, management area, and, for pot gear, by areas north and south of Point Chehalis, WA. All depths fished are included in the summaries. The sum of expanded weight is the sum of the estimated gross P. halibut discard across categories. The sampled discarded PHLB weight is the sum of sampled PHLB. The total discard (gross) is the sum of the PHLB in unsampled sets plus the sampled PHLB. All weights are metric tons (mt). (*) confidential data.

Area	Depth (fm)	Year	IFQ Flatfish				Mixed IFQ species				Non-IFQ Species				Unsorted				Sum of Exp. Discard Weight	Samp. Discarded PHLB	Total Discard	
			Samp. Weigh	Discard Ratio	Unsamp Weight	Est. Dis- card	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Dis- card	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Dis- card	Samp. Weight	Discard Ratio	Unsamp Weight	Est. Dis- card				
Hook and Line																						
North of 40°10' N. lat.		2011	7.19	0.84	0.00	0.00	22.01	0.28	0.00	0.00	56.74	0.00	0.00	0.00	78.76	0.08	0.00	0.00	0.00	6.06	6.06	
	South of 40°10' N. lat.	2011	0.18	0.00	0.00	0.00	3.72	0.00	0.00	0.00	21.06	0.00	0.00	0.00	24.78	0.00	0.00	0.00	0.00	0.00	0.00	
		Coastwide	2012	19.31	0.76	0.00	0.00	36.87	0.40	0.00	0.00	97.36	0.00	0.00	0.00	134.24	0.11	0.00	0.00	0.00	14.66	14.66
			2013	5.10	0.59	0.00	0.00	8.29	0.36	0.00	0.00	27.60	0.00	0.00	0.00	35.88	0.08	0.00	0.00	0.00	3.00	3.00
			2014	5.37	0.64	0.00	0.00	8.41	0.41	0.00	0.00	35.36	0.00	0.00	0.00	43.76	0.08	9.85	0.38	0.38	3.43	3.80
			2015	10.76	0.88	0.00	0.00	16.49	0.58	0.00	0.00	38.39	0.00	0.00	0.00	54.88	0.17	0.00	0.00	0.00	9.49	9.49
			2016	8.69	0.74	0.00	0.00	18.97	0.34	0.00	0.00	61.15	0.00	0.00	0.00	80.13	0.08	0.00	0.00	0.00	6.39	6.39
			2017	7.59	0.54	0.00	0.00	9.62	0.43	0.00	0.00	17.52	0.00	0.00	0.00	27.13	0.15	0.49	0.02	0.02	4.12	4.14
			Pot																			
North of Pt. Chehalis	2011	1.05	0.98	0.00	0.00	1.56	0.66	0.00	0.00	0.26	0.00	0.00	0.00	1.82	0.57	0.00	0.00	0.00	1.03	1.03		
	2012	2.46	0.52	0.00	0.00	9.15	0.14	0.00	0.00	2.27	0.00	0.01	0.00	11.42	0.11	0.00	0.00	0.00	1.27	1.27		
	2013	0.28	0.79	0.00	0.00	1.08	0.20	0.00	0.00	0.66	0.00	0.01	0.00	1.73	0.13	0.00	0.00	0.00	0.22	0.22		
	Pt. Chehalis to 40°10' N. lat.	2011	2.45	0.94	0.00	0.00	7.92	0.29	0.00	0.00	3.37	0.00	0.00	0.00	11.29	0.20	3.18	0.02	0.02	2.30	2.33	
2012		1.22	0.51	0.00	0.00	3.86	0.16	0.00	0.00	6.03	0.00	0.00	0.00	9.88	0.06	0.00	0.00	0.00	0.62	0.62		
2013		1.23	0.62	0.00	0.00	6.77	0.11	0.00	0.00	10.90	0.00	0.00	0.00	17.67	0.04	0.00	0.00	0.00	0.76	0.76		
2015		1.78	0.74	0.00	0.00	7.90	0.17	0.00	0.00	7.52	0.00	0.00	0.00	15.42	0.08	0.00	0.00	0.00	1.31	1.31		
South of 40°10' N. lat.		2011	0.30	0.00	0.00	0.00	6.49	0.00	0.00	0.00	6.91	0.00	0.00	0.00	13.41	0.00	0.00	0.00	0.00	0.00	0.00	
	2012	0.52	0.00	0.00	0.00	4.21	0.00	0.00	0.00	4.59	0.00	0.00	0.00	8.80	0.00	0.00	0.00	0.00	0.00	0.00		
	2013	0.03	0.00	0.00	0.00	3.01	0.00	0.00	0.00	3.62	0.00	0.00	0.00	6.64	0.00	0.00	0.00	0.00	0.00	0.00		
	2015	0.01	0.00	0.00	0.00	1.46	0.00	0.00	0.00	8.65	0.00	0.00	0.00	10.12	0.00	0.00	0.00	0.00	0.00	0.00		
	Coastwide	2014	0.58	0.55	0.00	0.00	11.53	0.03	0.00	0.00	16.58	0.00	0.01	0.00	28.11	0.01	0.00	0.00	0.00	0.32	0.32	
2016		2.20	0.77	0.00	0.00	6.54	0.26	0.00	0.00	5.50	0.00	0.00	0.00	12.04	0.14	0.00	0.00	0.00	1.70	1.70		
2017		1.91	0.57	0.00	0.00	7.30	0.15	0.00	0.00	7.92	0.00	0.00	0.00	15.23	0.07	0.00	0.00	0.00	1.09	1.09		

Table 14: Pacific halibut viabilities in the U.S. West Coast groundfish IFQ bottom trawl fishery by management area, depth, and year. The condition of sampled P. halibut was identified as Excellent (Exc), Poor, or Dead (see Appendices in WCGOP manual), consistent with IPHC protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. In addition, all years combined are also shown. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate provided, see text for explanation.

Bottom Trawl							
Area, Depth (fm)	Number				Weighted Percentages		
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead
North of Pt. Chehalis, 0-60							
2011	517	137	308	962	57.34%	14.21%	28.45%
2012	314	156	299	769	45.94%	20.28%	33.78%
2013	327	114	464	905	41.06%	13.61%	45.33%
2014	252	27	26	305	85.12%	8.02%	6.86%
2015	349	51	90	490	71.79%	12.54%	15.67%
2016	242	61	89	392	66.54%	14.69%	18.76%
2017	60	16	12	88	74.62%	18.98%	6.40%
All	2061	562	1288	3911	57.88%	14.64%	27.49%
North of Pt. Chehalis, 60+							
2011	1063	439	927	2429	46.75%	18.24%	35.01%
2012	1299	709	1368	3376	40.36%	20.82%	38.82%
2013	2100	534	984	3618	62.12%	14.22%	23.65%
2014	1669	595	1055	3319	52.59%	16.97%	30.43%
2015	1529	404	679	2612	59.53%	14.33%	26.14%
2016	837	326	630	1793	47.79%	16.90%	35.30%
2017	1057	327	890	2274	49.07%	13.35%	37.58%
All	9554	3334	6533	19421	51.40%	16.57%	32.03%
Pt. Chehalis to 40 deg. 10' N. lat., 0-60							
2011	1076	169	199	1444	80.30%	9.53%	10.17%
2012	791	174	229	1194	67.70%	13.85%	18.45%
2013	659	238	260	1157	59.12%	21.69%	19.19%
2014	1095	229	307	1631	68.69%	13.72%	17.59%
2015	778	232	426	1436	59.35%	15.05%	25.60%
2016	525	137	447	1109	49.51%	11.91%	38.58%
2017	395	42	150	587	72.18%	6.60%	21.22%
All	5319	1221	2018	8558	65.54%	13.55%	20.91%
Pt. Chehalis to 40 deg. 10' N. lat., 60+							

Table 14: Pacific halibut viabilities in the U.S. West Coast groundfish IFQ bottom trawl fishery by management area, depth, and year. The condition of sampled P. halibut was identified as Excellent (Exc), Poor, or Dead (see Appendices in WCGOP manual), consistent with IPHC protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. In addition, all years combined are also shown. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate provided, see text for explanation. *(continued)*

Bottom Trawl							
Area, Depth (fm)	Number				Weighted Percentages		
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead
2011	967	554	1188	2709	37.57%	20.22%	42.22%
2012	850	446	1201	2497	35.47%	17.55%	46.97%
2013	753	404	1100	2257	34.57%	18.55%	46.88%
2014	765	363	865	1993	42.04%	17.22%	40.74%
2015	1402	556	1513	3471	41.39%	17.07%	41.54%
2016	1319	515	1813	3647	38.09%	13.80%	48.11%
2017	1648	599	1575	3822	46.06%	15.04%	38.90%
All	7704	3437	9255	20396	39.93%	16.68%	43.39%
South of 40 deg. 10' N. lat., 0-60							
2011	0	0	10	10	0.00%	0.00%	100.00%
2012	*	*	*	*	*	*	*
2013‡	2	0	0	2	100.00%	0.00%	0.00%
2014‡	0	0	0	0	0.00%	0.00%	0.00%
2015‡	0	0	0	0	0.00%	0.00%	0.00%
2016‡	*	*	*	*	*	*	*
2017‡	0	0	0	0	0.00%	0.00%	0.00%
All‡	*	*	*	*	*	*	*
South of 40 deg. 10' N. lat., 60+							
2011	7	1	6	14	48.21%	6.06%	45.73%
2012	35	7	36	78	49.26%	9.18%	41.56%
2013‡	27	14	51	92	32.05%	16.05%	51.90%
2014‡	24	9	14	47	63.47%	13.76%	22.76%
2015‡	10	3	15	28	54.15%	9.94%	35.91%
2016‡	6	4	1	11	73.40%	22.25%	4.35%
2017‡	3	1	2	6	55.70%	22.93%	21.37%
All‡	112	39	125	276	48.32%	12.87%	38.80%
LE CA Halibut South of 40 deg. 10' N. lat., all depths							
2011	0	0	0	0	0.00%	0.00%	0.00%
2012	*	*	*	*	*	*	*

Table 14: Pacific halibut viabilities in the U.S. West Coast groundfish IFQ bottom trawl fishery by management area, depth, and year. The condition of sampled P. halibut was identified as Excellent (Exc), Poor, or Dead (see Appendices in WCGOP manual), consistent with IPHC protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. In addition, all years combined are also shown. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate provided, see text for explanation. *(continued)*

Bottom Trawl							
Area, Depth (fm)	Number				Weighted Percentages		
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead
All	*	*	*	*	*	*	*

Table 15: Pacific halibut viabilities in the U.S. West Coast groundfish IFQ midwater trawl fishery by management area and year. The condition of sampled P. halibut was identified as Excellent (Exc), Poor, or Dead (see Appendices in WC-GOP manual), consistent with IPHC protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. In addition, all years combined are also shown. (*) confidential data, (-) no estimate provided, see text for explanation.

Midwater Trawl							
Area	Number				Weighted Percentages		
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead
Catch Shares North of 40 deg. 10' N. lat.							
2011	*	*	*	*	*	*	*
2012	0	0	0	0	0.00%	0.00%	0.00%
2013	0	0	0	0	0.00%	0.00%	0.00%
2014	0	0	0	0	0.00%	0.00%	0.00%
All	*	*	*	*	*	*	*
Midwater Hake North of 40 deg. 10' N. lat.							
2015	0	0	0	0	0.00%	0.00%	0.00%
2016	0	0	0	0	0.00%	0.00%	0.00%
2017	3	0	2	5	67.97%	0.00%	32.03%
All	3	0	2	5	67.97%	0.00%	32.03%
Shoreside Hake North of 40 deg. 10' N. lat.							
2011	0	1	2	3	0.00%	46.01%	53.99%
2012	0	0	0	0	0.00%	0.00%	0.00%
2013	2	0	1	3	91.76%	0.00%	8.24%
2014	6	2	0	8	89.99%	10.01%	0.00%
All	8	3	3	14	78.15%	12.12%	9.73%
Midwater Rockfish North of 40 deg. 10' N. lat.							
2015	0	0	0	0	0.00%	0.00%	0.00%
2016	0	0	0	0	0.00%	0.00%	0.00%
2017	0	0	0	0	0.00%	0.00%	0.00%
All	0	0	0	0	0.00%	0.00%	0.00%

Table 16: Pacific halibut viabilities in the U.S. West Coast groundfish IFQ pot fishery by management area and year. The condition of sampled P. halibut was identified as Excellent (Exc), Poor, or Dead (see Appendices in WCGOP manual), consistent with IPHC protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. In addition, all years combined are also shown. (*) confidential data, (-) no estimate provided, see text for explanation.

Pot							
Area	Number				Weighted Percentages		
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead
North of Pt. Chehalis.							
2011	53	3	19	75	83.58%	2.14%	14.27%
2012	103	21	24	148	66.34%	16.72%	16.94%
2013	18	1	11	30	60.78%	1.83%	37.39%
All	174	25	54	253	71.77%	10.11%	18.12%
Pt. Chehalis to 40 deg. 10' N. lat.							
2011	149	10	65	224	69.06%	4.57%	26.37%
2012	58	4	3	65	86.97%	7.77%	5.27%
2013	76	7	8	91	83.18%	6.94%	9.88%
2015	145	7	17	169	83.65%	4.47%	11.88%
All	428	28	93	549	77.71%	5.29%	17.00%
South of 40 deg. 10' N. lat.							
2011	0	0	0	0	0.00%	0.00%	0.00%
2012	0	0	0	0	0.00%	0.00%	0.00%
2013	0	0	0	0	0.00%	0.00%	0.00%
2015	0	0	0	0	0.00%	0.00%	0.00%
All	0	0	0	0	0.00%	0.00%	0.00%
Coastwide							
2014	24	0	8	32	73.71%	0.00%	26.29%
2016	195	3	13	211	90.04%	1.73%	8.23%
2017	52	40	27	119	40.69%	35.31%	24.00%
All	271	43	48	362	70.92%	13.35%	15.73%

Table 17: Pacific halibut caught in the U.S. West Coast groundfish IFQ hook and line fishery by management area and year. The viabilities are not currently obtained on IFQ vessels. (*) confidential data, (-) no data collected

Hook and Line								
Area	Number				Weighted Percentages			
Year	Exc	Poor	Dead	Total	Exc	Poor	Dead	
North of 40 deg. 10' N. lat.								
2011	-	-	-	902	-	-	-	
All	-	-	-	902	-	-	-	
South of 40 deg. 10' N. lat.								
2011	-	-	-	0	-	-	-	
All	-	-	-	0	-	-	-	
Coastwide								
2012	-	-	-	1271	-	-	-	
2013	-	-	-	404	-	-	-	
2014	-	-	-	698	-	-	-	
2015	-	-	-	963	-	-	-	
2016	-	-	-	672	-	-	-	
2017	-	-	-	690	-	-	-	
All	-	-	-	4698	-	-	-	

Table 18: Estimated Pacific halibut gross at-sea discard (mt) and at-sea discard mortality (mt) in the U.S. West Coast groundfish IFQ bottom trawl fishery by management area, depth, and year. Estimates were allocated to three condition categories based on information presented in Table 14. DMR=Discard Mortality Rate. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate, see text for explanation.

Bottom Trawl									
Area, Depth (fm)	Estimated Gross Discard (mt)				Estimated Discard Mortality				DMR
Year	Exc	Poor	Dead	Total	m(Exc)	m(Poor)	m(Dead)	Total	
North of Pt. Chehalis, 0-60									
2011	4.58	1.13	2.27	7.98	0.92	0.62	2.04	3.58	0.45
2012	2.21	0.97	1.62	4.80	0.44	0.54	1.46	2.44	0.51
2013	2.24	0.74	2.47	5.46	0.45	0.41	2.23	3.08	0.56
2014	1.76	0.17	0.14	2.07	0.35	0.09	0.13	0.57	0.28
2015	2.79	0.49	0.61	3.89	0.56	0.27	0.55	1.37	0.35
2016	1.84	0.41	0.52	2.76	0.37	0.22	0.47	1.06	0.38
2017	0.62	0.16	0.05	0.84	0.12	0.09	0.05	0.26	0.31
North of Pt. Chehalis, 60+									
2011	10.42	4.07	7.80	22.29	2.08	2.24	7.02	11.34	0.51
2012	13.19	6.80	12.68	32.67	2.64	3.74	11.42	17.79	0.54
2013	18.58	4.25	7.07	29.91	3.72	2.34	6.37	12.42	0.42
2014	13.45	4.34	7.78	25.58	2.69	2.39	7.00	12.08	0.47
2015	14.80	3.56	6.50	24.86	2.96	1.96	5.85	10.77	0.43
2016	6.76	2.39	4.99	14.14	1.35	1.31	4.49	7.16	0.51
2017	8.54	2.33	6.54	17.41	1.71	1.28	5.89	8.88	0.51
Pt. Chehalis to 40 deg. 10' N. lat., 0-60									
2011	8.83	1.05	1.12	11.00	1.77	0.58	1.01	3.35	0.30
2012	5.46	1.12	1.49	8.06	1.09	0.61	1.34	3.04	0.38
2013	5.05	1.85	1.64	8.55	1.01	1.02	1.48	3.51	0.41
2014	7.20	1.44	1.84	10.48	1.44	0.79	1.66	3.89	0.37
2015	6.07	1.54	2.62	10.22	1.21	0.85	2.35	4.41	0.43
2016	3.62	0.87	2.82	7.32	0.72	0.48	2.54	3.75	0.51
2017	3.40	0.31	1.00	4.71	0.68	0.17	0.90	1.75	0.37
Pt. Chehalis to 40 deg. 10' N. lat., 60+									
2011	8.46	4.55	9.50	22.51	1.69	2.50	8.55	12.75	0.57
2012	7.26	3.59	9.61	20.46	1.45	1.98	8.65	12.08	0.59
2013	7.29	3.91	9.88	21.08	1.46	2.15	8.89	12.50	0.59
2014	7.19	2.95	6.97	17.11	1.44	1.62	6.28	9.33	0.55
2015	12.31	5.08	12.36	29.75	2.46	2.79	11.12	16.38	0.55

Table 18: Estimated Pacific halibut gross at-sea discard (mt) and at-sea discard mortality (mt) in the U.S. West Coast groundfish IFQ bottom trawl fishery by management area, depth, and year. Estimates were allocated to three condition categories based on information presented in Table 14. DMR=Discard Mortality Rate. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate, see text for explanation. *(continued)*

Bottom Trawl									
Area, Depth (fm)	Estimated Gross Discard (mt)				Estimated Discard Mortality				DMR
Year	Exc	Poor	Dead	Total	m(Exc)	m(Poor)	m(Dead)	Total	
2016	11.99	4.34	15.14	31.48	2.40	2.39	13.63	18.42	0.59
2017	16.66	5.44	14.07	36.17	3.33	2.99	12.66	18.99	0.52
South of 40 deg. 10' N. lat., 0-60									
2011	0.00	0.00	0.17	0.17	0.00	0.00	0.15	0.15	0.90
2012	*	*	*	*	*	*	*	*	*
2013‡	0.03	0.00	0.00	0.03	0.01	0.00	0.00	0.01	0.20
2014‡	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2015‡	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2016‡	*	*	*	*	*	*	*	*	*
2017‡	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
South of 40 deg. 10' N. lat., 60+									
2011	0.08	0.01	0.08	0.16	0.02	0.01	0.07	0.09	0.54
2012	0.41	0.08	0.35	0.84	0.08	0.04	0.31	0.44	0.52
2013‡	0.29	0.14	0.47	0.90	0.06	0.08	0.42	0.56	0.62
2014‡	0.36	0.08	0.13	0.57	0.07	0.04	0.12	0.23	0.41
2015‡	0.18	0.03	0.12	0.33	0.04	0.02	0.11	0.16	0.49
2016‡	0.07	0.02	0.00	0.10	0.01	0.01	0.00	0.03	0.31
2017‡	0.04	0.02	0.02	0.07	0.01	0.01	0.01	0.03	0.43
LE CA Halibut South of 40 deg. 10' N. lat., all depths									
2011	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2012	*	*	*	*	*	*	*	*	*

Table 19: Estimated Pacific halibut gross at-sea discard (mt) and at-sea discard mortality (mt) in the U.S. West Coast groundfish IFQ midwater trawl fishery by management area and year. Estimates were allocated to three condition categories based on information presented in Table 15. DMR=Discard Mortality Rate. (*) confidential data, (-) no estimate, see text for explanation.

Midwater Trawl									
Area	Estimated Gross Discard (mt)				Estimated Discard Mortality				DMR
Year	Exc	Poor	Dead	Total	m(Exc)	m(Poor)	m(Dead)	Total	
Catch Shares North of 40 deg. 10' N. lat.									
2011	*	*	*	*	*	*	*	*	*
2012	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2013	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2014	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Midwater Hake North of 40 deg. 10' N. lat.									
2015	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2016	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2017	0.02	0.00	0.01	0.03	0.00	0.00	0.00	0.03	1.00
Shoreside Hake North of 40 deg. 10' N. lat.									
2011	0.00	0.01	0.01	0.03	0.00	0.00	0.00	0.03	1.00
2012	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2013	0.05	0.00	0.00	0.05	0.00	0.00	0.00	0.05	1.00
2014	0.10	0.01	0.00	0.11	0.00	0.00	0.00	0.11	1.00
Midwater Rockfish North of 40 deg. 10' N. lat.									
2015	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2016	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2017	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 20: Estimated Pacific halibut gross at-sea discard (mt) and at-sea discard mortality (mt) in the U.S. West Coast groundfish IFQ pot fishery by management area and year. Estimates were allocated to three condition categories based on information presented in Table 16. DMR=Discard Mortality Rate. (*) confidential data, (-) no estimate, see text for explanation.

Pot									
Area	Estimated Gross Discard (mt)				Estimated Discard Mortality				DMR
Year	Exc	Poor	Dead	Total	m(Exc)	m(Poor)	m(Dead)	Total	
North of Pt. Chehalis.									
2011	0.86	0.02	0.15	1.03	0.00	0.02	0.15	0.17	0.16
2012	0.84	0.21	0.21	1.27	0.00	0.21	0.21	0.43	0.34
2013	0.13	0.00	0.08	0.22	0.00	0.00	0.08	0.09	0.39
Pt. Chehalis to 40 deg. 10' N. lat.									
2011	1.61	0.11	0.61	2.33	0.00	0.11	0.61	0.72	0.31
2012	0.54	0.05	0.03	0.62	0.00	0.05	0.03	0.08	0.13
2013	0.63	0.05	0.07	0.76	0.00	0.05	0.07	0.13	0.17
2015	1.10	0.06	0.16	1.31	0.00	0.06	0.16	0.21	0.16
South of 40 deg. 10' N. lat.									
2011	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2012	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2013	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2015	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Coastwide									
2014	0.23	0.00	0.08	0.32	0.00	0.00	0.08	0.08	0.26
2016	1.53	0.03	0.14	1.70	0.00	0.03	0.14	0.17	0.10
2017	0.44	0.38	0.26	1.09	0.00	0.38	0.26	0.64	0.59

Table 21: Estimated Pacific halibut gross at-sea discard (mt) and at-sea discard mortality (mt) in the U.S. West Coast groundfish IFQ hook and line fishery by management area, depth, and year. A single discard mortality rate (DMR) of 0.16 is applied in this fishery. Viabilities are not used to determine discard mortality, see text for explanation. (*) confidential data, (-) no data collected.

Hook and Line									
Area	Estimated Gross Discard (mt)				Estimated Discard Mortality				DMR
Year	Exc	Poor	Dead	Total	m(Exc)	m(Poor)	m(Dead)	Total	
North of 40 deg. 10' N. lat.									
2011	-	-	-	6.06	-	-	-	0.97	0.16
South of 40 deg. 10' N. lat.									
2011	-	-	-	0.00	-	-	-	0.00	0.00
Coastwide									
2012	-	-	-	14.66	-	-	-	2.34	0.16
2013	-	-	-	3.00	-	-	-	0.48	0.16
2014	-	-	-	3.80	-	-	-	0.61	0.16
2015	-	-	-	9.49	-	-	-	1.52	0.16
2016	-	-	-	6.39	-	-	-	1.02	0.16
2017	-	-	-	4.14	-	-	-	0.66	0.16

Table 22: Estimated Pacific halibut discard (mt), discard mortality (mt), legal-sized (82 cm) mortality (mt), and percent of legal-sized discard by weight in the U.S. West Coast groundfish IFQ bottom trawl fisheries by management area, depth, and year. The proportion of legal-sized P. halibut in the non-hake IFQ bottom trawl sector north of 40°10' N. lat. is 76%. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate, see text for explanation.

Bottom Trawl					
Area, Depth (fm)		Total		Estimated legal-size	
Year	Discards (mt)	Discard mortality (mt)	Mortality (mt)	% by weight	
North of Pt. Chehalis, 0-60					
2011	7.98	3.58	1.96	54.66%	
2012	4.80	2.44	1.14	46.94%	
2013	5.46	3.08	1.23	39.75%	
2014	2.07	0.57	0.27	47.56%	
2015	3.89	1.37	0.95	68.79%	
2016	2.76	1.06	0.64	60.15%	
2017	0.84	0.26	0.18	70.98%	
North of Pt. Chehalis, 60+					
2011	22.29	11.34	8.00	70.52%	
2012	32.67	17.79	12.31	69.19%	
2013	29.91	12.42	7.96	64.07%	
2014	25.58	12.08	6.46	53.50%	
2015	24.86	10.77	6.96	64.63%	
2016	14.14	7.16	4.30	60.06%	
2017	17.41	8.88	5.75	64.81%	
Pt. Chehalis to 40 deg. 10' N. lat., 0-60					
2011	11.00	3.35	2.08	62.17%	
2012	8.06	3.04	1.61	53.04%	
2013	8.55	3.51	2.18	62.10%	
2014	10.48	3.89	1.91	49.16%	
2015	10.22	4.41	2.54	57.52%	
2016	7.32	3.75	2.06	55.01%	
2017	4.71	1.75	1.26	71.91%	
Pt. Chehalis to 40 deg. 10' N. lat., 60+					
2011	22.51	12.75	8.78	68.87%	
2012	20.46	12.08	8.51	70.44%	
2013	21.08	12.50	8.81	70.48%	
2014	17.11	9.33	6.90	73.89%	
2015	29.75	16.38	11.84	72.32%	

Table 22: Estimated Pacific halibut discard (mt), discard mortality (mt), legal-sized (82 cm) mortality (mt), and percent of legal-sized discard by weight in the U.S. West Coast groundfish IFQ bottom trawl fisheries by management area, depth, and year. The proportion of legal-sized P. halibut in the non-hake IFQ bottom trawl sector north of 40°10' N. lat. is 76%. (‡) combined IFQ and LE CA Halibut, (*) confidential data, (-) no estimate, see text for explanation. (*continued*)

Bottom Trawl				
Area, Depth (fm)	Total		Estimated legal-size	
Year	Discards (mt)	Discard mortality (mt)	Mortality (mt)	% by weight
2016	31.48	18.42	13.43	72.95%
2017	36.17	18.99	15.39	81.06%
South of 40 deg. 10' N. lat., 0-60				
2011	0.17	0.15	0.15	100.00%
2012	*	*	*	*
2013‡	0.03	0.01	0.01	100.00%
2014‡	0.00	0.00	0.00	0.00%
2015‡	0.00	0.00	0.00	0.00%
2016‡	*	*	*	*
2017‡	0.00	0.00	0.00	0.00%
South of 40 deg. 10' N. lat., 60+				
2011	0.16	0.09	0.09	96.93%
2012	0.84	0.44	0.38	86.31%
2013‡	0.90	0.56	0.45	80.25%
2014‡	0.57	0.23	0.21	90.96%
2015‡	0.33	0.16	0.14	88.19%
2016‡	0.10	0.03	0.03	88.20%
2017‡	0.07	0.03	0.03	91.17%
LE CA Halibut South of 40 deg. 10' N. lat., all depths				
2011	0.00	0.00	0.00	0.00%
2012	*	*	*	*

Table 23: Estimated Pacific halibut discard (mt), discard mortality (mt), legal-sized (82 cm) mortality (mt), and percent of legal-sized discard by weight in the U.S. West Coast groundfish IFQ midwater trawl fisheries by area and year. (*) confidential data, (-) no estimate, see text for explanation.

Midwater Trawl				
Area	Total		Estimated legal-size	
Year	Discards (mt)	Discard mortality (mt)	Mortality (mt)	% by weight
Catch Shares North of 40 deg. 10' N. lat.				
2011	*	*	*	*
2012	0.00	0.00	0.00	0.00%
2013	0.00	0.00	0.00	0.00%
2014	0.00	0.00	0.00	0.00%
Midwater Hake North of 40 deg. 10' N. lat.				
2015	0.00	0.00	0.00	0.00%
2016	0.00	0.00	0.00	0.00%
2017	0.03	0.03	0.03	87.31%
Shoreside Hake North of 40 deg. 10' N. lat.				
2011	0.03	0.03	0.02	76.44%
2012	0.00	0.00	0.00	0.00%
2013	0.05	0.05	0.05	91.55%
2014	0.11	0.11	0.10	90.18%
Midwater Rockfish North of 40 deg. 10' N. lat.				
2015	0.00	0.00	0.00	0.00%
2016	0.00	0.00	0.00	0.00%
2017	0.00	0.00	0.00	0.00%

Table 24: Estimated Pacific halibut discard (mt), discard mortality (mt), legal-sized (82 cm) mortality (mt), and percent of legal-sized discard by weight in the U.S. West Coast groundfish IFQ pot fisheries by area and year. (*) confidential data, (-) no estimate, see text for explanation.

Pot				
Area	Total		Estimated legal-size	
Year	Discards (mt)	Discard mortality (mt)	Mortality (mt)	% by weight
North of Pt. Chehalis.				
2011	1.03	0.17	0.13	77.00%
2012	1.27	0.43	0.34	80.73%
2013	0.22	0.09	0.07	77.82%
Pt. Chehalis to 40 deg. 10' N. lat.				
2011	2.33	0.72	0.54	74.48%
2012	0.62	0.08	0.06	73.97%
2013	0.76	0.13	0.09	70.53%
2015	1.31	0.21	0.16	73.94%
South of 40 deg. 10' N. lat.				
2011	0.00	0.00	0.00	0.00%
2012	0.00	0.00	0.00	0.00%
2013	0.00	0.00	0.00	0.00%
2015	0.00	0.00	0.00	0.00%
Coastwide				
2014	0.32	0.08	0.07	84.94%
2016	1.70	0.17	0.12	72.61%
2017	1.09	0.64	0.51	79.23%

Table 25: Estimated Pacific halibut discard (mt), discard mortality (mt), legal-sized (82 cm) mortality (mt), and percent of legal-sized discard by weight in the U.S. West Coast groundfish IFQ hook and line fisheries by area and year. (*) confidential data, (-) no estimate, see text for explanation.

Hook and Line				
Area	Total		Estimated legal-size	
Year	Discards (mt)	Discard mortality (mt)	Mortality (mt)	% by weight
North of 40 deg. 10' N. lat.				
2011	6.06	0.97	0.43	44.66%
South of 40 deg. 10' N. lat.				
2011	0.00	0.00	0.00	0.00%
Coastwide				
2012	14.66	2.34	1.81	76.99%
2013	3.00	0.48	0.24	49.73%
2014	3.80	0.61	0.30	49.87%
2015	9.49	1.52	0.65	42.61%
2016	6.39	1.02	0.43	42.08%
2017	4.14	0.66	0.31	46.56%

Table 26: Pacific halibut bycatch by month for vessels fishing bottom trawl gear in the 2017 IFQ fishery. We present coastwide estimates across all depths to maintain confidentiality. Note that adding values across columns might give slightly different results because values are rounded to two decimals for reporting.

Month	Expanded Discard (mt)	Sampled Discard (mt)	Total Bycatch (mt)
Jan	0.00	2.60	2.60
Feb	0.00	3.79	3.79
Mar	0.03	6.52	6.55
Apr	0.00	8.66	8.67
May	0.01	8.20	8.21
Jun	0.02	3.14	3.15
Jul	0.00	5.25	5.25
Aug	0.01	2.60	2.61
Sep	0.00	4.37	4.37
Oct	0.63	5.20	5.83
Nov	0.00	3.38	3.38
Dec	0.03	4.76	4.79

Table 27: Physical measurements of *P. halibut* length (cm) in the U.S. west coast groundfish IFQ fishery (2011-2017) for vessels using bottom trawl gear. Length bins include the lower bound and exclude the upper bound.

IFQ Bottom Trawl physical lengths					
Length bin (cm)	Total No. Individuals	No. Dead Individuals	Length bin (cm)	Total No. Individuals	No. Dead Individuals
14-16	1	1	104-106	701	328
18-20	1	0	106-108	527	233
22-24	1	0	108-110	474	226
30-32	3	1	110-112	418	185
32-34	5	3	112-114	341	160
34-36	4	1	114-116	248	115
36-38	6	1	116-118	188	86
38-40	28	8	118-120	148	68
40-42	28	12	120-122	115	53
42-44	28	11	122-124	111	46
44-46	15	8	124-126	78	34
46-48	27	10	126-128	56	30
48-50	58	20	128-130	46	17
50-52	54	32	130-132	29	10
52-54	70	38	132-134	26	8
54-56	118	61	134-136	21	8
56-58	147	83	136-138	19	6
58-60	354	203	138-140	12	6
60-62	662	363	140-142	13	8
62-64	1161	619	142-144	9	5
64-66	1602	856	144-146	7	3
66-68	1979	1043	146-148	10	2
68-70	2611	1348	148-150	9	3
70-72	3027	1552	150-152	3	2
72-74	3365	1775	152-154	3	1
74-76	3373	1770	154-156	2	1
76-78	3279	1753	160-162	1	1
78-80	3270	1736	162-164	2	1
80-82	3160	1624	164-166	2	0
82-84	3066	1619	168-170	3	1
84-86	2739	1420	170-172	3	1
86-88	2370	1239	172-174	6	2
88-90	2184	1122	174-176	1	1
90-92	2119	1079	176-178	1	0
92-94	1840	925	178-180	3	2
94-96	1456	717	180-182	2	0
96-98	1239	614	182-184	3	2
98-100	1106	545	184-186	2	1
100-102	1044	486	186-188	2	2
102-104	883	425	188-190	1	0
104-106	701	328	192-194	3	1
106-108	527	233	198-200	1	0
108-110	474	226	212-214	1	1

Table 28: Physical measurements of *P. halibut* length (cm) in the U.S. west coast groundfish IFQ fishery (2011-2017) for vessels using pot gear. Length bins include the lower bound and exclude the upper bound.

IFQ Pot physical lengths		
Length bin (cm)	Total No. Individuals	No. Dead Individuals
44-46	1	0
46-48	1	0
50-52	3	0
54-56	4	1
56-58	2	0
58-60	4	1
60-62	14	4
62-64	12	5
64-66	12	3
66-68	8	1
68-70	27	4
70-72	45	10
72-74	59	7
74-76	56	13
76-78	48	16
78-80	74	19
80-82	97	22
82-84	111	28
84-86	98	21
86-88	80	20
88-90	65	12
90-92	71	15
92-94	49	19
94-96	45	15
96-98	37	15
98-100	19	5
100-102	28	6
102-104	18	5
104-106	17	7
106-108	14	5
108-110	9	3
110-112	6	1
112-114	4	0
114-116	7	4
116-118	3	1
118-120	3	2
120-122	2	0
122-124	3	0
128-130	1	0
130-132	2	1
134-136	1	0
136-138	1	0
138-140	1	0
166-168	1	0
200-202	1	0

Table 29: Visual estimates of *P. halibut* lengths (cm) from the U.S. West Coast groundfish IFQ fishery (2011-2017) for vessels using bottom trawl, pot, and hook and line gear. Length bins include the lower bound and exclude the upper bound. On IFQ hook & line vessels, only visual estimates are taken on *P. halibut*.

Length bin (cm)	IFQ visual lengths, no. of individuals		
	Bottom Trawl (no.)	Pot (no.)	Hook and Line (no.)
25-34	0	1	25
35-44	2	2	177
45-54	3	1	358
55-64	11	2	974
65-74	32	4	1480
75-84	34	14	1085
85-94	41	8	662
95-104	27	8	387
105-114	8	1	215
115-124	10	2	124
125-134	6	1	31
135-144	4	0	13
145-154	4	0	2
155-164	1	0	1
165-174	0	0	2
175-184	2	0	1

8.2 Tables: Non-Nearshore Fixed Gear Fisheries

Table 30: Number of observed vessels, trips, and sets by year and gear type in the LE Sablefish Endorsed fishery.

LE Sablefish Endorsed									
Year	Longline						Pot		
	North			South			vessels	trips	sets
	vessels	trips	sets	vessels	trips	sets			
2002	9	23	207	18	47	181	6	23	247
2003	8	25	191	9	26	160	6	35	362
2004	6	13	121	13	35	205	3	13	139
2005	10	31	402	18	73	276	7	39	492
2006	9	31	291	10	34	160	7	39	289
2007	9	36	381	14	40	136	4	30	154
2008	6	17	195	13	60	345	6	24	329
2009	3	12	177	6	34	110	3	27	67
2010	5	18	251	20	127	511	7	43	314
2011	7	18	284	20	84	389	3	22	227
2012	5	7	47	16	86	485	5	19	351
2013	6	12	135	14	49	218	3	14	47
2014	5	12	246	13	74	249	4	16	195
2015	6	15	174	24	87	458	9	36	308
2016	4	10	212	20	87	459	7	55	596
2017	7	15	209	21	98	492	3	14	186

Table 31: Number of observed vessels, trips, and sets by year and gear type in the LE Sablefish NonEndorsed fishery. The number of observed pot vessels in this fishery is too small to meet confidentiality and thus not reported.

LE Sablefish Non-Endorsed			
Year	Longline		
	vessels	trips	sets
2002	4	11	22
2003	17	130	219
2004	14	62	130
2005	11	35	60
2006	21	121	201
2007	36	158	304
2008	32	122	221
2009	34	138	273
2010	38	226	472
2011	38	201	426
2012	26	128	252
2013	22	124	248
2014	18	77	154
2015	21	65	144
2016	16	41	70
2017	12	34	71

Table 32: Number of observed vessels, trips, and sets by year and gear type in the OA Fixed Gear fishery. OA Fixed Gear fishery was not observed until 2003.

OA Fixed Gear						
Year	Longline			Pot		
	vessels	trips	sets	vessels	trips	sets
2002	—	—	—	—	—	—
2003	13	41	49	7	16	50
2004	14	42	52	17	96	185
2005	10	34	37	14	43	50
2006	7	10	11	15	38	39
2007	25	51	67	21	46	75
2008	33	58	68	20	55	75
2009	34	69	104	18	30	45
2010	37	70	105	26	40	71
2011	40	69	101	29	61	85
2012	24	34	53	19	35	70
2013	14	23	30	17	25	48
2014	21	28	39	21	41	63
2015	20	38	54	17	49	64
2016	30	55	78	27	55	73
2017	43	61	79	44	85	126

Table 33: Expansion factors and WCGOP observed discard rate by gear type for limited entry (LE) and open access (OA) non-nearshore fixed gear sectors used to expand discard estimates of *Pacific halibut* to the entire fleet.

Sector	Gear	Expansion Factor	Sector and Gear Type Rate Applied
LE Sablefish Endorsed	Longline	Sablefish	LE Sablefish Endorsed Longline
LE Sablefish Endorsed	Longline	Sablefish	LE Sablefish Endorsed Pot
LE Sablefish Non-Endorsed	Longline	Groundfish	LE Sablefish Non-Endorsed Longline
LE Sablefish Non-Endorsed	Pot	Sablefish	OA Fixed Gear ¹ Pot
OA Fixed Gear	Hook and Line	Groundfish	OA Fixed Gear ¹ Hook and Line
OA Fixed Gear	Pot	Groundfish	OA Fixed Gear ¹ Pot

¹A coast-wide discard ratio and coast-wide discard estimate could not be computed in the OA fixed gear sector for 2002-06 because the WCGOP only covered OA vessels in California during this time

Table 34: Total sablefish or groundfish landings (mt) and observed discard ratios (1SE) for each sector and gear type in the non-nearshore fixed gear fishery. Sablefish landings were used as the expansion factor in all cases except for the LE Sablefish Non-Endorsed and the OA Fixed Gear sectors, where target species include a variety of groundfish species.

Year	LE Sablefish Endorsed			LE Sablefish Non-Endorsed		OA Fixed Gear	
	Longline		Pot	Longline	Pot	Hook-and-Line	Pot
	North of Pt. Chehalis	South of Pt. Chehalis					
Expansion Factors							
2002	383	409	352	622	7	388	111
2003	482	570	604	542	7	545	190
2004	665	653	620	399	11	474	186
2005	668	674	615	548	3	624	379
2006	684	717	582	467	30	486	443
2007	495	609	428	516	2	263	258
2008	403	701	433	650	3	401	241
2009	429	1012	489	757	7	646	373
2010	266	1039	504	1000	17	757	318
2011	223	930	372	1246	24	434	256
2012	206	873	286	810	9	323	126
2013	217	531	283	813	15	193	72
2014	183	564	338	724	4	219	148
2015	234	706	358	725	4	364	234
2016	295	737	359	742	12	291	207
2017	314	746	375	784	12	348	211
Discard Ratios							
2002	0.3297 (0.05)	0.0283 (0.01)	0.0114 (0.00)	--	--	--	--
2003	0.3532 (0.05)	0.0467 (0.01)	0.0005 (0.00)	0.0003 (0.00)	--	--	--
2004	0.2369 (0.07)	0.0741 (0.01)	0.0526 (0.01)	--	--	--	--
2005	0.3318 (0.07)	0.0203 (0.00)	0.0043 (0.00)	--	--	--	--
2006	0.7827 (0.11)	0.1636 (0.05)	0.0271 (0.00)	--	--	--	--
2007	0.2184 (0.03)	0.0333 (0.01)	0.0092 (0.00)	0.0033 (0.00)	--	0.0789 (0.02)	0.0034 (0.00)
2008	0.3715 (0.07)	0.1473 (0.03)	0.0153 (0.00)	0.0046 (0.00)	--	0.0994 (0.04)	0.0010 (0.00)
2009	0.6497 (0.10)	0.0413 (0.01)	0.0017 (0.00)	0.0003 (0.00)	--	0.0541 (0.02)	0.0007 (0.00)
2010	0.2522 (0.06)	0.0631 (0.01)	0.0100 (0.00)	0.0004 (0.00)	--	0.0424 (0.03)	0.0016 (0.00)
2011	0.4780 (0.06)	0.0281 (0.00)	0.0110 (0.00)	0.0171 (0.01)	--	0.0299 (0.01)	0.0003 (0.00)
2012	0.4534 (0.16)	0.0594 (0.01)	0.0209 (0.00)	0.0204 (0.01)	--	0.0719 (0.03)	0.0032 (0.00)
2013	0.0871 (0.02)	0.0063 (0.00)	0.0000 (0.00)	--	--	0.0089 (0.00)	0.0008 (0.00)
2014	0.8892 (0.13)	0.0177 (0.00)	0.0060 (0.00)	--	--	0.0152 (0.00)	0.0011 (0.00)
2015	0.3685 (0.07)	0.0562 (0.01)	0.0046 (0.00)	0.0006 (0.00)	--	0.0278 (0.01)	0.0000 (0.00)
2016	0.4211 (0.06)	0.0596 (0.01)	0.0159 (0.00)	0.0069 (0.00)	--	0.1466 (0.03)	0.0003 (0.00)
2017	0.8062 (0.11)	0.0527 (0.01)	0.0024 (0.00)	0.0009 (0.00)	--	0.1361 (0.04)	0.0042 (0.00)

Table 35: Percent of observed trips that caught Pacific halibut by sector, gear, and area (where applicable). Observed average, minimum and maximum annual catch and discard weights and the percent of P. halibut catch weight discarded by year. n.o.c. No observed catch of P. halibut and thus a % discard calculation is not possible. – No WCGOP observers were deployed for the sector-year-gear type combination.

Year	LE Sablefish Endorsed			LE Sablefish Non-Endorsed		OA Fixed Gear	
	Longline		Pot	Longline	Pot	Hook-and-Line	Pot
	North of Pt. Chehalis	South of Pt. Chehalis					
% of observed trips that caught P. halibut							
2002	95.7%	46.8%	17.4%	0.0%	–	0.0%	0.0%
2003	100.0%	50.0%	8.6%	0.8%	–	0.0%	0.0%
2004	100.0%	71.4%	38.5%	0.0%	–	0.0%	0.0%
2005	96.8%	58.9%	33.3%	0.0%	–	0.0%	0.0%
2006	100.0%	76.5%	56.4%	0.0%	–	10.0%	0.0%
2007	94.4%	47.5%	33.3%	1.9%	–	25.5%	6.5%
2008	100.0%	78.3%	83.3%	3.3%	–	34.5%	5.5%
2009	91.7%	35.3%	33.3%	0.7%	–	37.7%	10.0%
2010	83.3%	47.2%	51.2%	1.3%	–	21.4%	2.5%
2011	88.9%	42.9%	45.5%	6.0%	–	30.4%	6.6%
2012	71.4%	58.1%	31.6%	7.0%	–	32.4%	8.6%
2013	83.3%	26.5%	21.4%	0.0%	–	13.0%	4.0%
2014	100.0%	24.3%	56.2%	0.0%	–	25.0%	9.8%
2015	100.0%	49.4%	61.1%	1.5%	–	34.2%	4.1%
2016	100.0%	60.9%	61.8%	7.3%	–	58.2%	7.3%
2017	93.3%	48.0%	57.1%	2.9%	–	31.1%	8.2%
Observed annaul catch (mt) of Pacific haibut							
Mean	43.4	11.4	1.8	0.3	–	0.9	0.0
Min	8.0	0.7	0.1	0.0	–	0.0	0.0
Max	118.4	36.6	5.4	1.4	–	2.3	0.1
Observed annaul discard (mt) of Pacific haibut							
Mean	38.2	11.3	1.8	0.3	–	0.9	0.0
Min	5.5	0.7	0.1	0.0	–	0.0	0.0
Max	109.6	36.6	5.4	1.4	–	2.3	0.1
% of Pacific haibut catch that was discarded							
2002	77.7%	95.5%	100.0%	n.o.c.	–	n.o.c.	n.o.c.
2003	80.1%	99.4%	100.0%	0.0%	–	n.o.c.	n.o.c.
2004	76.3%	97.3%	100.0%	n.o.c.	–	n.o.c.	n.o.c.
2005	82.7%	100.0%	100.0%	n.o.c.	–	n.o.c.	n.o.c.
2006	92.6%	97.5%	100.0%	n.o.c.	–	100.0%	n.o.c.
2007	78.0%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2008	87.4%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2009	100.0%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2010	100.0%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2011	100.0%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2012	96.6%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2013	69.0%	100.0%	0.0%	0.0%	–	100.0%	100.0%
2014	95.7%	100.0%	100.0%	0.0%	–	100.0%	100.0%
2015	95.5%	100.0%	100.0%	0.0%	–	100.0%	0.0%
2016	84.5%	100.0%	100.0%	67.0%	–	100.0%	100.0%
2017	89.1%	100.0%	97.7%	0.0%	–	100.0%	100.0%

Table 36: Pacific halibut viabilities caught by longline vessels in the U.S. west coast Limited Entry Sablefish Endorsed fishery by year and area north and south of Point Chehalis, WA. The condition of sampled P. halibut was identified as Minor (Mi), Moderate (Mo), Serious (Ser) or Dead (see Appendix in WCGOP manual), consistent with IPHC protocol. Sample sizes allowed for annual weighted percentages as described in the Methods.

Year	Limited Entry Sablefish																							
	North of Pt. Chehalis				South of Pt. Chehalis				Coastwide				North of Pt. Chehalis				South of Pt. Chehalis				Coastwide			
	no. of fish				no. of fish				no. of fish				weighted %				weighted %				weighted %			
	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead
2011	6039	157	95	124	2414	220	40	315	8453	377	135	439	95.6%	1.8%	1.3%	1.4%	80.1%	6.9%	1.0%	12.1%	91.2%	3.2%	1.2%	4.4%
2012	919	41	0	136	6197	798	190	461	7116	839	190	597	84.1%	3.5%	0.0%	12.4%	78.5%	11.3%	3.0%	7.3%	79.2%	10.2%	2.6%	8.0%
2013	2740	318	0	207	522	9	15	46	3262	327	15	253	83.1%	9.7%	0.0%	7.1%	90.3%	0.4%	4.0%	5.3%	84.5%	7.9%	0.8%	6.8%
2014	7238	296	144	921	1565	255	60	233	8803	551	204	1154	83.2%	3.3%	1.8%	11.6%	71.4%	12.1%	3.3%	13.2%	80.5%	5.4%	2.2%	12.0%
2015	7111	165	123	145	9513	663	298	249	16624	828	421	394	94.2%	2.1%	1.7%	2.0%	88.4%	6.1%	2.9%	2.5%	90.8%	4.4%	2.4%	2.3%
2016	7604	461	28	112	8919	1815	107	656	16523	2276	135	768	93.1%	5.3%	0.4%	1.2%	74.2%	17.4%	1.7%	6.8%	80.9%	13.1%	1.2%	4.8%
2017	9786	725	177	895	7202	744	54	471	16988	1469	231	1366	85.4%	6.2%	1.5%	6.9%	86.5%	7.4%	0.7%	5.5%	85.9%	6.7%	1.1%	6.3%

Table 37: Pacific halibut viabilities caught by longline vessels in the U.S. west coast Limited Entry Sablefish Non-Endorsed fishery, coastwide by year. The condition of sampled P. halibut was identified as Minor (Mi), Moderate (Mo), Serious (Ser) or Dead (see Appendix in WCGOP manual), consistent with IPHC protocol. Sample sizes necessitated the use of a five year rolling average to calculate weighted percentages, as described in the Methods.

Year	Limited Entry Sablefish Non-Endorsed							
	no. of fish				weighted %			
	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead
2011	496	214	0	0	64.8%	35.2%	0.0%	0.0%
2012	374	16	0	0	97.3%	2.7%	0.0%	0.0%
2013	0	0	0	0	0.0%	0.0%	0.0%	0.0%
2014	0	0	0	0	0.0%	0.0%	0.0%	0.0%
2015	24	0	0	0	100.0%	0.0%	0.0%	0.0%
2016	58	36	0	0	56.1%	43.9%	0.0%	0.0%
2017	11	0	0	0	100.0%	0.0%	0.0%	0.0%

Table 38: Pacific halibut viabilities caught by hook-&-line vessels in the U.S. west coast Open Access fixed gear fishery, coastwide by year. The condition of sampled P. halibut was identified as Minor (Mi), Moderate (Mo), Serious (Ser) or Dead (see Appendix in WCGOP manual), consistent with IPHC protocol. Sample sizes necessitated the use of a five year rolling average to calculate weighted percentages, as described in the Methods.

Year	Open Access Fixed Gear							
	no. of fish				weighted %			
	Mi	Mo	Ser	Dead	Mi	Mo	Ser	Dead
2011	286	76	31	6	72.0%	17.5%	10.0%	0.5%
2012	473	37	0	18	91.1%	8.6%	0.0%	0.3%
2013	53	0	0	0	100.0%	0.0%	0.0%	0.0%
2014	110	16	0	0	80.8%	19.2%	0.0%	0.0%
2015	473	25	0	0	96.9%	3.1%	0.0%	0.0%
2016	1878	89	0	0	92.5%	7.5%	0.0%	0.0%
2017	738	39	5	19	92.8%	4.4%	0.3%	2.5%

Table 39: Estimated gross discard (mt) and discard mortality (mt) in the limited entry (LE) sablefish endorsed, LE sablefish non-endorsed, and open access (OA) fixed gear sectors. Estimated discard mortality (mt) on longline and hook-&-line vessels was computed by two methods. For the 2002-2010 period, a 16 % discard mortality rate was applied to gross discard estimates because viability data was not available. Since 2011, mortality rates on longline vessels are based on the viability categories assigned to individuals. For all years, an 18% discard mortality rate was applied to gross discard estimates from pot vessels.

Year	LE Sablefish Endorsed				LE Sablefish Non-Endorsed		OA Fixed Gear	
	Longline			Pot	Longline	Pot	Hook-and-Line	Pot
	North of Pt. Chehalis	South of Pt. Chehalis	Coastwide					
Gross Discard Estimates								
2002	126.15	11.57	137.73	4.03	0.00	[0.02] *	[35.41] *	[0.23] *
2003	170.20	26.61	196.81	0.30	0.17	[0.01] *	[49.79] *	[0.40] *
2004	157.62	48.35	205.97	32.60	0.00	[0.02] *	[43.29] *	[0.39] *
2005	221.53	13.68	235.21	2.62	0.00	[0.01] *	[57.01] *	[0.79] *
2006	535.52	117.32	652.84	15.79	0.00	[0.06] *	[44.40] *	[0.92] *
2007	108.06	20.27	128.33	3.94	1.72	0.01	20.78	0.89
2008	149.62	103.22	252.84	6.62	2.99	0.00	39.89	0.23
2009	278.91	41.84	320.75	0.85	0.24	0.01	34.99	0.26
2010	66.99	65.58	132.56	5.04	0.37	0.03	32.06	0.50
2011	106.73	26.12	132.85	4.08	21.30	0.01	12.97	0.06
2012	93.39	51.87	145.27	5.99	16.54	0.03	23.22	0.41
2013	18.86	3.37	22.23	0.00	0.00	0.01	1.72	0.06
2014	162.77	9.98	172.76	2.03	0.00	0.00	3.33	0.16
2015	86.15	39.66	125.82	1.63	0.46	0.00	10.13	0.00
2016	124.22	43.94	168.17	5.72	5.09	0.00	42.73	0.05
2017	253.31	39.34	292.66	0.90	0.67	0.05	47.37	0.87
Estimated Discard Mortality								
2002	20.18	1.85	22.04	0.73	0.00	— *	— *	— *
2003	27.23	4.26	31.49	0.05	0.03	— *	— *	— *
2004	25.22	7.74	32.96	5.87	0.00	— *	— *	— *
2005	35.44	2.19	37.63	0.47	0.00	— *	— *	— *
2006	85.68	18.77	104.45	2.84	0.00	— *	— *	— *
2007	17.29	3.24	20.53	0.71	0.28	0.00	3.32	0.16
2008	23.94	16.52	40.45	1.19	0.48	0.00	6.38	0.04
2009	44.62	6.69	51.32	0.15	0.04	0.00	5.60	0.05
2010	10.72	10.49	21.21	0.91	0.06	0.00	5.13	0.09
2011	6.62	4.71	11.33	0.73	3.20	0.00	2.07	0.01
2012	15.51	8.35	23.86	1.08	0.72	0.00	1.54	0.07
2013	2.56	0.38	2.94	0.00	0.00	0.00	0.06	0.01
2014	27.57	2.22	29.79	0.37	0.00	0.00	0.33	0.03
2015	6.19	3.89	10.08	0.29	0.02	0.00	0.46	0.00
2016	8.21	7.37	15.59	1.03	0.91	0.00	2.55	0.01
2017	33.21	4.58	37.78	0.16	0.02	0.01	3.57	0.16

*The LE sablefish non-endorsed pot sector has not been observed by the WCGOP and therefore estimates are based on discard rates from observed OA fixed gear pot vessels. OA fixed gear vessels were not observed coastwide until 2007 and thus 2002-06 estimates are based on the 2007-08 coastwide discard rate, shown in brackets.

Table 40: Estimated P. halibut discard mortality (mt, mortality rate applied, see text for description) from each sector of the non-nearshore fixed gear fishery by year.

	Estimated discard mortality (mt)			All Sectors
	LE Sablefish Endorsed	LE Sablefish Non- Endorsed	OA Fixed Gear	
2002	22.76	0.00	0.00	22.76
2003	31.54	0.03	0.00	31.57
2004	38.82	0.00	0.00	38.82
2005	38.10	0.00	0.00	38.10
2006	107.30	0.00	0.00	107.30
2007	21.24	0.28	3.48	25.00
2008	41.65	0.48	6.42	48.55
2009	51.47	0.04	5.65	57.16
2010	22.12	0.06	5.22	27.40
2011	12.07	3.20	2.09	17.35
2012	24.94	0.73	1.61	27.28
2013	2.94	0.00	0.07	3.01
2014	30.16	0.00	0.35	30.51
2015	10.37	0.02	0.46	10.84
2016	16.62	0.91	2.56	20.09
2017	37.95	0.03	3.73	41.71

Table 41: Physical measurements of *P. halibut* length (cm) from the U.S. west coast LE Sablefish Endorsed fishery (2002-2017) for vessels using hook & line gear. Length bins include the lower bound and exclude the upper bound.

LE Sablefish Endorsed Hook & Line, physical lengths					
Length bin (cm)	Total No.	Dead No.	Length bin (cm)	Total No.	Dead No.
38-40	2	0	88-90	306	46
40-42	1	0	90-92	292	35
42-44	2	0	92-94	307	30
44-46	3	0	94-96	264	31
46-48	3	1	96-98	219	23
48-50	4	0	98-100	155	20
50-52	9	1	100-102	126	17
52-54	11	0	102-104	133	23
54-56	9	0	104-106	105	18
56-58	24	2	106-108	82	12
58-60	21	2	108-110	78	14
60-62	32	2	110-112	52	8
62-64	56	5	112-114	48	10
64-66	78	8	114-116	44	4
66-68	89	12	116-118	37	6
68-70	170	23	118-120	19	2
70-72	229	26	120-122	18	3
72-74	292	35	122-124	9	1
74-76	357	45	124-126	15	1
76-78	393	45	126-128	14	3
78-80	438	52	128-130	6	1
80-82	378	54	130-132	2	1
82-84	404	51	132-134	4	0
84-86	404	51	134-136	2	1
86-88	375	52	136-138	2	0
88-90	306	46	140-142	3	0
90-92	292	35	142-144	1	0
92-94	307	30	158-160	1	0

Table 42: Physical measurements of *P. halibut* length (cm) from the U.S. west coast LE Sablefish Endorsed fishery (2002-2017) for vessels using pot gear. Length bins include the lower bound and exclude the upper bound.

LE Sablefish Endorsed Pot, physical lengths					
Length bin (cm)	Total No.	Dead No.	Length bin (cm)	Total No.	Dead No.
42-44	1	1	82-84	85	19
44-46	1	0	84-86	83	12
46-48	1	1	86-88	78	17
48-50	1	0	88-90	64	10
50-52	1	0	90-92	51	10
52-54	3	0	92-94	36	7
58-60	1	0	94-96	40	7
60-62	5	1	96-98	32	4
62-64	4	1	98-100	23	8
64-66	10	1	100-102	16	4
66-68	16	3	102-104	12	2
68-70	18	2	104-106	7	2
70-72	37	6	106-108	6	0
72-74	42	8	108-110	9	1
74-76	70	9	110-112	7	2
76-78	70	12	114-116	1	0
78-80	76	13	116-118	3	1
80-82	78	16	118-120	2	0
82-84	85	19	122-124	6	2
84-86	83	12	130-132	1	0
86-88	78	17	146-148	1	0

Table 43: Physical measurements of *P. halibut* length (cm) from the U.S. west coast LE Sablefish Non-Endorsed fishery (2002-2017) for vessels using hook and line gear. The WCGOP does not cover vessels fishing pot gear in this fishery. Length bins include the lower bound and exclude the upper bound.

LE Sablefish Non-Endorsed Hook & Line, physical lengths		
Length bin (cm)	Total No.	Dead No.
66-68	1	0
68-70	3	0
72-74	5	1
74-76	4	0
76-78	6	0
78-80	4	0
80-82	3	0
82-84	3	0
84-86	3	0
86-88	5	1
88-90	6	0
90-92	5	1
92-94	4	0
94-96	2	0
96-98	3	0
98-100	6	1
100-102	1	0
102-104	3	0
104-106	3	0
106-108	2	0
110-112	1	0
112-114	4	1
118-120	2	0
122-124	1	0
134-136	1	0

Table 44: Physical measurements of *P. halibut* length (cm) from the U.S. west coast OA Fixed Gear fishery (2002-2017) for vessels using hook and line or pot gear.

Length bin (cm)	OA Fixed Gear, physical lengths			
	Hook and Line		Pot	
	Total (no.)	Dead (no.)	Total (no.)	Dead (no.)
34-36	1	0	0	0
40-42	1	0	0	0
44-46	3	1	0	0
48-50	1	0	0	0
50-52	2	0	0	0
52-54	1	0	0	0
54-56	1	0	0	0
56-58	2	0	0	0
58-60	4	0	0	0
60-62	5	0	0	0
62-64	6	0	1	0
64-66	9	2	1	1
66-68	9	0	1	1
68-70	16	1	1	0
70-72	14	0	3	0
72-74	26	3	0	0
74-76	26	1	1	0
76-78	22	1	1	0
78-80	22	1	2	0
80-82	26	1	2	1
82-84	27	2	0	0
84-86	35	2	1	0
86-88	29	3	2	0
88-90	15	1	1	0
90-92	17	1	0	0
92-94	21	2	2	1
94-96	13	1	1	0
96-98	12	0	0	0
98-100	6	0	0	0
100-102	8	0	0	0
102-104	2	0	0	0
104-106	6	1	0	0
106-108	4	0	0	0
108-110	6	2	2	0

Table 44: Physical measurements of *P. halibut* length (cm) from the U.S. west coast OA Fixed Gear fishery (2002-2017) for vessels using hook and line or pot gear. (*continued*)

Length bin (cm)	OA Fixed Gear, physical lengths			
	Hook and Line		Pot	
	Total (no.)	Dead (no.)	Total (no.)	Dead (no.)
110-112	4	0	1	0
112-114	3	0	0	0
114-116	5	1	0	0
116-118	1	0	0	0
118-120	2	0	0	0
120-122	1	0	0	0
122-124	2	0	0	0

Table 45: Visual estimates of *P. halibut* lengths (cm) from the U.S. West Coast groundfish Non-Nearshore fixed gear fisheries (2002-2017) for vessels using hook and line gear and pot gear. Numbers are the numbers of individuals caught with each gear type. The WCGOP does not observe LE Non-Endorsed Sablefish vessels fishing with pot gear. Length bins include the lower bound and exclude the upper bound.

Length bin (cm)	LE Sablefish Endorsed		LE Sablefish Non-Endorsed	OA Fixed Gear	
	No. Hook and Line	No. Pot	No. Hook and Line	No. Hook and Line	No. Pot
25-34	28	0	0	1	0
35-44	87	1	0	6	0
45-54	360	2	2	20	0
55-64	2357	6	11	54	0
65-74	6275	8	28	137	0
75-84	8367	25	37	156	2
85-94	6345	12	26	85	1
95-104	3604	6	15	38	0
105-114	1080	0	8	14	0
115-124	362	0	9	7	0
125-134	87	0	4	1	0
135-144	24	0	0	0	0
145-154	6	0	0	0	0
155-164	2	0	0	1	0

8.3 Tables: Legal-Sublegal P. halibut Lengths

Table 46: Pacific halibut physically measured lengths and visual estimates of lengths approximating legal (82 cm>) versus sublegal definitions (IPHC), collected by the WCGOP in the IFQ fishery (2011-2017), Non-Nearshore fixed gear fisheries (LE sablefish endorsed, LE non-endorsed, OA fixed gear; 2002-2017), and the At-sea Hake sectors (2002-2017). Note that visual length estimates are not taken in the At-sea Hake sectors.

Fishery	Type of Measurement	Length bin (cm)	No. of individuals	Percentage of Total
Non-Nearshore Fixed Gear	actual	0-82.0	3272	42.8%
Non-Nearshore Fixed Gear	actual	82.0>	4374	57.2%
Non-Nearshore Fixed Gear	visual	0-74.0	9452	20.6%
Non-Nearshore Fixed Gear	visual	75.0-84.0	12394	27.0%
Non-Nearshore Fixed Gear	visual	82.0>	24019	52.4%
Catch Shares	actual	0-82.0	28908	54.2%
Catch Shares	actual	82.0>	24404	45.8%
Catch Shares	visual	0-74.0	1556	27.0%
Catch Shares	visual	75.0-84.0	1133	19.6%
Catch Shares	visual	82.0>	3077	53.4%
At-Sea Hake	actual	0-82.0	201	27.8%
At-Sea Hake	actual	82.0>	521	72.2%

8.4 Tables: IPHC P. halibut Derby Fishery

Table 47: Observer coverage information for the Pacific halibut Derby fishery by year. The WCGOP began observing the Pacific halibut derby fishery in 2017 at the request of the International Pacific Halibut Commission.

P. halibut Derby Fishery									
Observed							Fleet-wide		
Year	Gear	Number of:					Retained (mt)	Retained (mt)	Coverage
		Vessels	Trips	Sets	Lost Hooks	Total Hooks			
2017	Hook-and-Line	13	22	62	2269	973150	10.21	137.06	7.0%

Table 48: Discard ratio and gross discard estimate of P. halibut for the P. halibut Derby fishery. LCL = lower confidence limit; UCL = upper confidence limit

P. halibut Derby Fishery						
Observed					Fleet-wide	
Year	Gear	Discard (mt)	Retained (mt)	Ratio (LCL-UCL)	Retained (mt)	Gross discard (mt, LCL-UCL)
2017	Hook-and-Line	1.9	10.2	0.19 (0.1 - 0.3)	137.1	25.7 (9.4-42)

Table 49: Pacific halibut viabilities from the Pacific halibut Derby fishery. The injury status of sampled discarded P. halibut was identified as minor, moderate, and serious injuries or dead (see Appendices in WCGOP manual), consistent with International Pacific Halibut Commission protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods.

P. halibut Derby Fishery									
Number						% Weighted			
Year	Gear	Minor	Moderate	Serious	Dead	Minor	Moderate	Serious	Dead
2017	Hook-and-Line	102	8	2	2	89.0%	7.0%	2.0%	2.0%

Table 50: Pacific halibut mortality from the Pacific halibut Derby fishery. The injury status of sampled discarded P. halibut was identified as minor, moderate, and serious injuries or dead (see Appendices in WCGOP manual), consistent with International Pacific Halibut Commission protocol. The number of fish in each category was weighted based on the length-weight relationship as described in the Methods. Mortality rates (Table 7) were applied to each injury category. LCL = lower confidence limit; UCL = upper confidence limit

P. halibut Derby Fishery							
Year	Gear	Gross (mt, LCL-UCL)	Discard Mortality (mt, LCL-UCL)				Total
		Discards	Minor	Moderate	Serious	Dead	
2017	Hook-and-Line	25.7 (9.4-42)	0.8 (0.3-1.3)	0.6 (0.2-1)	0.3 (0.1-0.6)	0.5 (0.2-0.8)	2.3 (0.8-3.7)

Table 51: Physical length measurements of discarded P. halibut from the Pacific halibut Derby fishery. Length bins include the lower bound and exclude the upper bound.

P. halibut Derby Fishery			
Hook-and-Line physical lengths			
Length bin (cm)	Total No. Individuals	No. Dead Individuals	
44-46	1	0	
54-56	1	0	
60-62	1	0	
62-64	2	0	
66-68	1	0	
68-70	4	0	
70-72	10	1	
72-74	7	0	
74-76	18	2	
76-78	22	1	
78-80	27	2	
80-82	14	2	
82-84	6	1	

Table 52: Visual length estimates of discarded P. halibut from the Pacific halibut Derby fishery. Length bins include the lower bound and exclude the upper bound.

P. halibut Derby Fishery	
Hook-and-Line visual estimates	
Length bin (cm)	Total No. Individuals
45-54	1
55-64	15
65-74	64
75-84	76
85-94	7
95-104	2

8.5 Tables: Observed State Fisheries

Table 53: Coverage information, bycatch rates, and bycatch estimates for Pacific halibut in the Oregon and California nearshore fixed gear groundfish fisheries by state and year. The WCGOP began observing the California nearshore fishery in 2003 and the Oregon nearshore fishery in 2004. Gear specific mortality rates cannot be applied to *P. halibut* bycatch in this fishery because of confidentiality issues. Coverage rate in the state nearshore fisheries is defined as the proportion of nearshore target species landings that were observed. Nearshore target species are listed in the WCGOP Data Processing Appendix. Washington does not allow a state nearshore fishery.

Nearshore fixed gear groundfish fishery											
State	Observed							Estimated			
Year	Fleet observer coverage rate	Number of observed sets	% of sets with <i>P. halibut</i>	<i>P. halibut</i> bycatch (mt)	Nearshore species retained (mt)	<i>P. halibut</i> bycatch rate	SE bycatch rate	Total fleet catch of nearshore species (mt)	<i>P. halibut</i> bycatch (mt)	Bycatch lower 95% CI (mt)	Bycatch upper 95% CI (mt)
Oregon											
2002	<i>not observed</i>	—	—	—	—	—	—	278.68	—	—	—
2003	<i>not observed</i>	—	—	—	—	—	—	207.77	—	—	—
2004	4.91%	210	1.90%	0.05	10.27	0.00	0.00	209.42	0.997	0.438	1.556
2005	6.61%	170	0.59%	0.03	11.84	0.00	0.00	179.04	0.492	0.395	0.589
2006	11.94%	385	1.30%	0.06	19.63	0.00	0.00	164.41	0.526	0.272	0.780
2007	9.01%	248	0.40%	0.01	16.23	0.00	0.00	180.22	0.086	0.072	0.100
2008	7.78%	185	0.54%	0.03	14.63	0.00	0.00	188.22	0.350	0.287	0.413
2009	6.26%	225	2.22%	0.08	13.95	0.01	0.00	222.83	1.281	0.745	1.816
2010	7.86%	213	0.47%	0.01	13.46	0.00	0.00	171.17	0.078	0.065	0.091
2011	8.19%	244	2.05%	0.09	15.88	0.01	0.00	193.85	1.093	0.400	1.787
2012	10.59%	290	1.38%	0.11	20.71	0.01	0.00	195.56	1.061	0.321	1.802
2013	7.80%	264	0.76%	0.02	16.08	0.00	0.00	206.05	0.290	0.196	0.383
2014	8.26%	197	2.03%	0.08	16.64	0.00	0.00	201.39	0.954	0.547	1.361
2015	8.53%	237	1.69%	0.12	18.43	0.01	0.00	216.05	1.436	0.098	2.774
2016	12.25%	265	4.53%	0.37	21.91	0.02	0.00	178.91	3.015	1.336	4.694
2017	12.33%	237	3.80%	0.19	24.77	0.01	0.00	200.95	1.554	0.974	2.133
California											
2002	<i>not observed</i>	—	—	—	—	—	—	381.26	—	—	—
2003	3.17%	208	0.00%	0.00	8.11	0.00	0.00	256.15	0.000	0.000	0.000
2004	8.00%	434	0.00%	0.00	23.26	0.00	0.00	290.62	0.000	0.000	0.000
2005	4.66%	219	0.91%	0.08	13.01	0.01	0.00	279.51	1.707	0.003	3.567
2006	3.24%	161	0.00%	0.00	8.34	0.00	0.00	257.52	0.000	0.000	0.000
2007	4.32%	227	0.00%	0.00	11.82	0.00	0.00	273.58	0.000	0.000	0.000
2008	2.28%	89	0.00%	0.00	6.69	0.00	0.00	293.75	0.000	0.000	0.000
2009	2.59%	123	0.00%	0.00	6.72	0.00	0.00	259.67	0.000	0.000	0.000
2010	3.23%	117	0.00%	0.00	7.07	0.00	0.00	218.85	0.000	0.000	0.000
2011	3.91%	214	0.47%	0.08	8.45	0.01	0.00	215.99	1.975	1.543	2.407
2012	5.95%	239	1.26%	0.07	11.91	0.01	0.00	200.19	1.187	0.174	2.200
2013	5.31%	194	1.55%	0.06	11.67	0.00	0.00	219.52	1.071	0.561	1.581
2014	4.69%	183	0.00%	0.00	11.46	0.00	0.00	244.51	0.000	0.000	0.000
2015	6.90%	277	0.00%	0.00	22.85	0.00	0.00	331.03	0.000	0.000	0.000
2016	5.34%	156	0.00%	0.00	13.23	0.00	0.00	247.64	0.000	0.000	0.000
2017	5.87%	163	0.00%	0.00	12.21	0.00	0.00	208.11	0.000	0.000	0.000

Table 54: Coverage information, bycatch rates, and bycatch estimates for Pacific halibut in the Washington and Oregon state pink shrimp fisheries by state and year. The WCGOP began observing the OR state pink shrimp fishery in 2004, but was unable to observe it in 2006. The WA state pink shrimp fishery was added for observation in 2010. Mortality rates are not applied to P. halibut bycatch in these fisheries because mortality rates for pink shrimp trawl gear have not been estimated. Coverage rate in the pink shrimp fisheries is defined as the proportion of pink shrimp landings that were observed. (*) = Confidential data; (–) = not observed; LCL/UCL = lower/upper 95% confidence limit.

Year	Observed							Estimated			
	Coverage rate	No. of sets	% sets w/P. halibut	P. halibut bycatch (kg)	Shrimp retained (kg)	Bycatch rate	SE	Shrimp landings (mt)	P. halibut bycatch (mt)	LCL	UCL
Washington											
2010	9.30%	341	0.00%	0.00	399484	0.00000	0.00000	4296	0.00	0.00	0.00
2011	16.17%	578	0.17%	7.66	697238	0.00001	0.00000	4312	0.05	0.04	0.05
2012	14.77%	522	0.00%	0.00	625952	0.00000	0.00000	4239	0.00	0.00	0.00
2013	10.18%	386	0.00%	0.00	626823	0.00000	0.00000	6158	0.00	0.00	0.00
2014	7.03%	401	0.00%	0.00	976192	0.00000	0.00000	13876	0.00	0.00	0.00
2015	11.37%	1454	0.00%	0.00	2139754	0.00000	0.00000	18814	0.00	0.00	0.00
2016	17.32%	973	0.00%	0.00	1107926	0.00000	0.00000	6396	0.00	0.00	0.00
2017	19.49%	915	0.00%	0.00	592624	0.00000	0.00000	3041	0.00	0.00	0.00
Oregon											
2002	<i>not observed</i>	–	–	–	–	–	–	18896	–	–	–
2003	<i>not observed</i>	–	–	–	–	–	–	9322	–	–	–
2004	7.72%	765	0.00%	0.00	427212	0.00000	0.00000	5537	0.00	0.00	0.00
2005	5.63%	533	0.19%	2.27	402886	0.00001	0.00000	7159	0.04	0.04	0.05
2006	<i>not observed</i>	–	–	–	–	–	–	5532	–	–	–
2007	7.12%	929	0.22%	15.26	649983	0.00002	0.00001	9129	0.21	0.03	0.39
2008	5.81%	785	0.00%	0.00	672491	0.00000	0.00000	11576	0.00	0.00	0.00
2009	7.48%	672	0.00%	0.00	751198	0.00000	0.00000	10049	0.00	0.00	0.00
2010	11.94%	1233	0.00%	0.00	1706840	0.00000	0.00000	14290	0.00	0.00	0.00
2011	13.63%	1892	0.11%	19.33	2985964	0.00001	0.00000	21915	0.14	0.05	0.24
2012	13.52%	2122	0.00%	0.00	3014219	0.00000	0.00000	22292	0.00	0.00	0.00
2013	10.74%	1403	0.00%	0.00	2313243	0.00000	0.00000	21538	0.00	0.00	0.00
2014	9.64%	1445	0.00%	0.00	2272045	0.00000	0.00000	23573	0.00	0.00	0.00
2015	9.38%	1984	0.00%	0.00	2275792	0.00000	0.00000	24274	0.00	0.00	0.00
2016	14.33%	2469	0.00%	0.00	2309357	0.00000	0.00000	16116	0.00	0.00	0.00
2017	13.91%	1623	0.00%	0.00	1454896	0.00000	0.00000	10459	0.00	0.00	0.00

Table 55: Coverage information, bycatch rates, and bycatch estimates for Pacific halibut in the California state pink shrimp fisheries by state and year. The WCGOP began observing the CA state pink shrimp fishery in 2004, but was unable to observe it in 2006. Mortality rates are not applied to P. halibut bycatch in this fishery because mortality rates for pink shrimp trawl gear have not been estimated. Coverage rate is defined as the proportion of pink shrimp landings that were observed. (*) = Confidential data; (-) = not observed; LCL/UCL = lower/upper 95% confidence limit.

Year	Observed							Estimated			
	Coverage rate	No. of sets	% sets w/P. halibut	P. halibut bycatch (kg)	Shrimp retained (kg)	Bycatch rate	SE	Shrimp landings (mt)	P. halibut bycatch (mt)	LCL	UCL
California											
2002	<i>not observed</i>	—	—	—	—	—	—	1849	—	—	—
2003	<i>not observed</i>	—	—	—	—	—	—	974	—	—	—
2004	*	*	*	*	*	*	*	992	*	*	*
2005	*	*	*	*	*	*	*	859	*	*	*
2006	<i>not observed</i>	—	—	—	—	—	—	63	—	—	—
2007	*	*	*	*	*	*	*	289	*	*	*
2008	*	*	*	*	*	*	*	945	*	*	*
2009	*	*	*	*	*	*	*	1183	*	*	*
2010	14.99%	134	0.00%	0.00	265531	0.00000	0.00000	1771	0.00	0.00	0.00
2011	12.62%	203	0.00%	0.00	420595	0.00000	0.00000	3333	0.00	0.00	0.00
2012	12.46%	175	0.00%	0.00	347598	0.00000	0.00000	2791	0.00	0.00	0.00
2013	9.19%	188	0.00%	0.00	359770	0.00000	0.00000	3915	0.00	0.00	0.00
2014	15.54%	337	0.00%	0.00	597530	0.00000	0.00000	3845	0.00	0.00	0.00
2015	9.69%	335	0.30%	0.91	334660	0.00000	0.00000	3453	0.01	0.01	0.01
2016	22.75%	405	0.00%	0.00	311723	0.00000	0.00000	1370	0.00	0.00	0.00
2017	16.29%	257	0.00%	0.00	241800	0.00000	0.00000	1484	0.00	0.00	0.00

Table 56: Coverage information, bycatch rates, and bycatch estimates for Pacific halibut in the state California halibut trawl fishery by sector and year. The WCGOP recognizes two sectors; a limited entry sector and an open access sector. In 2010, the LE and OA sectors are combined to maintain confidentiality. Beginning in 2011, the limited entry sector is observed under the IFQ groundfish fishery and estimates for this sector are included in the IFQ tables (above). Mortality rates are not applied to P. halibut bycatch in these fisheries because mortality rates for CA halibut trawl gear have not been estimated. Coverage rate in the CA halibut fishery is defined as the proportion of CA halibut landings that were observed.

California halibut trawl fishery											
Sector	Observed								Estimated		
Year	Fleet observer coverage rate	Number of observed tows	% of tows with P. halibut	P. halibut bycatch (kg)	CA halibut retained (kg)	P. halibut bycatch rate	SE bycatch rate	Total fleet catch of CA halibut (mt)	P. halibut bycatch (mt)	Bycatch lower 95% CI (mt)	Bycatch upper 95% CI (mt)
Limited Entry Sector											
2010	7.12%	153	0.00%	0.000	8798	0.00000	0.00000	124	0.000	0.000	0.000
2002	3.41%	52	0.00%	0.000	3590	0.00000	0.00000	105	0.000	0.000	0.000
2003	18.09%	207	0.00%	0.000	19093	0.00000	0.00000	106	0.000	0.000	0.000
2004	23.10%	171	0.58%	3.493	31488	0.00011	0.00001	136	0.015	0.012	0.018
2005	16.16%	235	0.43%	4.717	30514	0.00015	0.00001	189	0.029	0.024	0.034
2006	11.95%	224	0.89%	2.903	14286	0.00020	0.00007	120	0.024	0.007	0.042
2007	29.29%	81	1.23%	8.119	5447	0.00149	0.00023	19	0.028	0.019	0.036
2008	26.48%	118	8.47%	82.605	9637	0.00857	0.00162	36	0.312	0.196	0.428
LE & OA Sectors combined											
2009	6.14%	29	0.00%	0.000	2898	0.00000	0.00000	47	0.000	0.000	0.000
2011-2017					Observed under IFQ fisheries, see Table 18						
Open Access Sector											
2002	not observed	—	—	—	—	—	—	36	—	—	—
2003	7.68%	110	0.00%	0.000	1977	0.00000	0.00000	26	0.000	0.000	0.000
2004	7.21%	244	1.64%	49.351	5102	0.00967	0.00334	71	0.685	0.221	1.148
2005	11.52%	362	0.00%	0.000	7431	0.00000	0.00000	65	0.000	0.000	0.000
2006	not observed	—	—	—	—	—	—	55	—	—	—
2007	7.01%	227	0.00%	0.000	2745	0.00000	0.00000	39	0.000	0.000	0.000
2008	5.14%	199	0.00%	0.000	2666	0.00000	0.00000	52	0.000	0.000	0.000
2009	0.77%	30	0.00%	0.000	634	0.00000	0.00000	82	0.000	0.000	0.000
2011	15.57%	204	0.00%	0.000	12446	0.00000	0.00000	80	0.000	0.000	0.000
2012	6.35%	78	0.00%	0.000	3541	0.00000	0.00000	56	0.000	0.000	0.000
2013	6.22%	81	0.00%	0.000	4305	0.00000	0.00000	69	0.000	0.000	0.000
2014	22.27%	145	0.00%	0.000	18139	0.00000	0.00000	81	0.000	0.000	0.000
2015	33.26%	339	0.00%	0.000	30615	0.00000	0.00000	92	0.000	0.000	0.000
2016	30.49%	500	0.00%	0.000	27326	0.00000	0.00000	90	0.000	0.000	0.000
2017	29.42%	556	0.00%	0.000	28373	0.00000	0.00000	96	0.000	0.000	0.000

Table 57: Observer coverage information for the Sea Cucumber fishery by year. The WCGOP began observing the Sea Cucumber fishery in 2017.

Sea Cucumber Fishery									
Year	Gear	Observed					Fleet-wide		
		No. of Vessels	No. of Trips	No. of Sets	Cucumber Retained (mt)	P. halibut (mt) ¹	Cucumber Landed (mt)	Total Landed (mt)	Coverage
2017	Bottom Trawl	3	22	52	1.95	0	13.82	13.88	14.0%

¹ includes both discarded and retained

Table 58: Observer coverage information for the Ridgeback Prawn fishery by year. The WCGOP began observing the Ridgeback Prawn fishery in 2017.

Ridgeback Prawn Fishery									
Year	Gear	Observed					Fleet-wide		
		No. of Vessels	No. of Trips	No. of Sets	Prawn Retained (mt)	P. halibut (mt) ¹	Prawn Landed (mt)	Total Landed (mt)	Coverage
2017	Shrimp Trawl	9	67	297	17.01	0	148.45	185.92	11.0%

¹ includes both discarded and retained

8.6 Tables: At-Sea Hake Fisheries

Table 59: Coverage information and Pacific halibut bycatch in the At-sea Pacific hake fisheries by sector and year. Gear specific mortality rates cannot be applied to P. halibut bycatch in this fishery because mortality rates have not been determined for midwater trawl gear. (*) confidential

At-sea Pacific hake fishery				
Sector	Fleet	Number of	% of sets	P. halibut
Year	observer	observed	with P.	bycatch
	coverage	sets	halibut	(mt)
Catcher-Processor				
2002	100%	559	3.22%	1.013
2003	100%	768	4.04%	2.619
2004	100%	1501	1.07%	0.806
2005	100%	1337	1.72%	1.217
2006	100%	1497	0.27%	0.111
2007	100%	1577	1.65%	0.504
2008	100%	1886	5.51%	2.070
2009	100%	868	0.12%	0.014
2010	100%	1068	0.47%	0.143
2011	100%	1549	1.48%	0.488
2012	100%	1107	2.35%	0.542
2013	100%	1459	1.30%	0.667
2014	100%	1696	0.06%	0.039
2015	100%	1519	0.07%	0.012
2016	100%	2205	0.05%	0.028
2017	100%	2159	0.51%	0.264
Mothership Catcher Vessels				
2002	100%	574	0.17%	0.048
2003	100%	536	0.37%	0.035
2004	100%	571	1.23%	0.323
2005	100%	1040	1.25%	0.567
2006	100%	1283	1.95%	0.532
2007	100%	1147	2.01%	0.621
2008	100%	1349	2.82%	0.629
2009	100%	600	3.50%	0.255
2010	100%	908	3.41%	1.080
2011	100%	1248	0.48%	0.085
2012	100%	949	0.63%	0.099
2013	100%	1256	2.15%	0.397
2014	100%	1308	1.22%	0.332
2015	100%	640	0.31%	0.049
2016	100%	1565	0.51%	0.123
2017	100%	1309	0.84%	0.289

Table 60: Coverage information and Pacific halibut bycatch in the Tribal At-sea Pacific hake fishery by year. Tribal At-sea P. hake fishery has not operated since 2012. Gear specific mortality rates cannot be applied to P. halibut bycatch in this fishery because mortality rates have not been determined for midwater trawl gear. (*) confidential

At-sea Pacific hake fishery				
Sector				
Year	Fleet observer coverage	Number of observed sets	% of sets with P. halibut	P. halibut bycatch (mt)
Tribal Sector				
2002	100%	633	0.32%	0.079
2003	100%	540	0.00%	0.000
2004	100%	632	0.00%	0.000
2005	100%	633	0.79%	0.182
2006	100%	160	3.12%	0.192
2007	100%	156	0.64%	0.053
2008	100%	382	7.33%	1.280
2009	100%	404	0.99%	0.064
2010	100%	516	3.49%	0.349
2011	100%	228	0.88%	0.034
2012	100%	4	0.00%	0.000

Table 61: Physical P. halibut length frequencies (cm) collected by A-SHOP observers in the At-sea hake fishery (2002-2017). Length bins include the lower bound and exclude the upper bound.

At-sea Hake				
Length bin (cm)	No. of Excellent	No. of Poor	No. of Dead	No. of Unknown
58-60	0	0	2	0
60-62	0	1	3	0
62-64	0	0	2	0
64-66	0	2	6	0
66-68	0	0	9	0
68-70	1	0	10	0
70-72	0	1	21	0
72-74	1	0	23	0
74-76	0	1	19	0
76-78	1	1	21	1
78-80	0	0	27	0
80-82	1	0	46	1
82-84	2	0	21	0
84-86	0	1	31	0
86-88	0	2	28	0
88-90	2	0	40	2
90-92	0	0	37	1
92-94	0	0	31	0
94-96	1	2	39	0
96-98	0	0	23	0
98-100	1	0	31	0
100-102	0	4	32	0
102-104	1	1	21	0
104-106	0	1	22	0
106-108	0	0	23	0
108-110	2	1	22	1
110-112	1	0	19	0
112-114	1	0	9	0
114-116	0	0	8	0
116-118	0	1	10	1
118-120	0	0	7	0
120-122	0	0	6	0
122-124	0	0	7	0
124-126	0	0	4	0
126-128	0	1	1	0
128-130	0	0	4	0
130-132	0	0	6	0
132-134	0	0	2	0
136-138	0	0	2	0
138-140	0	0	1	0
140-142	0	0	2	0
142-144	0	0	1	0
154-156	0	0	1	0

8.7 Tables: IFQ Electronic Monitoring

Table 62: Number of vessels, trips, and tows (or sets) and gross metric tons of Pacific halibut discarded at-sea, P. halibut discarded at sea with mortality rate applied (bottom trawl = 0.90; pot = 0.18) and the P. halibut landed and discarded at the dock (mortality rate = 1.0) under the IFQ Electronic Monitoring Exempted Fishing Permit (EFP). All participating vessels carry electronic monitoring equipment on all fishing trips. Data are summarized from the EM program administered by Pacific States Marine Fisheries Commission

Area		Electronic Monitoring EFP (IFQ)					
Depth (fm)	Year	No. of vessels	No. of trips	No. of sets	discarded at sea (gross, mt)	discard mortality (mt)	discarded at dock (mt)
Bottom Trawl							
North of Pt. Chehalis							
All depths							
	2015	0	0	0	0.00	0.00	0.00
	2016	0	0	0	0.00	0.00	0.00
	2017	0	0	0	0.00	0.00	0.00
South of Pt. Chehalis							
All depths							
	2015	5	23	139	0.18	0.16	0.00
	2016	7	98	574	2.93	2.63	0.00
	2017	11	149	870	5.36	4.83	0.05
Pot							
North of 40°10' N. lat.							
All depths							
	2015	3	32	302	0.89	0.16	0.00
	2016	5	44	417	0.07	0.01	0.00
	2017	5	51	578	0.75	0.13	0.00
South of 40°10' N. lat.							
All depths							
	2015	5	26	398	0.00	0.00	0.00
	2016	4	27	603	0.00	0.00	0.00
	2017	7	31	648	0.00	0.00	0.00
Midwater Hake Trawl							
North of 40°10' N. lat.							
	2015	17	454	1178	0.00	0.00	0.56
	2016	20	648	1411	0.01	0.01	0.65
	2017	22	1104	2072	0.00	0.00	0.46
Midwater Rockfish Trawl							
North of 40°10' N. lat.							
	2015	8	26	81	0.00	0.00	0.00
	2016	6	30	74	0.00	0.00	0.00
	2017	10	53	88	0.00	0.00	0.00

8.8 Tables: Other EFP fleet and PHLB catch summaries

Table 63: Metric tons of Pacific halibut discarded at sea and landed and discarded at the dock on observed Exempted Fishing Permit (EFP) vessels. Note: This does not contain the Catch Shares Electronic Monitoring EFP data, see Table 62 for those data.

Observed							
Year	Gear	Sector	No. vessels	No. trips	No. hauls	P. halibut discarded (mt)	P. halibut landed (mt)
2002	EFP	Bottom & Midwater Trawl	7	38	279	53.36	0.00
2003	EFP	Bottom & Midwater Trawl	12	156	1491	50.79	0.00
2004	EFP	Bottom & Midwater Trawl	6	59	427	30.68	0.00
2005	EFP	—	0	0	0	0.00	0.00
2006	EFP	Bottom & Midwater Trawl	9	48	80	0.00	0.00
2007	EFP	—	0	0	0	0.00	0.00
2008	EFP	Fixed Gears	3	29	162	0.00	0.00
2009	EFP	Fixed Gears	5	83	141	0.00	0.00
2010	EFP	Fixed Gears	6	136	389	0.00	0.00
2011	EFP	—	0	0	0	0.00	0.00
2013	EFP	Fixed Gears	2	5	166	0.00	0.00
2014	EFP	Fixed Gears	3	11	21	0.00	0.00
2015	EFP	Fixed Gears	1	3	4	0.00	0.00
2016	EFP	Fixed Gears	1	3	3	0.00	0.00
2017	EFP	Fixed Gears	4	17	25	0.00	0.00

Table 64: Metric tons of Pacific halibut landed in non-groundfish fisheries that are not observed by the NWFSC Observer Program. Data are summarized from the PacFIN fish tickets and do not include any P. halibut landed under the IPHC P. halibut directed fishery.

Year	Sector	P. halibut landings (mt)
2002	Other Fisheries	24.85
2003	Other Fisheries	28.33
2004	Other Fisheries	26.63
2005	Other Fisheries	26.39
2006	Other Fisheries	20.73
2007	Other Fisheries	14.95
2008	Other Fisheries	9.14
2009	Other Fisheries	7.07
2010	Other Fisheries	17.21
2011	Other Fisheries	14.21
2012	Other Fisheries	17.99
2013	Other Fisheries	18.35
2014	Other Fisheries	21.33
2015	Other Fisheries	17.23
2016	Other Fisheries	15.42
2017	Other Fisheries	20.05

Table 65: Discard estimates for all fishery sectors observed by the NWFSC Groundfish Observer Program, 2002-2017. Total discard mortality estimates are also provided where discard mortality rates were applied. Estimates include individuals discarded at the dock (100% mortality). Ridgeback Prawn and Sea Cucumber fisheries had zero (0) observed P. halibut catch.

Year	LE bottom trawl 2002-10	IFQ Fishery ⁷					Non-Nearshore fixed gear			Nearshore Fixed Gear ³	Pink Shrimp ³	CA Halibut ^{3,6}	P. halibut Derby	At-sea Hake ³	Total
		Bottom Trawl ^{1,2,7}	LE CA Halibut ^{1,3}	Hook & Line	Pot ⁷	Midwater Rockfish ^{3,4,7}	Midwater Hake ^{2,3,5,7}	LE Endorsed	LE Non- Endsd.	OA					
Gross Discard Estimates (mt)															
2002	524.41							141.76	0.00	-	-	0.00		1.14	667.31
2003	186.65							197.11	0.17	-	0.00	0.00		2.65	386.58
2004	212.43							238.57	0.00	-	1.00	0.00		1.13	453.83
2005	460.35							237.82	0.00	-	2.20	0.04		1.97	702.41
2006	390.91							668.62	0.00	-	0.53	0.02		0.83	1060.91
2007	294.38							132.27	1.73	21.66	0.09	0.21		1.18	451.55
2008	305.21							259.46	2.99	40.12	0.35	0.00		3.98	612.42
2009	385.24							321.60	0.25	35.25	1.28	0.00		0.33	743.95
2010	265.08							137.60	0.39	32.56	0.08	0.00		1.57	437.28
2011		64.14	0	6.06	3.36	*	0.35	136.93	21.30	13.03	3.07	0.19		0.61	249.04
2012		67.13	*	14.66	1.89	0.0	0.62	151.25	16.57	23.63	2.25	0.00		0.64	278.64
2013		66.08	see ¹	3.00	0.98	0.0	1.34	22.23	0.01	1.78	1.36	0.00		1.06	97.84
2014		55.89	see ¹	3.80	0.32	0.0	1.36	174.79	0.00	3.49	0.95	0.00		0.37	240.97
2015		69.27	see ¹	9.49	2.21	0.0	0.70	127.45	0.46	10.13	1.44	0.01		0.06	221.22
2016		58.85	see ¹	6.39	1.77	0.0	0.68	173.88	5.09	42.78	3.01	0.00		0.15	292.60
2017		64.77	see ¹	4.14	1.84	0.0	0.51	293.56	0.72	48.24	1.55	0.00	25.74	0.55	441.62
Total Discard Mortality (mt)															
2002	344.82							22.76	0.00	-	-	0.00		1.14	368.72
2003	124.43							31.54	0.03	-	0.00	0.00		2.65	158.65
2004	133.12							38.82	0.00	-	1.00	0.00		1.13	174.77
2005	286.52							38.10	0.00	-	2.20	0.04		1.97	328.86
2006	242.47							107.30	0.00	-	0.53	0.02		0.83	351.15
2007	208.81							21.24	0.28	3.48	0.09	0.21		1.18	235.32
2008	207.81							41.65	0.48	6.42	0.35	0.00		3.98	261.00
2009	251.1							51.47	0.04	5.65	1.28	0.00		0.33	309.87
2010	180.97							22.12	0.06	5.22	0.08	0.00		1.57	210.02
2011		31.30	0	0.97	0.89	*	0.35	12.07	3.20	2.09	3.07	0.19		0.61	54.74
2012		36.13	*	2.34	0.51	0.0	0.62	24.94	0.73	1.61	2.25	0.00		0.64	69.77
2013		32.41	see ¹	0.48	0.21	0.0	1.34	2.94	0.00	0.07	1.36	0.00		1.06	39.87
2014		26.28	see ¹	0.61	0.08	0.0	1.36	30.16	0.00	0.35	0.95	0.00		0.37	60.16
2015		33.36	see ¹	1.52	0.38	0.0	0.70	10.37	0.02	0.46	1.44	0.01		0.06	48.32
2016		33.28	see ¹	1.02	0.18	0.0	0.68	16.62	0.91	2.56	3.01	0.00		0.15	58.41
2017		35.11	see ¹	0.66	0.78	0.0	0.51	37.95	0.03	3.73	1.55	0.00	2.26	0.55	83.13

¹Starting in 2013, LE CA Halibut estimates are combined with IFQ Bottom Trawl estimates.

²Includes a small amount landed and discarded at the dock.

³100% mortality rate

⁴from 2011-14, 'Midwater Trawl'

⁵from 2011-14, 'Shoreside Hake'

⁶Starting in 2011, this sector only includes OA CA halibut

⁷Includes P. halibut catch from IFQ electronic monitoring EFP

Note: Ridgeback Prawn and Sea Cucumber fisheries had zero (0) observed P. halibut catch

9 FIGURES




Figure 2: Number of vessels by month for IFQ bottom trawl vessels in 2017 (solid line) and averaged over the 2011–17 period (dotted line). Grey ribbon represents the monthly maximum and minimum across 2011-2017.




Figure 3: Number of tows by month for IFQ bottom trawl vessels in 2017 (solid line) and averaged over the 2011–17 period (dotted line). Grey ribbon represents the monthly maximum and minimum across 2011-2017.




Figure 4: Tow hours by month for IFQ bottom trawl vessels in 2017 (solid line) and averaged over the 2011–17 period (dotted line). Grey ribbon represents the monthly maximum and minimum across 2011-2017.




Figure 5: Estimated discard mortality of *P. halibut* in the non-nearshore fixed gear fishery by sector and year. We apply a fixed average discard rate from 2007-08 data to generate 2002-06 discard estimates for the OA sector because only a portion of the fishery was observed 2002-06. The 'Other fixed gear sectors' includes LE sablefish non-endorsed and OA fixed gear vessels fishing with pot gear. The inset is an expanded view of each of the sectors, except LE sablefish endorsed longline (LL) gear, during years with very small bycatch.





Figure 6: Length frequency distribution of discarded Pacific halibut on WCGOP observed Non-Nearshore Fixed Gear limited entry (LE) and open access (OA) groundfish vessels from September 2003 through December 2017. The majority of P. halibut lengths collected in this fishery were visual estimates (grey bars) which are only estimated in 10 cm bins. The sublegal-legal size cut-off (82 cm) is indicated by a vertical dashed line.

Figure 7: Spatial distribution of Pacific halibut bycatch (mt/km²) observed by West Coast Ground-fish Observer Program (2002-2017), off the U.S. West Coast. Gear types observed by the WCGOP include bottom trawl, midwater trawl, shrimp trawl, fixed gear hook-&-line and pot gear. The five catch classifications were defined by excluding any 0 values and then applying the Jenks natural breaks classification method. Cells (200 sq. km) with less than 3 vessels were omitted from the map to maintain confidentiality.






Figure 8: Number of sets, trips, and vessels by opening day for the P. halibut Derby fishery

A Appendices

A.1 IFQ Electronic Monitoring DMR comparison

PFMC staff, the NOAA Western Regional Office, and IPHC have requested a comparison of discard mortality rates (DMR) for bottom trawl and pot vessels in the IFQ program that carry electronic monitoring (EM) equipment versus those that carry observers on 100% of the fishing trips. When notified, EM vessels are required to carry observers for scientific observation, including collection of Pacific halibut viabilities. The WCGOP aims to observe approximately 30% of EM fishing trips. DMRs for EM vessels were calculated and compared using two methods:

1. Observer Viability Method
2. Time on Deck Model

The Observer Viability Method used human observer data collected on EM vessels. These data were stratified to match, as closely as possible, the current stratification used in the IFQ fishery while meeting confidentiality requirements. Confidentiality of EM data required combining strata across years, depths and areas. Mortality data from non-EM IFQ vessels is also shown for comparison purposes. Other than slight modification of stratification to maintain confidentiality, the observer viability method is identical to the method described in Section [3.2.1](#).

The Time-on-Deck model was developed in a collaborative process between PSMFC and the PFMC's Groundfish Management Team (GMT). The model measures the time each fish spends out of the water which correlates with P. halibut viability: the less time a fish spends out of the water the higher probability of the fish being in 'excellent' viability condition and therefore lower mortality rate. The Time-on-Deck model substitutes for a viability assessment on EM vessels when fisheries observers are not present on the vessel to assess viabilities. The model and discussion are detailed in a [PSMFC report](#) and a [PFMC GMT report](#).

The comparison below is for informational purposes only. Due to low sample sizes the NWFSC Observer Program cautions against using these estimates for management purposes. Data from EM pot vessels were obtained 2015-2017 on pot vessels, but only from 2016-2017 on bottom trawl EM vessels. The corresponding non-EM data (i.e., 2015-2017 pot; 2016-2017 bottom trawl) were used to allow direct comparison between vessels with and without EM. Confidentiality in the EM fleet precluded the use of the full stratification currently used in the Catch Shares fishery (see Tables [14](#) & [16](#)).

Table 66: Observed number of IFQ Electronic Monitoring bottom trawl vessels, trips, and sets that caught Pacific halibut and the number of P. halibut in each viability category. Exc = Excellent

Year	IFQ EM Bottom Trawl					
	Observed Number					
	Vessels	Trips	Sets	Exc	Poor	Dead
South of Pt. Chehalis						
2016	5	13	28	16	14	51
2017	3	4	6	7	4	9

Table 67: P. halibut observed weight (Obs), estimated total at-sea gross weight of discards (Grs), and estimated total discard weight (Dis) with the mortality rate applied based on observer viability, in each viability category from IFQ bottom trawl vessels. Discard mortality rates (DMR) are shown for the observer viability method (Obs) and the Time-on-Deck model (ToD) as applied by the Pacific States Marine Fisheries Commission. The ToD Discard is the weight of each fish multiplied by the mortality rate based on the time-on-deck model and summed across all individuals. Electronic Monitoring (EM) vessels carried electronic monitoring equipment. Viabilities on both EM and non-EM vessels were obtained by at-sea human observers. EM vessels only fished south of Pt. Chehalis. Values north of Pt. Chehalis represent non-EM vessels and are presented for comparison purposes only. All weights are metric tons (mt). Exc = Excellent = 0.10 mortality rate; Poor = 0.55 mortality rate; Dead = 0.90 mortality rate; All = all years combined

		Bottom Trawl														
		Exc (mt)			Poor (mt)			Dead (mt)			Total (mt)		Observer		ToD	
	Year	Obs	Grs	Dis	Obs	Grs	Dis	Obs	Grs	Dis	Obs	Grs	Dis	DMR	Dis	DMR
South of Pt. Chehalis																
EM	2016	0.17	0.66	0.13	0.14	0.53	0.29	0.46	1.74	1.56	0.77	2.93	1.99	0.68	—	—
	2017	0.09	2.31	0.46	0.03	0.90	0.50	0.08	2.15	1.94	0.20	5.36	2.89	0.54	3.45	0.65
	All	0.26	2.26	0.45	0.17	1.51	0.83	0.54	4.70	4.23	0.97	8.47	5.51	0.65	—	—
Non-EM	2016	14.64	15.71	3.14	4.88	5.24	2.88	16.74	17.96	16.16	36.26	38.90	22.18	0.57	—	—
	2017	18.36	20.16	4.03	5.24	5.75	3.16	13.70	15.04	13.54	37.30	40.95	20.73	0.51	—	—
	All	33.00	53.90	10.78	10.12	16.52	9.09	30.44	49.71	44.74	73.56	120.14	64.61	0.54	—	—
North of Pt. Chehalis																
	2016	7.55	8.64	1.73	2.44	2.79	1.54	4.78	5.47	4.93	14.77	16.90	8.19	0.48	—	—
	2017	8.03	9.14	1.83	2.18	2.48	1.36	5.82	6.63	5.97	16.03	18.25	9.16	0.50	—	—
	All	15.58	32.32	6.46	4.62	9.58	5.27	10.61	22.00	19.80	30.81	63.90	31.53	0.49	—	—

Table 68: Observed number of IFQ Electronic Monitoring pot vessels, trips, and sets that caught Pacific halibut and the number of P. halibut in each viability category. Exc = Excellent

Year	IFQ EM Pot					
	Observed Number					
	Vessels	Trips	Sets	Exc	Poor	Dead
Coastwide						
2015-17	3	10	16	19	4	11

Table 69: P. halibut observed weight (Obs), estimated total at-sea gross weight of discards (Grs), and estimated total discard weight (Dis) with the mortality rate applied based on observer viability, in each viability category from IFQ pot vessels. Total discard mortality rates (DMR) are shown for the observer viability method (Obs). Electronic Monitoring (EM) vessels carried electronic monitoring equipment. Viabilities on both EM and non-EM vessels were obtained by at-sea human observers. All weights are metric tons (mt). Exc = Excellent = 0.10 mortality rate; Poor = 0.55 mortality rate; Dead = 0.90 mortality rate; All = all years combined

		Pot												
		Exc (mt)			Poor (mt)			Dead (mt)			Total (mt)			DMR
		Year	Obs	Grs	Dis	Obs	Grs	Dis	Obs	Grs	Dis	Obs	Grs	
Coastwide														
EM	All	0.16	0.97	0	0.03	0.19	0.19	0.09	0.54	0.54	0.28	1.71	0.73	0.43
Non-EM	All	2.81	3.07	0	0.43	0.47	0.47	0.51	0.56	0.56	3.75	4.10	1.03	0.25

A.2 Catch Shares Weighted Length Frequencies

Weighted catch composition data from the IFQ fishery for bottom trawl and pot gears. The frequency within each length bin was weighted based on the following equation:

$$n_{w_l} = n_l \times \frac{W_{st}}{\sum_l w_{stl}} \times \frac{\sum_t W_{st}}{W_{st}} \times \frac{\hat{W}_s}{\sum_t W_{st}} = n_l \times \frac{\hat{W}_s}{\sum_l w_{stl}} \quad (9)$$

where:

s = stratum

t = tow

l = length bin

n = number of measured fish

w = total weight of fish, as determined through the IPHC length-weight relationship (Table 9 in Appendix A.4)

W = total observed discard weight of Pacific halibut

\hat{W} = estimated total discard weight of P. halibut

Table 70: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using bottom trawl gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. Since 2013, IFQ bottom trawl lengths could also include lengths taken on both IFQ and LE California halibut bottom trawl fisheries.

Bottom Trawl							
Length bin (cm)	2011	2012	2013	2014	2015	2016	2017
10	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
12	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
14	0.0000	0.0000	0.0000	0.0000	0.0000	0.0370	0.0000
16	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
18	0.0065	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
20	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
22	0.0000	0.0109	0.0000	0.0000	0.0000	0.0000	0.0000
24	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
26	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
28	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
30	0.0000	0.0076	0.0037	0.0000	0.0000	0.0000	0.0000
32	0.0000	0.0061	0.0030	0.0028	0.0000	0.0016	0.0000
34	0.0000	0.0101	0.0000	0.0000	0.0000	0.0000	0.0000
36	0.0000	0.0043	0.0000	0.0000	0.0009	0.0008	0.0000
38	0.0000	0.0109	0.0000	0.0000	0.0027	0.3872	0.0011
40	0.0014	0.0054	0.0019	0.0014	0.0091	0.3255	0.0009

Table 70: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using bottom trawl gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. Since 2013, IFQ bottom trawl lengths could also include lengths taken on both IFQ and LE California halibut bottom trawl fisheries. *(continued)*

Length bin (cm)	Bottom Trawl						
	2011	2012	2013	2014	2015	2016	2017
42	0.0023	0.0110	0.0000	0.0000	0.0057	0.0068	0.0025
44	0.0000	0.0024	0.0000	0.0000	0.0061	0.0022	0.0007
46	0.0003	0.0073	0.0006	0.0004	0.0023	0.0028	0.0013
48	0.0029	0.0064	0.0028	0.0011	0.0044	0.0101	0.0026
50	0.0034	0.0071	0.0032	0.0000	0.0030	0.0044	0.0047
52	0.0046	0.0072	0.0048	0.0021	0.0010	0.0035	0.0045
54	0.0079	0.0057	0.0482	0.0044	0.0052	0.0082	0.0104
56	0.0074	0.0062	0.0074	0.0050	0.0069	0.1125	0.0105
58	0.0194	0.0148	0.0474	0.0141	0.0120	0.0119	0.0152
60	0.0324	0.0294	0.0562	0.0305	0.0186	0.0151	0.0179
62	0.0441	0.0428	0.0553	0.0551	0.0334	0.0272	0.0260
64	0.0565	0.0529	0.0848	0.0740	0.0472	0.0273	0.0240
66	0.0589	0.0542	0.0710	0.0776	0.2292	0.0427	0.0276
68	0.0571	0.0623	0.1653	0.2307	0.0770	0.0734	0.0545
70	0.0762	0.0711	0.1995	0.1719	0.0817	0.0712	0.0484
72	0.0737	0.0708	0.1645	0.0848	0.2100	0.1733	0.0624
74	0.0858	0.0678	0.1020	0.1820	0.1822	0.0680	0.0658
76	0.0669	0.0629	0.1228	0.1326	0.1239	0.0659	0.0621
78	0.0561	0.0536	0.1091	0.0591	0.1132	0.0687	0.0712
80	0.0571	0.0486	0.1213	0.0760	0.0538	0.0587	0.3426
82	0.0478	0.0469	0.1003	0.1693	0.0528	0.0528	0.0676
84	0.0460	0.0376	0.0695	0.0821	0.1522	0.0703	0.0573
86	0.0309	0.0302	0.0610	0.1364	0.1048	0.0374	0.0471
88	0.0284	0.0255	0.0505	0.0822	0.0948	0.0315	0.0471
90	0.0258	0.0237	0.0487	0.0912	0.0267	0.0265	0.2364
92	0.0213	0.0214	0.0579	0.0162	0.0496	0.0432	0.0320
94	0.0167	0.0160	0.0429	0.0117	0.0665	0.0340	0.1988
96	0.0134	0.0110	0.0499	0.0408	0.0116	0.0131	0.0211
98	0.0096	0.0097	0.0156	0.0077	0.0109	0.0101	0.1695
100	0.0086	0.0084	0.0138	0.0075	0.0526	0.0263	0.0142
102	0.0070	0.0075	0.0228	0.0421	0.0072	0.0077	0.0121

Table 70: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using bottom trawl gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. Since 2013, IFQ bottom trawl lengths could also include lengths taken on both IFQ and LE California halibut bottom trawl fisheries. *(continued)*

Length bin (cm)	Bottom Trawl						
	2011	2012	2013	2014	2015	2016	2017
104	0.0054	0.0043	0.0102	0.0165	0.0258	0.0049	0.0088
106	0.0039	0.0036	0.0180	0.0025	0.0043	0.0043	0.0058
108	0.0030	0.0034	0.0118	0.0020	0.0205	0.0038	0.1117
110	0.0025	0.0033	0.0063	0.0312	0.0182	0.0025	0.1044
112	0.0021	0.0021	0.0180	0.0104	0.0024	0.0024	0.0029
114	0.0017	0.0015	0.0124	0.0009	0.0160	0.0010	0.0018
116	0.0011	0.0012	0.0043	0.0005	0.0146	0.0111	0.0014
118	0.0009	0.0007	0.0007	0.0004	0.0009	0.0008	0.0010
120	0.0005	0.0008	0.0066	0.0003	0.0248	0.0006	0.0008
122	0.0005	0.0005	0.0005	0.0073	0.0006	0.0006	0.0005
124	0.0006	0.0003	0.0002	0.0136	0.0113	0.0003	0.0004
126	0.0003	0.0004	0.0028	0.0063	0.0002	0.0003	0.0002
128	0.0003	0.0000	0.0001	0.0002	0.0003	0.0074	0.0000
130	0.0001	0.0000	0.0000	0.0057	0.0002	0.0001	0.0001
132	0.0002	0.0001	0.0000	0.0001	0.0001	0.0001	0.0001
134	0.0000	0.0000	0.0001	0.0053	0.0000	0.0001	0.0000
136	0.0001	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000
138	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000
140	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000
142	0.0001	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000
144	0.0001	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000
146	0.0000	0.0000	0.0000	0.0039	0.0000	0.0000	0.0000
148	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000

Table 71: Percentage of weighted length measurements in each viability category, for IFQ bottom trawl vessels by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. Since 2013, IFQ bottom trawl lengths could also include lengths taken on both IFQ and LE California halibut bottom trawl fisheries.

Length bin (cm)	Bottom Trawl																				
	Excellent							Poor							Dead						
	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017
10	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
12	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
14	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%
16	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
18	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
20	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
22	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
24	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
26	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
28	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
30	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%
32	0.0%	50.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	50.0%	100.0%	0.0%	0.0%	0.0%	0.0%
34	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
36	0.0%	100.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
38	0.0%	82.7%	0.0%	0.0%	92.7%	100.0%	0.0%	0.0%	15.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.2%	0.0%	0.0%	7.3%	0.0%	100.0%
40	0.0%	82.2%	22.2%	0.0%	32.3%	88.3%	0.0%	100.0%	0.0%	0.0%	0.0%	38.4%	0.0%	0.0%	0.0%	17.8%	77.8%	100.0%	29.3%	11.7%	100.0%
42	47.9%	68.0%	0.0%	0.0%	56.6%	100.0%	0.0%	52.1%	23.6%	0.0%	0.0%	19.7%	0.0%	69.1%	0.0%	8.5%	0.0%	0.0%	23.7%	0.0%	30.9%
44	0.0%	46.8%	0.0%	0.0%	50.6%	100.0%	0.0%	0.0%	0.0%	0.0%	14.3%	0.0%	0.0%	0.0%	0.0%	53.2%	0.0%	0.0%	35.1%	0.0%	100.0%
46	0.0%	83.4%	0.0%	0.0%	83.1%	82.9%	0.0%	0.0%	16.6%	0.0%	0.0%	16.9%	0.0%	100.0%	100.0%	0.0%	100.0%	100.0%	0.0%	17.1%	0.0%
48	24.9%	96.4%	34.3%	100.0%	78.4%	86.2%	0.0%	24.9%	0.0%	29.0%	0.0%	21.6%	13.8%	61.1%	50.1%	3.6%	36.7%	0.0%	0.0%	0.0%	38.9%
50	29.9%	66.0%	20.7%	0.0%	2.8%	54.5%	30.9%	0.0%	10.4%	22.1%	0.0%	0.0%	8.5%	18.8%	70.1%	23.6%	57.2%	0.0%	97.2%	37.1%	50.2%
52	23.1%	52.4%	29.7%	30.1%	100.0%	80.9%	0.0%	42.3%	15.3%	22.4%	11.6%	0.0%	14.0%	9.2%	34.6%	32.3%	48.0%	58.3%	0.0%	5.1%	90.8%
54	15.6%	59.9%	40.3%	50.8%	55.2%	67.9%	34.4%	43.2%	29.4%	18.2%	0.0%	34.6%	5.6%	13.9%	41.2%	10.7%	41.5%	49.2%	10.3%	26.5%	51.7%
56	21.0%	44.2%	54.8%	35.4%	38.1%	60.6%	19.8%	45.5%	13.9%	1.9%	0.8%	17.6%	17.0%	27.9%	33.5%	41.9%	43.2%	63.8%	44.3%	22.4%	52.4%
58	19.8%	41.0%	36.5%	32.9%	38.4%	57.2%	26.6%	31.2%	10.0%	23.0%	31.3%	23.4%	10.8%	5.9%	48.9%	49.0%	40.5%	35.8%	38.2%	32.0%	67.4%
60	32.8%	36.8%	39.5%	38.7%	52.4%	40.6%	36.8%	24.3%	21.8%	8.3%	23.6%	9.5%	14.3%	11.4%	42.9%	41.4%	52.1%	37.7%	38.1%	45.1%	51.9%
62	37.8%	40.0%	43.4%	43.4%	52.1%	34.2%	33.0%	22.7%	20.9%	18.7%	20.1%	13.4%	20.7%	25.4%	39.6%	39.2%	37.9%	36.5%	34.6%	45.1%	41.6%
64	39.6%	32.2%	46.1%	45.0%	47.9%	41.8%	36.8%	18.7%	20.9%	17.6%	19.7%	12.4%	14.8%	14.9%	41.7%	46.9%	36.3%	35.3%	39.7%	43.4%	48.3%
66	36.7%	35.9%	45.1%	48.9%	44.1%	32.5%	34.5%	21.0%	22.3%	14.3%	23.7%	20.3%	18.0%	17.8%	42.3%	41.9%	40.6%	27.5%	35.6%	49.5%	47.7%
68	42.6%	35.1%	50.5%	46.7%	47.7%	46.3%	43.6%	12.0%	21.5%	12.3%	20.8%	16.3%	12.9%	14.1%	45.3%	43.4%	37.2%	32.5%	35.9%	40.8%	42.3%
70	41.6%	39.6%	45.2%	53.6%	46.1%	37.6%	44.7%	20.8%	19.5%	17.1%	17.8%	16.5%	13.9%	15.4%	37.7%	40.9%	37.7%	28.6%	37.4%	48.5%	39.9%
72	38.6%	32.2%	48.6%	50.7%	49.3%	35.2%	39.9%	20.9%	18.8%	16.9%	18.4%	14.0%	17.7%	19.5%	40.5%	49.0%	34.5%	30.8%	36.7%	47.1%	40.6%
74	40.0%	32.5%	47.4%	53.7%	52.4%	36.3%	39.3%	17.4%	21.9%	19.1%	14.8%	14.2%	15.4%	14.0%	42.6%	45.7%	33.5%	31.6%	33.5%	48.3%	46.7%
76	45.5%	36.9%	45.0%	44.0%	47.1%	35.9%	36.8%	17.0%	17.2%	17.8%	18.2%	13.2%	17.4%	18.4%	37.5%	45.9%	37.2%	37.7%	39.7%	46.7%	44.8%
78	41.1%	33.3%	44.6%	52.3%	47.0%	35.8%	42.7%	19.0%	24.6%	16.0%	17.9%	17.0%	12.2%	15.8%	39.9%	42.1%	39.5%	29.8%	35.9%	52.0%	41.6%
80	45.7%	38.8%	53.9%	50.1%	47.4%	39.9%	37.8%	16.0%	18.5%	13.1%	16.6%	16.7%	14.3%	18.8%	38.3%	42.7%	33.0%	33.3%	35.9%	45.8%	43.4%
82	45.8%	36.6%	45.4%	50.6%	46.3%	34.6%	43.2%	19.9%	20.9%	18.3%	11.3%	13.5%	15.8%	16.5%	34.3%	42.5%	36.3%	38.1%	40.3%	49.6%	40.3%
84	50.1%	38.5%	50.6%	45.6%	45.4%	39.5%	39.0%	14.8%	18.9%	14.5%	13.3%	14.3%	14.3%	15.7%	35.1%	42.6%	34.9%	41.1%	40.3%	46.2%	45.3%
86	44.6%	36.4%	55.6%	48.8%	42.0%	35.2%	42.1%	14.6%	21.7%	15.5%	18.1%	20.1%	12.3%	18.5%	40.8%	41.8%	28.9%	33.2%	37.8%	52.5%	39.3%
88	41.7%	39.2%	52.9%	43.5%	51.2%	36.8%	48.1%	16.1%	21.5%	15.2%	22.0%	14.8%	14.4%	11.6%	42.2%	39.3%	31.9%	34.5%	34.0%	48.8%	40.3%
90	48.3%	40.9%	57.9%	43.1%	46.9%	35.1%	47.3%	17.0%	18.9%	13.8%	18.7%	16.6%	15.0%	12.0%	34.7%	40.1%	28.4%	38.2%	36.6%	49.9%	40.7%
92	46.6%	41.0%	58.4%	50.6%	49.1%	32.1%	45.8%	17.3%	20.2%	14.7%	14.0%	19.5%	18.7%	19.0%	36.1%	38.9%	26.9%	35.4%	31.3%	49.1%	35.2%
94	51.2%	46.4%	54.6%	49.4%	44.5%	42.2%	50.1%	20.1%	14.3%	15.6%	17.6%	17.3%	15.4%	12.0%	28.7%	39.3%	29.8%	33.1%	38.2%	42.4%	37.9%
96	49.4%	40.5%	58.5%	57.5%	50.5%	36.7%	48.8%	14.6%	16.9%	12.5%	14.6%	12.4%	15.2%	10.8%	36.0%	42.6%	29.0%	27.9%	37.1%	48.1%	40.4%
98	50.0%	39.7%	52.5%	43.5%	50.5%	41.8%	49.1%	18.2%	17.8%	19.6%	23.2%	16.8%	14.7%	14.4%	31.8%	42.4%	27.9%	33.3%	32.7%	43.5%	36.5%
100	53.8%	43.8%	60.9%	57.3%	60.4%	41.7%	49.0%	18.2%	21.0%	14.8%	5.6%	13.0%	12.9%	13.1%	28.0%	35.2%	24.3%	37.2%	26.6%	45.3%	38.0%
102	47.3%	51.1%	58.6%	52.2%	46.7%	48.0%	52.6%	16.1%	16.5%	14.3%	13.3%	16.9%	14.3%	8.2%	36.7%	32.4%	27.1%	34.5%	36.4%	37.8%	39.2%
104	53.0%	44.5%	55.6%	60.8%	54.6%	43.0%	50.9%	18.9%	10.3%	14.3%	17.8%	12.1%	12.7%	16.3%	28.1%	45.2%	30.1%	21.4%	33.3%	44.3%	32.8%
106	54.2%	39.6%	71.7%	66.2%	50.6%	54.9%	59.5%	18.4%	26.6%	12.7%	9.2%	19.4%	7.1%	13.8%	27.3%	33.8%	15.6%	24.6%	30.0%	38.0%	26.7%
108	53.4%	44.3%	58.5%	62.6%	45.3%	35.7%	60.3%	20.3%	16.4%	14.1%	23.2%	18.9%	20.1%	7.9%	26.3%	39.3%	27.4%	14.2%	35.8%	44.2%	31.8%
110	56.5%	51.4%	56.2%	60.9%	62.0%	33.2%	57.0%	11.2%	14.2%	26.9%	16.0%	12.9%	19.3%	6.7%	32.3%	34.4%	16.9%	23.1%	25.1%	47.5%	36.3%
112	56.6%	54.4%	58.0%	53.5%	30.3%	40.5%	58.3%	22.5%	22.4%	20.7%	14.9%	26.9%	8.0%	20.9%	20.9%	23.2%	21.3%	31.6%	42.8%	51.5%	20.8%
114	49.8%	43.9%	68.4%	64.7%	52.7%	23.2%	60.8%	25.2%	22.7%	12.7%	12.9%	12.0%	17.6%	9.6%	25.0%	33.4%	18.9%	22.3%	35.3%	59.3%	29.6%

Table 72: Table 71 continued for IFQ bottom trawl vessels. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

Length bin (cm)	Bottom Trawl																				
	Excellent							Poor							Dead						
	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017
116	60.6%	42.8%	59.7%	42.6%	57.3%	52.7%	47.0%	13.5%	20.0%	20.0%	37.1%	15.2%	11.0%	22.2%	25.9%	37.1%	20.2%	20.3%	27.5%	36.3%	30.9%
118	55.8%	58.4%	62.9%	62.3%	54.5%	25.4%	48.4%	9.6%	6.4%	17.3%	29.2%	21.4%	35.9%	18.5%	34.5%	35.2%	19.8%	8.5%	24.2%	38.7%	33.1%
120	47.5%	20.3%	79.4%	81.7%	58.1%	42.3%	67.3%	28.1%	16.5%	18.8%	0.0%	16.4%	0.8%	14.8%	24.3%	63.2%	1.8%	18.3%	25.4%	57.0%	18.1%
122	54.3%	58.9%	59.0%	80.1%	56.7%	52.0%	76.8%	8.0%	31.2%	14.5%	0.0%	7.5%	16.1%	17.8%	37.7%	9.9%	26.5%	19.9%	35.7%	31.9%	5.8%
124	39.8%	39.0%	47.7%	73.5%	29.7%	82.7%	62.5%	21.8%	48.5%	16.1%	16.0%	35.1%	0.0%	12.4%	38.4%	12.5%	36.1%	10.5%	35.1%	17.3%	25.1%
126	42.1%	29.4%	100.0%	0.0%	34.9%	31.2%	64.1%	19.0%	30.6%	0.0%	37.8%	0.0%	0.0%	35.9%	38.9%	40.1%	0.0%	62.2%	65.1%	68.8%	0.0%
128	52.5%	96.4%	49.5%	85.0%	84.8%	67.9%	0.0%	35.7%	0.0%	50.5%	0.0%	3.9%	20.7%	0.0%	11.7%	3.6%	0.0%	15.0%	11.5%	11.4%	0.0%
130	75.4%	0.0%	77.8%	100.0%	82.5%	79.0%	53.7%	24.6%	0.0%	0.0%	0.0%	0.0%	0.0%	46.3%	0.0%	100.0%	22.2%	0.0%	17.5%	21.0%	0.0%
132	45.2%	100.0%	22.2%	100.0%	100.0%	100.0%	52.4%	18.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	36.2%	0.0%	77.8%	0.0%	0.0%	0.0%	47.6%
134	79.3%	100.0%	67.0%	100.0%	25.6%	61.6%	0.0%	20.7%	0.0%	33.0%	0.0%	0.0%	26.5%	0.0%	0.0%	0.0%	0.0%	74.4%	11.9%	100.0%	0.0%
136	25.2%	100.0%	100.0%	100.0%	100.0%	74.7%	100.0%	49.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	25.2%	0.0%	0.0%	0.0%	0.0%	25.3%	0.0%
138	0.0%	8.2%	0.0%	0.0%	0.0%	90.3%	0.0%	100.0%	55.3%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	35.9%	0.0%	0.0%	100.0%	9.7%	0.0%
140	49.7%	0.0%	0.0%	0.0%	4.4%	0.0%	86.9%	50.3%	0.0%	0.0%	0.0%	46.5%	0.0%	13.1%	0.0%	0.0%	0.0%	49.0%	100.0%	0.0%	0.0%
142	25.1%	0.0%	0.0%	0.0%	10.3%	0.0%	0.0%	24.8%	100.0%	0.0%	0.0%	59.5%	0.0%	0.0%	50.1%	0.0%	0.0%	30.1%	0.0%	0.0%	0.0%
144	59.4%	0.0%	0.0%	59.9%	0.0%	50.6%	0.0%	40.6%	0.0%	0.0%	40.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	49.4%	0.0%	0.0%
146	100.0%	100.0%	0.0%	0.0%	100.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
148	50.2%	0.0%	0.0%	0.0%	100.0%	89.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	49.8%	0.0%	0.0%	0.0%	31.0%	0.0%	0.0%
150	0.0%	0.0%	0.0%	0.0%	45.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	54.5%	0.0%	0.0%
152	0.0%	100.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%
154	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
156	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
158	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
160	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
162	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
164	0.0%	0.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
166	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
168	0.0%	0.0%	0.0%	0.0%	91.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	9.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
170	0.0%	0.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
172	0.0%	0.0%	0.0%	0.0%	91.8%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	8.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
174	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%
176	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
178	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	50.0%	0.0%
180	0.0%	0.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
182	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%
184	0.0%	0.0%	0.0%	0.0%	45.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	54.9%	0.0%	0.0%
186	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%
188	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
190	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
192	0.0%	0.0%	0.0%	0.0%	100.0%	31.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	69.0%	0.0%	0.0%
194	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
196	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
198	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
200	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
202	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
204	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
206	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
208	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
210	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
212	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%
214	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
216	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
218	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
220	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
222	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Table 73: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using pot gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

Length bin (cm)	Pot						
	2011	2012	2013	2014	2015	2016	2017
0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
4	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
6	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
8	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Table 73: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using pot gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. *(continued)*

Length bin (cm)	Pot						
	2011	2012	2013	2014	2015	2016	2017
10	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
12	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
14	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
16	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
18	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
20	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
22	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
24	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
26	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
28	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
30	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
32	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
34	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
36	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
38	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
40	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
42	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
44	0.0248	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
46	0.0000	0.0000	0.0556	0.0000	0.0000	0.0000	0.0000
48	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
50	0.0000	0.0000	0.0000	0.0000	0.0255	0.0419	0.0000
52	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
54	0.0129	0.0000	0.0882	0.0000	0.0212	0.0000	0.0000
56	0.0054	0.0000	0.0000	0.0000	0.0000	0.0273	0.0000
58	0.0151	0.0000	0.0000	0.0000	0.0337	0.0000	0.0000
60	0.0672	0.0000	0.0148	0.0934	0.0151	0.0459	0.0000
62	0.0538	0.0000	0.0000	0.0000	0.0264	0.0203	0.0000
64	0.0217	0.0377	0.0000	0.0000	0.0238	0.0184	0.0000
66	0.0136	0.0113	0.0052	0.0000	0.0443	0.0162	0.0000
68	0.0215	0.0308	0.0531	0.0000	0.0584	0.0609	0.0000
70	0.0745	0.0239	0.0792	0.0000	0.0628	0.1038	0.0153
72	0.0908	0.0608	0.2634	0.0546	0.0980	0.1387	0.0292
74	0.0541	0.0595	0.2056	0.2002	0.0598	0.0750	0.0651

Table 73: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using pot gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. *(continued)*

Length bin (cm)	Pot						
	2011	2012	2013	2014	2015	2016	2017
76	0.0183	0.0295	0.1398	0.0918	0.0964	0.0477	0.1319
78	0.0744	0.0907	0.1474	0.0421	0.1261	0.0391	0.1217
80	0.1017	0.0891	0.1285	0.2270	0.1052	0.0675	0.1638
82	0.0631	0.1473	0.2159	0.1407	0.0862	0.0911	0.0752
84	0.0543	0.1230	0.0940	0.1990	0.1490	0.1379	0.0782
86	0.0411	0.0636	0.0759	0.2435	0.1113	0.0572	0.0724
88	0.0372	0.0659	0.0992	0.0550	0.1027	0.0199	0.0452
90	0.0473	0.0399	0.0716	0.0000	0.0476	0.0488	0.0831
92	0.0217	0.0337	0.0377	0.0238	0.0591	0.0285	0.0907
94	0.0187	0.0260	0.0300	0.0461	0.0345	0.0430	0.1096
96	0.0153	0.0259	0.0470	0.0416	0.0161	0.0074	0.0284
98	0.0123	0.0016	0.0000	0.0201	0.0091	0.0093	0.0215
100	0.0163	0.0062	0.0094	0.0188	0.0112	0.0396	0.0150
102	0.0025	0.0085	0.0206	0.1038	0.0027	0.0062	0.0137
104	0.0024	0.0054	0.0085	0.0000	0.0200	0.0096	0.0132
106	0.0000	0.0137	0.0340	0.0000	0.0023	0.0000	0.0082
108	0.0035	0.0012	0.0000	0.0000	0.0090	0.0017	0.0152
110	0.0014	0.0011	0.0090	0.0277	0.0042	0.0000	0.0000
112	0.0013	0.0010	0.0000	0.0000	0.0000	0.0030	0.0000
114	0.0028	0.0020	0.0000	0.0123	0.0000	0.0015	0.0000
116	0.0005	0.0000	0.0000	0.0233	0.0000	0.0000	0.0000
118	0.0011	0.0009	0.0028	0.0000	0.0000	0.0000	0.0000
120	0.0015	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
122	0.0029	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
124	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
126	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
128	0.0008	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
130	0.0004	0.0000	0.0000	0.0000	0.0012	0.0000	0.0000
132	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
134	0.0007	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
136	0.0007	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
138	0.0003	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
140	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Table 73: Weighted length frequency distributions for Pacific halibut in the IFQ fishery for vessels using pot gears, by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. *(continued)*

Length bin (cm)	Pot						
	2011	2012	2013	2014	2015	2016	2017
142	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
144	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
146	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
148	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
150	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
152	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
154	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
156	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
158	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
160	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
162	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
164	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
166	0.0004	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
168	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
170	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
172	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
174	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
176	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
178	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
180	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
182	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
184	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
186	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
188	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
190	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
192	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
194	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
196	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
198	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
200	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Table 74: Percentage of weighted length measurements in each viability category, for IFQ pot vessels by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

Length bin (cm)	Pot																				
	Excellent							Poor							Dead						
	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017	2011	2012	2013	2014	2015	2016	2017
40	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
42	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
44	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
46	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
48	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
50	0.0%	0.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
52	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
54	0.0%	0.0%	100.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
56	100.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
58	68.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	32.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
60	57.3%	0.0%	100.0%	100.0%	100.0%	76.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	42.7%	0.0%	0.0%	0.0%	0.0%	24.0%	0.0%
62	38.1%	0.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	61.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
64	34.6%	100.0%	0.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	65.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
66	50.0%	100.0%	100.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
68	69.8%	100.0%	36.2%	0.0%	66.4%	100.0%	0.0%	0.0%	0.0%	63.8%	0.0%	33.6%	0.0%	0.0%	30.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
70	62.3%	100.0%	77.9%	0.0%	86.1%	100.0%	100.0%	3.4%	0.0%	10.8%	0.0%	0.0%	0.0%	0.0%	34.3%	0.0%	11.3%	0.0%	13.9%	0.0%	0.0%
72	77.3%	85.9%	96.9%	100.0%	100.0%	100.0%	100.0%	0.0%	14.1%	0.0%	0.0%	0.0%	0.0%	0.0%	22.7%	0.0%	3.1%	0.0%	0.0%	0.0%	0.0%
74	69.2%	93.6%	64.1%	100.0%	100.0%	100.0%	0.0%	9.1%	6.4%	12.0%	0.0%	0.0%	0.0%	59.8%	21.7%	0.0%	23.9%	0.0%	0.0%	0.0%	40.2%
76	43.1%	49.7%	50.0%	100.0%	100.0%	100.0%	44.9%	0.0%	37.8%	33.1%	0.0%	0.0%	0.0%	36.8%	56.9%	12.4%	16.9%	0.0%	0.0%	0.0%	18.2%
78	59.1%	63.3%	100.0%	100.0%	90.2%	87.3%	64.1%	7.8%	14.6%	0.0%	0.0%	0.0%	0.0%	8.9%	33.1%	22.2%	0.0%	0.0%	9.8%	12.7%	27.0%
80	57.6%	100.0%	95.5%	65.8%	88.7%	86.5%	50.8%	1.7%	0.0%	0.0%	0.0%	0.0%	0.0%	24.9%	40.7%	0.0%	4.5%	34.2%	11.3%	13.5%	24.4%
82	86.4%	54.9%	61.6%	100.0%	87.5%	90.9%	62.3%	5.6%	9.6%	16.8%	0.0%	0.0%	4.6%	25.0%	8.0%	35.5%	21.6%	0.0%	12.5%	4.5%	12.7%
84	59.3%	73.6%	100.0%	100.0%	79.8%	100.0%	55.5%	6.0%	13.2%	0.0%	0.0%	6.8%	0.0%	33.6%	34.7%	13.2%	0.0%	0.0%	13.4%	0.0%	10.9%
86	85.3%	76.6%	87.9%	25.2%	75.0%	87.6%	44.5%	7.4%	7.6%	0.0%	0.0%	8.4%	6.3%	33.2%	7.4%	15.8%	12.1%	74.8%	16.6%	6.1%	22.3%
88	92.4%	79.3%	91.4%	100.0%	75.5%	100.0%	50.0%	0.0%	6.8%	0.0%	0.0%	8.2%	0.0%	50.0%	7.6%	13.9%	8.6%	0.0%	16.4%	0.0%	0.0%
90	70.5%	68.2%	100.0%	0.0%	75.4%	93.8%	65.9%	0.0%	21.4%	0.0%	0.0%	0.0%	0.0%	17.1%	29.5%	10.5%	0.0%	0.0%	24.6%	6.2%	17.1%
92	55.8%	59.0%	100.0%	0.0%	100.0%	100.0%	28.6%	22.1%	23.5%	0.0%	0.0%	0.0%	0.0%	57.0%	22.1%	17.4%	0.0%	100.0%	0.0%	0.0%	14.3%
94	52.1%	100.0%	88.9%	50.0%	79.6%	87.4%	22.5%	23.9%	0.0%	0.0%	0.0%	0.0%	0.0%	33.7%	23.9%	0.0%	11.1%	50.0%	20.4%	12.6%	43.8%
96	45.5%	80.2%	47.1%	0.0%	80.3%	100.0%	19.6%	13.4%	13.2%	0.0%	0.0%	19.7%	0.0%	19.6%	41.1%	6.7%	52.9%	100.0%	0.0%	0.0%	60.8%
98	53.1%	100.0%	0.0%	100.0%	100.0%	50.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	24.6%	0.0%	46.9%	0.0%	0.0%	0.0%	0.0%	25.4%	0.0%
100	77.6%	100.0%	100.0%	100.0%	100.0%	78.0%	33.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	33.7%	22.4%	0.0%	0.0%	0.0%	0.0%	22.0%	32.6%
102	100.0%	34.1%	100.0%	100.0%	100.0%	100.0%	0.0%	0.0%	33.0%	0.0%	0.0%	0.0%	0.0%	66.7%	0.0%	33.0%	0.0%	0.0%	0.0%	0.0%	33.3%
104	100.0%	0.0%	100.0%	0.0%	74.6%	79.8%	0.0%	0.0%	50.0%	0.0%	0.0%	25.4%	0.0%	66.3%	0.0%	50.0%	0.0%	0.0%	0.0%	20.2%	33.7%
106	0.0%	45.4%	76.4%	0.0%	100.0%	0.0%	50.8%	0.0%	54.6%	23.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	49.2%
108	18.4%	100.0%	0.0%	0.0%	100.0%	100.0%	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%	81.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
110	100.0%	100.0%	23.1%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	76.9%	0.0%	0.0%	0.0%	0.0%
112	100.0%	100.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
114	57.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	42.6%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%
116	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
118	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
120	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
122	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
124	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
126	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
128	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
130	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%

Table 75: Table 75 continued for IFQ pot vessels. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

[illegible]

Table 76: Weighted length frequency distributions for Pacific halibut in the limited entry bottom trawl fishery, 2002-10. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

Length bin (cm)	Weighted length frequency distribution							Length bin (cm)	Weighted length frequency distribution						
	2004	2005	2006	2007	2008	2009	2010		2004	2005	2006	2007	2008	2009	2010
22	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	94	0.0169	0.0108	0.0099	0.0148	0.0164	0.0151	0.0053
24	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	96	0.0062	0.0052	0.0066	0.0089	0.0143	0.0087	0.0066
26	0.0000	0.0125	0.0000	0.0000	0.0000	0.0000	0.0000	98	0.0034	0.0058	0.0066	0.0091	0.0110	0.0103	0.0067
28	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	100	0.0089	0.0045	0.0025	0.0053	0.0080	0.0088	0.0023
30	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	102	0.0060	0.0034	0.0029	0.0036	0.0061	0.0069	0.0018
32	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	104	0.0065	0.0023	0.0027	0.0041	0.0083	0.0062	0.0021
34	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	106	0.0043	0.0029	0.0032	0.0031	0.0059	0.0028	0.0013
36	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	108	0.0016	0.0014	0.0019	0.0018	0.0027	0.0025	0.0014
38	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	110	0.0048	0.0015	0.0004	0.0017	0.0018	0.0021	0.0009
40	0.0048	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	112	0.0015	0.0007	0.0020	0.0010	0.0016	0.0024	0.0013
42	0.0000	0.0044	0.0000	0.0000	0.0000	0.0000	0.0000	114	0.0020	0.0010	0.0007	0.0007	0.0020	0.0017	0.0001
44	0.0025	0.0012	0.0057	0.0000	0.0000	0.0010	0.0000	116	0.0026	0.0006	0.0002	0.0000	0.0010	0.0005	0.0005
46	0.0037	0.0000	0.0094	0.0000	0.0000	0.0009	0.0000	118	0.0007	0.0004	0.0003	0.0002	0.0004	0.0002	0.0002
48	0.0000	0.0034	0.0046	0.0000	0.0000	0.0000	0.0000	120	0.0013	0.0005	0.0002	0.0002	0.0005	0.0003	0.0002
50	0.0027	0.0068	0.0092	0.0000	0.0007	0.0010	0.0000	122	0.0008	0.0003	0.0000	0.0004	0.0003	0.0003	0.0002
52	0.0021	0.0069	0.0080	0.0041	0.0001	0.0053	0.0000	124	0.0010	0.0002	0.0001	0.0000	0.0003	0.0002	0.0003
54	0.0156	0.0076	0.0164	0.0042	0.0025	0.0004	0.0000	126	0.0000	0.0001	0.0002	0.0001	0.0001	0.0002	0.0002
56	0.0138	0.0211	0.0242	0.0071	0.0022	0.0019	0.0000	128	0.0002	0.0000	0.0002	0.0000	0.0000	0.0002	0.0000
58	0.0187	0.0331	0.0322	0.0293	0.0027	0.0091	0.0022	130	0.0003	0.0002	0.0001	0.0002	0.0000	0.0002	0.0000
60	0.0400	0.0431	0.0670	0.0593	0.0169	0.0175	0.0056	132	0.0005	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000
62	0.0329	0.0719	0.0751	0.0638	0.0285	0.0275	0.0121	134	0.0006	0.0000	0.0001	0.0000	0.0001	0.0001	0.0000
64	0.0428	0.0783	0.1001	0.0932	0.0614	0.0545	0.0155	136	0.0001	0.0001	0.0002	0.0000	0.0000	0.0001	0.0000
66	0.0532	0.0807	0.0979	0.1150	0.0705	0.0606	0.0185	138	0.0000	0.0001	0.0000	0.0000	0.0000	0.0001	0.0000
68	0.0757	0.0845	0.0870	0.0000	0.0599	0.0835	0.0256	140	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000
70	0.0672	0.0851	0.0986	0.1022	0.0871	0.0971	0.0154	142	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000
72	0.0774	0.0882	0.0478	0.1029	0.0973	0.0972	0.0314	144	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
74	0.0998	0.0746	0.0588	0.0840	0.1023	0.0941	0.0383	146	0.0001	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000
76	0.0890	0.0538	0.0461	0.0710	0.0743	0.0697	0.0284	148	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
78	0.0658	0.0506	0.0423	0.0539	0.0688	0.0744	0.0349	150	0.0001	0.0000	0.0001	0.0000	0.0000	0.0000	0.0000
80	0.0586	0.0427	0.0372	0.0460	0.0599	0.0527	0.0298	152	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
82	0.0486	0.0320	0.0258	0.0325	0.0443	0.0434	0.0239	154	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
84	0.0337	0.0255	0.0186	0.0316	0.0428	0.0335	0.0227	156	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
86	0.0221	0.0166	0.0130	0.0000	0.0300	0.0290	0.0141	158	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
88	0.0235	0.0115	0.0120	0.0154	0.0263	0.0290	0.0122	160	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
90	0.0193	0.0127	0.0115	0.0168	0.0225	0.0263	0.0100	162	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
92	0.0157	0.0092	0.0101	0.0122	0.0179	0.0204	0.0094	164	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

116

Length bin (cm)	2004			2005			2006			Length bin (cm)	2007			2008			2009		
	Exc	Poor	Dead	Exc	Poor	Dead	Exc	Poor	Dead		Exc	Poor	Dead	Exc	Poor	Dead	Exc	Poor	Dead
22	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	22	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
24	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	24	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
26	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	26	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
28	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	28	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
30	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	30	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
32	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	32	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
34	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	34	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%
36	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	36	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
38	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	38	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
40	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	40	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
42	0.0%	0.0%	0.0%	0.0%	88.4%	11.6%	0.0%	0.0%	0.0%	42	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
44	0.0%	0.0%	100.0%	0.0%	70.8%	29.2%	0.0%	0.0%	100.0%	44	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%
46	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	46	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%
48	0.0%	0.0%	0.0%	22.4%	0.0%	77.6%	0.0%	0.0%	100.0%	48	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
50	0.0%	0.0%	100.0%	61.1%	9.9%	29.0%	0.0%	0.0%	100.0%	50	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	100.0%	0.0%	0.0%
52	100.0%	0.0%	0.0%	23.6%	31.3%	45.2%	0.0%	0.0%	100.0%	52	33.4%	0.0%	66.6%	100.0%	0.0%	0.0%	99.5%	0.5%	0.0%
54	75.5%	11.9%	12.6%	10.0%	20.8%	69.2%	16.9%	0.0%	83.1%	54	35.6%	0.0%	64.4%	0.0%	4.4%	95.6%	42.3%	57.7%	0.0%
56	12.6%	37.9%	49.5%	25.1%	12.7%	62.2%	22.0%	15.2%	62.8%	56	33.9%	0.0%	66.1%	0.0%	0.0%	100.0%	15.7%	65.3%	19.0%
58	21.4%	25.6%	53.0%	15.1%	29.5%	55.4%	4.1%	20.2%	75.7%	58	9.4%	6.8%	83.8%	3.3%	3.3%	93.3%	51.0%	4.4%	44.6%
60	58.6%	14.4%	27.0%	18.2%	21.0%	60.8%	12.9%	25.5%	61.6%	60	5.3%	7.4%	87.2%	9.0%	14.3%	76.8%	28.7%	21.9%	49.4%
62	40.0%	21.6%	38.4%	18.5%	23.7%	57.8%	27.3%	22.3%	50.4%	62	20.8%	9.5%	69.7%	6.1%	15.7%	78.2%	19.3%	19.5%	61.2%
64	33.4%	18.4%	48.2%	25.2%	28.4%	46.4%	31.5%	21.0%	47.5%	64	18.9%	5.3%	75.8%	17.3%	7.5%	75.2%	38.0%	9.4%	52.6%
66	23.9%	24.7%	51.4%	20.9%	26.7%	52.3%	29.6%	17.3%	53.0%	66	9.1%	12.5%	78.4%	25.8%	8.9%	65.4%	26.7%	19.7%	53.6%

Table 78: Continuation of Table 77. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm.

Length bin (cm)	2010			Length bin	2010			Length bin	2010		
	Exc	Poor	Dead		Exc	Poor	Dead		Exc	Poor	Dead
10	0.0%	100.0%	0.0%	58	100.0%	0.0%	0.0%	106	2.4%	0.0%	97.6%
12	0.0%	0.0%	0.0%	60	33.4%	0.0%	66.6%	108	0.0%	20.1%	79.9%
14	0.0%	0.0%	0.0%	62	15.7%	29.4%	54.9%	110	14.2%	58.8%	27.0%
16	0.0%	0.0%	0.0%	64	30.1%	21.2%	48.7%	112	39.9%	0.0%	60.1%
18	0.0%	0.0%	0.0%	66	17.8%	15.4%	66.8%	114	0.0%	0.0%	100.0%
20	0.0%	0.0%	0.0%	68	15.0%	10.3%	74.8%	116	50.0%	0.0%	50.0%
22	0.0%	0.0%	0.0%	70	22.2%	7.4%	70.4%	118	0.0%	100.0%	0.0%
24	0.0%	0.0%	0.0%	72	23.6%	17.4%	59.0%	120	0.0%	0.0%	100.0%
26	0.0%	0.0%	0.0%	74	13.5%	24.8%	61.7%	122	0.0%	0.0%	100.0%
28	0.0%	0.0%	0.0%	76	20.1%	16.9%	63.0%	124	100.0%	0.0%	0.0%
30	0.0%	0.0%	0.0%	78	17.0%	17.4%	65.7%	126	0.0%	100.0%	0.0%
32	0.0%	0.0%	0.0%	80	10.6%	22.8%	66.6%	128	0.0%	0.0%	0.0%
34	0.0%	0.0%	0.0%	82	18.9%	19.9%	61.2%	130	0.0%	0.0%	0.0%
36	0.0%	0.0%	0.0%	84	21.9%	25.3%	52.8%	132	0.0%	0.0%	0.0%
38	0.0%	0.0%	0.0%	86	14.9%	16.4%	68.7%	134	0.0%	0.0%	0.0%
40	0.0%	0.0%	0.0%	88	24.8%	17.8%	57.4%	136	100.0%	0.0%	0.0%
42	0.0%	0.0%	0.0%	90	25.8%	24.2%	50.1%	138	0.0%	0.0%	0.0%
44	0.0%	0.0%	0.0%	92	5.0%	9.9%	85.1%	140	0.0%	0.0%	0.0%
46	0.0%	0.0%	0.0%	94	26.1%	29.2%	44.7%	142	0.0%	0.0%	0.0%
48	0.0%	0.0%	0.0%	96	17.4%	39.9%	42.7%	144	0.0%	0.0%	0.0%
50	0.0%	0.0%	0.0%	98	14.3%	23.3%	62.4%	146	0.0%	0.0%	0.0%
52	0.0%	0.0%	0.0%	100	2.2%	31.0%	66.8%	148	0.0%	0.0%	0.0%
54	0.0%	0.0%	0.0%	102	21.7%	20.6%	57.8%	150	0.0%	0.0%	0.0%
56	0.0%	0.0%	0.0%	104	18.3%	37.2%	44.6%	152	0.0%	100.0%	0.0%
								154	0.0%	0.0%	0.0%

Table 79: Number of dead P. halibut in each length bin, summed across viability categories, for IFQ bottom trawl vessels by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. Since 2013, IFQ bottom trawl lengths could also include lengths taken on both IFQ and LE California halibut bottom trawl fisheries. This analysis assumes that there is no size-dependent mortality within viability categories.

Bottom Trawl																
Length bin (cm)	2011	2012	2013	2014	2015	2016	2017	Length bin (cm)	2011	2012	2013	2014	2015	2016	2017	
10	0	0	0	0	0	0	0	110	29	41	44	36	35	27	56	
12	0	0	0	0	0	0	0	112	23	26	66	23	34	27	23	
14	0	0	0	0	0	1	0	114	24	27	60	11	34	14	15	
16	0	0	0	0	0	0	0	116	13	22	19	7	22	26	15	
18	1	0	0	0	0	0	0	118	15	14	9	4	14	10	10	
20	0	0	0	0	0	0	0	120	7	18	16	4	13	10	7	
22	0	1	0	0	0	0	0	122	9	7	8	10	11	7	4	
24	0	0	0	0	0	0	0	124	10	6	6	12	8	4	4	
26	0	0	0	0	0	0	0	126	7	8	2	3	5	7	2	
28	0	0	0	0	0	0	0	128	4	3	3	2	4	7	0	
30	0	1	1	0	0	0	0	130	2	1	2	2	4	2	1	
32	0	2	1	1	0	1	0	132	4	1	2	1	1	1	2	
34	0	1	0	0	0	0	0	134	1	1	2	2	2	3	1	
36	0	1	0	0	1	1	0	136	3	1	1	1	1	2	1	
38	0	3	0	0	2	4	1	138	1	2	1	0	1	2	0	
40	1	2	2	1	5	5	1	140	1	0	0	0	6	1	2	
42	1	5	0	0	4	2	2	142	3	1	0	0	3	0	0	
44	0	3	0	0	5	1	1	144	1	0	0	1	0	2	0	
46	1	3	1	1	3	2	1	146	1	1	0	1	1	1	1	
48	3	4	3	1	4	5	4	148	2	0	0	0	1	2	0	
50	6	7	5	0	7	6	7	150	0	1	0	0	2	0	0	
52	7	9	8	3	1	5	11	152	0	1	0	0	0	1	1	
54	12	11	13	5	5	9	16	154	0	0	0	0	1	1	0	
56	13	14	13	10	12	15	22	156	0	0	0	0	0	0	0	
58	44	39	49	26	23	18	35	158	0	0	0	0	0	0	0	
60	70	77	144	57	44	36	35	160	0	0	0	0	1	0	0	
62	103	115	136	109	83	68	55	162	0	0	0	0	1	1	0	
64	146	173	328	160	129	80	61	164	0	0	0	0	1	1	0	
66	175	188	208	176	245	131	73	166	0	0	0	0	0	0	0	
68	173	236	492	746	239	215	143	168	0	0	0	0	1	0	0	
70	240	270	1124	572	271	244	129	170	0	0	0	0	1	1	0	
72	262	310	1260	257	394	696	179	172	0	0	0	0	2	1	0	
74	334	325	391	602	327	259	212	174	0	0	0	0	1	0	0	
76	256	295	722	363	370	274	214	176	0	0	0	0	0	1	0	
78	236	277	664	232	511	311	247	178	0	0	0	0	1	2	0	
80	255	251	568	395	244	283	324	180	0	0	0	0	1	1	0	
82	218	264	990	864	264	280	266	182	0	0	0	0	2	1	0	
84	223	237	532	674	494	383	243	184	0	0	0	0	2	0	0	
86	172	204	262	616	478	214	212	186	0	0	0	0	2	0	0	
88	170	187	358	436	236	186	216	188	0	0	0	0	0	1	0	
90	155	177	620	298	171	177	287	190	0	0	0	0	0	0	0	
92	137	167	316	114	241	287	169	192	0	0	0	0	1	2	0	
94	105	126	456	85	262	177	255	194	0	0	0	0	0	0	0	
96	97	100	344	174	92	103	118	196	0	0	0	0	0	0	0	
98	74	95	116	68	85	78	201	198	0	0	0	0	1	0	0	
100	68	77	156	70	194	149	92	200	0	0	0	0	0	0	0	
102	66	71	152	159	71	60	84	202	0	0	0	0	0	0	0	
104	51	55	96	70	102	45	55	204	0	0	0	0	0	0	0	
106	37	44	126	25	44	35	39	206	0	0	0	0	0	0	0	
108	31	46	100	20	75	38	74	208	0	0	0	0	0	0	0	
110	29	41	44	36	35	27	56	210	0	0	0	0	0	0	0	
112	23	26	66	23	34	27	23	212	0	0	0	0	0	1	0	

Table 80: Number of dead P. halibut in each length bin, summed across viability categories, for IFQ pot vessels by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. This analysis assumes that there is no size-dependent mortality within viability categories.

Pot															
Length bin (cm)	2011	2012	2013	2014	2015	2016	2017	Length bin (cm)	2011	2012	2013	2014	2015	2016	2017
40	0	0	0	0	0	0	0	118	1	1	0	0	0	0	0
42	0	0	0	0	0	0	0	120	0	0	0	0	0	0	0
44	0	0	0	0	0	0	0	122	0	0	0	0	0	0	0
46	0	0	0	0	0	0	0	124	0	0	0	0	0	0	0
48	0	0	0	0	0	0	0	126	0	0	0	0	0	0	0
50	0	0	0	0	0	0	0	128	0	0	0	0	0	0	0
52	0	0	0	0	0	0	0	130	0	0	0	0	1	0	0
54	1	0	0	0	0	0	0	132	0	0	0	0	0	0	0
56	0	0	0	0	0	0	0	134	0	0	0	0	0	0	0
58	1	0	0	0	0	0	0	136	0	0	0	0	0	0	0
60	3	0	0	0	0	1	0	138	0	0	0	0	0	0	0
62	5	0	0	0	0	0	0	140	0	0	0	0	0	0	0
64	3	0	0	0	0	0	0	142	0	0	0	0	0	0	0
66	1	0	0	0	0	0	0	144	0	0	0	0	0	0	0
68	1	0	1	0	2	0	0	146	0	0	0	0	0	0	0
70	7	0	2	0	1	0	0	148	0	0	0	0	0	0	0
72	5	1	1	0	0	0	0	150	0	0	0	0	0	0	0
74	4	1	3	0	0	0	5	152	0	0	0	0	0	0	0
76	4	3	3	0	0	0	6	154	0	0	0	0	0	0	0
78	8	5	0	0	1	1	4	156	0	0	0	0	0	0	0
80	12	0	1	1	2	2	4	158	0	0	0	0	0	0	0
82	3	12	6	0	2	2	3	160	0	0	0	0	0	0	0
84	8	6	0	0	3	0	4	162	0	0	0	0	0	0	0
86	2	3	2	3	3	2	5	164	0	0	0	0	0	0	0
88	1	3	2	0	3	0	3	166	0	0	0	0	0	0	0
90	6	3	0	0	3	1	2	168	0	0	0	0	0	0	0
92	4	4	0	1	0	0	10	170	0	0	0	0	0	0	0
94	4	0	1	1	1	1	7	172	0	0	0	0	0	0	0
96	4	2	3	1	1	0	4	174	0	0	0	0	0	0	0
98	3	0	0	0	0	2	0	176	0	0	0	0	0	0	0
100	2	0	0	0	0	2	2	178	0	0	0	0	0	0	0
102	0	2	0	0	0	0	3	180	0	0	0	0	0	0	0
104	0	2	0	0	1	1	3	182	0	0	0	0	0	0	0
106	0	3	1	0	0	0	1	184	0	0	0	0	0	0	0
108	2	0	0	0	0	0	1	186	0	0	0	0	0	0	0
110	0	0	1	0	0	0	0	188	0	0	0	0	0	0	0
112	0	0	0	0	0	0	0	190	0	0	0	0	0	0	0
114	1	1	0	1	0	1	0	192	0	0	0	0	0	0	0
116	1	0	0	0	0	0	0	194	0	0	0	0	0	0	0
118	1	1	0	0	0	0	0	196	0	0	0	0	0	0	0
120	0	0	0	0	0	0	0	198	0	0	0	0	0	0	0
122	0	0	0	0	0	0	0	200	0	0	0	0	0	0	0

Table 81: Number of dead *P. halibut* in each length bin for Shoreside Hake vessels 2011-14. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. This analysis assumes 100% mortality of all individuals because viability has not been determined for *P. halibut* caught with midwater trawl nets. Starting in 2015, Shoreside Hake trips were sorted into Midwater Hake or Midwater Rockfish depending on landing amount of *P. hake*.

Shoreside Hake Midwater Trawl									
Length bin (cm)	2011	2012	2013	2014	Length bin (cm)	2011	2012	2013	2014
72	0	0	1	0	106	0	0	0	1
73	0	0	0	0	107	0	0	0	0
74	0	0	0	1	108	0	0	0	0
75	0	0	0	0	109	0	0	0	0
76	0	0	0	0	110	0	0	0	1
77	0	0	0	0	111	0	0	0	0
78	0	0	0	0	112	0	0	0	0
79	0	0	0	0	113	0	0	0	0
80	1	0	0	1	114	0	0	0	0
81	0	0	0	0	115	0	0	0	0
82	0	0	0	0	116	0	0	0	0
83	0	0	0	0	117	0	0	0	0
84	0	0	0	0	118	0	0	0	0
85	0	0	0	0	119	0	0	0	0
86	1	0	0	0	120	0	0	0	0
87	0	0	0	0	121	0	0	0	0
88	0	0	0	0	122	0	0	0	0
89	0	0	0	0	123	0	0	0	0
90	0	0	0	0	124	0	0	0	0
91	0	0	0	0	125	0	0	0	0
92	0	0	0	0	126	0	0	0	0
93	0	0	0	0	127	0	0	0	0
94	0	0	0	1	128	0	0	0	1
95	0	0	0	0	129	0	0	0	0
96	0	0	0	0	130	0	0	0	0
97	0	0	0	0	131	0	0	0	0
98	1	0	0	0	132	0	0	0	0
99	0	0	0	0	133	0	0	0	0
100	0	0	0	1	134	0	0	0	0
101	0	0	0	0	135	0	0	0	0
102	0	0	0	0	136	0	0	1	0
103	0	0	0	0	137	0	0	0	0
104	0	0	1	1					
105	0	0	0	0					

Table 82: Number of dead P. halibut in each length bin for Midwater Rockfish vessels by year. Length bins are inclusive of the bin value (lower) and exclude the upper value, e.g., 10 = lengths 10.0 to 11.99 cm. This analysis assumes 100% mortality of all individuals because viability has not been determined for P. halibut caught with midwater trawl nets.

Midwater Rockfish			
Length bin (cm)	2017	Length bin (cm)	2017
72	1	80	0
74	0	82	1
76	0	84	2
78	0	86	0
80	0	88	0
82	1	90	1

A.3 Pacific Halibut IBQ Expansions for In-Season Management, Special Cases

A.3.1 In season reporting to the Vessel Account System

The Vessel Account System (VAS) is a NOAA, West Coast Region database that allows fishers to manage their IFQ quota pounds. On a weekly basis, the WCGOP provides trip-level estimates of discarded P. halibut IBQ to the Pacific States Marine Fisheries Commission (PSMFC). The PSMFC then uploads the data to the VAS. Occasionally, special circumstances required alternative calculations of P. halibut IBQ. Alternative calculations of P. halibut IBQ were identified by observer program staff and incorporated into the VAS. Scenarios triggering an alternative calculation and the equations used for those calculations are given in Table 83 below.

The WCGOP database calculates IBQ weight at the haul-level when the observer collects all the required data elements. The calculation is dependent on the gear fished.

A.3.2 In season IBQ Weight Calculations for Bottom Trawl Gear

The sampled P. halibut lengths are converted to weight using the IPHC length-weight conversion table (Table 9 in Appendix A.4). The total weight of P. halibut in the haul is calculated as:

$$W = \frac{w}{n} \times N \quad (10)$$

where, for each haul:

W = total weight of P. halibut

w = sampled weight of P. halibut

n = sampled number of P. halibut

N = total number of P. halibut

IBQ weight for each haul is then calculated as:

$$W_{IBQ} = \sum_c \left(\frac{w_c}{\sum_c w_c} \times W \times m_c \right) \quad (11)$$

where, for each haul:

c = viability condition category

W_{IBQ} = IBQ weight (mortality rate applied) of P. halibut

W = total weight of P. halibut in haul

w = sampled weight of P. halibut

m = mortality rate (Table 5)

A.3.3 In season IBQ Weight Calculations for Pot Gear

The sampled P. halibut lengths are converted to weight using the IPHC length-weight conversion table (Table 9 in Appendix A.4). Observers are not always able to sample 100% of all gear units due to time constraints and logistics, therefore sample weights need to be expanded to the haul/set level. The total weight of P. halibut in the set is calculated as:

$$W = \left(\frac{w}{n} \times N \right) \times \left(\frac{P}{p} \right) \quad (12)$$

where, for each set:

W = total weight of P. halibut

w = sampled weight of P. halibut

n = sampled number of P. halibut

N = total number of P. halibut

P = total number of pots fished

p = sampled number of pots

IBQ weight for each haul is then calculated as:

$$W_{IBQ} = \sum_c \left(\frac{w_c}{\sum_c w_c} \times W \times m_c \right) \quad (13)$$

where, for each set:

c = viability condition category

W_{IBQ} = IBQ weight (mortality rate applied) of P. halibut

W = total weight of P. halibut in set

w = sampled weight of P. halibut

m = mortality rate (Table 6)

A.3.4 In season IBQ Weight Calculations for Hook-&-Line Gear

The visual estimates of Pacific halibut length (10 cm increments) are converted to weight using the IPHC length-weight conversion table (Table 9 in Appendix A.4). Observers are not always

able to sample 100% of all gear units due to time constraints and logistics, therefore sample weights need to be expanded to the haul/set level. The total weight of P. halibut in the set is calculated as:

$$W_{IBQ} = \left(\frac{H}{h} \times w \right) \times 0.16 \quad (14)$$

where, for each set:

W_{IBQ} = IBQ weight (mortality rate applied) of P. halibut

w = sampled weight of P. halibut

H = total number of hooks fished

h = sampled number of hooks

0.16 = IPHC mortality rate applied to hook-&-line gear

A.3.5 In season IBQ Weight Alternative Calculation Scenarios

The most prevalent causes for alternative IBQ calculations were due to pre-sorting of P. halibut by the crew and improper sampling. In these scenarios, observer program staff reviewed the trip and calculated IBQ weight manually.

To determine the most appropriate method to calculate IBQ weight, the observer program data management team consulted with the IPHC. For bottom trawl and pot gear, the IPHC preferred the use of manually measured fish from other properly sampled hauls within the same trip, rather than the use of visually estimated lengths from the haul. All calculations utilized data from the same trip or a different trip from the same vessel. In other words, there was never a circumstance where data from Vessel A was used to calculate IBQ weight for Vessel B.

In addition to scenarios where the observer did not collect all required data, there were also instances of hauls where P. halibut was not sampled by the observer or all the gear was lost. In these instances, properly sampled hauls were used to estimate IBQ weight for the unsampled haul. Methods for expanding P. halibut weight to unsampled or partially sampled hauls varied by gear type.

To calculate P. halibut IBQ weight for unsampled trawl hauls, the sum of all IBQ weight from other properly sampled hauls is divided by the sum of tow duration (hours) from sampled hauls and multiplied by tow duration of the unsampled haul.

$$W_{IBQ} = \left(\frac{\sum_t w_{IBQ}}{\sum_t d} \right) \times D \quad (15)$$

where, for each tow:

t = tow

W_{IBQ} = unsampled IBQ weight (mortality rate applied) of P. halibut

w_{IBQ} = sampled IBQ weight (mortality rate applied) of P. halibut

d = tow duration (hr) of sampled haul

D = tow duration (hr) of unsampled haul

To calculate P. halibut IBQ weight when trawl gear is lost (i.e., entire net or codend is lost), the sum of all P. halibut expanded species weight from other properly sampled hauls is divided by the sum of tow durations from sampled hauls, multiplied by the tow duration of the unsampled haul. For lost trawl gear, a mortality rate for the “dead” P. halibut viability condition (0.90) is applied.

$$W_{IBQ} = \left(\frac{\sum_t w}{\sum_t d} \right) \times D \times 0.90 \quad (16)$$

where, for each tow with lost gear:

t = tow

W_{IBQ} = unsampled IBQ weight (mortality rate applied) of P. halibut

w_{IBQ} = sampled IBQ weight (mortality rate applied) of P. halibut

d = tow duration (hr) of sampled haul

D = tow duration (hr) of unsampled haul

To calculate P. halibut IBQ weight in unsampled fixed gear sets, the sum of all P. halibut IBQ weight from sets with similar properties (i.e., date, depth, target, gear type, area; determined by WCGOP data managers) is divided by the sum of the number of gear units sampled, and the result is multiplied by the total number of gear units fished from the unsampled set.

$$W_{IBQ} = \left(\frac{\sum_t w_{IBQ}}{\sum_t g} \right) \times G \quad (17)$$

where, for each set:

t = tow

W_{IBQ} = unsampled IBQ weight (mortality rate applied) of P. halibut

w_{IBQ} = sampled IBQ weight (mortality rate applied) of P. halibut

g = number of sampled gear units (e.g., hooks, pots)

G = total number of gear units (e.g., hooks, pots) fished in the unsampled set

To calculate P. halibut IBQ weight when fixed gear is lost, the sum of P. halibut weight from the sampled portion of the set, or, if all gear is lost, from sets with similar properties is divided by the sum of units sampled, and the result is multiplied by the total hooks from the unsampled set. For any lost fixed gear, a mortality rate for the “dead” P. halibut viability condition (1.0) is applied.

$$W_{IBQ} = \left(\frac{\sum_t w_{IBQ}}{\sum_t g} \right) \times G \times 1.0 \quad (18)$$

where, for each set with lost gear:

t = tow

W_{IBQ} = unsampled IBQ weight (mortality rate applied) of P. halibut

w_{IBQ} = sampled IBQ weight (mortality rate applied) of P. halibut

g = number of sampled gear units (e.g., hooks, pots)

G = total number of gear units (e.g., hooks, pots) fished in the unsampled set

Scenario 1: Total count of P. halibut exists with no length or viability data.

Resolution: Determine an average mortality weight per individual P. halibut in the trip from all sampled hauls. Multiply that average by the total count of P. halibut to determine an IBQ.

Scenario 2: *Total count of P. halibut exists with actual lengths and no viability data.*

Resolution: Determine catch weight for P. halibut using the lengths in the haul and then apply that to the total count for a total weight. Determine CATCH_WEIGHT_MORT for all viabilities (E, P, D) from all other properly sampled hauls in the trip and apply to the CATCH_WEIGHT for IBQ estimate.

Scenario 3: *Total count of P. halibut exists with visual estimates of P. halibut lengths and no viabilities.*

Resolution: The use of visual lengths was discouraged by the IPHC so the most appropriate method is to determine an average IBQ per individual P. halibut in the trip from all sampled hauls. Multiply that average by the total count of P. halibut to determine an IBQ.

Scenario 4: *Total count of P. halibut exists with visual estimates of P. halibut lengths and proper in-hand viabilities.*

Resolution: The use of visual lengths was discouraged by the IPHC, so the most appropriate method here would be to determine an average IBQ per individual P. halibut in the trip from all sampled hauls. Multiply that average by the total count of P. halibut to determine an IBQ.

Scenario 5: *P. halibut not sampled or only visual estimates of length are available.*

Resolution: Confirm P. halibut was present in the haul, and no data was collected on them. Determine an average IBQ per haul for all sampled hauls in the trip. This scenario is unlikely and, to date, has never occurred.

Scenario 6: *Total count of P. halibut does not exist with length and no viability data.*

Resolution: Catch weight of the haul will be determined by taking the measured P. halibut sample, converted to weight, divided by the number of fish sampled, multiplied by the average number of P. halibut for all sampled hauls in the trip. Then the average mortality rates from the sampled hauls are applied to the calculated P. halibut weight and, to date, has never occurred.

Scenario 7: *Total count of P. halibut does not exist with length and viability data.*

Resolution: P. halibut catch weight for the haul will be determined by taking the length of the P. halibut sample, converted to weight, divided by the number of fish sampled, multiplied by the average number of P. halibut for all sampled hauls in the trip. Because viabilities and lengths exist, IBQ can be determined using normal protocols and the calculated catch weight and, to date, has never occurred.

Scenario 8: *Total count of P. halibut does not exist with visual length and viability data.*

Resolution: Determine an average IBQ per haul for all sampled hauls in the trip and apply to the unsampled haul(s).

Scenario 9: *Observer encounters predated fish that are dead and badly damaged so that accurate biological data cannot be collected.*

Resolution: If properly sampled P. halibut exist in the haul they can be used to determine the portion of the catch weight attributed to the predated and non-predated fish. The IBQ for the P. halibut not predated would be calculated separately using the data collected in the haul. The IBQ for the predated fish would be the portion of the P. halibut catch weight attributed to the predated fish multiplied by the mortality rate for “dead” from the IPHC viability tables for that gear.

If all P. halibut in the haul are heavily predated then a catch weight for the haul will need to be determined. This can be done by taking the total count of P. halibut in the haul times an average catch weight (not IBQ estimates) per P. halibut from other hauls in the trip (or like “sets” if P. halibut doesn’t exist in any other hauls). The estimated catch weight will then be multiplied by the mortality rate for “dead” from the IPHC viability tables for that gear to determine IBQ. In 2011, there were two instances where a P. halibut IBQ was manually calculated due to sand flea predation.

Table 83: Calculations used by the Vessel Account System (VAS) to determine Pacific halibut IBQ weight for unsampled or partially sampled fishing events in the U.S. West Coast groundfish IFQ fishery. The calculated values, $\hat{w}_{IBQ_{u,p}}$, are added to the sampled P. halibut to obtain total IBQ weight. Note that these calculations differ slightly from the methods used in this report.

Scenario(s)	Calculation
1,3,4	$\hat{w}_{IBQ_u} = \left(\frac{\sum_{h,v} (l_{h,v} \times r_v)}{\sum_h c_h} \right) \times c_u$
2	$\hat{w}_{IBQ_u} = \left(\frac{\sum_{h,v} l_{h,v}}{\sum_h l_h} \times r_v \right) \times \left(\frac{\sum_f l_f}{\sum_f c_f} \right)$
6,7	$\hat{w}_{IBQ_u} = \left[\left(\frac{\sum_f l_f}{\sum_f c_f} \times r_v \right) \times \frac{\sum_h c_h}{h} \right] \times \left(\frac{\sum_{h,v} l_{h,v}}{\sum_h l_h} \right)$
5,8	$\hat{w}_{IBQ_u} = \frac{\sum_h w_{IBQ_h}}{\sum_h t_h} \times \sum_u t_u$
9	$\hat{w}_{IBQ_p} = \frac{\sum_h l_h}{\sum_h c_h} \times c_p$

where:

c = count of P. halibut

w = weight of P. halibut

l = length of P. halibut, converted to weight via IPHC length-weight table

v = viability of P. halibut, Excellent, Poor, or Dead

r = mortality rate applied for a given viability and gear combination, see Tables 5 & 6

h = sampled hauls

u = unsampled hauls

f = individual sampled P. halibut

t = tow time

p = predated fish

A.4 IPHC Length-Weight Table

Figure 9: IPHC length-weight conversion table for Pacific halibut.

Centimeter	Pounds	Kilograms	Centimeter	Pounds	Kilograms	Centimeter	Pounds	Kilograms	Centimeter	Pounds	Kilograms
10	0.02	0.01	71	9.19	4.17	131	66.82	30.31	191	226.70	102.83
11	0.02	0.01	72	9.61	4.36	132	68.48	31.06	192	230.56	104.58
12	0.02	0.01	73	10.05	4.56	133	70.17	31.83	193	234.48	106.36
13	0.04	0.02	74	10.49	4.76	134	71.89	32.61	194	238.45	108.16
14	0.04	0.02	75	10.98	4.98	135	73.66	33.41	195	242.44	109.97
15	0.07	0.03	76	11.44	5.19	136	75.44	34.22	196	246.50	111.81
16	0.07	0.03	77	11.95	5.42	137	77.25	35.04	197	250.60	113.67
17	0.09	0.04	78	12.46	5.65	138	79.08	35.87	198	255.74	116.00
18	0.11	0.05	79	12.99	5.89	139	80.95	36.72	199	258.93	117.45
19	0.13	0.06	80	13.51	6.13	140	82.87	37.59	200	263.17	119.37
20	0.15	0.07	81	14.07	6.38	141	84.79	38.46	201	267.46	121.32
21	0.18	0.08	82	14.64	6.64	142	86.75	39.35	202	271.79	123.28
22	0.20	0.09	83	15.23	6.91	143	88.76	40.26	203	276.17	125.27
23	0.24	0.11	84	15.83	7.18	144	90.79	41.18	204	280.60	127.28
24	0.26	0.12	85	16.45	7.46	145	92.84	42.11	205	285.10	129.32
25	0.31	0.14	86	17.09	7.75	146	94.93	43.06	206	289.62	131.37
26	0.35	0.16	87	17.75	8.05	147	97.05	44.02	207	294.21	133.45
27	0.40	0.18	88	18.41	8.35	148	99.21	45.00	208	298.84	135.55
28	0.46	0.21	89	19.09	8.66	149	101.39	45.99	209	303.51	137.67
29	0.51	0.23	90	19.80	8.98	150	103.62	47.00	210	308.25	139.82
30	0.57	0.26	91	20.53	9.31	151	105.87	48.02	211	313.03	141.99
31	0.62	0.28	92	21.25	9.64	152	108.16	49.06	212	317.86	144.18
32	0.71	0.32	93	22.02	9.99	153	110.50	50.12	213	322.73	146.39
33	0.77	0.35	94	22.80	10.34	154	112.83	51.18	214	327.67	148.63
34	0.84	0.38	95	23.59	10.70	155	115.24	52.27	215	332.65	150.89
35	0.93	0.42	96	24.41	11.07	156	117.66	53.37	216	337.70	153.18
36	1.01	0.46	97	25.24	11.45	157	120.13	54.49	217	342.79	155.49
37	1.10	0.50	98	26.08	11.83	158	122.62	55.62	218	347.93	157.82
38	1.21	0.55	99	26.96	12.23	159	125.16	56.77	219	353.13	160.18
39	1.32	0.60	100	27.87	12.64	160	127.71	57.93	220	358.38	162.56
40	1.43	0.65	101	28.77	13.05	161	130.32	59.11	221	363.69	164.97
41	1.59	0.72	102	29.70	13.47	162	132.96	60.31	222	369.05	167.40
42	1.68	0.76	103	30.67	13.91	163	135.65	61.53	223	374.45	169.85
43	1.81	0.82	104	31.64	14.35	164	138.36	62.76	224	379.92	172.33
44	1.94	0.88	105	32.63	14.80	165	141.12	64.01	225	385.45	174.84
45	2.09	0.95	106	33.64	15.26	166	143.90	65.27	226	391.03	177.37
46	2.25	1.02	107	34.68	15.73	167	146.72	66.55	227	396.67	179.93
47	2.43	1.10	108	35.74	16.21	168	149.54	67.83	228	402.36	182.51
48	2.58	1.17	109	36.84	16.71	169	152.49	69.17	229	408.09	185.11
49	2.76	1.25	110	37.94	17.21	170	155.45	70.51	230	413.91	187.75
50	2.95	1.34	111	39.07	17.72	171	158.42	71.86	231	419.76	190.40
51	3.15	1.43	112	40.21	18.24	172	161.44	73.23	232	425.69	193.09
52	3.35	1.52	113	41.38	18.77	173	164.51	74.62	233	431.66	195.80
53	3.57	1.62	114	42.59	19.32	174	167.60	76.02	234	437.68	198.53
54	3.79	1.72	115	43.81	19.87	175	170.75	77.45	235	443.76	201.29
55	4.01	1.82	116	45.06	20.44	176	173.92	78.89	236	449.91	204.08
56	4.25	1.93	117	46.32	21.01	177	177.14	80.35	237	456.13	206.90
57	4.52	2.05	118	47.62	21.60	178	180.40	81.83	238	462.39	209.74
58	4.76	2.16	119	48.94	22.20	179	183.71	83.33	239	468.72	212.61
59	5.05	2.29	120	50.29	22.81	180	187.06	84.85	240	475.09	215.50
60	5.31	2.41	121	51.65	23.43	181	190.46	86.39	241	481.55	218.43
61	5.62	2.55	122	53.07	24.07	182	193.87	87.94	242	488.05	221.38
62	5.93	2.69	123	54.48	24.71	183	197.36	89.52	243	494.60	224.35
63	6.24	2.83	124	55.93	25.37	184	200.86	91.11	244	501.24	227.36
64	6.57	2.98	125	57.41	26.04	185	204.43	92.73	245	507.92	230.39
65	6.90	3.13	126	58.91	26.72	186	208.03	94.36	246	514.66	233.45
66	7.25	3.29	127	60.43	27.41	187	211.67	96.01	247	521.48	236.54
67	7.61	3.45	128	61.99	28.12	188	214.71	97.39	248	528.36	239.66
68	7.98	3.62	129	63.56	28.83	189	218.50	99.11	249	535.28	242.80
69	8.38	3.80	130	65.17	29.56	190	222.89	101.10	250	542.29	245.98
70	8.77	3.98									

A.5 Data flow

Figure 10: IFQ groundfish fishery data flow from the Northwest Fisheries Science Center Observer Program to the Vessel Account System (VAS) of the NMFS Western Regional Office.

