

1 DISTRIBUTION, BIOMASS, AND DEMOGRAPHICS OF  
2 COASTAL PELAGIC FISHES IN THE CALIFORNIA  
3 CURRENT ECOSYSTEM DURING SUMMER 2022 BASED ON  
4 ACOUSTIC-TRAWL SAMPLING

5 K.L. Stierhoff, NOAA Fisheries-Southwest Fisheries Science Center (NMFS-SWFSC)  
6 J.P. Zwolinski, UCSC, Coop. Inst. for Mar., Earth, & Atm. Sys. and NMFS-SWFSC  
7 J.S. Renfree, NMFS-SWFSC D.A. Demer, NMFS-SWFSC

8 2023-02-27 20:34:23 UTC

9 **Contents**

10	<b>Executive Summary</b>	<b>1</b>
11	<b>1 Introduction</b>	<b>3</b>
12	<b>2 Methods</b>	<b>6</b>
13	2.1 Sampling . . . . .	6
14	2.1.1 Design . . . . .	6
15	2.1.2 Acoustic . . . . .	8
16	2.1.3 Oceanographic . . . . .	13
17	2.1.4 Fish-eggs . . . . .	13
18	2.1.5 Species and Demographics . . . . .	14
19	2.2 Data processing . . . . .	19
20	2.2.1 Acoustic and oceanographic data . . . . .	19
21	2.2.2 Sound speed and absorption calculation . . . . .	19
22	2.2.3 Echo classification . . . . .	19
23	2.2.4 Removal of non-CPS backscatter . . . . .	20
24	2.2.5 Extraction of nearshore backscatter . . . . .	22
25	2.2.6 Quality Assurance and Quality Control . . . . .	22
26	2.2.7 Echo integral partitioning and acoustic inversion . . . . .	22
27	2.2.8 Trawl clustering and species proportion . . . . .	24
28	2.3 Data analysis . . . . .	26
29	2.3.1 Post-stratification . . . . .	26

30	2.3.2	Biomass and sampling precision estimation . . . . .	29
31	2.3.3	Abundance- and biomass-at-length estimation . . . . .	29
32	2.3.4	Percent biomass per cluster contribution . . . . .	29
33	<b>3</b>	<b>Results</b>	<b>30</b>
34	3.1	Sampling effort and allocation . . . . .	30
35	3.2	Acoustic backscatter . . . . .	31
36	3.3	Egg densities and distributions . . . . .	31
37	3.4	Trawl catch . . . . .	32
38	3.5	Purse-seine catch . . . . .	32
39	3.5.1	<i>Lisa Marie</i> . . . . .	32
40	3.5.2	<i>Long Beach Carnage</i> . . . . .	32
41	3.5.3	Combined catch . . . . .	32
42	3.6	Biomass distribution and demographics . . . . .	39
43	3.6.1	Northern Anchovy . . . . .	39
44	3.6.2	Pacific Sardine . . . . .	47
45	3.6.3	Pacific Mackerel . . . . .	55
46	3.6.4	Jack Mackerel . . . . .	59
47	3.6.5	Pacific Herring . . . . .	64
48	<b>4</b>	<b>Discussion</b>	<b>68</b>
49	4.1	Biomass and abundance . . . . .	68
50	4.1.1	Northern Anchovy . . . . .	68
51	4.1.2	Pacific Sardine . . . . .	68
52	4.1.3	Pacific Mackerel . . . . .	69
53	4.1.4	Jack Mackerel . . . . .	69
54	4.1.5	Pacific Herring . . . . .	69
55	4.2	Ecosystem dynamics: Forage fish community . . . . .	71
56		<b>Acknowledgements</b>	<b>75</b>
57		<b>References</b>	<b>76</b>
58		<b>Appendix</b>	<b>80</b>
59	<b>A</b>	<b>Length distributions and percent biomass by cluster</b>	<b>80</b>
60	A.1	Northern Anchovy . . . . .	80
61	A.2	Pacific Sardine . . . . .	81
62	A.3	Pacific Mackerel . . . . .	82
63	A.4	Jack Mackerel . . . . .	83
64	A.5	Pacific Herring . . . . .	84

## 65 List of Tables

66	1	Wide-Bandwidth Transceiver (Simrad EK80 WBT; Kongsberg) information, pre-calibration settings, and post-calibration beam model results (below the horizontal line). Prior to the survey, on-axis gain ( $G_0$ ), beam angles, angle offsets, and $S_A$ Correction ( $S_{A\text{corr}}$ ) values from calibration results were entered into the WBT control software (Simrad EK80; Kongsberg). . . . .	10
67			
68			
69			
70	2	Wideband Transceiver (Simrad EK80 WBT; Kongsberg) and transducer information (above horizontal line) and beam model results (below horizontal line) estimated from calibration of the echosounders aboard <i>Lisa Marie</i> using a WC38.1 standard sphere. . . . .	11
71			
72			
73	3	General Purpose Transceiver (Simrad EK60 GPT; Kongsberg) and transducer information (above horizontal line) and beam model results (below horizontal line) estimated from a tank calibration, using a WC38.1 standard sphere, of the echosounders later installed and used aboard <i>Long Beach Carnage</i> . . . . .	11
74			
75			
76			
77	4	Miniature Wideband Transceiver (Simrad-Kongsberg WBT Mini) beam model results estimated from calibrations of echosounders using a WC38.1 standard sphere, of the echosounders aboard the two USVs. . . . .	12
78			
79			
80	5	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for the northern stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	39
81			
82			
83	6	Abundance versus standard length ( $L_S$ , cm) for the northern stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. . . . .	40
84			
85	7	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for the central stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	43
86			
87			
88	8	Abundance versus standard length ( $L_S$ , cm) for the central stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. . . . .	44
89			
90	9	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for the northern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	47
91			
92			
93	10	Abundance versus standard length ( $L_S$ , cm) for the northern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. . . . .	48
94			
95	11	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for the southern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	51
96			
97			
98	12	Abundance versus standard length ( $L_S$ , cm) for the southern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. . . . .	52
99			
100	13	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for Pacific Mackerel ( <i>Scomber japonicus</i> ) in nearshore survey region. Stratum areas are $\text{nmi}^2$ . . . . .	55
101			
102			
103	14	Abundance versus fork length ( $L_F$ , cm) for Pacific Mackerel ( <i>Scomber japonicus</i> ) in the core and nearshore survey regions. . . . .	56
104			
105	15	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for Jack Mackerel ( <i>Trachurus symmetricus</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	59
106			
107			
108	16	Abundance versus fork length ( $L_F$ , cm) for Jack Mackerel ( <i>Trachurus symmetricus</i> ) in the core and nearshore survey regions. . . . .	60
109			

110	17	Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals, $CI_{95\%}$ ; and coefficients of variation, CVs) for Pacific Herring ( <i>Clupea pallasii</i> ) in the core and nearshore survey regions. Stratum areas are $\text{nmi}^2$ . . . . .	64
111			
112			
113	18	Abundance versus fork length ( $L_F$ , cm) for Pacific Herring ( <i>Clupea pallasii</i> ) in the core and nearshore survey regions. . . . .	65
114			

DRAFT



## 115 List of Figures

116	1	Conceptual spring (shaded region) and summer (hashed region) distributions of potential habitat for the northern stock of Pacific Sardine along the west coasts of Mexico, the United States, and Canada. The dashed and dotted lines represent, respectively, the approximate summer and spring positions of the $0.2 \text{ mg m}^{-3}$ chlorophyll-a concentration isoline. This isoline appears to oscillate in synchrony with the transition zone chlorophyll front (TZCF, Polovina <i>et al.</i> , 2001) and the offshore limit of the northern stock Pacific Sardine potential habitat (Zwolinski <i>et al.</i> , 2011). Mackerels are found within and on the edge of the same oceanographic habitat (e.g., Demer <i>et al.</i> , 2012; Zwolinski <i>et al.</i> , 2012). The TZCF may delineate the offshore and southern limit of both Pacific Sardine and Pacific Mackerel distributions, and juveniles may have nursery areas in the SCB, downstream of upwelling regions. . . . .	4
126	2	Planned compulsory and adaptive transects sampled by <i>Lasker</i> and <i>Carranza</i> ; interstitial and offshore transects sampled by USVs; and nearshore transects sampled by <i>Lisa Marie</i> and <i>Long Beach Carnage</i> . Isobaths (light gray lines) are 50, 200, 500, and 2,000 m. . . . .	7
129	3	Echosounder transducers mounted on the bottom of the retractable centerboard on <i>Lasker</i> . During the survey, the centerboard was extended, typically positioning the transducers $\sim 2$ m below the keel at a water depth of $\sim 7$ m. . . . .	9
132	4	Transducers (Top-bottom: Simrad ES200-7C, ES120-7C, ES38-12, and ES70-7C, Kongsberg) in a pole-mounted multi-transducer array (MTA4) installed on <i>Long Beach Carnage</i> . . . . .	9
134	5	Schematic drawings of the Nordic 264 rope trawl a) net and b) cod-end. . . . .	15
135	6	Example depths (m) of the trawl headrope (red line) and footrope (blue line) measured using temperature-depth recorders (TDRs) during the net deployment (dashed box) and when actively fishing (shaded region). The vessel speed over ground (kn, black line) was measured using the ship's GPS. . . . .	16
139	7	Specimen length versus weight from the current survey (colored points, by sex) compared to those from previous SWFSC surveys during the same season (gray points, all sexes) and models [dashed lines; Palance <i>et al.</i> (2019)]. . . . .	18
142	8	Two examples of echograms depicting CPS schools (red) and plankton aggregations (blue and green) at 38 kHz (top) and 120 kHz (bottom). Example data processing steps include the original echogram (a, d), after noise subtraction and bin-averaging (b, e), and after filtering to retain only putative CPS echoes (c, f). . . . .	21
146	9	Echoes from fishes with swimbladders (blue points, scaled by backscatter intensity) along an example acoustic transect (top) and the corresponding echogram image (bottom). In this example, the upper (blue) and lower lines (green) indicate boundaries within which echoes were retained. When the lower boundary is deeper than the seabed (black line), echoes above the seabed are retained. Echoes from deep, bottom-dwelling schools of non-CPS fishes with swimbladders, and from diffuse scatterers near the surface were excluded. The proximity of the echoes to the seabed was also used to define the lower limit for vertical integration. . . .	21
153	10	a) Polygons enclosing 100-m acoustic intervals from <i>Lasker</i> , <i>Lisa Marie</i> , and USVs assigned to catches from each trawl cluster or purse-seine set, and b) the acoustic proportions of CPS in catches from trawl clusters or purse-seine sets. See <b>Section 3.5.1</b> and <b>Fig. 13</b> for a description of the method used for this survey to estimate species proportions and Jack Mackerel lengths in the area between Cape Flattery and Cape Mendocino. The numbers inside each polygon in panel a) are the cluster or purse-seine numbers, which are located at the average latitude and longitude of all trawls in that cluster or each individual purse-seine set. Black points in panel b) indicate trawl clusters or purse-seine sets with no CPS present in the catch. . . . .	25

162	11	Biomass density ( $\log_{10}(t \text{ nmi}^2 + 1)$ ) versus latitude (easternmost portion of each transect) and strata used to estimate biomass and abundance (shaded regions; outline indicates stratum number) for each species in the core survey region. Data labels (blue numbers) correspond to transects with positive biomass ( $\log_{10}(t + 1) > 0.01$ ). Transect spacing (nmi; point color), and stock breaks for Northern Anchovy and Pacific Sardine (red dashed lines and text) are indicated. . . . .	27
163			
164			
165			
166			
167			
168	12	Summary of all transects sampled throughout the survey by <i>Lasker</i> (red) and <i>Lisa Marie</i> (yellow) in relation to the probability thresholds of the updated model of potential habitat for the northern stock of Pacific Sardine (Zwolinski and Demer, In prep.). The shaded regions with probabilities higher than 0.29 and 0.18 contained 82% and 95% of all spring northern stock Pacific Sardine spawning stock biomass, respectively. The habitat-model output is averaged in areas, $\pm 2^\circ$ latitude and longitude, centered around the daytime location of each vessel throughout the survey. Areas without data (white), occurred in the analysis domain where and when clouds prevented satellite-sensed observations. . . . .	28
169			
170			
171			
172			
173			
174			
175			
176	13	Maps of species proportions in <i>Lasker's</i> nighttime trawl-catch clusters during summer a) 2018, b) 2019, and c) 2021 depicting an analysis region between the Columbia River and Cape Mendocino (dashed line); d) a model of the acoustic proportions of Pacific Sardine versus latitude in <i>Lasker's</i> nighttime trawl-catch clusters between 2018 and 2021; e) species proportions in <i>Lisa Marie's</i> summer 2022 purse-seine catches; and f) the species proportions used in the estimation of summer 2022 CPS biomasses north of Cape Mendocino, replacing <i>Lisa Marie's</i> catches containing Pacific Sardine, in the analysis region, with the modeled proportion of Pacific Sardine, with its complement assigned to Jack Mackerel. . . . .	33
177			
178			
179			
180			
181			
182			
183			
184	14	Proportion (top) and cumulative proportion (bottom) of biomass versus distance to the nearest positive trawl cluster. Dashed vertical lines (bottom) represent the cluster distance where cumulative biomass equals 90%. . . . .	34
185			
186			
187	15	Spatial distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ , $\text{m}^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; b) CUFES egg density (eggs $\text{m}^{-3}$ ) for Northern Anchovy, Pacific Sardine, and Jack Mackerel; and c) acoustic proportions of CPS in trawl clusters (black outline) and purse-seine sets (white outline). Black points indicate trawl clusters or purse-seine sets with no CPS. . . . .	35
188			
189			
190			
191			
192	16	Nearshore survey transects sampled by <i>Lisa Marie</i> overlaid with the distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ , $\text{m}^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; and b) the proportions, by weight, of CPS in each purse-seine catch. Species with low catch weights may not be visible at this scale. . . . .	36
193			
194			
195			
196	17	Nearshore transects sampled by <i>Long Beach Carnage</i> overlaid with the distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ , $\text{m}^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; and b) the proportions, by weight, of CPS in each purse-seine catch. Species with low catch weights may not be visible at this scale. . . . .	37
197			
198			
199			
200	18	Spatial distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ , $\text{m}^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS from nearshore sampling; and b) acoustic proportions of CPS in trawl clusters (black outline) and purse-seine sets (white outline). . . . .	38
201			
202			
203	19	Biomass densities (colored points) of the northern stock of Northern Anchovy ( <i>Engraulis mordax</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Northern Anchovy (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	41
204			
205			
206			

207	20	Abundance versus standard length ( $L_S$ , upper panels) and biomass (t) versus $L_S$ (lower panels) for the northern stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. This plot correctly shows the abundances and biomasses in the core region, which is based on one specimen resulting in the mode at 10 cm (see Cluster 60 in <b>Appendix A.1</b> ). Abundance and biomass in the nearshore region is negligible relative to the core region and not visible at this scale. . . . .	42
208			
209			
210			
211			
212			
213	21	Biomass densities (colored points) of central stock of Northern Anchovy ( <i>Engraulis mordax</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Northern Anchovy (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	45
214			
215			
216			
217	22	Abundance versus standard length ( $L_S$ , upper panels) and biomass (t) versus $L_S$ (lower panels) for the central stock of Northern Anchovy ( <i>Engraulis mordax</i> ) in the core and nearshore survey regions. . . . .	46
218			
219			
220	23	Biomass densities (colored points) of the northern stock of Pacific Sardine ( <i>Sardinops sagax</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Sardine (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	49
221			
222			
223			
224	24	Estimated abundance (upper panel) and biomass (lower panel) versus standard length ( $L_S$ , cm) for the northern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. . . . .	50
225			
226			
227	25	Biomass densities (colored points) of the southern stock of Pacific Sardine ( <i>Sardinops sagax</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Sardine (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	53
228			
229			
230			
231	26	Estimated abundance (upper panels) and biomass (lower panels) versus standard length ( $L_S$ , cm) for the southern stock of Pacific Sardine ( <i>Sardinops sagax</i> ) in the core and nearshore survey regions. . . . .	54
232			
233			
234	27	Biomass densities (colored points) of Pacific Mackerel ( <i>Scomber japonicus</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Mackerel (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	57
235			
236			
237			
238	28	Estimated abundance (upper panels) and biomass (lower panels) versus fork length ( $L_F$ , cm) for Pacific Mackerel ( <i>Scomber japonicus</i> ) in the core and nearshore survey regions. . . . .	58
239			
240	29	Biomass densities (colored points) of Jack Mackerel ( <i>Trachurus symmetricus</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Jack Mackerel (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	62
241			
242			
243			
244	30	Estimated abundance (upper panel) and biomass (lower panel) versus fork length ( $L_F$ , cm) for Jack Mackerel ( <i>Trachurus symmetricus</i> ) in the core and nearshore survey regions. . . . .	63
245			
246	31	Biomass densities (colored points) of Pacific Herring ( <i>Clupea pallasii</i> ), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Herring (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects. . . . .	66
247			
248			
249			
250	32	Estimated abundance (upper panel) and biomass (lower panel) versus fork length ( $L_F$ , cm) for Pacific Herring ( <i>Clupea pallasii</i> ) in the core and nearshore survey regions. . . . .	67
251			

252	33	Differentiation of northern (blue) and southern (red) stocks of Pacific Sardine by: a) length	
253		distributions; b) individual (grey points) and catch-mean (colored points) lengths at the lat-	
254		itudes of their respective trawls; and c) geographic locations of trawls catches with (colored	
255		points) and without (black points) Pacific Sardine. . . . .	70
256	34	Distributions of species proportions in <i>Lasker's</i> nighttime trawl catches, summer 2015 through	
257		2022. In 2015, the integrated CPS-hake survey sample northward of Vancouver Island. In	
258		2017, there was no sampling in the SCB. In 2020, there was no survey due to the COVID-19	
259		pandemic. In 2021, through a collaboration with Mexico, the CPS survey extended farther	
260		south into Baja California. In 2022, there was no nighttime trawl sampling north of Cape	
261		Mendocino, California. . . . .	73
262	35	a) Estimated and b) cumulative estimated biomasses ( $t$ ) of the eight most abundant CPS	
263		stocks of six species in the CCE during summer since 2008. Surveys typically span the area	
264		between Cape Flattery and San Diego, but in some years also include Vancouver Island,	
265		Canada (2015-2019) and portions of Baja CA (2021-2022). . . . .	74

DRAFT

## Executive Summary

This report provides: 1) a detailed description of the acoustic-trawl method (ATM) used by NOAA's Southwest Fisheries Science Center for direct assessments of the dominant coastal pelagic species (CPS; i.e.: Pacific Sardine *Sardinops sagax*, Northern Anchovy *Engraulis mordax*, Pacific Mackerel *Scomber japonicus*, Jack Mackerel *Trachurus symmetricus*, and Pacific Herring *Clupea pallasii* in the California Current Ecosystem off the west coast of the United States (U.S.) and portions of Baja CA, Mexico (MX); and 2) estimates of the biomasses, distributions, and demographics of those CPS encountered in the survey area between 27 June and 30 September 2022.

The core survey region, which was to be sampled by NOAA ship *Reuben Lasker* (hereafter, *Lasker*) and two wind-powered uncrewed surface vehicles (Explorer USVs; Sairdron, Inc.), spanned most of the continental shelf between Cape Flattery, WA and Punta Baja, Baja CA Norte. Planned transects were oriented approximately perpendicular to the coast, from the shallowest navigable depth (~20 m) to either a distance of 35 nmi or to the 1,000 fth (~1830 m) isobath, whichever is farthest. In the SCB, transects in the core region were extended to approximately 75 nmi. However, after losing roughly half of the scheduled sea days aboard *Lasker*, the plan was modified for chartered fishing vessel *Lisa Marie* to survey *Lasker*'s transects, 20 nmi apart, between Cape Flattery, WA and Cape Mendocino, but extended into the nearshore region to ~5 m depth. Both *Lisa Marie* and *Lasker* sampled in the area between Cape Mendocino and Bodega Bay, and then *Lasker* sampled farther south, ending at Punta Baja. North of Cape Mendocino, where *Lasker* did not sample, species composition and CPS length distributions were estimated from *Lisa Marie*'s daytime purse-seine catches, but adjusted to reflect the associations between Pacific Sardine and Jack Mackerel in this region during summer 2018-2021 (see **Section 3.5.1**). Between Cape Mendocino and Punta Baja, species composition and CPS length distributions were estimated, as usual, by the catches from nighttime surface trawls.

Because sampling by *Lasker* and the USVs in water shallower than ~20 m was deemed inefficient, unsafe, or both, fishing vessel *Long Beach Carnage* sampled CPS in the U.S. nearshore region, along 2.5 to 5 nmi-long transects spaced 5 nmi apart off the mainland coast of the U.S., between Bodega Bay and San Diego, as well as around Santa Cruz and Santa Catalina Islands in the Southern CA Bight. In the nearshore region, the species composition and CPS length distributions were estimated using catches daytime purse-seine sets by *Long Beach Carnage* or nighttime surface trawls by *Lasker*, whichever was nearest to the acoustically sampled CPS.

The biomasses, distributions, and demographics for each species and stock are for the survey area and period, and therefore may not represent their entire population or stock. Nearshore sampling was not conducted off Baja CA, so nearshore biomass estimates are for U.S. waters only.

The estimated biomass of the northern stock of Northern Anchovy was 16,432 t ( $CI_{95\%} = 5,646 - 27,680$  t,  $CV = 34\%$ ). In the core region, the biomass was 16,432 t ( $CI_{95\%} = 5,646 - 27,680$  t,  $CV = 34\%$ ), and in the nearshore region, biomass was 0.0934 t ( $CI_{95\%} = 0 - 0.285$  t,  $CV = 94\%$ ), or 0.00057% of the total biomass. The estimated nearshore biomass includes uncertainty associated with the assumed nearshore-area (see **Section 3.5.1**). The northern stock ranged from approximately Westport, WA to Cape Blanco, OR, and the distribution of their standard length ( $L_S$ ) ranged from 10 to 15 cm with a mode at 10 cm in both the core and nearshore regions.

The estimated biomass of the central stock of Northern Anchovy was 2,235,996 t ( $CI_{95\%} = 1,248,956 - 3,051,863$  t,  $CV = 20\%$ ), of which 6% was observed in Mexican waters. In the core region, the biomass was 2,197,812 t ( $CI_{95\%} = 1,231,227 - 3,002,630$  t,  $CV = 21\%$ ), and in the nearshore region, the biomass was 38,184 t ( $CI_{95\%} = 17,729 - 49,233$  t,  $CV = 21\%$ ), or 1.7% of the total biomass. The central stock ranged from approximately Bodega Bay to El Rosario, and the distribution of their  $L_S$  ranged from 5 to 16 cm with modes at 9 and 12 cm in the core region and 9 cm in the nearshore region.

The estimated biomass of the northern stock of Pacific Sardine was 69,506 t ( $CI_{95\%} = 30,484 - 99,021$  t,  $CV = 21\%$ ). In the core region, the biomass was 53,741 t ( $CI_{95\%} = 29,672 - 84,749$  t,  $CV = 26\%$ ), and in the nearshore region, the biomass was 15,765 t ( $CI_{95\%} = 812 - 14,272$  t,  $CV = 23\%$ ), or 23% of the total biomass. The nearshore and total biomasses include uncertainties associated with the assumed nearshore-area (see

316 **Section 2.2.5**) and the estimated species proportions from the daytime purse-seine sets (see **Section 3.5.1**),  
317 respectively. Within the survey area, the northern stock ranged from approximately Westport, WA to Point  
318 Conception. The distribution of  $L_S$  ranged from 11 to 27 cm with a mode at 16 cm in the core region and  
319 modes at 11 and 14 cm in the nearshore region.

320 The estimated biomass of the southern stock of Pacific Sardine in the surveyed area was 107,468 t ( $CI_{95\%}$   
321 = 47,994 - 178,947 t, CV = 23%), of which 0.9% was observed in Mexican waters. In the core region, the  
322 biomass was 40,206 t ( $CI_{95\%}$  = 4,741 - 79,328 t, CV = 48%), and in the nearshore region, the biomass was  
323 67,262 t ( $CI_{95\%}$  = 43,253 - 99,620 t, CV = 23%), or 63% of the total biomass. Within the survey area, the  
324 southern stock ranged from approximately Point Conception to El Rosario. The distribution of  $L_S$  ranged  
325 from 9 to 21 cm with modes at 13 and 18 cm in the core region and at 12 and 16 cm in the nearshore region.

326 The estimated biomass of Pacific Mackerel was 7,968 t ( $CI_{95\%}$  = 3,741 - 12,662 t, CV = 22%), of which 27%  
327 was observed in Mexican waters. In the core region, the biomass was 5,619 t ( $CI_{95\%}$  = 2,851 - 9,108 t, CV  
328 = 29%), and in the nearshore region, the biomass was 2,349 t ( $CI_{95\%}$  = 890 - 3,553 t, CV = 30%), or 29%  
329 of the total biomass. The estimated nearshore biomass includes uncertainty associated with the assumed  
330 nearshore-area (see **Section 2.2.5**). Pacific Mackerel ranged from approximately Cape Mendocino to El  
331 Rosario, but was mostly south of Point Conception and around Santa Cruz and Santa Catalina Islands. The  
332 distribution of fork length ( $L_F$ ) ranged from 8 to 38 cm with modes at 11 and 15 cm in the core region, and  
333 at 18 and 27 cm in the nearshore region.

334 The estimated biomass of Jack Mackerel was 807,090 t ( $CI_{95\%}$  = 515,560 - 1,145,812 t, CV = 20%), of  
335 which 0.06% was observed in Mexican waters. In the core region, biomass was 799,082 t ( $CI_{95\%}$  = 512,231 -  
336 1,132,052 t, CV = 20%), and, in the nearshore region, biomass was 8,009 t ( $CI_{95\%}$  = 3,328 - 13,761 t, CV =  
337 35%), or 0.99% of the total biomass. The nearshore and total biomasses include uncertainties associated with  
338 the assumed nearshore-area (see **Section 2.2.5**) and the estimated species proportions from the daytime  
339 purse-seine sets (see **Section 3.5.1**), respectively. Jack Mackerel were present throughout the survey area  
340 from the Columbia River to El Rosario, but were most abundant in the core region between Astoria and  
341 Bodega Bay and around Santa Cruz Island in the nearshore region. The distribution of  $L_F$  ranged from 3  
342 to 51 cm with modes at 8 and 34 cm in the core region, and at 19 cm in the nearshore region.

343 The total estimated biomass of Pacific Herring was 50,718 t ( $CI_{95\%}$  = 14,460 - 99,700 t, CV = 41%).  
344 In the core region, biomass was 47,024 t ( $CI_{95\%}$  = 13,306 - 93,207 t, CV = 44%), and was distributed  
345 from approximately Cape Flattery to Cape Mendocino, but was most abundant from Cape Flattery to the  
346 Columbia River, and between Crescent City and Cape Mendocino. The distribution of  $L_F$  ranged from 13  
347 to 17 cm, with modes at 13 and 17 cm. In the nearshore region, biomass was 3,694 t ( $CI_{95\%}$  = 1,154 - 6,493  
348 t, CV = 36%), or 7.3% of the total biomass. The distribution of  $L_F$  ranged from 13 to 17 cm, and had a  
349 mode at 14 cm.

350 The total estimated biomass of seven stocks of five species within the survey area was 3,295,179 t. Of this  
351 68% (2,235,996 t) was from the central stock of Northern Anchovy. Proportions of other stocks, in decreasing  
352 order, were Jack Mackerel (24%), southern stock of Pacific Sardine (3%), northern stock of Pacific Sardine  
353 (2.1%), Pacific Herring (1.5%), northern stock of Northern Anchovy (0.5%), and Pacific Mackerel (0.2%).  
354 The biomass of the central stock of Northern Anchovy, which had been growing exponentially since 2015,  
355 decreased ~20% from 2,721,689 t estimated in summer 2021 (Stierhoff *et al.*, 2023). Jack Mackerel, which  
356 were found mostly north of Cape Mendocino, included an abundance of apparently age-0 fish farther south,  
357 suggesting a strong recruitment. The southern stock of Pacific Sardine was found mostly north of the  
358 U.S.-Mexico border and more were found nearshore than in the core area. Even considering the additional  
359 uncertainties in the biomass estimates north of Cape Mendocino (see **Sections 2.2.5** and **3.5.1**) there is  
360 no indication that the biomasses of the northern stock of Pacific Sardine have changed significantly since  
361 summer 2021.



# 1 Introduction

In the California Current Ecosystem (CCE), multiple coastal pelagic fish species (CPS; i.e.: Pacific Sardine *Sardinops sagax*, Northern Anchovy *Engraulis mordax*, Jack Mackerel *Trachurus symmetricus*, Pacific Mackerel *Scomber japonicus*, and Pacific Herring *Clupea pallasii*) comprise the bulk of the forage fish assemblage. These populations, which can change by an order of magnitude within a few years, represent important prey for marine mammals, birds, and larger migratory fishes (Field *et al.*, 2001), and are targets of commercial fisheries.

During summer and fall, the northern stock of Pacific Sardine typically migrates north to feed in the productive coastal upwelling off OR, WA, and Vancouver Island (Zwolinski *et al.*, 2012, and references therein, **Fig. 1**). In synchrony, but separately, the southern stock of Pacific Sardine migrates from Northern Baja CA, Mexico to the Southern CA Bight (SCB) (Smith, 2005). The predominantly piscivorous adult Pacific and Jack Mackerels also migrate north in summer, but go farther offshore to feed (Zwolinski *et al.*, 2014 and references therein). In the winter and spring, the northern stock of Pacific Sardine typically migrates south to its spawning grounds, generally off Central and Southern CA (Demer *et al.*, 2012) and occasionally off OR and WA (Lo *et al.*, 2011). These migrations vary in extent with population size; fish age and length; and oceanographic conditions (Zwolinski *et al.*, 2012). For example, the transition zone chlorophyll front [TZCF; Polovina *et al.* (2001)] may delineate the offshore and southern limit of both Pacific Sardine and Pacific Mackerel habitat (e.g., Demer *et al.*, 2012; Zwolinski *et al.*, 2012), and juveniles may have nursery areas in the SCB, downstream of upwelling regions. In contrast, Northern Anchovy spawn predominantly during winter and closer to the coast where seasonal down-welling increases retention of their eggs and larvae (Bakun and Parrish, 1982). Pacific Herring spawn in intertidal beach areas (Love, 1996). The northern stock of Northern Anchovy is located off WA and OR and the central stock is located off Central and Southern CA and northern Baja CA. Whether a species migrates or remains in an area depends on its reproductive and feeding behaviors, affinity to certain oceanographic or seabed habitats, and its population size.

Acoustic-trawl method (ATM) surveys, which combine information collected with echosounders and nets, were introduced to the CCE more than 48 years ago to survey CPS off the west coast of the United States (U.S.) (Mais, 1974, 1977; Smith, 1978). Following a two-decade hiatus, the ATM was reintroduced in the CCE in spring 2006 to sample the then-abundant Pacific Sardine population (Cutter and Demer, 2008). Since then, this sampling effort has continued and expanded through annual or semi-annual surveys (Demer *et al.*, 2012; Zwolinski *et al.*, 2014). Beginning in 2011, the ATM estimates of Pacific Sardine abundance, age structure, and distribution have been incorporated in the annual assessments of the northern stock (Hill *et al.*, 2017). ATM estimates are used in assessments of Pacific Mackerel (Crone *et al.*, 2019; Crone and Hill, 2015) and the central stock of Northern Anchovy (Kuriyama *et al.*, 2022). Additionally, ATM survey results have yielded estimated abundances, demographics, and distributions of epipelagic and semi-demersal fishes (e.g., Swartzman, 1997; Williams *et al.*, 2013; Zwolinski *et al.*, 2014) and zooplankton (Hewitt and Demer, 2000).

This document, and references herein, describes in detail the ATM as presently used by NOAA's Southwest Fisheries Science Center (SWFSC) to survey the distributions and abundances of CPS and their oceanographic environments (e.g., Cutter and Demer, 2008; Demer *et al.*, 2012; Zwolinski *et al.*, 2014). In general terms, the contemporary ATM combines information from satellite-sensed oceanographic conditions, multi-frequency echosounders, probe-sampled oceanographic conditions, pumped samples of fish eggs, and trawl-net catches of juvenile and adult CPS. The survey area is initially defined with consideration to the potential habitat of the northern stock of Pacific Sardine for summer surveys (**Fig. 1**) or the central stock of Northern Anchovy for spring surveys, when they occur. The survey area is further expanded to encompass as much of the potential habitat as possible for other CPS present off the West Coast of the U.S. and Baja CA, as time permits.

Along transects in the survey area, multi-frequency split-beam echosounders transmit sound pulses downward beneath the ship and receive echoes from animals and the seabed in the path of the sound waves. Measurements of sound speed and absorption from conductivity-temperature-depth (CTD) probes allow accurate compensation of these echoes for propagation losses. The calibrated echo intensities, normalized to the range-dependent observational volume, provide indications of the target type and behavior (e.g., Demer




Figure 1: Conceptual spring (shaded region) and summer (hashed region) distributions of potential habitat for the northern stock of Pacific Sardine along the west coasts of Mexico, the United States, and Canada. The dashed and dotted lines represent, respectively, the approximate summer and spring positions of the 0.2 mg m<sup>-3</sup> chlorophyll-a concentration isoline. This isoline appears to oscillate in synchrony with the transition zone chlorophyll front (TZCF, Polovina *et al.*, 2001) and the offshore limit of the northern stock Pacific Sardine potential habitat (Zwolinski *et al.*, 2011). Mackerels are found within and on the edge of the same oceanographic habitat (e.g., Demer *et al.*, 2012; Zwolinski *et al.*, 2012). The TZCF may delineate the offshore and southern limit of both Pacific Sardine and Pacific Mackerel distributions, and juveniles may have nursery areas in the SCB, downstream of upwelling regions.



414 Echoes from marine organisms are a function of their body composition, shape, and size relative to the  
415 sensing-sound wavelength, and their orientation relative to the incident sound waves (Cutter *et al.*, 2009;  
416 Demer *et al.*, 2009b; Renfree *et al.*, 2009). Variations in echo intensity across frequencies, known as echo  
417 spectra, indicate the taxonomic groups contributing to the echoes. The CPS, with highly reflective swim  
418 bladders, create high intensity echoes of sound pulses at all echosounder frequencies (e.g., Conti and Demer,  
419 2003). In contrast, krill, with acoustic properties closer to those of the surrounding seawater, produce lower  
420 intensity echoes, particularly at lower frequencies (e.g., Demer *et al.*, 2003). The echo energy attributed  
421 to CPS, based on empirical echo spectra (Demer *et al.*, 2012), are apportioned to species using trawl-catch  
422 proportions (Zwolinski *et al.*, 2014).

423 Animal densities are estimated by dividing the summed intensities attributed to a species by the length-  
424 weighted average echo intensity, i.e., the mean backscattering cross-section, from animals of that species (e.g.,  
425 Demer *et al.*, 2012). Transects with similar densities are grouped into post-sampling strata that mimic the  
426 natural patchiness of the target species (e.g., Zwolinski *et al.*, 2014). An estimate of abundance is obtained  
427 by multiplying the average estimated density in the stratum by the stratum area (Demer *et al.*, 2012). The  
428 associated sampling variance is calculated using non-parametric bootstrap of the mean transect densities.  
429 The total abundance estimate in the survey area is the sum of abundances in all strata. Similarly, the total  
430 variance estimate is the sum of the variance in each stratum.

431 The primary objectives of the SWFSC's ATM surveys are to survey the distributions and abundances of  
432 CPS, krill, and their abiotic environments in the CCE. Typically, spring surveys are conducted during 25-40  
433 days-at-sea (DAS) between March and May, and summer surveys are conducted during 50-90 DAS between  
434 June and October. In spring, the ATM surveys focus primarily on the northern stock of Pacific Sardine  
435 and the central stock of Northern Anchovy. In summer, the ATM surveys also include the northern stock of  
436 Northern Anchovy and Pacific Herring. During spring and summer, biomasses are also estimated for other  
437 CPS (e.g., Pacific Mackerel, Jack Mackerel, and Round Herring) present in the survey area.

438 In summer 2022, the ATM survey, spanning U.S. and Mexican waters, was conducted by fishing vessel *Lisa*  
439 *Marie*, between Cape Flattery and Cape Mendocino, and by *Lasker* from Cape Mendocino to Punta Baja.  
440 Between Newport, OR and Bodega Bay, along adaptive transects not sampled by *Lisa Marie* or *Lasker*, two  
441 wind-powered uncrewed surface vessels (Explorer USVs; Sailable, Inc.) conducted acoustic sampling. From  
442 Cape Flattery to San Diego, sampling from fishing vessels *Lisa Marie* and *Long Beach Carnage* was used to  
443 estimate the biomasses of CPS in the nearshore regions, where sampling by *Lasker* was not possible or safe.

444 Presented here are: 1) a detailed description of the ATM used to survey CPS in the California Current  
445 Ecosystem (CCE) off the west coast of the U.S. and portions of Baja CA; and 2) estimates of the abundances,  
446 biomasses, size structures, and distributions of CPS, specifically the northern and southern stocks of Pacific  
447 Sardine; the central and northern stocks of Northern Anchovy, Pacific Mackerel, Jack Mackerel, and Pacific  
448 Herring for the core and nearshore survey regions in which they were sampled. Additional details about the  
449 survey may be found in the survey report (Renfree *et al.*, 2023).

450 This survey was conducted with the approval of the Secretaria de Relaciones Exteriores (SRE, Diplomatic  
451 note UAN0807/2022), the Instituto Nacional de Estadística y Geografía (INEGI; Authorization: EG0022022,  
452 through official letters 400./58/2022 and 400./59/2022), Unidad de Planeación y Coordinación Estratégica de  
453 la Secretaría de Marina (SEMAR; Letter no AI/1223/22), Unidad Coordinadora de Asuntos Internacionales  
454 (UCAI) de la Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT; Letter UCAI/01314/2022),  
455 Universidad Nacional Autónoma de México (UNAM; Letter ICML/DIR/127/2022), Unidad de Concesiones  
456 y Servicios del Instituto Federal de Telecomunicaciones (IFT; Letter IFT/223/UCS/DG-AUSE/3059/2022),  
457 and the Comisión Nacional de Acuacultura y Pesca (CONAPESCA; Permit: PPFE/DGPPE.-05325/120722).

## 2 Methods

### 2.1 Sampling

#### 2.1.1 Design

The summer 2022 survey was conducted principally using *Lasker* and *Lisa Marie*, but was augmented with acoustic sampling by two uncrewed surface vehicles (USVs), and nearshore sampling by *Long Beach Carnage*. The sampling domain, or core region, between Cape Flattery, WA and Punta Baja, Baja CA Norte, was defined by the conceptual distribution of potential habitat for the northern stock of Pacific Sardine in summer (Fig. 1), but also encompassed the anticipated distributions of the southern stock of Pacific Sardine and the central and northern stocks of Northern Anchovy off the west coasts of the U.S. and portion of Baja CA, Mexico. It also spanned portions of the Pacific Mackerel, Jack Mackerel, Pacific Herring, and Round Herring populations. East to west, the sampling domain extended from the coast to at least the 1,000 ftm (~1830 m) isobath (Fig. 2). Considering the expected distribution of the target species, the acceptable uncertainty in biomass estimates, and the available ship time (75 days at sea, DAS), the principal survey objectives were to estimate the biomasses of the northern and southern stocks of Pacific Sardine and the northern and central stocks of Northern Anchovy in the survey region. Secondary objectives were to estimate the population biomasses of Pacific Mackerel, Jack Mackerel, Pacific Herring, and Round Herring in the survey region.

The core region transects were perpendicular to the coast, extending from the shallowest navigable depth (~20 m) to either a distance of 35 nmi or to the 1,000 ftm isobath, whichever was farthest (Fig. 2). Compulsory transects were spaced 10 nmi-apart in areas of historic CPS abundance (e.g., between Cape Flattery and Newport, and between San Francisco and San Diego) and 20-nmi apart elsewhere. When CPS were observed within the westernmost 3 nmi of a transect, that transect and the next one to the south were extended in 5-nmi increments until no CPS were observed in the last 3 nmi of the extension, to a maximum extension of 50 nmi. If a transect was extended, the ensuing transect was extended by the same amount.

Leg I on *Lasker* was cancelled for reasons outside of the SWFSC's control, so *Lisa Marie* was directed to sample the transects between Cape Flattery, WA to Bodega Bay, CA using 20-nmi spacing, including the nearshore portions of those transects. Meanwhile USVs (SD-1076 and SD-1077) were directed to sample the interstitial transects, also spaced 20-nmi apart, from Newport, OR to San Francisco (cyan lines, Fig. 2). Leg III on *Lasker* was delayed by 16 days for similar reasons, so the remaining four days were used to sample transects spaced 20-nmi apart, from San Diego northward into the Southern California Bight (SCB). During Leg IV on *Lasker*, transects were sampled southward from Monterey, CA to Punta Baja, Baja CA Norte. To progress as far south as possible, compulsory transects off Baja CA were shortened to 30-nmi, and adaptive transects were omitted.

To estimate the abundances and biomasses of CPS between Cape Flattery and San Diego, in the nearshore area where *Lasker* and the USVs could not efficiently or safely navigate or trawl, two fishing vessels conducted acoustic and purse-seine sampling (magenta lines, Fig. 2). *Lisa Marie* sampled transects to ~5-m depth, spaced 20 nmi apart between Cape Flattery and Bodega Bay. *Long Beach Carnage* sampled 5-nmi-long transects spaced 5 nmi apart between Bodega Bay and San Diego, and 2.5-nmi-long transects spaced 2.5 nmi apart around Santa Cruz and Santa Catalina Islands in the SCB.




Figure 2: Planned compulsory and adaptive transects sampled by *Lasker* and *Carranza*; interstitial and off-shore transects sampled by USVs; and nearshore transects sampled by *Lisa Marie* and *Long Beach Carnage*. Isobaths (light gray lines) are 50, 200, 500, and 2,000 m.

496 **2.1.2 Acoustic**

497 **2.1.2.1 Acoustic equipment**

498 **2.1.2.1.1 *Lasker*** Multi-frequency Wide-Bandwidth Transceivers (18-, 38-, 70-, 120-, 200-, and 333-kHz  
499 Simrad EK80 WBTs; Kongsberg) were configured with split-beam transducers (Simrad ES18-11, ES38B,  
500 ES70-7C, ES120-7C, ES200-7C, and ES333-7C, respectively; Kongsberg). The transducers were mounted on  
501 the bottom of a retractable keel or “centerboard” (**Fig. 3**). The keel was retracted (transducers at ~5-m  
502 depth) during calibration, and extended to the intermediate position (transducers at ~7-m depth) during  
503 the survey. Exceptions were made during shallow water operations, when the keel was retracted; or during  
504 times of heavy weather, when the keel was extended (transducers at ~9-m depth) to provide extra stability  
505 and reduce the effect of weather-generated noise. In addition, acoustic data were also collected using a  
506 multibeam echosounder (Simrad ME70; Kongsberg), multibeam sonar (Simrad MS70; Kongsberg), scanning  
507 sonar (Simrad SX90; Kongsberg), acoustic Doppler current profiler and echosounder (Simrad EC150-3C,  
508 Kongsberg), and a separate ADCP (Ocean Surveyor OS75; Teledyne RD Instruments). Transducer position  
509 and motion were measured at 5 Hz using an inertial motion unit (Applanix POS-MV; Trimble).

510 **2.1.2.1.2 *Lisa Marie*** On *Lisa Marie*, multi-frequency Wideband Transceivers (Simrad 38- and 200-  
511 kHz EK80 WBTs; Kongsberg) were connected to the vessel’s hull-mounted split-beam transducers (Simrad  
512 ES38-7 and ES200-7C; Kongsberg). The transducers were at a water depth of ~4 m.

513 **2.1.2.1.3 *Long Beach Carnage*** On *Long Beach Carnage*, the SWFSC’s multi-frequency General Pur-  
514 pose Transceivers (38-, 70-, 120-, and 200-kHz Simrad EK60 GPTs; Kongsberg) were configured with the  
515 SWFSC’s split-beam transducers (Simrad ES38-12, ES70-7C, ES120-7C and ES200-7C; Kongsberg) mounted  
516 in a multi-frequency transducer array (MTA4) on the bottom of a retractable pole (**Fig. 4**). The transducers  
517 were at a water depth of roughly 2 m.

518 **2.1.2.1.4 USVs** On the two USVs (SD-1076 and SD-1077), miniature Wide-Bandwidth Transceivers  
519 (Simrad WBT-Mini; Kongsberg) were configured with gimbaled, keel-mounted, dual-frequency transducers  
520 (Simrad ES38-18|200-18C; Kongsberg) containing a split-beam 38-kHz transducer and single-beam 200-kHz  
521 transducer with nominally 18° beamwidths. The transducers were at a water depth of ~1.9 m.




Figure 3: Echosounder transducers mounted on the bottom of the retractable centerboard on *Lasker*. During the survey, the centerboard was extended, typically positioning the transducers ~2 m below the keel at a water depth of ~7 m.



Figure 4: Transducers (Top-bottom: Simrad ES200-7C, ES120-7C, ES38-12, and ES70-7C, Kongsberg) in a pole-mounted multi-transducer array (MTA4) installed on *Long Beach Carnage*.

522 **2.1.2.2 Echosounder calibrations**

523 **2.1.2.2.1 Lasker** The echosounder systems aboard *Lasker* were calibrated on 23 June while the vessel  
524 was docked at 10th Avenue Marine Terminal, San Diego Bay (32.6956 °N, -117.15278 °W) using the standard  
525 sphere technique (Demer *et al.*, 2015; Foote *et al.*, 1987). Each WBT was calibrated in both CW (i.e.,  
526 continuous wave or narrowband mode) and FM mode (i.e., frequency modulation or broadband mode). The  
527 reference target was a 38.1-mm diameter sphere made from tungsten carbide (WC) with 6% cobalt binder  
528 material (WC38.1; *Lasker* sphere #1); for FM mode, additional calibrations were conducted for the 120,  
529 200, and 333-kHz echosounders using a 25-mm WC sphere (WC25). Prior to the calibrations, temperature  
530 and salinity were measured to a depth of 10 m using a handheld probe (Pro2030, YSI) to estimate sound  
531 speeds at the transducer and sphere depths, and the time-averaged sound speed and absorption coefficients  
532 for the range between them. The theoretical target strength ( $TS$ ; dB re 1 m<sup>2</sup>) of the sphere was calculated  
533 using values for the sphere, sound-pulse, and seawater properties. The sphere was positioned throughout  
534 the main lobe of each of the transducer beams using three motorized downriggers, two on one side of the  
535 vessel and one on the other. The WBTs were configured using the calibration results via the control software  
536 (Simrad EK80 v21.15.1; Kongsberg; **Table 1**). Calibration results for WBTs in FM mode are presented in  
537 the survey report (Renfree *et al.*, 2023).

Table 1: Wide-Bandwidth Transceiver (Simrad EK80 WBT; Kongsberg) information, pre-calibration settings, and post-calibration beam model results (below the horizontal line). Prior to the survey, on-axis gain ( $G_0$ ), beam angles, angle offsets, and  $S_A$  Correction ( $S_{A\text{corr}}$ ) values from calibration results were entered into the WBT control software (Simrad EK80; Kongsberg).

	Units	Frequency (kHz)					
		18	38	70	120	200	333
Model		ES18	ES38-7	ES70-7C	ES120-7C	ES200-7C	ES333-7C
Serial Number		2106	337	233	783	513	124
Transmit Power ( $p_{\text{et}}$ )	W	1000	2000	600	200	90	35
Pulse Duration ( $\tau$ )	ms	1.024	1.024	1.024	1.024	1.024	1.024
Temperature	C	23.4	23.4	23.4	23.4	23.4	23.4
Salinity	ppt	35.0	35.0	35.0	35.0	35.0	35.0
Sound speed	m s <sup>-1</sup>	1530.5	1530.5	1530.5	1530.5	1530.5	1530.5
On-axis Gain ( $G_0$ )	dB re 1	22.96	26.07	27.24	26.54	26.57	26.40
$S_a$ Correction ( $S_{a\text{corr}}$ )	dB re 1	0.00	-0.37	-0.08	-0.07	-0.05	-0.15
3-dB Beamwidth Along. ( $\alpha_{-3\text{dB}}$ )	deg	10.29	6.55	6.82	6.59	6.46	6.47
3-dB Beamwidth Athw. ( $\beta_{-3\text{dB}}$ )	deg	10.47	6.76	6.73	6.54	6.48	6.46
Angle Offset Along. ( $\alpha_0$ )	deg	-0.01	0.03	-0.00	-0.00	0.01	-0.00
Angle Offset Athw. ( $\beta_0$ )	deg	-0.02	-0.06	-0.03	-0.01	-0.01	0.01
Equivalent Two-way Beam Angle ( $\Psi$ )	dB re 1 sr	-16.90	-20.19	-20.17	-20.09	-20.07	-19.55



538 **2.1.2.2.2 *Lisa Marie*** The 38- and 200-kHz WBTs aboard *Lisa Marie* were calibrated on 15 June 2022,  
 539 using the standard sphere technique, while the vessel was anchored in Grays Harbor near Westport, WA  
 540 (46.9202 N, 124.1090 W). Calibration results for *Lisa Marie* are presented in **Table 2**.

Table 2: Wideband Transceiver (Simrad EK80 WBT; Kongsberg) and transducer information (above horizontal line) and beam model results (below horizontal line) estimated from calibration of the echosounders aboard *Lisa Marie* using a WC38.1 standard sphere.

	Units	Frequency (kHz)	
		38	200
Model		ES38-7	ES200-7C
Serial Number		448	899
Transmit Power ( $p_{et}$ )	W	1000	90
Pulse Duration ( $\tau$ )	ms	1.024	1.024
Temperature	C	13.1	13.1
Salinity	ppt	27.5	27.5
Sound speed	m s <sup>-1</sup>	1491.6	1491.6
On-axis Gain ( $G_0$ )	dB re 1	26.76	26.48
$S_a$ Correction ( $S_{a,corr}$ )	dB re 1	-0.05	-0.05
3-dB Beamwidth Along. ( $\alpha_{-3dB}$ )	deg	6.49	6.71
3-dB Beamwidth Athw. ( $\beta_{-3dB}$ )	deg	6.40	7.25
Angle Offset Along. ( $\alpha_0$ )	deg	-0.03	-0.16
Angle Offset Athw. ( $\beta_0$ )	deg	0.00	0.01
Equivalent Two-way Beam Angle ( $\Psi$ )	dB re 1 sr	-20.35	-20.46

541 **2.1.2.2.3 *Long Beach Carnage*** The 38, 70, 120, and 200 kHz EK60 GPTs aboard *Long Beach Carnage*  
 542 were calibrated on 7 July 2022, using the standard sphere technique, in a tank at the SWFSC (Demer *et al.*,  
 543 2015). Calibration results for *Long Beach Carnage* are presented in **Table 3**.

Table 3: General Purpose Transceiver (Simrad EK60 GPT; Kongsberg) and transducer information (above horizontal line) and beam model results (below horizontal line) estimated from a tank calibration, using a WC38.1 standard sphere, of the echosounders later installed and used aboard *Long Beach Carnage* .

	Units	Frequency (kHz)			
		38	70	120	200
Model		ES38-12	ES70-7C	ES120-7C	ES200-7C
Serial Number		28075	234	813	616
Transmit Power ( $p_{et}$ )	W	1000	600	200	90
Pulse Duration ( $\tau$ )	ms	1.024	1.024	1.024	1.024
Temperature	C	19.1	19.1	19.1	19.1
Salinity	ppt	35.9	35.9	35.9	35.9
Sound speed	m s <sup>-1</sup>	1520.1	1520.1	1520.1	1520.1
On-axis Gain ( $G_0$ )	dB re 1	21.77	26.21	26.03	26.70
$S_a$ Correction ( $S_{a,corr}$ )	dB re 1	-0.71	-0.29	-0.39	-0.24
3-dB Beamwidth Along. ( $\alpha_{-3dB}$ )	deg	12.50	7.08	7.17	6.92
3-dB Beamwidth Athw. ( $\beta_{-3dB}$ )	deg	12.49	7.10	7.33	6.96
Angle Offset Along. ( $\alpha_0$ )	deg	-0.04	0.04	0.11	-0.05
Angle Offset Athw. ( $\beta_0$ )	deg	0.15	0.03	-0.01	0.02
Equivalent Two-way Beam Angle ( $\Psi$ )	dB re 1 sr	-15.64	-20.23	-20.13	-20.06

544 **2.1.2.2.4 USVs** For the two USVs, the echosounders were calibrated by Sairdron, Inc., using the stan-  
 545 dard sphere technique, while dockside. The results, processed and derived by the SWFSC (Renfree *et al.*,  
 546 2019), are presented in **Table 4**.

Table 4: Miniature Wideband Transceiver (Simrad-Kongsberg WBT Mini) beam model results estimated from calibrations of echosounders using a WC38.1 standard sphere, of the echosounders aboard the two USVs.

	Units	Sairdron (Frequency)			
		1076 (38)	1076 (200)	1077 (38)	1077 (200)
Echosounder SN		266961-07	266961-08	268632-07	268632-08
Transducer SN		136	136	131	131
Temperature	C	20.2	20.2	20.1	20.1
Salinity	ppt	31.0	31.0	31.1	31.1
Sound speed	m s <sup>-1</sup>	1517.5	1517.5	1517.5	1517.5
Eq. Two-way Beam Angle ( $\Psi$ )	dB re 1 sr	-12.9	-11.7	-12.4	-11.6
On-axis Gain ( $G_0$ )	dB re 1	19.18	19.45	18.87	19.00
$S_a$ Correction ( $S_{acorr}$ )	dB re 1	0.08	0.08	0.02	0.09
3-dB Beamwidth Along. ( $\alpha_{-3dB}$ )	deg	17.3	19.4	18.2	20.1
3-dB Beamwidth Athw. ( $\beta_{-3dB}$ )	deg	17.0	20.2	18.4	19.9
Angle Offset Along. ( $\alpha_0$ )	deg	0.1	0.5	0.3	0.2
Angle Offset Athw. ( $\beta_0$ )	deg	-0.5	0.2	-0.6	-0.4
RMS	dB	0.27	0.46	0.32	0.41

### 547 2.1.2.3 Data collection

548

549 On *Lasker*, the computer clocks were synchronized with the GPS clock (UTC) using synchronization software  
 550 (NetTime<sup>1</sup>). The 18-kHz WBT, operated by a separate PC from the other echosounders, was programmed  
 551 to track the seabed and output the detected depth to the ship’s Scientific Computing System (SCS). The  
 552 38-, 70-, 120-, 200-, and 333-kHz echosounders were controlled by the EK80 Adaptive Logger (EAL<sup>2</sup>, Renfree  
 553 and Demer, 2016). The EAL optimizes the pulse interval based on the seabed depth, while avoiding aliased  
 554 seabed echoes, and was programmed such that once an hour the echosounders would record three pings in  
 555 passive mode, for obtaining estimates of the background noise level. Acoustic sampling for CPS-density  
 556 estimation along the pre-determined transects was limited to daylight hours (approximately between sunrise  
 557 and sunset).

558 During daytime aboard *Lasker*, measurements of volume backscattering strength ( $S_v$ ; dB re 1 m<sup>2</sup> m<sup>-3</sup>) and  
 559 target strength ( $TS$ ; dB re 1 m<sup>2</sup>), indexed by time and geographic positions provided by GPS receivers, were  
 560 logged to 60 m beyond the detected seabed range or to a maximum range of 500, 500, 500, 300, and 150 m  
 561 for 38, 70, 120, 200, and 333 kHz, respectively, and stored, with a 50-GB maximum file size, in Simrad-EK80  
 562 .raw format. At nighttime, echosounders were set to FM mode to improve target strength estimation and  
 563 species differentiation for CPS near the surface, and logged to 100 m to reduce data volume. For each  
 564 acoustic instrument, the prefix for the file names is a concatenation of the survey name (e.g., 2207RL), the  
 565 operational mode (CW or FM), and the logging commencement date and time from the EK80 software.  
 566 For example, file generated by the EK80 software (v21.15.1) for a WBT operated in CW mode is named  
 567 2207RL-CW-D20220801-T125901.raw.

568 To minimize acoustic interference, transmit pulses from the EK80, ME70, MS70, SX90, EC150-3C, and  
 569 ADCP were triggered using a synchronization system (Simrad K-Sync; Kongsberg). The K-Sync trigger  
 570 rate, and thus echosounder ping interval, was modulated by the EAL (Renfree and Demer, 2016) using the  
 571 seabed depth measured using the 18-kHz echosounder. During daytime, the ME70, MS70, SX90, and ADCP  
 572 were operated continuously, but only recorded at the discretion of the acoustician during times when CPS

<sup>1</sup><http://timesynctool.com>

<sup>2</sup><https://www.fisheries.noaa.gov/west-coast/science-data/ek80-adaptive-logger/>



573 were present. At nighttime, only the EK80 and ADCP were operated. All other instruments that produce  
574 sound within the echosounder bandwidths were secured during daytime survey operations. Exceptions were  
575 made during stations (e.g., plankton sampling and fish trawling) or in shallow water when the vessel's  
576 command occasionally operated the bridge's 50- and 200-kHz echosounders (Furuno), the Doppler velocity  
577 log (SRD-500A; Sperry Marine), or both. Data from the ME70, MS70, and SX90 are not presented in this  
578 report.

579 On *Lisa Marie* and *Long Beach Carnage*, the EAL was used to control the EK80 software to modulate the  
580 echosounder recording ranges and ping intervals to avoid aliased seabed echoes. When the EAL was not  
581 utilized, the EK80 software recorded to 1000 m and used the maximum ping rate. Transmit pulses from the  
582 EK60s and fishing sonars were not synchronized. Therefore, the latter was secured during daytime acoustic  
583 transects.

584 On the USVs, the echosounders were programmed to transmit CW pulses to different ranges, dependent  
585 on the seabed depth. For deeper seabed depths, the ping interval was 2 s and the 38 and 200-kHz  
586 echosounders recorded to 1000 and 400 m, respectively. For shallower depths, the ping interval was 1 s  
587 and both echosounders recorded to 250 m. Once an hour, the echosounders operated in passive mode and  
588 recorded data from three pings to obtain estimates of the background noise levels.

### 589 2.1.3 Oceanographic

#### 590 2.1.3.1 Conductivity and temperature versus depth (CTD)

591 Conductivity and temperature were measured versus depth to 350 m (or to within ~10 m of the shallower  
592 than 350 m) with calibrated sensors on a CTD rosette (Model SBE911+, Seabird) or underway probe [Un-  
593 derwayCTD (UCTD); Oceanscience] cast from the vessel. At least one cast was planned along each acoustic  
594 transect. These data were used to calculate the harmonic mean sound speed (Demer *et al.*, 2015) for esti-  
595 mating ranges to the sound scatterers, and frequency-specific sound absorption coefficients for compensating  
596 signal attenuation of the sound pulse between the transducer and scatterers (Simmonds and MacLennan,  
597 2005) (see **Section 2.2.2**).

#### 598 2.1.3.2 Scientific Computer System

599 While underway, information about the position and direction (e.g., latitude, longitude, speed, course over  
600 ground, and heading), weather (air temperature, humidity, wind speed and direction, and barometric pres-  
601 sure), and sea-surface oceanography (e.g., temperature, salinity, and fluorescence) were measured continu-  
602 ously and logged using *Lasker's* Scientific Computer System (SCS). During and after the survey, data from a  
603 subset of these sensors, logged with a standardized format at 1-min resolution, are available on the internet  
604 via NOAA's ERDDAP data server<sup>3</sup>.

### 605 2.1.4 Fish-eggs

606 On *Lasker*, fish eggs were sampled during the day using a continuous underway fish egg sampler (CUFES,  
607 Checkley *et al.*, 1997), which collects water and plankton at a rate of ~640 l min<sup>-1</sup> from an intake at ~3-m  
608 depth on the hull of the ship. The particles in the sampled water were sieved by a 505- $\mu$ m mesh. Pacific  
609 Sardine, Northern Anchovy, Jack Mackerel, and Pacific Hake (*Merluccius productus*) eggs were identified to  
610 species, counted, and logged. Eggs from other species (e.g., Pacific Mackerel and flatfishes) were also counted  
611 and logged as "other fish eggs." Typically, the duration of each CUFES sample was 30 min, corresponding  
612 to a distance of 5 nmi at a speed of 10 kn. Because the durations of the early egg stages are short for most  
613 fish species, the egg distributions inferred from CUFES indicated the nearby presence of actively spawning  
614 fish, and were used in combination with CPS echoes to select trawl locations.

<sup>3</sup><http://coastwatch.pfeg.noaa.gov/erddap/tabledap/fsuNoaaShipWTEG.html>

## 615 2.1.5 Species and Demographics

616 The net catches provide information about species composition, lengths, weights and ages of CPS sampled  
617 acoustically during the day. Nighttime trawls were conducted to sample the fish dispersed near the sea  
618 surface, because after sunset, schools of CPS and other fish tend to ascend and disperse and are less likely to  
619 avoid a trawl net (Mais, 1977). Daytime purse-seine nets were set nearshore to sample CPS schools where  
620 their depth is constrained by the seabed, and their vision is obscured by non-transparent, light-scattering  
621 water, due to primary production and suspended particulates. In summer 2022, *Lisa Marie* used daytime  
622 purse-seine sets to sample CPS schools nearshore, but also offshore in deeper, clearer water, and with mixed  
623 success (see **Section 3.5.1**). For example, fast swimming Jack Mackerel often avoided capture, resulting in  
624 unquantified species selectivity.

### 625 2.1.5.1 Trawl gear

626 **2.1.5.1.1 *Lasker*** A Nordic 264 rope trawl (NET Systems, Bainbridge Island, WA; **Figs. 5a,b**), was  
627 towed at the surface for 45 min at a speed of 3.5-4.5 kn. The net has a rectangular opening with an area of  
628 approximately 300 m<sup>2</sup> (~15-m tall x 20-m wide), a throat with variable-sized mesh and a “marine mammal  
629 excluder device” to prevent the capture of large animals, such as dolphins, turtles, or sharks while retaining  
630 target species (Dotson *et al.*, 2010), and an 8-mm square-mesh cod-end liner (to retain a large range of  
631 animal sizes). The trawl doors were foam-filled and the trawl headrope was lined with floats so the trawl  
632 towed at the surface. Temperature-depth recorders (TDRs; RBRduet<sup>3</sup> T.D., RBR) were attached to the  
633 kite and footrope to evaluate trawl performance (**Fig. 6**).

### 634 2.1.5.2 Purse-seine gear

635 **2.1.5.2.1 *Lisa Marie* and *Long Beach Carnage*** *Lisa Marie* used an approximately 440-m-long and  
636 40-m-deep net with 17-mm-wide mesh (A. Blair, pers. comm.). *Long Beach Carnage* used an approximately  
637 200-m-long and 27-m-deep net with 17-mm-wide mesh; a small section on the back end of the net had 25-  
638 mm-wide mesh (R. Ashley, pers. comm.). Specimens collected by *Lisa Marie* and *Long Beach Carnage* were  
639 processed aboard the vessel by the WA Department of Fish and Wildlife (WDFW) and ashore by the CA  
640 Department of Fish and Wildlife (CDFW), respectively.





Figure 5: Schematic drawings of the Nordic 264 rope trawl a) net and b) cod-end.




Figure 6: Example depths (m) of the trawl headrope (red line) and footrope (blue line) measured using temperature-depth recorders (TDRs) during the net deployment (dashed box) and when actively fishing (shaded region). The vessel speed over ground (kn, black line) was measured using the ship's GPS.

DRAFT

### 641 2.1.5.3 Sampling locations

642 **2.1.5.3.1 *Lasker*** Up to three nighttime (i.e., 30 min after sunset to 30 min before sunrise) surface  
643 trawls, typically spaced at least 10-nmi apart, were conducted in areas where echoes from putative CPS  
644 schools were observed earlier that day. Trawl locations were selected using one or more of the following  
645 criteria, in descending priority: CPS schools in echograms that day; CPS eggs in CUFES that day; and the  
646 trawl locations and catches during the previous night. Each evening, trawl locations were selected by an  
647 acoustician, who monitored CPS echoes, and a biologist, who measured the densities of CPS eggs in the  
648 CUFES. The locations were provided to the watch officers who charted the proposed trawl sites.

649 If no CPS echoes or CPS eggs were observed along a transect that day, the trawls were alternately placed  
650 nearshore one night and offshore the next night, with consideration given to the seabed depth and the  
651 modeled distribution of CPS habitat. Each morning, after the last trawl or 30 min prior to sunrise, *Lasker*  
652 resumed sampling at the location where the acoustic sampling stopped the previous day.

653 **2.1.5.3.2 *Lisa Marie* and *Long Beach Carnage*** On *Lisa Marie*, as many as three purse-seine sets  
654 were conducted each day. For each set, three dip-net samples, were collected, spatially separated as much  
655 as possible.

656 On *Long Beach Carnage*, as many as three purse-seine sets were conducted each day, including evenings.  
657 For each set, three dip-net samples were collected, spatially separated as much as possible.

### 658 2.1.5.4 Sample processing

659 **2.1.5.4.1 *Lasker*** If the total volume of the trawl catch was five 35-l baskets (~175 l) or less, all target  
660 species were separated from the catch, sorted by species, weighed, and enumerated. If the volume of the  
661 entire catch was more than five baskets, a five-basket random subsample that included non-target species  
662 was collected, sorted by species, weighed, and enumerated; the remainder of the total catch was weighed.  
663 In these cases, the weight of the entire catch was calculated as the sum of the subsample and remainder  
664 weights. The weight of the  $e$ -th species in the total catch ( $C_{T,e}$ ) was obtained by summing the catch weight  
665 of the respective species in the subsample ( $C_{S,e}$ ) and the corresponding catch in the remainder ( $C_{R,e}$ ), which  
666 was calculated as:

$$C_{R,e} = C_R * P_{w,e}, \quad (1)$$

667 where  $P_{w,e} = C_{S,e} / \sum_1^s C_{S,e}$ , is the proportion in weight of the  $e$ -th species in the subsample. The number  
668 of specimens of the  $e$ -th species in the total catch ( $N_{T,e}$ ) was estimated by:

$$N_{T,e} = \frac{C_{T,e}}{\bar{w}_e}, \quad (2)$$

669 where  $\bar{w}_e$  is the mean weight of the  $e$ -th species in the subsample. For Pacific Sardine and Northern Anchovy  
670 with 75 specimens or less, individual measurements of standard length ( $L_S$ ) in mm and weight ( $w$ ) in g were  
671 recorded. For Jack Mackerel, Pacific Mackerel, and Pacific Herring with 50 specimens or less, individual  
672 measurements of fork length ( $L_F$ ) and  $w$  were recorded. In addition, sex and maturity were recorded for up  
673 to 75 Pacific Sardine and Northern Anchovy and up to 25 Jack and Pacific Mackerel. Ovaries were preserved  
674 for up to 10 specimens of each CPS species except Pacific Herring. Fin clips were removed from 50 Pacific  
675 Sardine and Northern Anchovy specimens from seven geographic zones (with boundaries at the Columbia  
676 River, Cape Mendocino, San Francisco Bay, Point Conception, San Diego, and San Quentin, Baja CA) and  
677 preserved in ethanol for genetic analysis. Otoliths were removed from all 50 Pacific Sardine in the subsample;  
678 for other CPS species except Pacific Herring, 25 otoliths were removed as equally as possible from the range  
679 of sizes present. The combined catches in up to three trawls per night (i.e., trawl cluster) were used to  
680 estimate the proportions of species contributing to the nearest samples of acoustic backscatter.

681 **2.1.5.4.2 Lisa Marie** For each dip-net sample, all specimens were sorted, weighed, and counted to  
 682 provide a combined weight and count for each. Next, all three dip net samples were combined and up to 50  
 683 specimens of each CPS species were randomly sampled to provide a combined weight for each set. Length  
 684 (mm),  $L_S$  for Pacific Sardine and Northern Anchovy and  $L_F$  for all others, and weight (g) were measured  
 685 for up to 50 randomly selected specimens of each species. Otoliths were extracted, macroscopic maturity  
 686 stage was determined visually, and gonads were collected and preserved from female specimens.

687 **2.1.5.4.3 Long Beach Carnage** For each dip net sample, all specimens were sorted, weighed, and  
 688 counted to provide a combined weight and count for each, and as many as 20 fish of each CPS species were  
 689 chosen randomly throughout the sample, and combined for a random sample of 50 fish collected throughout  
 690 the catch. The fish were frozen for later analysis by CDFW biologists, yielding measures of individual fish  
 691 and total sample weights (g); length (mm),  $L_S$  for Pacific Sardine and Northern Anchovy and  $L_F$  for all  
 692 others; maturity; and otolith-derived ages. No female gonad samples were analyzed.

693 **2.1.5.5 Quality Assurance and Quality Control** At sea, trawl data were entered into a database  
 694 (Microsoft Access). During and following the survey, data were further scrutinized and verified, or corrected.  
 695 Missing length ( $L_{miss}$ ) and weight ( $W_{miss}$ ) measurements were estimated as  $W_{miss} = \beta_0 L^{\beta_1}$  and  $L_{miss} =$   
 696  $(W/\beta_0)^{1/\beta_1}$ , respectively, where values for  $\beta_0$  and  $\beta_1$  are species- and season-specific parameters of the  
 697 length-versus-weight relationships described in Palance et al. (Palance et al., 2019). To identify measurement  
 698 or data-entry errors, length and weight data were graphically compared (Fig. 7) to measurements from  
 699 previous surveys and models of season-specific length-versus-weight from previous surveys (Palance et al.,  
 700 2019). Outliers were flagged, reviewed by the trawl team, and mitigated. Catch data were removed from  
 701 aborted trawl hauls, or hauls otherwise deemed unacceptable.




Figure 7: Specimen length versus weight from the current survey (colored points, by sex) compared to those from previous SWFSC surveys during the same season (gray points, all sexes) and models [dashed lines; Palance et al. (2019)].

## 2.2 Data processing

### 2.2.1 Acoustic and oceanographic data

The calibrated echosounder data from each transect were processed using commercial software (Echoview v12.1; Echoview Software Pty Ltd.) and estimates of the sound speed and absorption coefficient calculated with contemporaneous data from CTD probes cast while stationary or underway (UCTD, see Section 2.1.3.1). Data collected along the daytime transects at speeds  $\geq 5$  kn were used to estimate CPS densities. Nighttime acoustic data were not used for biomass estimations because they are assumed to be negatively biased due to diel-vertical migration and disaggregation of the target species' schools (Cutter and Demer, 2008).

### 2.2.2 Sound speed and absorption calculation

Depth derived from pressure in CTD casts was used to average samples in 1-m depth bins. Sound speed in each bin ( $c_{w,i}$ , m s<sup>-1</sup>) was estimated from the average salinity, density, and pH [if measured, else pH = 8; Chen and Millero (1977); Seabird (2013)]. The harmonic sound speed in the water column ( $\bar{c}_w$ , m s<sup>-1</sup>) was calculated over the upper 70 m as:

$$\bar{c}_w = \frac{\sum_{i=1}^N \Delta r_i}{\sum_{i=1}^N \Delta r_i / c_{w,i}}, \quad (3)$$

where  $\Delta r$  is the depth of increment  $i$  (Seabird, 2013). Measurements of seawater temperature ( $t_w$ , °C), salinity ( $s_w$ , psu), depth, pH, and  $\bar{c}_w$  are also used to calculate the mean species-specific absorption coefficients ( $\bar{\alpha}_a$ , dB m<sup>-1</sup>) over the entire profile using equations in Francois and Garrison (1982), Ainslie and McColm (1998), and Doonan et al. (2003). Both  $\bar{c}_w$  and  $\bar{\alpha}_a$  are later used to estimate ranges to the sound scatterers to compensate the echo signal for spherical spreading and attenuation during propagation of the sound pulse from the transducer to the scatterer range and back (Simmonds and MacLennan, 2005). The CTD rosette, when cast, also provides measures of fluorescence and dissolved oxygen concentration versus depth, which may be used to estimate the vertical dimension of Pacific Sardine potential habitat (Zwolinski et al., 2011), particularly the depth of the upper-mixed layer where most epipelagic CPS reside. The latter information is used to inform echo classification (see Section 2.2.3).

### 2.2.3 Echo classification

Echoes from schooling CPS and plankton (Figs. 8a,d) were identified using a semi-automated data processing algorithm implemented using Echoview software (v12.1; Echoview Software Pty Ltd). The filters and thresholds were based on a subsample of echoes from randomly selected CPS schools. The aim of the filter criteria is to retain at least 95% of the noise-free backscatter from CPS while rejecting at least 95% of the non-CPS backscatter (Fig. 8). Data from *Lasker* and *Long Beach Carnage* were processed using the following steps:

1. Match geometry of all  $S_v$  variables to the 38-kHz  $S_v$ ;
2. Remove passive-mode pings;
3. Estimate and subtract background noise using the background noise removal function (De Robertis and Higginbottom, 2007) in Echoview (Figs. 8b,e);
4. Average the noise-free  $S_v$  echograms using non-overlapping 11-sample by 3-ping bins;
5. Expand the averaged, noise-reduced  $S_v$  echograms with a 7 pixel x 7 pixel dilation;
6. For each pixel, compute:  $S_{v,200\text{kHz}} - S_{v,38\text{kHz}}$ ,  $S_{v,120\text{kHz}} - S_{v,38\text{kHz}}$ , and  $S_{v,70\text{kHz}} - S_{v,38\text{kHz}}$ ;
7. Create a Boolean echogram for  $S_v$  differences in the CPS range:  $-13.85 < S_{v,70\text{kHz}} - S_{v,38\text{kHz}} < 9.89$  and  $-13.5 < S_{v,120\text{kHz}} - S_{v,38\text{kHz}} < 9.37$  and  $-13.51 < S_{v,200\text{kHz}} - S_{v,38\text{kHz}} < 12.53$ ;



- 742 8. Compute the 120- and 200-kHz Variance-to-Mean Ratios ( $VMR_{120\text{kHz}}$  and  $VMR_{200\text{kHz}}$ , respectively,  
743 Demer *et al.*, 2009a) using the difference between noise-filtered  $S_v$  (Step 3) and averaged  $S_v$  (Step 4);  
744 9. Expand the  $VMR_{120\text{kHz}}$  and  $VMR_{200\text{kHz}}$  echograms with a 7 pixel x 7 pixel dilation;  
745 10. Create a Boolean echogram based on the  $VMR$ s in the CPS range:  $VMR_{120\text{kHz}} > -65$  dB and  
746  $VMR_{200\text{kHz}} > -65$  dB. Diffuse backscattering layers have low  $VMR$  (Zwolinski *et al.*, 2010) whereas  
747 fish schools have high  $VMR$  (Demer *et al.*, 2009a);  
748 11. Intersect the two Boolean echograms to create an echogram with “TRUE” samples for candidate CPS  
749 schools and “FALSE” elsewhere;  
750 12. Mask the noise-reduced echograms using the CPS Boolean echogram (**Figs. 8c,f**);  
751 13. Create an integration-start line 5 m below the transducer (~10 m depth);  
752 14. Create an integration-stop line 3 m above the estimated seabed (Demer *et al.*, 2009a), or to the  
753 maximum logging range (e.g., 350 m), whichever is shallowest;  
754 15. Set the minimum  $S_v$  threshold to -60 dB (corresponding to a density of approximately three 20-cm-long  
755 Pacific Sardine per 100 m<sup>3</sup>);  
756 16. Integrate the volume backscattering coefficients ( $s_v$ , m<sup>2</sup> m<sup>-3</sup>) attributed to CPS over 5-m depths and  
757 averaged over 100-m distances;  
758 17. Output the resulting nautical area scattering coefficients ( $s_A$ ; m<sup>2</sup> nmi<sup>-2</sup>) and associated information  
759 from each transect and frequency to comma-delimited text (.csv) files.

760 Data from *Lisa Marie* and the USVs were processed using the following steps:

- 761 1. Match geometry of the  $S_{v,200\text{kHz}}$  to the  $S_{v,38\text{kHz}}$ ;  
762 2. Remove passive-mode pings;  
763 3. Perform Steps 3-5 from *Lasker* processing;  
764 4. For each pixel, compute:  $S_{v,200\text{kHz}} - S_{v,38\text{kHz}}$ ;  
765 5. Create a Boolean echogram for  $S_v$  differences in the CPS range:  $-13.5 < S_{v,200\text{kHz}} - S_{v,38\text{kHz}} < 9.37$   
766 6. Perform Steps 8-9 from *Lasker* processing;  
767 7. Create a Boolean echogram mask using  $VMR > -57$  dB;  
768 8. Performs Steps 11-17 from *Lasker* processing.

769 When necessary, the start and stop integration lines were manually edited to exclude reverberation due to  
770 bubbles, to include the entirety of shallow CPS aggregations, or to exclude seabed echoes. Also, echoes  
771 from putative rockfish schools were excluded based on their aggregation shapes and proximity to the rocky  
772 seabed.

#### 773 2.2.4 Removal of non-CPS backscatter

774 In addition to echoes from target CPS, echoes may also be present from other pelagic fish species (Pacific  
775 Saury, *Cololabis saira*), or semi-demersal fish such as Pacific Hake and rockfishes (*Sebastes* spp.). When  
776 analyzing the acoustic-survey data, it was therefore necessary to filter “acoustic by-catch,” i.e., backscatter  
777 not from the target species. To exclude echoes from mid-water, demersal, and benthic fishes, echograms were  
778 visually examined to exclude fish echoes where the seabed was hard and rugose, or where diffuse schools are  
779 observed offshore either near the surface or deeper than ~250 m (**Fig. 9**). In areas dominated by Pacific  
780 Herring, for example off Vancouver Island, backscatter was integrated to a maximum depth of 75 m.






Figure 8: Two examples of echograms depicting CPS schools (red) and plankton aggregations (blue and green) at 38 kHz (top) and 120 kHz (bottom). Example data processing steps include the original echogram (a, d), after noise subtraction and bin-averaging (b, e), and after filtering to retain only putative CPS echoes (c, f).




Figure 9: Echoes from fishes with swimbladders (blue points, scaled by backscatter intensity) along an example acoustic transect (top) and the corresponding echogram image (bottom). In this example, the upper (blue) and lower lines (green) indicate boundaries within which echoes were retained. When the lower boundary is deeper than the seabed (black line), echoes above the seabed are retained. Echoes from deep, bottom-dwelling schools of non-CPS fishes with swimbladders, and from diffuse scatterers near the surface were excluded. The proximity of the echoes to the seabed was also used to define the lower limit for vertical integration.

781 **2.2.5 Extraction of nearshore backscatter**

782 In summer 2022, between Cape Flattery and Bodega Bay, *Lisa Marie* sampled along acoustic transects to a  
 783 depth of ~5 m. Because there was not a separate nearshore survey in this area, as there was between 2016 and  
 784 2021, acoustic intervals in water shallower than the 20-m isobath were assigned to nearshore strata, else to  
 785 core strata, and biomasses were estimated as usual. The 20-m isobath roughly corresponds to the shallowest  
 786 depth in which *Lasker* can safely navigate, so this nearshore area roughly approximates those from previous  
 787 surveys. However, a study of inter-annual variation in nearshore biomasses should use a standardized area  
 788 across years.

789 **2.2.6 Quality Assurance and Quality Control**

790 The largest 38-kHz integrated backscattering coefficients ( $s_A$ ,  $m^2 \text{ nmi}^{-2}$ ) were graphically examined to iden-  
 791 tify potential errors in the integrated data (e.g., when a portion of the seabed was accidentally integrated,  
 792 not shown). If found, errors were corrected and data were re-integrated prior to use for biomass estimation.

793 **2.2.7 Echo integral partitioning and acoustic inversion**

794 For fishes with swimbladders, the acoustic backscattering cross-section of an individual ( $\sigma_{bs}$ ,  $m^2$ ) depends  
 795 on many factors but mostly on the acoustic wavelength and the swimbladder size and orientation relative  
 796 to the incident sound pulse. For echosounder sampling conducted in this survey,  $\sigma_{bs}$  is a function of the  
 797 dorsal-surface area of the swimbladder and was approximated by a function of fish length ( $L$ ), i.e.:

$$\sigma_{bs} = 10^{\frac{m \log_{10}(L)+b}{10}}, \quad (4)$$

798 where  $m$  and  $b$  are frequency and species-specific parameters that are obtained theoretically or experimentally  
 799 (see references below).  $TS$ , a logarithmic representation of  $\sigma_{bs}$ , is defined as:

$$TS = 10 \log_{10}(\sigma_{bs}) = m \log_{10}(L) + b. \quad (5)$$

800  $TS$  has units of dB re  $1 \text{ m}^2$  if defined for an individual, or dB re  $1 \text{ m}^2 \text{ kg}^{-1}$  if defined by weight. The following  
 801 equations for  $TS_{38\text{kHz}}$ , were used in this analysis:

$$TS_{38\text{kHz}} = -14.90 \times \log_{10}(L_T) - 13.21, \text{ for Pacific Sardine;} \quad (6)$$

$$TS_{38\text{kHz}} = -11.97 \times \log_{10}(L_T) - 11.58561, \text{ for Pacific and Round Herrings;} \quad (7)$$

$$TS_{38\text{kHz}} = -13.87 \times \log_{10}(L_T) - 11.797, \text{ for Northern Anchovy; and} \quad (8)$$

$$TS_{38\text{kHz}} = -15.44 \times \log_{10}(L_T) - 7.75, \text{ for Pacific and Jack Mackerels,} \quad (9)$$

802 where the units for total length ( $L_T$ ) is cm and  $TS$  is dB re  $1 \text{ m}^2 \text{ kg}^{-1}$ .

803 Equations (6) and (9) were derived from echosounder measurements of  $\sigma_{bs}$  for in situ fish and measures of  
 804  $L_T$  and  $W$  from concomitant catches of South American Pilchard (*Sardinops ocellatus*) and Horse Mackerel  
 805 (*Trachurus trachurus*) off South Africa (Barange *et al.*, 1996). Because mackerels have similar  $TS$  (Peña,  
 806 2008), Equation (9) is used for both Pacific and Jack Mackerels. For Pacific Herring and Round Herring,  
 807 Equation (7) was derived from that of Thomas *et al.* (2002) measured at 120 kHz with the following  
 808 modifications: 1) the intercept used here was calculated as the average intercept of Thomas *et al.*'s spring

809 and fall regressions; 2) the intercept was compensated for swimbladder compression after Zhao *et al.* (2008)  
810 using the average depth for Pacific Herring of 44 m; 3) the intercept was increased by 2.98 dB to account  
811 for the change of frequency from 120 to 38 kHz (Saunders *et al.*, 2012). For Northern Anchovy, Equation  
812 (8) was derived from that of Kang *et al.* (2009), after compensation of the swimbladder volume (Ona, 2003;  
813 Zhao *et al.*, 2008) for the average depth of Northern Anchovy observed in summer 2016 (19 m, Zwolinski *et*  
814 *al.*, 2017).

815 To calculate  $TS_{38\text{kHz}}$ ,  $L_T$  was estimated from measurements of  $L_S$  or  $L_F$  using linear relationships between  
816 length and weight derived from specimens collected in the CCE (Palance *et al.*, 2019): for Pacific Sardine,  
817  $L_T = 0.3574 + 1.149L_S$ ; for Northern Anchovy,  $L_T = 0.2056 + 1.1646L_S$ ; for Pacific Mackerel,  $L_T =$   
818  $0.2994 + 1.092L_F$ ; for Jack Mackerel  $L_T = 0.7295 + 1.078L_F$ ; and for Pacific Herring  $L_T = -0.105 + 1.2L_F$ .  
819 Since a conversion does not exist for Round Herring, the equation for Pacific Herring was used to estimate  
820  $L_T$ .

821 The proportions of species in a trawl cluster were considered representative of the proportions of species in  
822 the vicinity of the cluster. Therefore, the proportion of the echo-integral from the  $e$ -th species ( $P_e$ ) in an  
823 ensemble of  $s$  species can be calculated from the species catches  $N_1, N_2, \dots, N_s$  and the respective average  
824 backscattering cross-sections  $\sigma_{bs1}, \sigma_{bs2}, \dots, \sigma_{bss}$  (Nakken and Dommasnes, 1975). The acoustic proportion for  
825 the  $e$ -th species in the  $a$ -th trawl ( $P_{ae}$ ) is:

$$P_{ae} = \frac{N_{ae} \times \bar{w}_{ae} \times \bar{\sigma}_{bs,ae}}{\sum_{e=1}^{s_a} (N_{ae} \times \bar{w}_{ae} \times \bar{\sigma}_{bs,ae})}, \quad (10)$$

826 where  $\bar{\sigma}_{bs,ae}$  is the arithmetic counterpart of the average target strength ( $\overline{TS}_{ae}$ ) for all  $n_{ae}$  individuals of  
827 species  $e$  in the random sample of trawl  $a$ :

$$\bar{\sigma}_{bs,ae} = \frac{\sum_{i=1}^{n_{ae}} 10^{(TS_i/10)}}{n_{ae}}, \quad (11)$$

828 and  $\bar{w}_{ae}$  is the average weight:  $\bar{w}_{ae} = \sum_{i=1}^{n_{ae}} w_{aei}/n_{ae}$ . The total number of individuals of species  $e$  in a  
829 trawl  $a$  ( $N_{ae}$ ) is obtained by:  $N_{ae} = \frac{n_{ae}}{w_{s,ae}} \times w_{t,ae}$ , where  $w_{s,ae}$  is the weight of the  $n_{ae}$  individuals sampled  
830 randomly, and  $w_{t,ae}$  is the total weight of the respective species' catch.

831 The trawls within a cluster were combined to reduce sampling variability (see **Section 2.2.8**), and the  
832 number of individuals caught from the  $e$ -th species in a cluster  $g$  ( $N_{ge}$ ) was obtained by summing the  
833 catches across the  $h$  trawls in the cluster:  $N_{ge} = \sum_{a=1}^{h_g} N_{ae}$ . The backscattering cross-section for species  $e$   
834 in the  $g$ -th cluster with  $a$  trawls is then given by:

$$\bar{\sigma}_{bs,ge} = \frac{\sum_{a=1}^{h_g} N_{ae} \times \bar{w}_{ae} \times \bar{\sigma}_{bs,ae}}{\sum_{a=1}^{h_g} N_{ae} \times \bar{w}_{ae}}, \quad (12)$$

835 where:

$$\bar{w}_{ge} = \frac{\sum_{a=1}^{h_g} N_{ae} \times \bar{w}_{ae}}{\sum_{a=1}^{h_g} N_{ae}}, \quad (13)$$

836 and the proportion ( $P_{ge}$ ) is;

$$P_{ge} = \frac{N_{ge} \times \bar{w}_{ge} \times \bar{\sigma}_{bs,ge}}{\sum_{e=1}^s (N_{ge} \times \bar{w}_{ge} \times \bar{\sigma}_{bs,ge})}. \quad (14)$$

### 837 2.2.8 Trawl clustering and species proportion

838 Trawls that occurred on the same night were assigned to a trawl cluster. Biomass densities ( $\rho$ ) were calculated  
839 for 100-m transect intervals by dividing the integrated area-backscatter coefficients for each CPS species by  
840 the mean backscattering cross-sectional area (MacLennan *et al.*, 2002) estimated in the trawl cluster nearest  
841 in space. Survey data were post-stratified to account for spatial heterogeneity in sampling effort and biomass  
842 density in a similar way to that performed for Pacific Sardine (Zwolinski *et al.*, 2016).

843 For a generic 100-m long acoustic interval, the area-backscattering coefficient for species  $e$ :  $s_{A,e} = s_{A,cps} \times P_{ge}$ ,  
844 where  $P_{ge}$  is the species acoustic proportion of the nearest trawl cluster (Equation (14)), was used to estimate  
845 the biomass density ( $\rho_{w,e}$ ) (MacLennan *et al.*, 2002; Simmonds and MacLennan, 2005) for every 100-m  
846 interval, using the size and species composition of the nearest (space and time) trawl cluster (**Fig. 10**):

$$\rho_{w,e} = \frac{s_{A,e}}{4\pi\bar{\sigma}_{bs,e}}. \quad (15)$$

847 See **Section 3.5.1** and **Fig. 13** for a description of the method used for this survey to estimate species  
848 proportions and Jack Mackerel lengths in the area between Cape Flattery and Cape Mendocino. The biomass  
849 densities were converted to numerical densities using:  $\rho_{n,e} = \rho_{w,e}/\bar{w}_e$ , where  $\bar{w}_e$  is the corresponding mean  
850 weight. Also, for each acoustic interval, the biomass or numeric densities are partitioned into length classes  
851 according to the species' length distribution in the respective trawl cluster.




Figure 10: a) Polygons enclosing 100-m acoustic intervals from *Lasker*, *Lisa Marie*, and USVs assigned to catches from each trawl cluster or purse-seine set, and b) the acoustic proportions of CPS in catches from trawl clusters or purse-seine sets. See **Section 3.5.1** and **Fig. 13** for a description of the method used for this survey to estimate species proportions and Jack Mackerel lengths in the area between Cape Flattery and Cape Mendocino. The numbers inside each polygon in panel a) are the cluster or purse-seine numbers, which are located at the average latitude and longitude of all trawls in that cluster or each individual purse-seine set. Black points in panel b) indicate trawl clusters or purse-seine sets with no CPS present in the catch.

## 852 2.3 Data analysis

### 853 2.3.1 Post-stratification

854 The transects were sampling units (Simmonds and Fryer, 1996). Because each species does not generally  
855 span the entire survey area (Demer and Zwolinski, 2017; Zwolinski *et al.*, 2014), the sampling domain was  
856 stratified for each species and stock. Strata were defined by uniform transect spacing (sampling intensity)  
857 and either presences (positive densities and potentially structural zeros) or absences (real zeros) of species  
858 biomass. Each stratum has: 1) at least three transects, with approximately equal spacing, 2) fewer than  
859 three consecutive transects with zero-biomass density, and 3) bounding transects with zero-biomass density  
860 (**Fig. 11**). This approach tracks stock patchiness and creates statistically-independent, stationary, post-  
861 sampling strata (Johannesson and Mitson, 1983; Simmonds *et al.*, 1992). For Northern Anchovy, we define  
862 the separation between the northern and central stock at Cape Mendocino (40.5 °N). For Pacific Sardine,  
863 the northern and southern stocks present in the survey area (Felix-Uraga *et al.*, 2004; Felix-Uraga *et al.*,  
864 2005; Garcia-Morales *et al.*, 2012; Hill *et al.*, 2014) were separated using the Pacific Sardine potential habitat  
865 during the survey (**Fig. 12**). This separation is further supported by different distributions of  $L_S$  and a  
866 break in the distribution of Pacific Sardine biomass, which, in this survey, coincided geographically with Big  
867 Sur, CA (36.2 °N, **Fig. 11**).




Figure 11: Biomass density ( $\log_{10}(t + 1) \text{ nmi}^{-2}$ ) versus latitude (easternmost portion of each transect) and strata used to estimate biomass and abundance (shaded regions; outline indicates stratum number) for each species in the core survey region. Data labels (blue numbers) correspond to transects with positive biomass ( $\log_{10}(t + 1) > 0.01$ ). Transect spacing (nmi; point color), and stock breaks for Northern Anchovy and Pacific Sardine (red dashed lines and text) are indicated.






Figure 12: Summary of all transects sampled throughout the survey by *Lasker* (red) and *Lisa Marie* (yellow) in relation to the probability thresholds of the updated model of potential habitat for the northern stock of Pacific Sardine (Zwolinski and Demer, In prep.). The shaded regions with probabilities higher than 0.29 and 0.18 contained 82% and 95% of all spring northern stock Pacific Sardine spawning stock biomass, respectively. The habitat-model output is averaged in areas,  $\pm 2^\circ$  latitude and longitude, centered around the daytime location of each vessel throughout the survey. Areas without data (white), occurred in the analysis domain where and when clouds prevented satellite-sensed observations.



### 868 2.3.2 Biomass and sampling precision estimation

869 For each stratum and stock, the biomass ( $\hat{B}$ ; kg) of each species was estimated by:

$$\hat{B} = A \times \hat{D}, \quad (16)$$

870 where  $A$  is the stratum area ( $\text{nmi}^2$ ) and  $\hat{D}$  is the estimated mean biomass density ( $\text{kg nmi}^{-2}$ ):

$$\hat{D} = \frac{\sum_{l=1}^k \bar{\rho}_{w,l} c_l}{\sum_{l=1}^k c_l}, \quad (17)$$

871 where  $\bar{\rho}_{w,l}$  is the mean biomass density of the species on transect  $l$ ,  $c_l$  is the transect length, and  $k$  is the  
872 total number of transects. The variance of  $\hat{B}$  is a function of the variability of the transect-mean densities  
873 and associated lengths. Treating transects as replicate samples of the underlying population (Simmonds and  
874 Fryer, 1996), the variance was calculated using bootstrap resampling (Efron, 1981) based on transects as  
875 sampling units. Provided that each stratum has independent and identically-distributed transect means (i.e.,  
876 densities on nearby transects are not correlated, and they share the same statistical distribution), bootstrap  
877 or other random-sampling estimators provide unbiased estimates of variance.

878 The 95% confidence intervals ( $\text{CI}_{95\%}$ ) for the mean biomass densities ( $\hat{D}$ ) were estimated as the 0.025 and  
879 0.975 percentiles of the distribution of 1000 bootstrap survey-mean biomass densities. Coefficient of variation  
880 (CV, %) values were obtained by dividing the bootstrapped standard error by the mean estimate (Efron,  
881 1981). Total biomass in the survey area was estimated as the sum of the biomasses in each stratum, and the  
882 associated sampling variance was calculated as the sum of the variances across strata.

### 883 2.3.3 Abundance- and biomass-at-length estimation

884 The numerical densities by length class (**Section 2.2.8**) were averaged for each stratum in a similar way for  
885 that used for biomass (Equation (17)), and multiplied by the stratum area to obtain abundance per length  
886 class.

### 887 2.3.4 Percent biomass per cluster contribution

888 The percent contribution of each cluster to the estimated abundance in a stratum (**Appendix A**) was  
889 calculated as:

$$\frac{\sum_{i=1}^l \bar{\rho}_{ci}}{\sum_{c=1}^C \sum_{i=1}^l \bar{\rho}_{ci}}, \quad (18)$$

890 where  $\bar{\rho}_{ci}$  is the numerical density in interval  $i$  represented by the nearest trawl cluster  $c$ .

## 3 Results

### 3.1 Sampling effort and allocation

The summer 2022 survey spanned the area from Cape Flattery and El Rosario between 27 June and 30 September 2022, and included most of the potential habitat for the northern stock of Pacific Sardine at the time of the survey<sup>4</sup>. In the core survey region that spanned this area (**Fig. 15**), *Lasker* (40 days at sea, DAS), *Lisa Marie* (26 DAS), and the two USVs (108 mission days) sampled 96 east-west transects totaling 4,264 nmi. Catches from a total of 86 nighttime surface trawls and 41 purse-seine sets from *Lisa Marie* were combined into 73 trawl clusters. In the core area, one to six post-survey strata were defined by their transect spacing and the densities of echoes attributed to each species.

The nearshore region spanned an area from approximately Cape Flattery to San Diego, including around Santa Cruz and Santa Catalina Islands. *Lisa Marie* (26 DAS) surveyed from approximately Cape Flattery, WA to Stewarts Point, CA with 24 east-west transects totaling **18 nmi** and 41 purse-seine sets. *Long Beach Carnage* (22 DAS) surveyed from approximately Stewarts Point to San Diego, and around the Santa Cruz and Santa Catalina Islands, with 129 east-west transects totaling 511 nmi and 53 purse-seine sets (**Fig. 17**). In the nearshore area, one to fourteen post-survey strata were defined by their transect spacing and the biomass densities.

Biomasses and abundances were estimated for each species in both the core and nearshore survey areas. The total biomass for each stock within the survey region was estimated as the sum of its biomasses in the core and nearshore areas.

#### Leg I

Leg I on *Lasker* was canceled. Therefore, *Lisa Marie* was directed to sample the 20-nmi-spaced compulsory Transects, 178 to 143, between Cape Flattery and Port Orford, OR, but extending them shoreward to ~5 m depth. Two USVs (SD-1076 and SD-1077) sampled Transects 170 to 160, between Copalis Beach, WA to Tillamook Bay, OR, from 9 to 22 July.

#### Leg II

On 21 July, *Lasker* departed from the 10th Avenue Marine Terminal in San Diego, CA at ~1745 (all times GMT). Prior to the transit north, a calibration of the Simrad-Kongsberg EC150-3C ADCP-echosounder was attempted northwest of the sea buoy outside San Diego Bay (32.6598 N, 117.3833 W), but was not completed, due to GPS data-format incompatibilities. Throughout the northward transit, daytime sampling was conducted with CUFES, EK80s, ME70, MS70 and SX90 while personnel continued to troubleshoot GPS issues and test equipment. On 25 July, *Lasker* arrived at the waypoint offshore of Cape Mendocino, CA at ~1930 and conducted one tow before initiating acoustic sampling on transect 129 at sunrise. After encountering CPS echoes on the first transect, adaptive sampling was initiated. On 26 July, Simrad-Kongsberg representative David Barbee resolved the issues with GPS-attitude data inputs to the EK80s. ADCP calibrations were successfully completed on 27 July, taking advantage of good weather conditions, close proximity to the intended trawl locations, and the last two hours of daylight. On 29 July, after completing transect 119, *Lasker* ceased adaptive sampling. Small boat operations were conducted near Point Arena on 30 July at ~0230 to embark acoustician Scott Mau, biologist Rachel Backman, and Junior Officer Daniel Stofka. On 4 August, *Lasker* completed transect 103 off Monterey, CA, ceased acoustic sampling, and transited to Point Conception, CA to recover an acoustic lander. The lander was recovered on 5 August at ~1400 before *Lasker* continued south to San Diego. On 6 August, *Lasker* arrived at the 10th Avenue Marine Terminal in San Diego, CA at ~1400 to complete Leg II.

Meanwhile, *Lisa Marie* sampled core-region Transects 141 to 114, between Cape Sebastian, OR to Bodega Bay, CA, from 21 July to 1 August. On 2 August, on its transit north, *Lisa Marie* resampled Transect 133, which she had previously been sampled during Leg II, but without recording EK80 data. On 3 August, *Lisa Marie* returned to Westport, WA to conclude its portion of the 2022 Summer CCE survey.

<sup>4</sup>[https://coastwatch.pfeg.noaa.gov/erddap/griddap/sardine\\_habitat\\_modis.html](https://coastwatch.pfeg.noaa.gov/erddap/griddap/sardine_habitat_modis.html)

937 The two USVs (SD-1076 and SD-1077) sampled core-region Transects 158 to 146, between Neskowin, OR to  
938 Bandon, OR, from 23 July to 13 August.

939 *Long Beach Carnage* sampled nearshore Transects 227 to 186, between Bodega Bay, CA to Morro Bay, CA,  
940 from 30 July to 5 August.

### 941 **Leg III**

942 All but four days on Leg III were lost, so the remaining time was used for CPS reconnaissance in the SCB.  
943 At ~1400 on 26 August, after a 16-day delay, *Lasker* departed from the 10th Avenue Marine Terminal Pier  
944 in San Diego, CA. At ~1930 on 26 August, acoustic sampling was conducted along transect 72. A total of  
945 five transects were completed in the SCB. On 29 August, acoustic sampling ceased after the completion of  
946 transect 78 off Long Beach, CA. On 30 August, after completing a compass calibration outside of the San  
947 Diego sea buoy, *Lasker* arrived at the 10th Avenue Marine Terminal in San Diego, CA at ~1900 to complete  
948 Leg III.

949 The two USVs (SD-1076 and SD-1077) sampled core-region Transects 144 to 122, between Cape Blanco, OR  
950 to Beaver Point, CA, from 9 to 31 August. *Long Beach Carnage* sampled nearshore Transects 185 to 138,  
951 between Point Buchon, CA to the U.S.-Mexico border, and the Santa Cruz and Santa Catalina Islands, from  
952 20 August to 8 September.

### 953 **Leg IV**

954 At ~1615 on 9 September, *Lasker* departed from 10th Avenue Marine Terminal in San Diego, CA. At ~1430  
955 on 12 September, *Lasker* resumed acoustic sampling along transect 103 off Monterey, CA. At ~0330 on 16  
956 September, an acoustic lander was deployed at 34.438635 N, 120.54667 W. The lander, consisting of an  
957 autonomous echosounder (WBAT; Simrad-Kongsberg) and passive-acoustic recorder (AURAL-M2; Multi-  
958 Electronique), is part of an ongoing project to utilize stationary platforms for monitoring the ecosystem off  
959 Point Conception, CA, and stocks of CPS that migrate past there. At 0330 on 17 September, scientists  
960 Dayv Lowry and Daniel Hernandez Cruz embarked via small craft in Santa Barbara Harbor, CA. At 0330  
961 on 24 September, scientist Brittany Schwarzkopf disembarked and *Lasker's* XO embarked via small craft in  
962 Mission Bay, CA. At 0200 on 29 September, acoustic sampling ceased at sunset along transect 54 off Punta  
963 Baja, Baja California. At 0700 on 30 September, *Lasker* arrived at the 10th Avenue Marine Terminal in San  
964 Diego, CA to conclude the 2022 Summer CCE survey.

## 965 **3.2 Acoustic backscatter**

966 Acoustic backscatter ascribed to CPS was observed throughout the latitudinal range of the core survey area  
967 (**Fig. 15a**) and was present nearshore to the shelf break. Zero-biomass intervals were observed at the  
968 offshore end of each transect in the core region. The majority (greater than 90%) of the biomass for each  
969 species was apportioned using catch data from trawl clusters conducted within 30 nmi (**Fig. 14**).

970 Acoustic backscatter ascribed to CPS was also observed throughout the nearshore survey area (**Figs. 16a**  
971 and **17a**), but was most prevalent in transects sampled by *Lisa Marie* between Crescent City and Bodega Bay  
972 (**Fig. 16a**) and along transects sampled by *Long Beach Carnage* between Bodega Bay and San Francisco,  
973 between Big Sur and Long Beach, and around Santa Cruz Island (**Fig. 17a**).

## 974 **3.3 Egg densities and distributions**

975 Jack Mackerel eggs were predominant in CUFES samples collected north of Monterey, but were most abun-  
976 dant between Cape Mendocino and Fort Bragg (**Fig. 15b**). Northern Anchovy eggs were predominant in  
977 positive samples collected between Monterey and El Rosario (**Fig. 15b**). Some Pacific Sardine eggs were  
978 present in samples collected offshore between San Francisco and Half Moon Bay, CA and in a few samples  
979 throughout the SCB (**Fig. 15b**).

### 3.4 Trawl catch

Trawl catches from *Lasker* were comprised of mostly Jack Mackerel between Cape Mendocino and Point Arena, CA, and Northern Anchovy farther south (Fig. 15c). Pacific Herring were present in several trawl clusters off the coast of WA and OR. Pacific Sardine were caught in relatively small numbers between Cape Mendocino and Fort Bragg (Fig. 15c). Relatively few Pacific Mackerel and Pacific Herring were collected (Fig. 15c). Overall, the 86 trawls and 41 purse-seine sets captured a combined 7,356 kg of CPS (6,887 kg of Northern Anchovy, 213 kg of Pacific Sardine, 18 kg of Pacific Mackerel, 217 kg of Jack Mackerel, and 20.2 kg of Pacific Herring).

### 3.5 Purse-seine catch

#### 3.5.1 *Lisa Marie*

North of the Columbia River, Pacific Herring were predominant, by weight, in the purse-seine samples (Fig. 13e). Between the Columbia River and Cape Mendocino, Pacific Sardine were predominant (Fig. 13e). On numerous occasions, however, Jack Mackerel were reportedly schooling with Pacific Sardine, but eluded capture, thereby biasing the species composition in those samples (K. Hinton, pers. comm.). Therefore, for the region between Astoria and Cape Mendocino, the acoustic proportions of Pacific Sardine in nighttime trawl clusters from *Lasker* between 2018 and 2021 (Fig. 13a-c) were used to create a generalized additive model (GAM) describing their acoustic proportion versus latitude (Fig. 13d), which was used to estimate the acoustic proportion of Pacific Sardine in the purse-seine catch in 2022 (Fig. 13f). The complement acoustic proportion in the 2022 purse-seine catch was assumed to be Jack Mackerel, based on the association of Jack Mackerel and Pacific Sardine in the summer 2021 nighttime trawl catches (Fig. 13f). Other species (e.g., Pacific Herring, Pacific Mackerel, saury, and smelt) that could have contributed to the CPS backscatter in this area were also observed or suspected to be avoiding the net (K. Hinton, pers. comm.), but those species comprised a small fraction (always <5%, <1.4% on average, and mostly <1%) of the acoustic proportions in clusters that contained both Pacific Sardine and Jack Mackerel in 2021. The Jack Mackerel lengths in this region are those from the summer 2021 nighttime trawl clusters nearest to the CPS backscatter observed in summer 2022. The purse-seine was only deployed when schools were observed, so purse-seine sampling was sparse along portions of the WA and OR coast. Overall, the 41 seines captured a combined 62.4 kg of CPS (37.5 kg of Pacific Sardine, 12.1 kg of Jack Mackerel, 4.75 kg of Pacific Herring, 4.08 t of Pacific Mackerel, and 3.9 kg of Northern Anchovy).

#### 3.5.2 *Long Beach Carnage*

Northern Anchovy were predominant, by weight, in purse-seine samples collected by *Long Beach Carnage* nearshore off central CA between Cape Mendocino and Monterey (Fig. 17b). Pacific Sardine were predominant between Big Sur and San Diego (Fig. 17b). Some Pacific Mackerel were collected between Oceanside, CA and San Diego and around Santa Cruz and Santa Catalina Islands. Jack Mackerel were collected between Big Sur and Morro Bay, and off San Diego (Fig. 17b). Overall, dip net samples from 53 seines totaled 151 kg of CPS (87 kg of Pacific Sardine, 46 kg of Pacific Mackerel, kg of Jack Mackerel, and 10 kg of Northern Anchovy; and no Pacific Herring).

#### 3.5.3 Combined catch

In some areas, purse-seine sets were sparse (Figs. 16b and 17b). To estimate biomass in the nearshore region, acoustic intervals were assigned the species proportions from the nearest purse-seine set or trawl cluster, whichever was closest (Fig. 18b).




Figure 13: Maps of species proportions in *Lasker's* nighttime trawl-catch clusters during summer a) 2018, b) 2019, and c) 2021 depicting an analysis region between the Columbia River and Cape Mendocino (dashed line); d) a model of the acoustic proportions of Pacific Sardine versus latitude in *Lasker's* nighttime trawl-catch clusters between 2018 and 2021; e) species proportions in *Lisa Marie's* summer 2022 purse-seine catches; and f) the species proportions used in the estimation of summer 2022 CPS biomasses north of Cape Mendocino, replacing *Lisa Marie's* catches containing Pacific Sardine, in the analysis region, with the modeled proportion of Pacific Sardine, with its complement assigned to Jack Mackerel.




Figure 14: Proportion (top) and cumulative proportion (bottom) of biomass versus distance to the nearest positive trawl cluster. Dashed vertical lines (bottom) represent the cluster distance where cumulative biomass equals 90%.





Figure 15: Spatial distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ ,  $m^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; b) CUFES egg density ( $\text{eggs m}^{-3}$ ) for Northern Anchovy, Pacific Sardine, and Jack Mackerel; and c) acoustic proportions of CPS in trawl clusters (black outline) and purse-seine sets (white outline). Black points indicate trawl clusters or purse-seine sets with no CPS.



a)



b)




Figure 16: Nearshore survey transects sampled by *Lisa Marie* overlaid with the distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ ,  $m^2 \text{ nmi}^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; and b) the proportions, by weight, of CPS in each purse-seine catch. Species with low catch weights may not be visible at this scale.




Figure 17: Nearshore transects sampled by *Long Beach Carnage* overlaid with the distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ ,  $m^2 nmi^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS; and b) the proportions, by weight, of CPS in each purse-seine catch. Species with low catch weights may not be visible at this scale.




Figure 18: Spatial distributions of: a) 38-kHz integrated backscattering coefficients ( $s_A$ ,  $m^2 nmi^{-2}$ ; averaged over 2000-m distance intervals) ascribed to CPS from nearshore sampling; and b) acoustic proportions of CPS in trawl clusters (black outline) and purse-seine sets (white outline).

1021 **3.6 Biomass distribution and demographics**

1022 The biomasses, distributions, and demographics for each species and stock are for the survey area and period  
 1023 and therefore may not represent the entire population. No nearshore sampling was conducted off Baja CA,  
 1024 so nearshore biomass estimates are for U.S. waters only.

1025 **3.6.1 Northern Anchovy**

1026 **3.6.1.1 Northern stock**

1027 The total estimated biomass of the northern stock of Northern Anchovy was 16,432 t ( $CI_{95\%} = 5,646 -$   
 1028  $27,680$  t,  $CV = 34\%$ ; **Table 5**). In the core region, biomass was 16,432 t ( $CI_{95\%} = 5,646 - 27,680$  t,  $CV =$   
 1029  $34\%$ ; **Table 5**); the stock was distributed throughout the survey area from approximately Westport to Cape  
 1030 Blanco (**Fig. 19a**).  $L_S$  ranged from 10 to 15 cm with modes at 10 and 13 cm (**Table 6, Fig. 20**). In the  
 1031 nearshore region, biomass was 0.0934 t ( $CI_{95\%} = 0 - 0.285$  t,  $CV = 94\%$ ; **Table 5**), comprising 0.00057% of  
 1032 the total biomass, and was located near the entrance to the Columbia River (**Fig. 19b**).  $L_S$  had a single  
 1033 mode at  $\sim 13$  cm (**Table 6**; not visible in **Fig. 20**).

Table 5: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for the northern stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions. Stratum areas are  $nmi^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\hat{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	4	3,287	8	352	1	1	16,321	5,556	27,583	35
	5	6,558	13	659	2	9	111	2	319	76
	All	9,844	21	1,011	3	10	16,432	5,646	27,680	34
Nearshore	10	43	2	4	1	7	0	0	0	94
	All	43	2	4	1	7	0	0	0	94
<b>All</b>	-	<b>9,887</b>	<b>23</b>	<b>1,014</b>	<b>4</b>	<b>17</b>	<b>16,432</b>	<b>5,646</b>	<b>27,680</b>	<b>34</b>

Table 6: Abundance versus standard length ( $L_S$ , cm) for the northern stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions.

$L_S$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	0	0
6	0	0
7	0	0
8	0	0
9	0	0
10	1,704,924,599	0
11	0	0
12	0	0
13	2,091,901	2,070
14	1,568,926	1,553
15	428,218	0
16	0	0
17	0	0
18	0	0
19	0	0
20	0	0




Figure 19: Biomass densities (colored points) of the northern stock of Northern Anchovy (*Engraulis mordax*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Northern Anchovy (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 20: Abundance versus standard length ( $L_S$ , upper panels) and biomass (t) versus  $L_S$  (lower panels) for the northern stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions. This plot correctly shows the abundances and biomasses in the core region, which is based on one specimen resulting in the mode at 10 cm (see Cluster 60 in **Appendix A.1**). Abundance and biomass in the nearshore region is negligible relative to the core region and not visible at this scale.



### 3.6.1.2 Central stock

The total estimated biomass of the central stock of Northern Anchovy was 2,235,996 t ( $CI_{95\%} = 1,248,956 - 3,051,863$  t,  $CV = 20\%$ ; **Table 7**), of which 6% was observed in Mexican waters. In the core region, biomass was 2,197,812 t ( $CI_{95\%} = 1,231,227 - 3,002,630$  t,  $CV = 21\%$ ; **Table 7**); the stock was distributed throughout most of the survey area from Bodega Bay to El Rosario (**Fig. 21a**).  $L_S$  ranged from 5 to 16 cm with modes at 9 and 12 cm (**Table 8, Fig. 22**). In the nearshore region, biomass was 38,184 t ( $CI_{95\%} = 17,729 - 49,233$  t,  $CV = 21\%$ ; **Table 7**), comprising 1.7% of the total biomass, and was distributed between Bodega Bay and Los Angeles, CA (**Fig. 21b**). The nearshore length distribution had a single mode at 11 cm (**Table 8, Fig. 22**).

Table 7: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for the central stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions. Stratum areas are  $nmi^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\hat{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	1	4,744	8	237	4	149,276	141,459	32,781	328,023	56
	2	19,805	16	1,020	15	148,855	1,030,667	336,141	1,637,019	34
	3	6,319	16	600	6	197,764	1,025,686	572,423	1,545,177	26
	All	30,867	40	1,857	23	495,895	2,197,812	1,231,227	3,002,630	21
Nearshore	1	46	5	7	1	7	0	0	0	28
	2	142	11	37	5	925	7,872	2,632	13,536	37
	3	279	28	60	9	95,212	24,435	6,546	32,490	28
	4	293	21	66	4	55,065	5,023	1,589	10,124	46
	5	99	2	4	1	462	529	0	1,088	73
	6	99	10	19	2	54	38	11	79	49
	7	99	2	4	1	462	3	0	5	73
	8	85	2	4	1	8,416	53	0	108	74
	9	8	3	1	2	11,729	232	2	551	86
	All	1,149	84	203	24	172,334	38,184	17,729	49,233	21
	All	-	<b>32,017</b>	<b>124</b>	<b>2,060</b>	<b>47</b>	<b>668,229</b>	<b>2,235,996</b>	<b>1,248,956</b>	<b>3,051,863</b>

Table 8: Abundance versus standard length ( $L_S$ , cm) for the central stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions.

$L_S$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	667,895	0
6	296,968,038	4,728,472
7	3,554,272,853	39,943,859
8	12,432,116,940	51,948,230
9	29,065,634,300	79,734,397
10	25,632,382,890	166,702,066
11	11,650,326,085	893,956,011
12	30,509,134,701	860,733,228
13	26,142,957,433	250,610,771
14	9,673,296,890	56,452,414
15	715,539,669	257,977
16	136,425,482	125,663
17	0	0
18	0	0
19	0	0
20	0	0




Figure 21: Biomass densities (colored points) of central stock of Northern Anchovy (*Engraulis mordax*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Northern Anchovy (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 22: Abundance versus standard length ( $L_S$ , upper panels) and biomass (t) versus  $L_S$  (lower panels) for the central stock of Northern Anchovy (*Engraulis mordax*) in the core and nearshore survey regions.

1043 **3.6.2 Pacific Sardine**

1044 **3.6.2.1 Northern stock**

1045 The total estimated biomass of the northern stock of Pacific Sardine was 69,506 t ( $CI_{95\%} = 30,484 - 99,021$  t,  
 1046  $CV = 21\%$ ; **Table 9**). In the core region, biomass was 53,741 t ( $CI_{95\%} = 29,672 - 84,749$  t,  $CV = 26\%$ ; **Table**  
 1047 **9**), was distributed from Westport to Point Conception, but was most abundant between the Columbia River  
 1048 and Newport (**Fig. 23a**).  $L_S$  ranged from 11 to 27 cm with a mode at 16 cm (**Table 10, Fig. 24**). In  
 1049 the nearshore region, biomass was 15,765 t ( $CI_{95\%} = 812 - 14,272$  t,  $CV = 23\%$ ; **Table 9**), comprising 23%  
 1050 of the total biomass. It was distributed mostly between San Francisco and Point Conception, with some  
 1051 present near Crescent City (**Fig. 23b**).  $L_S$  had two modes at 11 and 15 cm (**Table 10, Fig. 24**).

Table 9: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for the northern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions. Stratum areas are  $\text{nm}^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\hat{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	4	4,985	11	410	4	401	4,838	1,145	9,466	45
	5	5,665	15	570	5	179	5,956	2,024	10,094	36
	6	16,113	36	1,672	13	1,056	42,946	20,342	72,916	32
	All	26,764	62	2,651	22	1,636	53,741	29,672	84,749	26
Nearshore	1	297	22	62	5	423	15,764	811	14,270	23
	2	9	3	2	2	52	0	0	1	61
	3	9	3	1	3	20	1	0	2	51
	All	315	28	65	10	495	15,765	812	14,272	23
<b>All</b>	-	<b>27,078</b>	<b>90</b>	<b>2,716</b>	<b>32</b>	<b>2,130</b>	<b>69,506</b>	<b>30,484</b>	<b>99,021</b>	<b>21</b>

Table 10: Abundance versus standard length ( $L_S$ , cm) for the northern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions.

$L_S$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	0	0
6	0	0
7	0	0
8	0	0
9	0	0
10	0	1,924,590
11	318,297	1,511,625
12	444,361	413,140
13	636,594	619,448
14	13,944,275	3,850,443
15	105,679,270	3,607,983
16	268,105,820	1,026,615
17	219,040,205	20,715
18	47,775,938	4,864
19	13,509,188	3,188
20	20,696,254	1,063
21	10,463,389	1,063
22	11,311,389	0
23	20,900,885	0
24	16,335,566	0
25	13,274,355	0
26	7,290,532	0
27	4,915,285	0
28	0	0
29	0	0
30	0	0




Figure 23: Biomass densities (colored points) of the northern stock of Pacific Sardine (*Sardinops sagax*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Sardine (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.






Figure 24: Estimated abundance (upper panel) and biomass (lower panel) versus standard length ( $L_S$ , cm) for the northern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions.

### 3.6.2.2 Southern stock

The total estimated biomass of the southern stock of Pacific Sardine was 107,468 t ( $CI_{95\%} = 47,994 - 178,947$  t,  $CV = 23\%$ ; **Table 11**), of which 0.9% was observed in Mexican waters. In the core region, biomass was 40,206 t ( $CI_{95\%} = 4,741 - 79,328$  t,  $CV = 48\%$ ; **Table 11**), and was distributed from approximately San Francisco to El Rosario (**Fig. 25a**).  $L_S$  ranged from 9 to 21 cm with two modes, at 13 and 18 cm (**Table 12, Fig. 26**). In the nearshore region, biomass was 67,262 t ( $CI_{95\%} = 43,253 - 99,620$  t,  $CV = 23\%$ ; **Table 11**), comprising 63% of the total biomass. The nearshore biomass was distributed between Point Conception and San Diego, but was greatest near Santa Barbara, San Diego, and around Santa Cruz Island. The  $L_S$  in the nearshore region ranged from 10 to 21 cm and had modes at 12 and 16 cm (**Table 12, Fig. 26**).

Table 11: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for the southern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions. Stratum areas are  $nmi^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\hat{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	1	2,611	5	131	2	200	376	68	674	40
	2	13,960	9	730	10	3,770	39,668	4,085	78,502	49
	3	3,555	4	177	1	2	161	7	425	75
	All	20,127	18	1,037	13	3,972	40,206	4,741	79,328	48
Nearshore	4	126	10	25	6	3,864	6,095	8,568	27,070	76
	5	491	45	114	20	713	51,182	19,167	70,237	28
	6	99	10	19	5	293	7,268	2,873	13,701	40
	7	99	7	15	6	343	1,958	765	3,855	43
	8	85	7	14	3	150	352	7	917	73
	9	85	9	19	5	214	408	79	870	50
	All	984	88	204	37	5,578	67,262	43,253	99,620	23
All	-	21,111	106	1,241	50	9,549	107,468	47,994	178,947	23

Table 12: Abundance versus standard length ( $L_S$ , cm) for the southern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions.

$L_S$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	0	0
6	0	0
7	0	0
8	0	0
9	2,070,865	0
10	4,439,182	8,383,644
11	54,646,303	158,236,963
12	52,549,062	275,597,032
13	306,314,473	62,485,385
14	121,021,559	181,015,402
15	11,453,420	348,793,820
16	11,453,420	446,726,568
17	120,610,666	335,727,403
18	195,066,323	49,621,103
19	34,484,274	3,238,436
20	111,371	1,305,083
21	5,726,710	1,338,380
22	0	0
23	0	0
24	0	0
25	0	0
26	0	0
27	0	0
28	0	0
29	0	0
30	0	0




Figure 25: Biomass densities (colored points) of the southern stock of Pacific Sardine (*Sardinops sagax*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Sardine (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 26: Estimated abundance (upper panels) and biomass (lower panels) versus standard length ( $L_S$ , cm) for the southern stock of Pacific Sardine (*Sardinops sagax*) in the core and nearshore survey regions.

1061 **3.6.3 Pacific Mackerel**

1062 The total estimated biomass of Pacific Mackerel was 7,968 t ( $CI_{95\%} = 3,741 - 12,662$  t,  $CV = 22\%$ ; **Table**  
 1063 **13**), of which 27% was observed in Mexican waters. In the core region, biomass was 5,619 t ( $CI_{95\%} = 2,851$   
 1064 - 9,108 t,  $CV = 29\%$ ) and was distributed from approximately Cape Mendocino to El Rosario, but was  
 1065 primarily located south of Point Conception (**Fig. 27a**). The distribution of  $L_F$  ranged from 8 to 38 cm  
 1066 with two modes at 11 and 15 cm (**Table 14, Fig. 28**). In the nearshore region, biomass was 2,349 t ( $CI_{95\%}$   
 1067 = 890 - 3,553 t,  $CV = 30\%$ ; **Table 13, Fig. 27b**), comprising 29% of the total biomass. It was distributed  
 1068 from Point Conception to San Diego, but was most abundant around Santa Cruz and Santa Catalina Islands.  
 1069 The distribution of  $L_F$  ranged from 8 to 35 cm and had modes at 18, 27, and 29 cm.

Table 13: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for Pacific Mackerel (*Scomber japonicus*) in nearshore survey region. Stratum areas are  $nmi^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\bar{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	1	4,744	8	237	4	197	1,495	159	3,577	61
	2	16,400	12	851	10	122	3,981	1,772	6,926	34
	3	2,327	6	232	1	2	144	0	288	55
	All	23,470	26	1,321	14	321	5,619	2,851	9,108	29
Nearshore	1	185	17	40	5	61	328	17	505	41
	2	75	6	21	2	10	50	0	107	56
	3	37	3	11	1	4	0	0	0	49
	4	99	20	39	5	71	1,802	525	3,216	39
	5	85	19	39	9	142	168	51	325	41
	All	481	65	149	21	288	2,349	890	3,553	30
<b>All</b>	-	<b>23,950</b>	<b>91</b>	<b>1,470</b>	<b>35</b>	<b>610</b>	<b>7,968</b>	<b>3,741</b>	<b>12,662</b>	<b>22</b>

Table 14: Abundance versus fork length ( $L_F$ , cm) for Pacific Mackerel (*Scomber japonicus*) in the core and nearshore survey regions.

$L_F$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	0	0
6	0	0
7	0	0
8	41,652,930	161,497
9	35,946,448	322,994
10	25,256,609	0
11	77,660,304	646,051
12	35,867,089	322,994
13	4,933,957	62
14	9,969,569	607
15	16,472,665	1,523,075
16	8,636,305	5,467,389
17	5,295,350	5,797,579
18	6,909,912	7,201,154
19	2,968,101	1,145,259
20	1,677,292	165,231
21	2,633,347	94,314
22	1,210,878	107,018
23	948,000	28,320
24	215,911	196,712
25	107,943	393,425
26	0	575,014
27	136,551	1,522,636
28	0	693,934
29	0	1,009,929
30	0	30,972
31	152,226	331,481
32	0	46,458
33	0	15,486
34	0	15,486
35	0	46,458
36	0	0
37	0	0
38	152,226	0
39	0	0
40	0	0






Figure 27: Biomass densities (colored points) of Pacific Mackerel (*Scomber japonicus*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Mackerel (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 28: Estimated abundance (upper panels) and biomass (lower panels) versus fork length ( $L_F$ , cm) for Pacific Mackerel (*Scomber japonicus*) in the core and nearshore survey regions.

1070 **3.6.4 Jack Mackerel**

1071 The total estimated biomass of Jack Mackerel was 807,090 t ( $CI_{95\%} = 515,560 - 1,145,812$  t,  $CV = 20\%$ ;  
 1072 **Table 15**), of which 0.06% was observed in Mexican waters. In the core region, the biomass was 799,082 t  
 1073 ( $CI_{95\%} = 512,231 - 1,132,052$  t,  $CV = 20\%$ ; **Table 15**). It was distributed throughout the survey area from  
 1074 the Columbia River to El Rosario, but were most abundant between Astoria and Bodega Bay. (**Fig. 29a**).  
 1075  $L_F$  ranged from 3 to 51 cm, with modes at 8 and 34 cm. (**Table 16, Fig. 30**). In the nearshore region,  
 1076 the biomass was 8,009 t ( $CI_{95\%} = 3,328 - 13,761$  t,  $CV = 35\%$ ; **Table 15**), comprising 0.99% of the total  
 1077 biomass. It was distributed from Point Conception to San Diego, but was most abundant near Long Beach  
 1078 and around Santa Cruz Island (**Fig. 29b**), and had a length mode at 19 cm (**Table 16, Fig. 30**). A small  
 1079 amount of biomass was present in the nearshore region near Fort Bragg.

Table 15: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for Jack Mackerel (*Trachurus symmetricus*) in the core and nearshore survey regions. Stratum areas are  $\text{nm}^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\bar{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	1	4,744	8	237	3	16	465	3	1,034	57
	2	19,805	16	1,020	18	1,478	25,551	12,637	39,938	27
	3	2,504	7	256	2	4	94	12	185	49
	4	24,712	59	2,539	11	333	772,972	488,717	1,104,653	21
	All	51,764	90	4,052	32	1,831	799,082	512,231	1,132,052	20
Nearshore	1	185	17	40	6	16	318	55	467	34
	2	85	2	4	1	1	5,589	0	0	0
	3	85	3	6	1	28	1,835	0	160	2
	5	85	2	4	2	20	55	0	2	1
	6	71	5	19	1	10	20	0	76	102
	7	37	3	11	1	81	1	0	1	16
	8	66	10	17	2	34	36	1	8	6
	9	69	6	15	1	3	0	0	0	12
	10	99	7	13	2	45	4	1,317	11,409	64,364
	11	99	12	24	4	120	0	604	3,165	769,833
	12	8	3	1	1	68	24	0	58	87
	13	3	2	0	1	146	75	0	105	40
	14	9	3	1	1	1	52	1	97	51
	All	900	75	155	21	573	8,009	3,328	13,761	35
	All	-	52,664	165	4,207	53	2,404	807,090	515,560	1,145,812

Table 16: Abundance versus fork length ( $L_F$ , cm) for Jack Mackerel (*Trachurus symmetricus*) in the core and nearshore survey regions.

$L_F$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	241,794	0
4	1,419,851	0
5	629,808	0
6	86,626,388	9,501
7	413,381,593	2,696,403
8	561,261,791	602,429
9	297,229,109	252,294
10	116,464,804	108,161
11	116,535,707	1,333,800
12	161,670,180	5,154,860
13	30,428,028	5,290,971
14	39,476,650	3,717,705
15	24,562,868	16,354,113
16	29,829,944	11,423,912
17	10,578,404	19,428,435
18	11,748,959	20,718,304
19	18,663,528	38,024,306
20	12,779,069	5,405,898
21	8,948,724	314,591
22	5,960,163	543,747
23	13,276,594	816,105
24	3,864,777	813,187
25	2,233,347	271,022
26	0	0
27	8,614,274	2,372
28	15,158,763	2,667
29	18,108,237	2,805
30	24,446,401	10,388
31	51,218,487	6,277
32	51,901,530	13,830
33	96,540,428	24,229
34	215,042,403	27,179
35	160,101,796	39,669
36	187,761,602	55,110
37	148,046,781	30,005
38	45,002,461	26,162
39	36,409,165	30,295
40	7,764,248	8,287
41	17,836,083	4,482
42	13,856,657	2,667
43	4,961,080	0
44	3,917,073	1,704
45	6,795,860	0
46	260,788	0
47	260,788	0

Table 16: Abundance versus fork length ( $L_F$ , cm) for Jack Mackerel (*Trachurus symmetricus*) in the core and nearshore survey regions. (*continued*)

$L_F$	Core	Nearshore
48	4,166,132	0
49	3,027,970	0
50	1,538,823	0
51	139,570,655	0
52	0	0
53	0	0
54	0	0
55	0	0
56	0	0
57	0	0
58	0	0
59	0	0
60	0	0

DRAFT




Figure 29: Biomass densities (colored points) of Jack Mackerel (*Trachurus symmetricus*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Jack Mackerel (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 30: Estimated abundance (upper panel) and biomass (lower panel) versus fork length ( $L_F$ , cm) for Jack Mackerel (*Trachurus symmetricus*) in the core and nearshore survey regions.



1080 **3.6.5 Pacific Herring**

1081 The total estimated biomass of Pacific Herring was 50,718 t ( $CI_{95\%} = 14,460 - 99,700$  t,  $CV = 41\%$ ; **Table**  
 1082 **17**). In the core region, biomass was 47,024 t ( $CI_{95\%} = 13,306 - 93,207$  t,  $CV = 44\%$ ; **Table 17**). It  
 1083 was distributed from approximately Cape Flattery to Cape Mendocino, but was most abundant from Cape  
 1084 Flattery to the Columbia River, and between Crescent City and Cape Mendocino (**Fig. 31a**).  $L_F$  ranged  
 1085 from 13 to 17 cm, with modes at 13 and 17 cm (**Table 18, Fig. 32**). In the nearshore region, biomass was  
 1086 3,694 t ( $CI_{95\%} = 1,154 - 6,493$  t,  $CV = 36\%$ ; **Table 17, Fig. 31b**), or 7.3% of the total biomass. It was  
 1087 distributed from Cape Flattery to Cape Mendocino (**Fig. 32**), and the distribution of  $L_F$  ranged from 13  
 1088 to 17 cm and had a mode at 14 cm (**Table 18, Fig. 32**).

Table 17: Biomass estimates (metric tons, t) and their precisions (upper and lower 95% confidence intervals,  $CI_{95\%}$ ; and coefficients of variation, CVs) for Pacific Herring (*Clupea pallasii*) in the core and nearshore survey regions. Stratum areas are  $\text{nm}^2$ .

Region	Stratum				Trawl		Biomass			
	Number	Area	Transects	Distance	Clusters	Individuals	$\hat{B}$	$CI_{L,95\%}$	$CI_{U,95\%}$	CV
Core	1	3,308	9	351	1	6	5,130	1,206	10,319	48
	2	2,087	4	227	1	1	23,177	0	70,776	85
	3	6,911	13	562	3	699	18,717	6,630	26,433	26
	All	12,305	26	1,140	5	706	47,024	13,306	93,207	44
Nearshore	1	26	5	3	1	6	163	2	356	60
	2	146	5	9	3	699	3,531	916	6,295	37
	All	172	10	13	4	705	3,694	1,154	6,493	36
<b>All</b>	-	<b>12,478</b>	<b>36</b>	<b>1,153</b>	<b>9</b>	<b>1,411</b>	<b>50,718</b>	<b>14,460</b>	<b>99,700</b>	<b>41</b>

Table 18: Abundance versus fork length ( $L_F$ , cm) for Pacific Herring (*Clupea pallasii*) in the core and nearshore survey regions.

$L_F$	Region	
	Core	Nearshore
1	0	0
2	0	0
3	0	0
4	0	0
5	0	0
6	0	0
7	0	0
8	0	0
9	0	0
10	0	0
11	0	0
12	0	0
13	37,385,309	10,751,591
14	201,859,877	48,768,596
15	284,781,511	37,136,356
16	115,944,703	15,043,225
17	481,156,110	1,427,246
18	0	0
19	0	0
20	0	0
21	0	0
22	0	0
23	0	0
24	0	0
25	0	0
26	0	0
27	0	0
28	0	0
29	0	0
30	0	0




Figure 31: Biomass densities (colored points) of Pacific Herring (*Clupea pallasii*), per stratum, in the a) core and b) nearshore survey regions. Overlaid are the locations of trawl clusters with at least one Pacific Herring clusters (blue numbers) in each stratum (colored polygons). Thick gray lines represent acoustic transects.




Figure 32: Estimated abundance (upper panel) and biomass (lower panel) versus fork length ( $L_F$ , cm) for Pacific Herring (*Clupea pallasii*) in the core and nearshore survey regions.

## 1089 4 Discussion

1090 The principal objectives of the 75-day, summer 2022 CCE Survey were to estimate the biomasses and  
1091 distributions of the northern and southern stocks of Pacific Sardine and the northern and central stocks of  
1092 Northern Anchovy. Secondary objectives were to produce estimates for Pacific Mackerel, Jack Mackerel, and  
1093 Pacific Herring within the survey area at the time of the survey.

1094 Despite inclement weather conditions, mechanical limitations, staffing issues, and logistical challenges related  
1095 to the COVID-19 pandemic, and the loss of nearly half of the allocated sea days aboard *Lasker*, the core  
1096 region was surveyed by *Lisa Marie* and two USVs from Cape Flattery to Bodega Bay, and by *Lasker* between  
1097 Cape Mendocino and El Rosario off Baja CA. Following the cancellation of Leg 1, the decision was made to  
1098 forgo sampling off Vancouver Island to maximize the likelihood of surveying all of the transects in U.S. waters.  
1099 To mitigate the loss of acoustic sampling effort north of Cape Mendocino, the decision was made to have  
1100 *Lisa Marie* and two USVs acoustically sample along *Lasker*'s compulsory transects in tandem, and daytime  
1101 purse-seine sampling was conducted by *Lisa Marie* to provide information about the distribution, species  
1102 composition, and length distribution of CPS. This approach deviated from the standard ATM surveys, which  
1103 created additional challenges and uncertainty (see **Section 3.5.1** and **Fig. 13**). For example, net avoidance  
1104 by fast-swimming species, such as Jack Mackerel, required a novel method to adjust species proportions in  
1105 that region to minimize uncertainty in the biomass estimates.

### 1106 4.1 Biomass and abundance

#### 1107 4.1.1 Northern Anchovy

1108 **4.1.1.1 Northern stock** The estimated biomass of the northern stock of Northern Anchovy in the survey  
1109 region north of Cape Mendocino was 16,432 t ( $CI_{95\%} = 5,646 - 27,680$  t) in summer 2022. The northern stock  
1110 biomass has comprised a small fraction (0.1 to 5.4%) of the total biomass in the ATM surveys conducted in  
1111 the CCE since at least 2015 (Stierhoff *et al.*, 2021a).

1112 **4.1.1.2 Central stock** The estimated biomass of the central stock of Northern Anchovy in the survey  
1113 region was 2,235,996 t ( $CI_{95\%} = 1,248,956 - 3,051,863$  t) and comprised 68% of the total CPS biomass in  
1114 summer 2022. The biomass represents a ~20% decrease from the 2,721,689 t estimated in summer 2021  
1115 (Stierhoff *et al.*, 2023). In summer 2022, 6% of the central stock Northern Anchovy biomass was observed in  
1116 Mexican waters. In 2015, the ATM survey documented a large recruitment to the central stock of Northern  
1117 Anchovy, and since 2018, the central stock of Northern Anchovy has been the dominant forage fish species  
1118 in the survey area (**Figs. 35a,b**).

#### 1119 4.1.2 Pacific Sardine

1120 **4.1.2.1 Northern stock** The boundary between the northern and southern stocks of Pacific Sardine was  
1121 Point Conception, based foremost on associations with potential habitat but corroborated by the distributions  
1122 of biomass density north and south of Point Conception, and differences in length distribution (**Fig. 33**).  
1123 The estimated biomass of 69,506 t ( $CI_{95\%} = 30,484 - 99,021$  t) in the survey region was a 46% increase  
1124 in biomass compared to the 47,721 t estimated in summer 2021 (Stierhoff *et al.*, 2023). Since 2014, the  
1125 ATM biomass of the northern stock of Pacific Sardine has remained less than the 150,000 t rebuilding target  
1126 adopted by the Pacific Fishery Management Council in 2020<sup>5</sup> (**Figs. 35a,b**).

<sup>5</sup><https://www.pcouncil.org/documents/2020/08/g-1-attachment-1-pacific-sardine-rebuilding-plan-preliminary-environmental-analysis.pdf/>

1127 **4.1.2.2 Southern stock** The estimated biomass of the southern stock of Pacific Sardine in the survey  
1128 region was 107,468 t ( $CI_{95\%} = 47,994 - 178,947$  t). In summer 2022, 106,930 t (core stratum 2 and all  
1129 nearshore strata; 99% of the total biomass) of southern stock biomass was observed in U.S. waters, and the  
1130 remaining 376 t (stratum 1; 0.9% of the total biomass) was observed off Baja CA. Of the portion in U.S.  
1131 waters in summer 2022, 67,262 t (63%) occurred in the nearshore region.

1132 The southern stock was first observed in U.S. waters by the SWFSC's ATM surveys in 2016 (323 t, Stierhoff  
1133 *et al.*, 2021b). Since then, the southern stock biomass in U.S. waters has been increasing, from 33,093 t in  
1134 summer 2018 (Stierhoff *et al.*, 2021b) to 107,468 t in summer 2022. In summer 2017, the summer survey  
1135 did not extend into the SCB (Zwolinski *et al.*, 2019), and no summer survey was conducted in 2020 due to  
1136 COVID-19.

#### 1137 **4.1.3 Pacific Mackerel**

1138 In summer 2022, the estimated biomass of Pacific Mackerel in the survey region was 7,968 t ( $CI_{95\%} = 3,741$   
1139 - 12,662 t), which is lower than recent estimates (21,998 - 42,423 between 2016 and 2021, and the lowest  
1140 biomass observed since 2015 (1,224 t, Stierhoff *et al.*, 2021a).

#### 1141 **4.1.4 Jack Mackerel**

1142 In summer 2022, the estimated biomass of Jack Mackerel in the survey region, south of Cape Flattery, was  
1143 807,090 t ( $CI_{95\%} = 515,560 - 1,145,812$  t), which is 1.4-fold higher than 569,793 t estimated in summer 2021  
1144 (Stierhoff *et al.*, 2023). In summer 2022, Jack Mackerel was the second most abundant CPS overall, and  
1145 comprised 24% of the total CPS biomass (Figs. 35a,b).

#### 1146 **4.1.5 Pacific Herring**

1147 In summer 2022, the estimated biomass of Pacific Herring in U.S. waters south of Cape Flattery, was 50,718  
1148 t ( $CI_{95\%} = 14,460 - 99,700$  t), which was 75% of the 67,920 t estimated in summer 2021 (Stierhoff *et al.*,  
1149 2023).




Figure 33: Differentiation of northern (blue) and southern (red) stocks of Pacific Sardine by: a) length distributions; b) individual (grey points) and catch-mean (colored points) lengths at the latitudes of their respective trawls; and c) geographic locations of trawls catches with (colored points) and without (black points) Pacific Sardine.

## 1150 4.2 Ecosystem dynamics: Forage fish community

1151 The acoustic-trawl method (ATM) has been used to monitor the biomasses and distributions of pelagic and  
1152 mid-water fish stocks worldwide (e.g., Coetzee *et al.*, 2008; Karp and Walters, 1994; Simmonds *et al.*, 2009).  
1153 In 2006, the SWFSC’s ATM survey in the CCE focused on Pacific Sardine (Cutter and Demer, 2008), but  
1154 evolved to assess the five most abundant CPS (Zwolinski *et al.*, 2014): Pacific Sardine, Northern Anchovy,  
1155 Jack Mackerel, Pacific Mackerel, and Pacific Herring. In the CCE, ATM surveys have been used to directly  
1156 assess Pacific Hake (Edwards *et al.*, 2018; JTC, 2014); rockfishes (Demer, 2012a, 2012b, 2012c; Starr *et*  
1157 *al.*, 1996); Pacific Herring (Thomas and Thorne, 2003); northern stock of Pacific Sardine (Hill *et al.*, 2017);  
1158 northern (Mais, 1974, 1977) and central stocks (Kuriyama *et al.*, 2022) of Northern Anchovy; and Pacific  
1159 Mackerel (Crone *et al.*, 2019; Crone and Hill, 2015). The proportions of these stocks that are in water too  
1160 shallow to be sampled by NOAA ships are estimated using samples collected from fishing vessels and USVs.  
1161 Also, concurrent satellite- and ship-based measures of their biotic and abiotic habitats are used to provide  
1162 an ecosystem perspective.

1163 Collectively, these annual or bi-annual ATM surveys provide a unique insight into the dynamics of forage  
1164 fishes in the CCE, including their distributions, abundances, interactions, and environments. For example,  
1165 results from 2006 through 2013 indicate that Pacific Sardine dominated the CPS assemblage, but their  
1166 biomass was declining (Demer and Zwolinski, 2012; Zwolinski and Demer, 2012) and their seasonal migration  
1167 was contracting (Zwolinski *et al.*, 2014). Meanwhile, harvest rates for the declining stock increased (Demer  
1168 and Zwolinski, 2017), and the total forage-fish biomass decreased to less than 200,000 t in 2014 and 2015  
1169 (**Figs. 35a,b**). The U.S. fishery for Pacific Sardine was closed in 2015 (National Marine Fisheries Service,  
1170 2015), and there were reports of mass strandings, deaths, and reproductive failures in Brown Pelicans  
1171 (*Pelecanus occidentalis*<sup>6</sup>), Common Murres (*Uria aalge*), Brandt’s Cormorants (*Phalacrocorax penicillatus*),  
1172 and California sea lions (*Zalophus californianus*<sup>7</sup>) (McClatchie *et al.*, 2016), all of which depend on forage  
1173 species. The National Marine Fisheries Service deemed the stock ‘overfished’ in 2019.

1174 The biomass of the central stock of Northern Anchovy, which had been growing exponentially since 2015,  
1175 decreased ~20% from the 2,721,689 t estimated in summer 2021 (Stierhoff *et al.*, 2023). Jack Mackerel,  
1176 which were found mostly north of Cape Mendocino, included an abundance of apparently age-0 fish farther  
1177 south, suggesting a strong recruitment. The southern stock of Pacific Sardine was found mostly north of the  
1178 U.S.-Mexico border and nearshore. Even considering the additional uncertainties in the biomass estimates  
1179 north of Cape Mendocino (see **Section 2.2.5**, **Section 3.5.1**, and **Fig. 13**) there is no indication that the  
1180 biomasses of the northern stock of Pacific Sardine and the northern stock of Northern Anchovy have changed  
1181 significantly since summer 2021.

1182 The survey time series of estimated CPS biomasses from summer 2008 to 2022 shows that the forage fish  
1183 assemblage in the CCE was dominated by the northern stock of Pacific Sardine until 2013 and a low biomass  
1184 of Jack Mackerel in 2014 and 2015 (**Figs. 35a,b**). Since 2016, the forage-fish biomass has increased, mainly  
1185 due to resurgences of Jack Mackerel and the now dominant central stock of Northern Anchovy (**Figs. 35a,b**),  
1186 whose biomass primarily (1,025,686 t, or 47% of the total estimate biomass) occurred in U.S. waters. In 2022,  
1187 as it was a half century ago (Mais, 1974, 1977), the CPS assemblage is now mostly composed of Northern  
1188 Anchovy and to a lesser extent Jack Mackerel. Between the summers of 2018 and 2021, the biomass of the  
1189 southern stock of Pacific Sardine in U.S. waters has increased from 33,093 to 106,930 t.

1190 Since the resurgence of the central stock of Northern Anchovy, beginning in 2015, there has been consistency  
1191 in the regional distributions of the three dominant species: Northern Anchovy, Jack Mackerel and Pacific  
1192 Herring (**Fig. 34**). Pacific Herring are caught mostly north of central Washington. Lower biomasses of  
1193 northern stock Pacific Sardine and northern stock Northern Anchovy are resident off Oregon and Northern  
1194 California. Jack mackerel are caught between central Washington and Cape Mendocino, often along with  
1195 fewer northern stock Pacific Sardine in recent years. Central stock Northern Anchovy are caught south  
1196 of Cape Mendocino and, with the exception of summer 2021, mostly south of Bodega Bay. The smaller  
1197 northern stock is resident from central Washington to northern California. The summer 2022 distribution

<sup>6</sup>[https://e360.yale.edu/features/brown\\_pelicans\\_a\\_test\\_case\\_for\\_the\\_endangered\\_species\\_act](https://e360.yale.edu/features/brown_pelicans_a_test_case_for_the_endangered_species_act)

<sup>7</sup><https://www.fisheries.noaa.gov/national/marine-life-distress/2013-2017-california-sea-lion-unusual-mortality-event-california>



<sup>1198</sup> of Northern Anchovy appears to have shifted south, better aligning with its distributions during 2015-2019.  
<sup>1199</sup> In contrast to earlier surveys in the time series, the southern stock of Pacific sardine has been persistently  
<sup>1200</sup> present during summer surveys in U.S. waters, mostly in the Southern California Bight (**Fig. 34**).

DRAFT




Figure 34: Distributions of species proportions in *Lasker's* nighttime trawl catches, summer 2015 through 2022. In 2015, the integrated CPS-hake survey sample northward of Vancouver Island. In 2017, there was no sampling in the SCB. In 2020, there was no survey due to the COVID-19 pandemic. In 2021, through a collaboration with Mexico, the CPS survey extended farther south into Baja California. In 2022, there was no nighttime trawl sampling north of Cape Mendocino, California.




Figure 35: a) Estimated and b) cumulative estimated biomasses (t) of the eight most abundant CPS stocks of six species in the CCE during summer since 2008. Surveys typically span the area between Cape Flattery and San Diego, but in some years also include Vancouver Island, Canada (2015-2019) and portions of Baja CA (2021-2022).

## Acknowledgements

1202 The authors greatly appreciate that the ATM surveys require an enormous effort by multiple groups of peo-  
1203 ple, particularly the SWFSC's Advanced Survey Technologies group (Alice Beittel, Scott Mau, David Murfin,  
1204 and Steve Sessions); CalCOFI and Ship Operations group (Noelle Bowlin, Nicolas Concha Saiz, Anne Freire,  
1205 Emily Gardner, Amy Hays, Megan Human, Bryan Overcash, Andrew Thompson, Lanora Vasquez de Mer-  
1206 cado, Bill Watson, and Ed Weber); Life History Group (Brad Erisman, Kelsey James, Brittany Schwartzkopf,  
1207 Owyn Snodgrass, and Jon Walker); Stock Assessment (Peter Kuriyama); and their volunteers (Rachel Back-  
1208 man, Shannon Dolan, Morgan Illman, Dayv Lowry, Dan Palance, Kamran Walsh, and Jake Wood). We  
1209 also thank biologists Daniel Hernandez Cruz and Selene del Carmen Morales Gutierrez (INAPESCA), the  
1210 officers and crew of *Lasker*, and the SWFSC's Fisheries Resources Division administrative staff. The authors  
1211 acknowledge that the methods used are the culmination of more than a half century of development efforts  
1212 from numerous researchers from around the globe. We thank Capts. Rick Blair (*Lisa Marie*) and Rich Ashley  
1213 and Tom Brinton (*Long Beach Carnage*), along with all the F/V crew members, for their coordination and  
1214 cooperation during the nearshore sampling, and to Joel van Noord (CWPA) and Kristen Hinton and Patrick  
1215 Biondo (WDFW) for leading the at-sea processing of purse seine specimens. We thank Diane Pleschner-  
1216 Steele (CWPA) for contracting the *Long Beach Carnage* to conduct the nearshore survey off central CA and  
1217 in the SCB, and Greg Shaughnessy (Ocean Gold Seafoods) for contracting the *Lisa Marie* to conduct the  
1218 nearshore survey of WA, OR, and northern CA. We thank Kelly Kloos, Trevor Stocking, Diego Aceituno,  
1219 Dane McDermott, Kyle Mooers, Jax Mikkelsen, Leeanne Laughlin, Chelsea Protasio, and Michelle Horeczko  
1220 (CDFW) for their efforts to coordinate with and process samples from *Long Beach Carnage*. Finally, reviews  
1221 by [LIST OF REVIEWERS] and **Annie Yau** improved this report.

## References

- 1222
- 1223 Ainslie, M. A., and McCole, J. G. 1998. [A simplified formula for viscous and chemical absorption in sea](#)  
1224 [water](#). Journal of the Acoustical Society of America, 103: 1671–1672.
- 1225 Bakun, A., and Parrish, R. H. 1982. Turbulence, transport, and pelagic fish in the California and Peru  
1226 current systems. California Cooperative Oceanic Fisheries Investigations Reports, 23: 99–112.
- 1227 Barange, M., Hampton, I., and Soule, M. 1996. [Empirical determination of the in situ target strengths of](#)  
1228 [three loosely aggregated pelagic fish species](#). ICES Journal of Marine Science, 53: 225–232.
- 1229 Checkley, D. M., Ortner, P. B., Settle, L. R., and Cummings, S. R. 1997. A continuous, underway fish egg  
1230 sampler. Fisheries Oceanography, 6: 58–73.
- 1231 Chen, C. T., and Millero, F. J. 1977. [Speed of sound in seawater at high pressures](#). Journal of the Acoustical  
1232 Society of America, 62: 1129–1135.
- 1233 Coetzee, J. C., Merkle, D., Moor, C. L. de, Twatwa, N. M., Barange, M., and Butterworth, D. S. 2008.  
1234 [Refined estimates of South African pelagic fish biomass from hydro-acoustic surveys: Quantifying the](#)  
1235 [effects of target strength, signal attenuation and receiver saturation](#). African Journal of Marine Science,  
1236 30: 205–217.
- 1237 Conti, S. G., and Demer, D. A. 2003. [Wide-bandwidth acoustical characterization of anchovy and sardine](#)  
1238 [from reverberation measurements in an echoic tank](#). ICES Journal of Marine Science, 60: 617–624.
- 1239 Crone, P. R., and Hill, K. T. 2015. Pacific mackerel (*Scomber japonicus*) stock assessment for USA man-  
1240 agement in the 2015-16 fishing year. Pacific Fishery Management Council, Pacific Fishery Management  
1241 Council, 7700 NE Ambassador Place, Suite 101, Portland, Oregon 97220: 131 p.
- 1242 Crone, P. R., Hill, K. T., Zwolinski, J. P., and Kinney, M. J. 2019. Pacific mackerel (*Scomber japonicus*) stock  
1243 assessment for U.S. management in the 2019-20 and 2020-21 fishing years. Pacific Fishery Management  
1244 Council, Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, Oregon  
1245 97220: 112 p.
- 1246 Cutter, G. R., and Demer, D. A. 2008. California Current Ecosystem Survey 2006. Acoustic cruise reports  
1247 for NOAA FSV *Oscar Dyson* and NOAA FRV *David Starr Jordan*. U.S. Dep. Commer., NOAA Tech.  
1248 Memo., NOAA-SWFSC-415: 98 pp.
- 1249 Cutter, G. R., Renfree, J. S., Cox, M. J., Brierley, A. S., and Demer, D. A. 2009. [Modelling three-dimensional](#)  
1250 [directivity of sound scattering by Antarctic krill: Progress towards biomass estimation using multibeam](#)  
1251 [sonar](#). ICES Journal of Marine Science, 66: 1245–1251.
- 1252 De Robertis, A., and Higginbottom, I. 2007. [A post-processing technique to estimate the signal-to-noise](#)  
1253 [ratio and remove echosounder background noise](#). ICES Journal of Marine Science, 64: 1282–1291.
- 1254 Demer, D. A. 2012a. 2007 survey of rockfishes in the Southern California Bight using the collaborative  
1255 optical-acoustic survey technique. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-SWFSC-498: 110.
- 1256 Demer, D. A. 2012b. 2004 survey of rockfishes in the Southern California Bight using the collaborative  
1257 optical-acoustic survey technique. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-SWFSC-497: 96.
- 1258 Demer, D. A. 2012c. 2003 survey of rockfishes in the Southern California Bight using the collaborative  
1259 optical-acoustic survey technique. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-SWFSC-496: 82.
- 1260 Demer, D. A., Berger, L., Bernasconi, M., Bethke, E., Boswell, K., Chu, D., Domokos, R., *et al.* 2015.  
1261 Calibration of acoustic instruments. ICES Cooperative Research Report No. 326: 133 pp.
- 1262 Demer, D. A., Conti, S. G., De Rosny, J., and Roux, P. 2003. [Absolute measurements of total target strength](#)  
1263 [from reverberation in a cavity](#). Journal of the Acoustical Society of America, 113: 1387–1394.
- 1264 Demer, D. A., Cutter, G. R., Renfree, J. S., and Butler, J. L. 2009a. [A statistical-spectral method for echo](#)  
1265 [classification](#). ICES Journal of Marine Science, 66: 1081–1090.
- 1266 Demer, D. A., Kloser, R. J., MacLennan, D. N., and Ona, E. 2009b. [An introduction to the proceedings](#)  
1267 [and a synthesis of the 2008 ICES Symposium on the Ecosystem Approach with Fisheries Acoustics and](#)  
1268 [Complementary Technologies \(SEAFACETS\)](#). ICES Journal of Marine Science, 66: 961–965.
- 1269 Demer, D. A., and Zwolinski, J. P. 2012. [Reply to MacCall et al.: Acoustic-trawl survey results provide](#)  
1270 [unique insight to sardine stock decline](#). Proceedings of the National Academy of Sciences of the United  
1271 States of America, 109: E1132–E1133.
- 1272 Demer, D. A., and Zwolinski, J. P. 2017. [A method to consistently approach the target total fishing fraction](#)  
1273 [of Pacific sardine and other internationally exploited fish stocks](#). North American Journal of Fisheries  
1274 Management, 37: 284–293.

- 1275 Demer, D. A., Zwolinski, J. P., Byers, K. A., Cutter, G. R., Renfree, J. S., Sessions, T. S., and Macewicz,  
1276 B. J. 2012. Prediction and confirmation of seasonal migration of Pacific sardine (*Sardinops sagax*) in the  
1277 California Current Ecosystem. *Fishery Bulletin*, 110: 52–70.
- 1278 Doonan, I. J., Coombs, R. F., and McClatchie, S. 2003. [The absorption of sound in seawater in relation to  
1279 the estimation of deep-water fish biomass](#). *ICES Journal of Marine Science*, 60: 1047–1055.
- 1280 Dotson, R. C., Griffith, D. A., King, D. L., and Emmett, R. L. 2010. Evaluation of a marine mammal  
1281 excluder device (MMED) for a Nordic 264 midwater rope trawl. U.S. Dep. Commer., NOAA Tech.  
1282 Memo., NOAA-SWFSC-455: 19.
- 1283 Edwards, A. M., Taylor, I. G., Grandin, C. J., and Berger, A. M. 2018. Status of the Pacific hake (whiting)  
1284 stock in U.S. and Canadian waters in 2018. Prepared by the Joint Technical Committee of the U.S. and  
1285 Canada Pacific Hake/Whiting Agreement, National Marine Fisheries Service and Fisheries and Oceans  
1286 Canada. Report. Pacific Fishery Management Council.
- 1287 Efron, B. 1981. Nonparametric standard errors and confidence intervals. *Canadian Journal of Statistics*, 9:  
1288 139–158.
- 1289 Felix-Uraga, R., Gomez-Mu noz, V. M., Quinonez-Velazquez, C., Melo-Barrera, F. N., and Garcia-Franco,  
1290 W. 2004. On the existence of Pacific sardine groups off the west coast of Baja California and southern  
1291 California. *California Cooperative Oceanic Fisheries Investigations Reports*, 45: 146–151.
- 1292 Felix-Uraga, R., Gomez-Mu noz, V., Hill, K., and Garcia-Franco, W. 2005. Pacific sardine (*Sardinops  
1293 sagax*) stock discrimination off the west coast of Baja California and southern California using otolith  
1294 morphometry. *California Cooperative Oceanic Fisheries Investigations Reports*, 46: 113–121.
- 1295 Field, J. C., Francis, R. C., and Strom, A. 2001. Toward a fisheries ecosystem plan for the northern California  
1296 Current. *California Cooperative Oceanic Fisheries Investigations Reports*, 42: 74–87.
- 1297 Foote, K. G., Knudsen, H. P., Vestnes, G., MacLennan, D. N., and Simmonds, E., J. 1987. Calibration of  
1298 acoustic instruments for fish density estimation: A practical guide. *ICES Cooperative Research Report*,  
1299 144: 69 pp.
- 1300 Francois, R. E., and Garrison, G. R. 1982. [Sound-absorption based on ocean measurements. Part 1: Pure  
1301 water and magnesium-sulfate contributions](#). *Journal of the Acoustical Society of America*, 72: 896–907.
- 1302 Garcia-Morales, R., Shirasago, B., Felix-Uraga, R., and Perez-Lezama, E. 2012. Conceptual models of Pacific  
1303 sardine distribution in the California Current System. *Current Developments in Oceanography*, 5: 23–47.
- 1304 Hewitt, R. P., and Demer, D. A. 2000. [The use of acoustic sampling to estimate the dispersion and abundance  
1305 of euphausiids, with an emphasis on Antarctic krill, \*Euphausia superba\*](#). *Fisheries Research*, 47: 215–229.
- 1306 Hill, K. T., Crone, P. R., Demer, D. A., Zwolinski, J., Dorval, E., and Macewicz, B. J. 2014. Assessment of  
1307 the Pacific sardine resource in 2014 for U.S. management in 2014-15. U.S. Dep. Commer., NOAA Tech.  
1308 Memo., NMFS-SWFSC-531.
- 1309 Hill, K. T., Crone, P. R., and Zwolinski, J. P. 2017. Assessment of the Pacific sardine resource in 2017 for  
1310 U.S. Management in 2017-18. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-576: 264 pp.
- 1311 Johannesson, K., and Mitson, R. 1983. Fisheries acoustics. A practical manual for aquatic biomass estima-  
1312 tion. *FAO Fisheries Technical Paper*.
- 1313 JTC. 2014. Status of the Pacific Hake (whiting) stock in U.S. and Canadian waters in 2014 with a manage-  
1314 ment strategy evaluation. Report.
- 1315 Kang, D., Cho, S., Lee, C., Myoung, J. G., and Na, J. 2009. [Ex situ target-strength measurements of  
1316 Japanese anchovy \(\*Engraulis japonicus\*\) in the coastal Northwest Pacific](#). *ICES Journal of Marine Science*,  
1317 66: 1219–1224.
- 1318 Karp, W. A., and Walters, G. E. 1994. Survey assessment of semi-pelagic Gadoids: the example of walleye  
1319 pollock, *Theragra chalcogramma*, in the Eastern Bering Sea. *Marine Fisheries Review*, 56: 8–22.
- 1320 Kuriyama, P. T., Zwolinski, J. P., Teo, S. L. H., and Hill, K. T. 2022. [Assessment of the Northern Anchovy  
1321 \(\*Engraulis mordax\*\) central subpopulation in 2021 for U.S. management](#). U.S. Dep. Commer., NOAA  
1322 Tech. Memo., NMFS-SWFSC-665: 132 pp.
- 1323 Lo, N. C. H., Macewicz, B. J., and Griffith, D. A. 2011. [Migration of Pacific sardine \(\*Sardinops sagax\*\) off  
1324 the West Coast of United States in 2003-2005](#). *Bulletin of Marine Science*, 87: 395–412.
- 1325 Love, M. S. 1996. Probably More Than You Want to Know About the Fishes of the Pacific Coast. Really  
1326 Big Press, Santa Barbara, CA.
- 1327 MacLennan, D. N., Fernandes, P. G., and Dalen, J. 2002. [A consistent approach to definitions and symbols  
1328 in fisheries acoustics](#). *ICES Journal of Marine Science*, 59: 365–369.

1329 Mais, K. F. 1974. Pelagic fish surveys in the California Current. State of California, Resources Agency,  
1330 Dept. of Fish and Game, Sacramento, CA: 79 pp.

1331 Mais, K. F. 1977. Acoustic surveys of Northern anchovies in the California Current System, 1966-1972.  
1332 International Council for the Exploration of the Sea, 170: 287–295.

1333 McClatchie, S., Goericke, R., Leising, A., Auth, T. D., Bjorkstedt, E., Robertson, R. R., Brodeur, R. D., *et*  
1334 *al.* 2016. State of the California Current 2015-16: Comparisons with the 1997-98 El Niño. California  
1335 Cooperative Ocean and Fisheries Investigations Reports, 57: 5–61.

1336 Nakken, O., and Dommasnes, A. 1975. The application of an echo integration system in investigations of  
1337 the stock strength of the Barents Sea capelin 1971-1974. ICES C.M., B:25: 20.

1338 National Marine Fisheries Service. 2015. Fisheries Off West Coast States; Coastal Pelagic Species Fisheries;  
1339 Closure. U.S. Federal Register, 80: 50 CFR Part 660.

1340 Ona, E. 2003. An expanded target-strength relationship for herring. ICES Journal of Marine Science, 60:  
1341 493–499.

1342 Palance, D., Macewicz, B., Stierhoff, K. L., Demer, D. A., and Zwolinski, J. P. 2019. Length conversions  
1343 and mass-length relationships of five forage-fish species in the California current ecosystem. Journal of  
1344 Fish Biology, 95: 1116–1124.

1345 Peña, H. 2008. [In situ target-strength measurements of Chilean jack mackerel \(\*Trachurus symmetricus\*  
1346 \*murphyi\*\) collected with a scientific echosounder installed on a fishing vessel.](#) ICES Journal of Marine  
1347 Science, 65: 594–604.

1348 Polovina, J. J., Howell, E., Kobayashi, D. R., and Seki, M. P. 2001. [The transition zone chlorophyll front, a  
1349 dynamic global feature defining migration and forage habitat for marine resources.](#) Progress in Oceanog-  
1350 raphy, 49: 469–483.

1351 Renfree, J. S., Beittel, A., Bowlin, N. M., Erisman, B. E., James, K., Mau, S. A., Murfin, D. W., *et al.* 2023.  
1352 [Report on the 2022 California Current Ecosystem \(CCE\) Survey \(2207RL\), 27 June to 30 September](#)  
1353 [2022, conducted aboard NOAA Ship \*Reuben Lasker\*, fishing vessels \*Lisa Marie\* and \*Long Beach Carnage\*,](#)  
1354 [and uncrewed surface vehicles.](#) U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-678: 42 pp.

1355 Renfree, J. S., and Demer, D. A. 2016. Optimising transmit interval and logging range while avoiding aliased  
1356 seabed echoes. ICES Journal of Marine Science, 73: 1955–1964.

1357 Renfree, J. S., Hayes, S. A., and Demer, D. A. 2009. [Sound-scattering spectra of steelhead \(\*Oncorhynchus\*  
1358 \*mykiss\*\), coho \(\*O. kisutch\*\), and chinook \(\*O. tshawytscha\*\) salmonids.](#) ICES Journal of Marine Science,  
1359 66: 1091–1099.

1360 Renfree, J. S., Sessions, T. S., Murfin, D. W., Palance, D., and Demer, D. A. 2019. Calibrations of Wide-  
1361 Bandwidth Transceivers (WBT Mini) with Dual-frequency Transducers (ES38-18/200-18C) for SAILDRONE  
1362 Surveys of the California Current Ecosystem During Summer 2018. U.S. Dep. Commer., NOAA Tech.  
1363 Memo., NMFS-SWFSC-608: 29 pp.

1364 Saunders, R. A., O'Donnell, C., Korneliussen, R. J., Fassler, S. M. M., Clarke, M. W., Egan, A., and Reid,  
1365 D. 2012. [Utility of 18-kHz acoustic data for abundance estimation of Atlantic herring \(\*Clupea harengus\*\).](#)  
1366 ICES Journal of Marine Science, 69: 1086–1098.

1367 Seabird. 2013. Seasoft V2 - SBE Data Processing Manual Revision 7.22.4. Sea-Bird Electronics, Washington,  
1368 USA.

1369 Simmonds, E. J., and Fryer, R. J. 1996. [Which are better, random or systematic acoustic surveys? A  
1370 simulation using North Sea herring as an example.](#) ICES Journal of Marine Science, 53: 39–50.

1371 Simmonds, E. J., Gutierrez, M., Chipollini, A., Gerlotto, F., Woillez, M., and Bertrand, A. 2009. [Optimizing  
1372 the design of acoustic surveys of Peruvian Anchoveta.](#) ICES Journal of Marine Science, 66: 1341–1348.

1373 Simmonds, E. J., and MacLennan, D. N. 2005. Fisheries Acoustics: Theory and Practice, 2nd Edition.  
1374 Blackwell Publishing, Oxford.

1375 Simmonds, E. J., Williamson, N. J., Gerlotto, F., and Aglen, A. 1992. Acoustic survey design and analysis  
1376 procedures: A comprehensive review of good practice. ICES Cooperative Research Report, 187: 1–127.

1377 Smith, P. E. 1978. Precision of sonar mapping for pelagic fish assessment in the California Current. ICES  
1378 Journal of Marine Science, 38: 33–40.

1379 Smith, P. E. 2005. A history of proposals for subpopulation structure in the Pacific sardine (*Sardinops*  
1380 *sagax*) population off Western North America. California Cooperative Oceanic Fisheries Investigations  
1381 Reports, 46: 75–82.

1382 Starr, R. M., Fox, D. S., Hixon, M. A., Tissot, B. N., Johnson, G. E., and Barss, W. H. 1996. Comparison of



1383 submersible-survey and hydroacoustic-survey estimates of fish density on a rocky bank. *Fishery Bulletin*,  
1384 94: 113–123.

1385 Stierhoff, K. L., Renfree, J. S., Rojas-González, R. I., Vallarta-Zárate, J. R. F., Zwolinski, J. P., and Demer,  
1386 D. A. 2023. [Distribution, biomass, and demographics of coastal pelagic fishes in the California Current  
1387 Ecosystem during summer 2021 based on acoustic-trawl sampling](#). U.S. Dep. Commer., NOAA Tech.  
1388 Memo., NMFS-SWFSC-676: 86 pp.

1389 Stierhoff, K. L., Zwolinski, J. P., and Demer, D. A. 2021a. Distribution, biomass, and demography of coastal  
1390 pelagic fishes in the California Current Ecosystem during summer 2015 based on acoustic-trawl sampling.  
1391 U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-648: 74 pp.

1392 Stierhoff, K. L., Zwolinski, J. P., and Demer, D. A. 2021b. Distribution, biomass, and demography of coastal  
1393 pelagic fishes in the California Current Ecosystem during summer 2016 based on acoustic-trawl sampling.  
1394 U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-649: 79 pp.

1395 Swartzman, G. 1997. [Analysis of the summer distribution of fish schools in the Pacific Eastern Boundary  
1396 Current](#). *ICES Journal of Marine Science*, 54: 105–116.

1397 Thomas, G. L., Kirsch, J., and Thorne, R. E. 2002. Ex situ target strength measurements of Pacific herring  
1398 and Pacific sand lance. *North American Journal of Fisheries Management*, 22: 1136–1145.

1399 Thomas, G. L., and Thorne, R. E. 2003. [Acoustical-optical assessment of Pacific Herring and their predator  
1400 assemblage in Prince William Sound, Alaska](#). *Aquatic Living Resources*, 16: 247–253.

1401 Williams, K., Wilson, C. D., and Horne, J. K. 2013. [Walleye pollock \(\*Theragra chalcogramma\*\) behavior in  
1402 midwater trawls](#). *Fisheries Research*, 143: 109–118.

1403 Zhao, X., Wang, Y., and Dai, F. 2008. Depth-dependent target strength of anchovy (*Engraulis japonicus*)  
1404 measured in situ. *ICES Journal of Marine Science*, 65: 882–888.

1405 Zwolinski, J. P., and Demer, D. A. In prep. An updated model of potential habitat for northern stock Pacific  
1406 Sardine (*Sardinops sagax*) and its use for attributing survey observations and fishery landings.

1407 Zwolinski, J. P., and Demer, D. A. 2012. [A cold oceanographic regime with high exploitation rates in  
1408 the northeast pacific forecasts a collapse of the sardine stock](#). *Proceedings of the National Academy of  
1409 Sciences of the United States of America*, 109: 4175–4180.

1410 Zwolinski, J. P., Demer, D. A., Byers, K. A., Cutter, G. R., Renfree, J. S., Sessions, T. S., and Macewicz,  
1411 B. J. 2012. Distributions and abundances of Pacific sardine (*Sardinops sagax*) and other pelagic fishes  
1412 in the California Current Ecosystem during spring 2006, 2008, and 2010, estimated from acoustic-trawl  
1413 surveys. *Fishery Bulletin*, 110: 110–122.

1414 Zwolinski, J. P., Demer, D. A., Cutter Jr., G. R., Stierhoff, K., and Macewicz, B. J. 2014. Building on  
1415 Fisheries Acoustics for Marine Ecosystem Surveys. *Oceanography*, 27: 68–79.

1416 Zwolinski, J. P., Demer, D. A., Macewicz, B. J., Cutter, G. R., Elliot, B. E., Mau, S. A., Murfin, D. W.,  
1417 *et al.* 2016. Acoustic-trawl estimates of northern-stock Pacific sardine biomass during 2015. U.S. Dep.  
1418 Commer., NOAA Tech. Memo., NMFS-SWFSC-559: 15 pp.

1419 Zwolinski, J. P., Demer, D. A., Macewicz, B. J., Mau, S. A., Murfin, D. W., Palance, D., Renfree, J. S.,  
1420 *et al.* 2017. Distribution, biomass and demography of the central-stock of Northern anchovy during  
1421 summer 2016, estimated from acoustic-trawl sampling. U.S. Dep. Commer., NOAA Tech. Memo.,  
1422 NMFS-SWFSC-572: 18 pp.

1423 Zwolinski, J. P., Emmett, R. L., and Demer, D. A. 2011. [Predicting habitat to optimize sampling of Pacific  
1424 sardine \(\*Sardinops sagax\*\)](#). *ICES Journal of Marine Science*, 68: 867–879.

1425 Zwolinski, J. P., Oliveira, P. B., Quintino, V., and Stratoudakis, Y. 2010. [Sardine potential habitat and  
1426 environmental forcing off western Portugal](#). *ICES Journal of Marine Science*, 67: 1553–1564.

1427 Zwolinski, J. P., Stierhoff, K. L., and Demer, D. A. 2019. [Distribution, biomass, and demography of coastal  
1428 pelagic fishes in the California Current Ecosystem during summer 2017 based on acoustic-trawl sampling](#).  
1429 U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-610: 76 pp.




1430 Appendix

1431 A Length distributions and percent biomass by cluster

1432 A.1 Northern Anchovy


1433 Standard length ( $L_S$ ) frequency distributions of Northern Anchovy (*Engraulis mordax*) per nighttime trawl  
 1434 cluster, annotated with the number of individuals caught and their percentage contributions to the abundance  
 1435 in each stratum.



1436


## A.2 Pacific Sardine

1438 Standard length ( $L_S$ ) frequency distributions of Pacific Sardine (*Sardinops sagax*) per nighttime trawl cluster,  
 1439 annotated with the number of individuals caught and their percentage contributions to the abundance in  
 1440 each stratum.



1442 **A.3 Pacific Mackerel**


1443 Fork length ( $L_F$ ) frequency distributions of Pacific Mackerel (*Scomber japonicus*) per nighttime trawl cluster,  
 1444 annotated with the number of individuals caught and their percentage contributions to the abundance in  
 1445 each stratum.



1446

1447 **A.4 Jack Mackerel**


1448 Fork length ( $L_F$ ) frequency distributions of Jack Mackerel (*Trachurus symmetricus*) per nighttime trawl  
 1449 cluster, annotated with the number of individuals caught and their percentage contributions to the abundance  
 1450 in each stratum.



1451

1452 **A.5 Pacific Herring**

1453 Fork length ( $L_F$ ) frequency distributions of Pacific Herring (*Clupea pallasii*) per nighttime trawl cluster,  
 1454 annotated with the number of individuals caught and their percentage contributions to the abundance in  
 1455 each stratum.



1456