

2012 OFLs (mt) and recommended 2013 and 2014 OFLs (mt) for west coast groundfish stocks (overfished stocks in CAPS; stocks with new assessments in bold; component stocks in status quo stock complexes in italics).

Stock	2012 OFL	2013 OFL	2014 OFL	Comments
OVERFISHED STOCKS				
BOCACCIO S. of 40°10' N latitude	732	TBD a/	TBD a/	Assessment will be reviewed at the mop-up panel. OFLs to be determined in Nov.
CANARY	622	592	597	Projected using a 50% SPR from the 2011 update assessment.
COWCOD S. of 40°10' N latitude	13	11	12	Sum of Conception and Monterey OFLs.
<i>COWCOD (Conception)</i>	6	7	7	Projected using a 50% SPR from the 2009 update assessment.
<i>COWCOD (Monterey)</i>	7	5	5	Revised DB-SRA estimate.
DARKBLOTCHED	497	TBD a/	TBD a/	Assessment will be reviewed at the mop-up panel. OFLs to be determined in Nov.
PACIFIC OCEAN PERCH	1,007	844	861	Projected using a 50% SPR from the 2011 full assessment.
PETRALE SOLE	1,279	2,711	2,774	Projected using a 30% SPR from the 2011 full assessment.
WIDOW	4,923	TBD a/	TBD a/	Assessment will be reviewed at the mop-up panel. OFLs to be determined in Nov.
YELLOWEYE	48	51	51	Projected using a 50% SPR from the 2011 update assessment.
NON-OVERFISHED STOCKS				
Arrowtooth Flounder	14,460	7,391	6,912	Projected using a 30% SPR from the 2007 full assessment.
Black Rockfish (OR-CA)	1,169	1,159	1,166	Projected using a 50% SPR from the 2007 full assessment with the addition of the northern OFL 3% reduction to account for the portion of the stock estimated between Cape Falcon and the Columbia River.
Black Rockfish (WA)	435	430	428	Projected using a 50% SPR from the 2007 full assessment with a 3% reduction to account for the portion of the stock estimated between Cape Falcon and the Columbia River.
Cabezon (CA)	176	170	165	Projected using a 45% SPR from the 2009 full assessment.
Cabezon (OR)	50	49	49	Projected using a 45% SPR from the 2009 full assessment.
California scorpionfish	132	126	122	Projected using a 45% SPR from the 2005 full assessment.
Chilipepper S. of 40°10' N latitude	1,872	1,768	1,722	Projected using a 50% SPR from the 2007 full assessment. The portion of the coastwide stock south of 40°10' N. lat. (93%) is based on average historical landings.
Dover Sole	44,826	92,955	77,774	Projected using a 30% SPR from the 2011 full assessment.
English Sole	10,620	7,129	5,906	Projected using a 30% SPR from the 2009 update assessment.
Lingcod N. of 42° N latitude (OR & WA)	2,251	2,102	1,984	Projected using a 45% SPR from the 2009 full assessment.
Lingcod S. of 42° N latitude (CA)	2,597	2,566	2,454	Projected using a 45% SPR from the 2009 full assessment.
Longnose skate	3,006	2,902	2,816	Projected using a 45% SPR from the 2007 full assessment.
Longspine Thornyhead (coastwide)	3,483	3,391	3,304	Projected using a 50% SPR from the 2005 full assessment.
Pacific Cod	3,200	3,200	3,200	Status quo OFL.
Sablefish (coastwide)	8,623	6,621	7,158	Projected using a 45% SPR from the 2011 full assessment.

Stock	2012 OFL	2013 OFL	2014 OFL	Comments
Shortbelly	6,950	6,950	6,950	MSY estimated from 2007 assessment.
Shortspine Thornyhead (coastwide)	2,358	2,333	2,310	Projected using a 50% SPR from the 2005 full assessment.
Splitnose S. of 40°10' N latitude	1,610	1,684	1,747	Projected using a 50% SPR from the 2009 full assessment. The portion of the coastwide stock south of 40°10' N. lat. (64.2%) is based on average historical (1916-2008) landings.
Starry Flounder	1,813	1,825	1,834	Projected using a 30% SPR from the 2005 full assessment.
Yellowtail N. of 40°10' N latitude	4,573	4,579	4,584	Projected using a 50% SPR from the 2005 full assessment.
STOCK COMPLEXES				
Minor Nearshore Rockfish North	116	110	110	Sum of OFL contributions of component stocks in the complex.
<i>Black and yellow</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Blue (CA)</i>	27.5	27.4	27.4	<i>Projected using a 50% SPR from the 2007 full assessment. The portion of the assessed stock in CA north of 40°10' N. lat. (12.7%) is based on average historical landings.</i>
<i>Blue (OR & WA)</i>	33.1	32.3	32.3	<i>Revised DCAC estimate.</i>
<i>Brown</i>	5.3	5.5	5.5	<i>Revised DB-SRA estimate.</i>
<i>Calico</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>China</i>	11.7	9.8	9.8	<i>Revised DB-SRA estimate.</i>
<i>Copper</i>	28.6	26.0	26.0	<i>Revised DB-SRA estimate.</i>
<i>Gopher</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Grass</i>	0.6	0.7	0.7	<i>Revised DB-SRA estimate.</i>
<i>Kelp</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Olive</i>	0.3	0.3	0.3	<i>Revised DB-SRA estimate.</i>
<i>Quillback</i>	8.7	7.4	7.4	<i>Revised DB-SRA estimate.</i>
<i>Treefish</i>	0.2	0.2	0.2	<i>Revised DB-SRA estimate.</i>
Minor Shelf Rockfish North	2,197	2,183	2,195	Sum of OFL contributions of component stocks in the complex.
<i>Bronzespotted</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Bocaccio</i>	268.2	284.0	284.0	<i>Revised DB-SRA estimate.</i>
<i>Chameleon</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Chilipepper</i>	140.9	133.1	129.6	<i>Projected using a 50% SPR from the 2007 full assessment. The portion of the coastwide stock north of 40°10' N. lat. (7%) is based on average historical landings.</i>
<i>Cowcod</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Flag</i>	0.1	0.1	0.1	<i>Revised DB-SRA estimate.</i>
<i>Freckled</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Greenblotched</i>	1.4	1.3	1.3	<i>Revised DB-SRA estimate.</i>
<i>Greenspotted 40°10' to 42° N latitude</i>	20.9	9.4	9.4	<i>Catch-based allocation of 50% SPR from the full 2011 assessment (22.2% of OFL from northern California model).</i>
<i>Greenspotted N. of 42 N latitude (OR & WA)</i>		6.1	6.1	<i>DCAC estimate.</i>

Stock	2012 OFL	2013 OFL	2014 OFL	Comments
<i>Greenstriped</i>	1,232.0	1,252.3	1,268.3	<i>Projected using a 50% SPR from the full 2009 assessment. The portion of the coastwide stock north of 40°10' N. lat. (84.5%) is based on the mean of the 2003-2008 swept area biomass estimates from the NMFS trawl survey.</i>
<i>Halfbanded</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Harlequin</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Honeycomb</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Mexican</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Pink</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Pinkrose</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Puget Sound</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Pygmy</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Redstripe</i>	288.3	269.9	269.9	<i>Revised DB-SRA estimate.</i>
<i>Rosethorn</i>	15.2	12.9	12.9	<i>Revised DB-SRA estimate.</i>
<i>Rosy</i>	2.5	3.0	3.0	<i>Revised DB-SRA estimate.</i>
<i>Silvergray</i>	180.0	159.4	159.4	<i>Revised DB-SRA estimate.</i>
<i>Speckled</i>	0.2	0.2	0.2	<i>Revised DB-SRA estimate.</i>
<i>Squarespot</i>	0.1	0.2	0.2	<i>Revised DB-SRA estimate.</i>
<i>Starry</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Stripetail</i>	35.3	40.4	40.4	<i>Revised DB-SRA estimate.</i>
<i>Swordspine</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Tiger</i>	1.1	1.0	1.0	<i>Revised DB-SRA estimate.</i>
<i>Vermilion</i>	11.1	9.7	9.7	<i>Revised DB-SRA estimate.</i>
Minor Slope Rockfish North	1,507	1,518	1,553	Sum of OFL contributions of component stocks in the complex.
<i>Aurora</i>	17	15.4	15.4	<i>Revised DB-SRA estimate.</i>
<i>Bank</i>	20	17.2	17.2	<i>Revised DB-SRA estimate.</i>
<i>Blackgill</i>	5	4.7	4.7	<i>Revised DCAC estimate.</i>
<i>Redbanded</i>	52	45.3	45.3	<i>Revised DB-SRA estimate.</i>
<i>Rougheye</i>	78	71.1	71.1	<i>Revised DB-SRA estimate.</i>
<i>Sharpchin</i>	232	214.5	214.5	<i>Revised DB-SRA estimate.</i>
<i>Shortraker</i>	22	18.7	18.7	<i>Revised DB-SRA estimate.</i>
<i>Splitnose</i>	897	939.0	974.1	<i>Projected using a 50% SPR from the 2009 full assessment. The portion of the coastwide stock north of 40°10' N. lat. (35.8%) is based on average historical (1916-2008) landings.</i>
<i>Yellowmouth</i>	185	192.4	192.4	<i>Revised DB-SRA estimate.</i>
Minor Nearshore Rockfish South	1,145	1,164	1,160	Sum of OFL contributions of component stocks in the complex.
<i>Shallow Nearshore Species</i>	NA	NA	NA	
<i>Black and yellow</i>	26.8	27.5	27.5	<i>Revised DB-SRA estimate.</i>
<i>China</i>	19.8	16.6	16.6	<i>Revised DB-SRA estimate.</i>
<i>Gopher (N of Pt. Conception)</i>	165.0	157.0	153.0	<i>Projected using a 50% SPR from the 2005 full assessment.</i>
<i>Gopher (S of Pt. Conception)</i>	26.0	25.6	25.6	<i>Revised DCAC estimate.</i>
<i>Grass</i>	55.6	59.6	59.6	<i>Revised DB-SRA estimate.</i>

Stock	2012 OFL	2013 OFL	2014 OFL	Comments
<i>Kelp</i>	25.9	27.7	27.7	<i>Revised DB-SRA estimate.</i>
<i>Deeper Nearshore Species</i>	NA	NA	NA	
<i>Blue (assessed area)</i>	190	187.8	187.8	<i>Projected using a 50% SPR from the 2007 full assessment. The portion of the assessed stock in CA south of 40°10' N. lat. (87.3%) is based on average historical landings.</i>
<i>Blue (S of 34°27' N latitude)</i>	74.0	72.9	72.9	<i>Revised DCAC estimate.</i>
<i>Brown</i>	197.4	204.6	204.6	<i>Revised DB-SRA estimate.</i>
<i>Calico</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Copper</i>	156.0	141.5	141.5	<i>Revised DB-SRA estimate.</i>
<i>Olive</i>	189.5	224.6	224.6	<i>Revised DB-SRA estimate.</i>
<i>Quillback</i>	6.3	5.4	5.4	<i>Revised DB-SRA estimate.</i>
<i>Treefish</i>	12.9	13.2	13.2	<i>Revised DB-SRA estimate.</i>
Minor Shelf Rockfish South	2,243.3	1,910.0	1,912.9	Sum of OFL contributions of component stocks in the complex.
<i>Bronzespotted</i>	6.7	3.6	3.6	<i>Revised DB-SRA estimate.</i>
<i>Chameleon</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Flag</i>	26.6	23.4	23.4	<i>Revised DB-SRA estimate.</i>
<i>Freckled</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Greenblotched</i>	24.6	23.1	23.1	<i>Revised DB-SRA estimate.</i>
<i>Greenspotted</i>	195.3	80.3	80.3	<i>Catch-based allocation of 50% SPR from the full 2011 assessment (77.8% of OFL from northern California model + the OFL from the southern California model).</i>
<i>Greenstriped</i>	226.0	229.7	232.7	<i>Projected using a 50% SPR from the full 2009 assessment. The portion of the coastwide stock south of 40°10' N. lat. (15.5%) is based on the mean of the 2003-2008 swept area biomass estimates from the NMFS trawl survey.</i>
<i>Halfbanded</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Harlequin</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Honeycomb</i>	7.8	9.9	9.9	<i>Revised DB-SRA estimate.</i>
<i>Mexican</i>	2.8	5.1	5.1	<i>Revised DB-SRA estimate.</i>
<i>Pink</i>	2.8	2.5	2.5	<i>Revised DB-SRA estimate.</i>
<i>Pinkrose</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Pygmy</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Redstripe</i>	0.5	0.5	0.5	<i>Revised DB-SRA estimate.</i>
<i>Rosethorn</i>	2.5	2.1	2.1	<i>Revised DB-SRA estimate.</i>
<i>Rosy</i>	36.9	44.5	44.5	<i>Revised DB-SRA estimate.</i>
<i>Silvergray</i>	0.6	0.5	0.5	<i>Revised DB-SRA estimate.</i>
<i>Speckled</i>	42.9	39.4	39.4	<i>Revised DB-SRA estimate.</i>
<i>Squarespot</i>	5.8	11.1	11.1	<i>Revised DB-SRA estimate.</i>
<i>Starry</i>	70.5	62.6	62.6	<i>Revised DB-SRA estimate.</i>
<i>Stripetail</i>	20.6	23.6	23.6	<i>Revised DB-SRA estimate.</i>
<i>Swordspine</i>	12.9	14.2	14.2	<i>Revised DB-SRA estimate.</i>
<i>Tiger</i>	0.0	0.0	0.0	<i>Revised DB-SRA estimate.</i>
<i>Vermilion</i>	308.4	269.3	269.3	<i>Revised DB-SRA estimate.</i>

Stock	2012 OFL	2013 OFL	2014 OFL	Comments
<i>Yellowtail</i>	1,248.9	1,064.4	1,064.4	<i>Revised DB-SRA estimate.</i>
Minor Slope Rockfish South	903	681	685	Sum of OFL contributions of component stocks in the complex.
<i>Aurora</i>	29.4	26.1	26.1	<i>Revised DB-SRA estimate.</i>
<i>Bank</i>	574.8	503.2	503.2	<i>Revised DB-SRA estimate.</i>
<i>Blackgill</i>	275.0	130.0	134.0	<i>Projected using a 50% SPR from the 2011 full assessment.</i>
<i>Pacific ocean perch</i>	0.0	0.0	0.0	<i>No harvest contribution (3a stock).</i>
<i>Redbanded</i>	11.9	10.4	10.4	<i>Revised DB-SRA estimate.</i>
<i>Rougheye</i>	0.5	0.4	0.4	<i>Revised DB-SRA estimate.</i>
<i>Sharpchin</i>	10.6	9.8	9.8	<i>Revised DB-SRA estimate.</i>
<i>Shortraker</i>	0.1	0.1	0.1	<i>Revised DB-SRA estimate.</i>
<i>Yellowmouth</i>	0.8	0.8	0.8	<i>Revised DB-SRA estimate.</i>
Other Flatfish	10,146	10,060	10,060	Sum of OFL contributions of component stocks in the complex.
<i>Butter sole</i>	4.6	4.6	4.6	<i>Based on the average catch during 1994-1998 + a 60% discard rate estimated from the EDCP study.</i>
<i>Curlfin sole</i>	8.2	8.2	8.2	<i>Based on the average catch during 1994-1998 + a 60% discard rate estimated from the EDCP study.</i>
<i>Flathead sole</i>	35.0	35.0	35.0	<i>Max. catch = 35 mt in 2005.</i>
<i>Pacific sanddab</i>	4,942.5	4,801.0	4,801.0	<i>Revised DB-SRA estimate.</i>
<i>Rex sole</i>	4,308.6	4,371.5	4,371.5	<i>Revised DB-SRA estimate.</i>
<i>Rock sole</i>	66.0	66.7	66.7	<i>Revised DB-SRA estimate.</i>
<i>Sand sole</i>	780.8	773.2	773.2	<i>Revised DB-SRA estimate.</i>
Other Fish	11,150	TBD c/	TBD c/	No analytical basis for the status quo OFL.
<i>Big skate</i>				
<i>Cabezon (WA)</i>				
<i>California skate</i>				
<i>Finescale codling</i>				
<i>Kelp greenling (CA)</i>	110.6	118.9	118.9	<i>Revised DB-SRA estimate.</i>
<i>Kelp greenling (OR & WA)</i>				
<i>Leopard shark</i>	164.0	167.1	167.1	<i>Revised DB-SRA estimate.</i>
<i>Pacific rattail</i>	1,178.1			
<i>Ratfish</i>				
<i>Southern shark</i>	62.4	61.6	61.6	<i>Revised DCAC estimate.</i>
<i>Spiny dogfish</i>	2,200.2	2,980.0	2,950.0	<i>Projected using a 45% SPR from the 2011 full assessment.</i>

a/ Values to be determined in November pending review of 2011 assessments at the mop-up panel..

b/ Values to be provided supplementally.

c/ Values for these specifications require supplemental harvest-based OFLs and/or the addition of new species to the complex (e.g., non-FMP skates and grenadiers).