

CHAPTER II: AFFECTED ENVIRONMENT - CHINOOK SALMON ASSESSMENT

CHINOOK STOCKS SOUTH OF CAPE FALCON

Sacramento River Fall Chinook

Predictor Description

The Council's Salmon FMP sets the escapement goal for SRFC as a range from 122,000 to 180,000 hatchery and natural area adults. This stock comprises a large proportion of the Chinook spawners returning to Central Valley streams and hatcheries. The Sacramento Index (SI) is the sum of (1) SRFC ocean fishery harvest south of Cape Falcon between September 1 and August 31, (2) SRFC impacts from non-retention ocean fisheries when they occur, (3) the recreational harvest of SRFC in the Sacramento River Basin, and (4) the SRFC adult spawner escapement (Table II-1, Figure II-1). The SRFC exploitation rate is the ocean and river harvest (and non-retention impacts) divided by the SI, and has varied significantly since 1983 (Table II-1). Since 1990, the SRFC exploitation rate has generally declined over time. In 2010 the SRFC exploitation rate was 18 percent.

The STT based the forecast of the SI on a zero-intercept linear model relating the previous year ($t-1$) SRFC jack escapement to the SI in year t (Figure II-2). In addition to the mean SI predictor model, 95 percent prediction intervals for the SI are displayed in Figure II-2. To estimate prediction intervals, errors in the zero-intercept linear model were assumed to be additive and normally distributed. The interpretation of this interval is that a single future observation would be expected to be contained within this interval with 95 percent probability. The additive error structure of this model results in the lower bound of the prediction interval including zero for SI forecasts made at low jack escapement levels. While this may be somewhat unrealistic (one would expect a nonzero SI in year t to result from a nonzero jack escapement in year $t-1$), the STT concludes that the error structure used for this model best approximates the true uncertainty in this forecast at the current time.

Predictor Performance

In 2010 the SI preseason forecast of 245,483 was 1.6 times its postseason value of 152,857.

2011 Stock Status

A total of 27,483 SRFC jacks were estimated to have escaped to Sacramento River basin hatcheries and natural spawning areas in 2010. The resulting 2011 SI forecast is 729,893 adult SRFC (Figure II-2). For the 2011 SI forecast, the upper bound of the 95 percent prediction interval is 1,228,114 and the lower bound is 231,671.

Klamath River Fall Chinook

Predictor Description

For Klamath River fall Chinook, linear regressions are used to relate September 1 ocean abundance estimates of age-3, age-4, and age-5 fish to that year's river run size estimates of age-2, age-3, and age-4 fish, respectively (Table II-2). Historical abundance estimates were derived from a cohort analysis of CWT information (brood years 1979-2006). The y-intercept of the regressions is constrained to zero, which gives the biologically reasonable expectation that a river run size of zero predicts an ocean abundance remainder of zero for the same cohort. The abundance of age-2 fish is not forecasted because no precursor to age-2 fish of that brood is available. Ocean fisheries harvest small numbers of age-2 KRFC.

Predictor Performance

Since 1985, the preseason ocean abundance forecasts for age-3 fish have ranged from 0.33 to 2.72 times the postseason estimates; for age-4 fish from 0.47 to 2.60 times the postseason estimates; and for the adult stock as a whole from 0.34 to 2.03 times the postseason estimates (Table II-3). The September 1, 2009 age-3 forecast (223,400) was 1.03 times its postseason estimate (217,062). The age-4 forecast (106,300) was 1.60 times its postseason estimate (66,452); and the age-5 forecast (1,800) was 3.51 times its postseason estimate (513) (Table II-3). The preseason forecast of the adult stock as a whole was 1.17 times the postseason estimate.

Management of KRFC harvest since 1986 has attempted to achieve specific harvest rates on fully-vulnerable age-4 and age-5 fish in ocean and river fisheries (Table II-4). The Council has used a combination of quotas and time/area restrictions in ocean fisheries in an attempt to meet the harvest rate objective set each year. Since 1992, fisheries have been managed to achieve 50/50 allocation between tribal and non-tribal fisheries. Tribal and recreational river fisheries have been managed on the basis of adult Chinook quotas.

The Council's FMP conservation objective for KRFC (Amendment 9) permits an average natural spawner reduction rate via fisheries of no more than 0.67, with a minimum escapement of 35,000 natural spawning adults. The plan allows for any ocean and river harvest allocation that meets the spawner reduction rate constraint, provided it also meets the minimum escapement goal. The regulations adopted in 2010 were expected to result in 40,700 natural area spawning adults and an age-4 ocean harvest rate of 12.3 percent. Postseason estimates of these quantities were 37,222 natural area adult spawners and an age-4 ocean harvest rate of 3.9 percent (Table II-5).

2011 Stock Status

The 2011 forecast for the ocean abundance of KRFC as of September 1, 2010 (preseason) is 304,600 age-3 fish, the age-4 forecast is 61,600, and the age-5 forecast is 5,000 fish.

Late-season ocean fisheries in 2010 (September through November) were estimated to have harvested zero adult KRFC; hence, no harvest will be deducted from the ocean fishery's allocation in determining the 2011 allowable ocean harvest.

Amendment 15 to the Salmon FMP (implemented March 26, 2008) provides for potential limited harvest of KRFC in ocean salmon fisheries during years that might otherwise be closed due to a projected shortfall in meeting the 35,000 natural spawner conservation objective, as long as this would not jeopardize the long term capacity of the stock to produce maximum sustainable yield on continuing basis. In 2011, there is no basis for invoking *de minimis* fishing under Amendment 15 because KRFC is not projected to fall short of the 35,000 floor.

KRFC triggered an overfishing concern in 2006, after failing to achieve a minimum of 35,000 natural area adult spawners in 2004, 2005, and 2006. The Council recommended that the overfishing concern be ended when KRFC exceeded 35,000 natural area adult spawners in three out of four consecutive years, or when escapement exceeded 40,700 natural area adults in two consecutive years. In 2010, KRFC satisfied the Council adopted criteria for ending the overfishing concern by exceeding 35,000 natural area adult spawners for the third time in four years (2007, 2009, and 2010).

Other California Coastal Chinook Stocks

Other California coastal streams that support fall Chinook stocks which contribute to ocean fisheries off Oregon and California, include the Smith, Little, Mad, Eel, and Mattole rivers, and Redwood Creek. Except for the Smith River, these stocks are included in the California coastal Chinook ESU, which is

listed as threatened under the ESA. Current information is insufficient to forecast the ocean abundance of these stocks, however, the NMFS ESA consultation standard restricts the Klamath River fall Chinook age-4 ocean harvest rate to no more than 16.0 percent to limit impacts on these stocks. In 2010 the age-4 ocean harvest rate was 4.0 percent.

Oregon Coast Chinook Stocks

Oregon coast Chinook stocks are categorized into three major subgroups based on ocean migration patterns; the North Oregon Coast (NOC) Chinook aggregate, the Mid Oregon Coast (MOC) Chinook aggregate, and the South Oregon Coast (SOC) Chinook aggregate. Although their ocean harvest distributions overlap somewhat, they have been labeled as far-north, north, or south/local migrating, respectively.

Far-North and North Migrating Chinook (NOC and MOC groups)

Far-north and north migrating Chinook stocks include spring and fall stocks north of and including the Elk River, with the exception of Umpqua River spring Chinook. Based on CWT analysis, the populations from ten major NOC river systems from the Nehalem through the Siuslaw Rivers are harvested primarily in ocean fisheries off British Columbia and Southeast Alaska, and to a much lesser degree in Council area and terminal area (state waters) fisheries off Washington and Oregon. CWT analysis indicates populations from five major MOC systems, from the Coos through the Elk Rivers, are harvested primarily in ocean fisheries off British Columbia, Canada, Washington, Oregon, and in terminal area fisheries. Minor catches occur in California fisheries and variable catches have been observed in southeast Alaska troll fisheries.

South/Local Migrating Chinook (SOC group)

South/local migrating Chinook stocks include Rogue River spring and fall Chinook, fall Chinook from smaller rivers south of the Elk River, and Umpqua River spring Chinook. These stocks are important contributors to ocean fisheries off Oregon and northern California. Umpqua River spring Chinook contribute to a lesser degree to fisheries off Washington, British Columbia, and southeast Alaska.

Rogue River fall Chinook contribute to ocean fisheries principally as age-3 through age-5 fish. Mature fish enter the river each year from mid-July through October, with the peak of the run occurring during August and September.

Umpqua and Rogue spring Chinook contribute to ocean fisheries primarily as age-3 fish. Mature Chinook enter the rivers primarily during April and May and generally prior to annual ocean fisheries. Quantitative abundance predictions are not made for these stocks.

Natural fall Chinook stocks from river systems south of the Elk River and spring Chinook stocks from the Rogue and Umpqua rivers dominate production from this subgroup. Substantial releases of hatchery spring Chinook occur in both the Rogue and Umpqua rivers, although also present in lesser numbers are hatchery fall Chinook, primarily from the Chetco River.

Predictor Description and 2011 Stock Status for NOC and MOC Groups

Quantitative abundance predictions are made for all three of the coastal Chinook groups (NOC, MOC, and SOC), but are not used in annual development of Council area fishery regulations. Quantitative forecasts of abundance are based on sibling regression analyses from individual basin's escapement assessment data and scale sampling, which occurs coast-wide. Forecast data for the NOC are used in the PSC management process in addition to terminal area management actions.

Natural spawner escapement is assessed yearly from the Nehalem through Sixes rivers. Peak spawning counts of adults are obtained from standard index areas on these rivers and monitored to assess stock trends (PFMC 2011, Chapter II, Table II-4 and Figure II-3). Natural fall Chinook stocks from both the NOC and MOC dominate production from this subgroup. Also present in lesser numbers are naturally-produced spring Chinook stocks from several rivers, and hatchery fall and/or spring Chinook released in the Trask, Nestucca, Salmon, Alsea, and Elk rivers.

Basin-specific forecasts constitute the overall aggregate forecasts and are derived in conjunction with annual PSC Chinook model input and calibration activities; however they were not available at publication time.

North Oregon Coast

Since 1977, the Salmon River Hatchery production has been tagged for use primarily as a PSC indicator stock for the NOC stock component. Because these fish are primarily harvested in fisheries north of the Council management area, the STT has not reviewed the procedure by which this indicator stock is used in estimating annual stock status. The annual spawner counts decreased from 2002 through 2008 despite excellent parental escapements indices in 2001 to 2004. The 2010 spawner counts were a 45 percent increase from 2009 (PFMC 2011, Appendix B, Table B-11). The 2011 NOC stock abundance is expected to be greater than the 2010 abundance.

Based on the density index of total spawners, the generalized expectation for NOC stocks in 2011 is above recent year's average abundance. Specifically, the 2010 spawner density in standard survey areas for the NOC averaged 87 spawners per mile.

Mid Oregon Coast

Since 1977, the Elk River Hatchery production has been tagged for potential use as a PSC indicator stock for the MOC stock aggregate. Age specific ocean abundance forecasts for 2011 are not currently available, but are being developed. The STT has not undertaken a review of the methods used by Oregon Department of Fish and Wildlife (ODFW) staff in developing these abundance forecasts.

The 2010 MOC density from standard survey areas was 92 adult spawners per mile, the highest since 2003 (PFMC 2011, Appendix B, Table B-11). Fall Chinook escapement goals are currently under development for the South Umpqua and Coquille basins of the MOC.

Predictor Description and 2011 Stock Status for South/Local Migrating Chinook

Quantitative abundance predictions are not made for all of these stocks, although an abundance index for Rogue River fall Chinook has been developed. General trends in stock abundance for SOC Chinook stocks are assessed through escapement indices (PFMC 2011, Chapter II, Table II-4 and Figure II-3).

Carcass recoveries in Rogue River index surveys covering a large proportion of the total spawning area were available for 1977-2004. Using Klamath Ocean Harvest Model (KOHM) methodology, these carcass numbers, allocated into age-classes from scale data, were used to estimate the Rogue Ocean Population Index (ROPI) for age-3 to age-5 fish. A linear regression was developed using the escapement estimates (all ages) in year t based on seining at Huntley Park (1976-2004) to predict the ROPI in year $t+1$ (1977-2005). The 2010 Huntley Park escapement estimate and the resulting 2011 ROPI forecast was then scaled to the historical carcass survey-based ROPI. The 2011 ROPI forecast (16,800) consisting of age-3 (9,500), age-4 (6,300) and age-5 (1,000) are based on the average annual age-class strengths of the carcass-based ROPIs from 1991-2005. This data-set was truncated at 1991 because significant harvest restrictions that could affect age structure began that year. The 2011 ROPI is slightly higher than the recent three-year average of 13,400 (Table II-6).

Other Stocks

Fall Chinook escapement goals and forecasts are currently under development for stocks south of the Elk River. These stocks are minor contributors to general season mixed stock ocean fisheries. Standard fall Chinook spawning index escapement data were available for the smaller SOC rivers (Winchuck, Chetco, and Pistol rivers). The average density from standard survey areas was 52 adult spawners per mile, the second highest observed since 2004 (PFMC 2011, Appendix B, Table B-8).

CHINOOK STOCKS NORTH OF CAPE FALCON

Columbia River Fall Chinook

Predictor Description and Past Performance

Columbia River fall Chinook stocks typically form the largest contributing stock group to Council Chinook fisheries north of Cape Falcon. Abundance of these stocks is a major factor in determining impacts of fisheries on weak natural stocks critical to Council area management. Abundance predictions are made for five major fall stock units characterized as being hatchery or natural production, and originating above or below Bonneville Dam. The upriver brights (URB) and lower river wild (LRW) are primarily naturally-produced stocks, although the upriver brights do have a significant hatchery component. The lower river hatchery (LRH) tule, Spring Creek Hatchery (SCH) tule, and mid-Columbia brights (MCB) are primarily hatchery-produced stocks. The MCB include the lower river bright (LRB) stock as a small naturally-produced component. LRB spawn in the mainstem Columbia River near Beacon Rock and are believed to have originated from MCB hatchery strays. The tule stocks generally mature at an earlier age than the bright fall stocks and do not migrate as far north. Minor stocks include the Select Area brights (SAB), a stock originally from the Rogue River.

Preseason forecasts of Columbia River fall Chinook stock abundance, used by the STT to assess the Council's adopted fishery regulations, are based on age-specific and stock-specific forecasts of annual ocean escapement (return to the Columbia River). These forecasts are developed by WDFW and a subgroup of the *U.S. v Oregon* Technical Advisory Committee (TAC). Columbia River return forecast methodologies used for Council management are identical to those used for planning Columbia River fall season fisheries, although minor updates to Council estimates of inriver run size may occur prior to finalization of the inriver fishery plans, based on results of planned ocean fisheries.

The 2011 return of each fall Chinook stock group is forecasted using relationships between successive age groups within a cohort. The database for these relationships was constructed by combining age-specific estimates of escapement and inriver fishery catches for years since 1964 (except for MCB, which started in the 1980's). Typically, only the more recent broods are used in the current predictions. Fall Chinook stock identification in the Columbia River mixed stock fisheries is determined by sampling catch and escapement for CWTs and visual stock identification (VSI). Age composition estimates are based on CWT data and scale reading of fishery and escapement samples, where available. These stock and age data for Columbia River fall Chinook are the basis for the return data presented in the *Review of 2010 Ocean Salmon Fisheries* (Appendix B, Tables B-15 through B-20). The 2010 returns for the five fall Chinook stocks listed in this report may differ somewhat from those provided in the *Review of 2010 Ocean Salmon Fisheries*, since ocean escapement estimates may have been updated after that report was printed.

Performance of the preliminary inriver run size estimation methodology can be assessed, in part, by examining the differences between preseason forecasts and postseason estimates (Table II-7). The recent 10-year average March preliminary preseason forecasts as a percentage of the postseason estimates for the URB, LRW, LRH, SCH, and MCB are 1.01, 1.09, 0.93, 1.06, and 0.94 respectively. None of the stocks had a bias in the recent time series of March preliminary forecasts.

Ocean escapement forecasts developed for the March Council meeting do not take into account variations in marine harvest. The STT combines the initial inriver run size (ocean escapement; Table II-7) with expected Council area fishery harvest levels and stock distribution patterns to produce adjusted ocean escapement forecasts based on the proposed ocean fishing regulations. These revised forecasts are available at the end of the Council preseason planning process in April and are used for preseason fishery modeling in the Columbia River.

2011 Stock Status

The preliminary forecast for 2011 URB fall Chinook ocean escapement is 398,200 adults, about 123 percent of last year's return and about 153 percent of the recent 10-year average of 259,610.

The preliminary forecast for 2011 ocean escapement of ESA-listed Snake River wild fall Chinook is 17,500, about 114 percent of last year's preliminary return estimate.

Ocean escapement of LRW fall Chinook in 2011 is forecast at 12,500 adults, about 115 percent of last year's forecast, and about 81 percent of the recent 10-year average return of 15,360. The forecast is greater than last year's actual return and the spawning escapement goal of 5,700 in the North Fork Lewis River should be achieved this year depending on fishing regulations.

The preliminary forecast for 2011 ocean escapement of LRH fall Chinook is for a return of 133,500 adults, about 130 percent of last year's return and 144 percent of the recent 10-year average of 92,500.

The preliminary ocean escapement forecast of SCH fall Chinook in 2011 is 116,400 adults, about 89 percent of last year's return and 111 percent of the 10-year average of 104,900.

The preliminary forecast for the 2011 ocean escapement of MCB fall Chinook is 100,000 adults, about 127 percent of last year's return and about 110 percent of the recent 10-year average of 90,560.

Washington Coast Chinook

Predictor Description and Past Performance

Council fisheries have negligible impacts on Washington coast Chinook stocks, and except for Willapa Bay fall Chinook, Hoh River Chinook and Quillayute River Chinook, forecast data is unavailable in time for publication of this report; therefore, preseason abundance estimates are not presented. However, abundance estimates are provided for Washington Coastal fall stocks in subsequent preseason fishery impact assessment reports prepared by the STT.

2011 Stock Status

The 2011 Willapa Bay hatchery fall Chinook ocean escapement forecast is 32,476, which is higher than the 2010 prediction of 31,135. The 2011 natural fall Chinook ocean escapement forecast is 4,341, which is higher than last year's prediction of 2,023, but still slightly less than the WDFW spawning escapement goal of 4,350.

For the Hoh River, the 2011 natural spring/summer Chinook ocean escapement forecast is 1,037, above the FMP conservation objective of 900. The natural fall Chinook forecast is 2,880, above the FMP conservation objective of 1,200.

The 2011 Quillayute hatchery spring Chinook ocean escapement forecast is 1,399 and the natural summer/fall Chinook forecast is 8,842 (1,299 summer and 7,543 fall). The FMP conservation objectives are spawning escapements of 1,200 summer Chinook and 3,000 fall Chinook.

Puget Sound Chinook

Run-size expectations for various Puget Sound stock management units are listed in Table I-1. A comparison of preseason and postseason forecasts for recent years is detailed in Table II-8. The STT has not undertaken a review of the methods employed by state and tribal staffs in preparing these abundance forecasts. Methodologies for estimates are described in the annual Puget Sound management reports (starting in 1993, reports are available by Puget Sound management unit, not by individual species). Forecasts for Puget Sound stocks generally assume production is dominated by age-4 adults. Puget Sound Chinook were listed as threatened under the ESA in March 1999. Southern U.S. fisheries that impact Puget Sound Chinook are constrained by terms of a Resource Management Plan (RMP), and are exempted from ESA Section 9 take prohibitions under Limit 6 of the 4(d) rule.

2011 Stock Status

Spring Chinook

Spring Chinook originating in Puget Sound are expected to remain depressed. Runs in the Nooksack, Skagit, White, and Dungeness rivers are of particular concern.

Summer/Fall Chinook

The 2011 preliminary forecast for Puget Sound summer/fall stocks is for a return of 244,377 Chinook, slightly higher than the 2010 preseason forecast of 225,664. The 2011 natural Chinook return forecast of 39,333 (includes supplemental category forecasts) is lower than the 2010 forecast of 42,981.

Since ESA listing and development of the RMP, fishery management for Puget Sound Chinook has changed from an escapement goal basis to the use of stock specific exploitation rates and “critical abundance thresholds.” This new approach is evaluated on an annual basis through the RMP.

SELECTIVE FISHERY CONSIDERATIONS FOR CHINOOK

As the North of Falcon region has moved forward with mass marking of hatchery Chinook salmon stocks, the first mark selective fishery for Chinook salmon in Council waters was implemented from June 12 through June 30, 2010 in the recreational fishery north of Cape Falcon. As in 2010, selective fishing options for non-Indian fisheries are under consideration in the ocean area from Cape Falcon, Oregon to the U. S./Canada border. Observed mark rates on Chinook in 2010 ocean fisheries in this area ranged from 50 to 70 percent. Based on preseason abundance forecasts, the expected mark rate for Chinook in this area for 2011 should be similar to those observed in 2010.

TABLE II-1. Harvest and abundance indices for Sacramento River fall Chinook (SRFC) in thousands of fish.

Year	SRFC Ocean Harvest				River Harvest	Spawning Escapement			Sacramento Index (SI) ^{c/}	Exploitation Rate (%) ^{d/}
	South of Cape Falcon ^{a/}					Natural	Hatchery	Total		
	Troll	Sport	Non-Ret ^{b/}	Total						
1983	248.0	86.7	0.0	334.7	18.1	91.4	18.8	110.2	463.0	76
1984	266.9	87.2	0.0	354.1	26.1	119.5	39.5	159.0	539.1	71
1985	359.5	160.2	0.0	519.7	39.3	209.5	29.9	239.3	798.3	70
1986	620.6	138.4	0.0	759.0	39.4	216.3	23.8	240.1	1,038.5	77
1987	687.9	176.3	0.0	864.2	32.0	174.8	20.3	195.1	1,091.3	82
1988	1,163.2	188.8	0.0	1,352.0	37.3	198.0	29.5	227.5	1,616.8	86
1989	606.5	160.1	0.0	766.5	25.0	126.7	25.9	152.6	944.1	84
1990	508.0	152.4	0.0	660.4	17.2	83.2	21.9	105.1	782.7	87
1991	301.0	90.8	0.0	391.8	26.0 ^{e/}	91.4	27.5	118.9	536.6	78
1992	233.3	70.5	0.0	303.8	13.3 ^{e/}	59.5	22.1	81.5	398.7	80
1993	342.8	115.8	0.0	458.6	27.7 ^{e/}	110.6	26.8	137.4	623.6	78
1994	303.4	165.7	0.0	469.1	28.9 ^{e/}	133.0	32.6	165.6	663.6	75
1995	730.7	390.3	0.0	1,121.1	48.5	253.5	41.8	295.3	1,464.8	80
1996	426.8	157.1	0.0	584.0	49.5	267.1	34.6	301.6	935.1	68
1997	582.4	219.5	0.0	801.9	56.6	279.6	65.2	344.8	1,203.3	71
1998	293.7	117.5	0.0	411.1	69.8 ^{e/}	168.1	77.8	245.9	726.8	66
1999	308.8	78.0	0.0	386.8	68.9 ^{e/}	353.7	46.1	399.8	855.4	53
2000	432.8	154.6	0.0	587.4	59.5 ^{e/}	369.2	48.3	417.5	1,064.4	61
2001	285.8	95.8	0.0	381.6	97.9	537.4	59.4	596.8	1,076.3	45
2002	458.2	194.6	0.0	652.8	89.2 ^{e/}	682.7	87.2	769.9	1,511.9	49
2003	508.3	109.0	0.0	617.3	85.8	413.4	109.6	523.0	1,226.1	57
2004	622.3	214.0	0.0	836.4	47.1	203.5	83.4	286.9	1,170.3	75
2005	370.4	128.0	0.0	498.3	65.0	210.7	185.3	396.0	959.3	59
2006	150.0	107.9	0.0	257.9	45.1	195.1	79.9	275.0	578.1	52
2007	120.0	32.2	0.0	152.1	14.3 ^{e/}	70.0	21.4	91.4	257.8	65
2008	3.2	0.9	0.0	4.1	0.1 ^{e/}	46.9	18.5	65.4	69.6	6
2009	0.0	0.2	0.1	0.3	0.0 ^{e/}	23.3	17.5	40.9	41.2	1
2010 ^{f/}	12.7	12.5	0.3	25.5	2.0 ^{e/}	85.7	39.7	125.4	152.9	18

a/ Ocean harvest for the period September 1 (t-1) through August 31 (t).

b/ Mortalities estimated from non-retention ocean fisheries (e.g., coho-only fisheries, non-retention GSI sampling).

c/ The SI is the sum of (1) SRFC ocean fishery harvest south of Cape Falcon between September 1 and August 31, (2) SRFC impacts from non-retention ocean fisheries when they occur, (3) the recreational harvest of SRFC in the Sacramento River Basin, and (4) the SRFC adult spawner escapement.

d/ Total ocean harvest, non-retention ocean fishery mortalities, and river harvest of SRFC as a percent of the SI.

e/ Estimates derived from CDFG Sacramento River Basin angler survey. Estimates not marked with a footnote are inferred from escapement data and the mean river harvest rate estimate.

f/ Preliminary.

TABLE II-2. Klamath River fall Chinook ocean abundance (thousands), harvest rate, and river run size estimates (thousands) by age.

Year (t)	Ocean Abundance Sept. 1 (t-1)			Annual Ocean Harvest Rate Sept. 1 (t-1) - Aug. 31 (t)		Klamath Basin River Run (t)				
	Age-3	Age-4	Total	Age-3	Age-4	Age-2	Age-3	Age-4	Age-5	Total Adults
1981	493.2	57.0	550.2	0.21	0.53	28.2	64.1	14.4	1.8	80.3
1982	561.1	133.4	694.5	0.30	0.52	39.4	30.1	33.9	2.6	66.6
1983	313.3	114.2	427.5	0.19	0.60	3.8	35.9	20.7	0.9	57.5
1984	157.3	82.8	240.1	0.08	0.38	8.3	21.7	24.4	1.1	47.2
1985	374.8	56.9	431.7	0.11	0.24	69.4	32.9	25.7	5.8	64.4
1986	1,304.4	140.8	1,445.2	0.18	0.46	44.6	162.9	29.8	2.3	195.0
1987	781.2	341.9	1,123.1	0.16	0.43	19.1	89.7	112.6	6.8	209.1
1988	756.3	234.8	991.0	0.20	0.39	24.1	101.2	86.5	3.9	191.6
1989	369.8	177.2	547.1	0.15	0.36	9.1	50.4	69.6	4.3	124.3
1990	176.1	104.0	280.1	0.30	0.55	4.4	11.6	22.9	1.3	35.9
1991	69.4	37.2	106.6	0.03	0.18	1.8	10.0	21.6	1.1	32.7
1992	39.5	28.2	67.7	0.02	0.07	13.7	6.9	18.8	1.0	26.7
1993	168.5	15.0	183.5	0.05	0.16	7.6	48.3	8.2	0.7	57.2
1994	119.9	41.7	161.6	0.03	0.09	14.4	37.0	26.0	1.0	64.0
1995	784.3	28.7	813.0	0.04	0.14	22.8	201.9	18.3	2.6	222.8
1996	192.3	225.5	417.8	0.05	0.16	9.5	38.8	136.7	0.3	175.8
1997	140.2	62.8	203.0	0.01	0.06	8.0	35.0	44.2	4.6	83.7
1998	154.8	44.7	199.5	0.00	0.09	4.6	59.2	29.7	1.7	90.6
1999	129.1	30.5	159.5	0.02	0.09	19.2	29.2	20.5	1.3	51.0
2000	617.1	44.2	661.3	0.06	0.10	10.2	187.1	30.5	0.5	218.1
2001	356.1	133.8	489.9	0.03	0.09	11.3	99.1	88.2	0.2	187.4
2002	513.6	98.9	612.5	0.02	0.15	9.2	94.6	62.5	3.7	160.8
2003	400.2	192.2	592.4	0.08	0.21	3.8	94.3	96.8	0.9	191.9
2004	159.6	105.1	264.6	0.12	0.34	9.7	33.2	40.7	5.3	79.2
2005	190.0	38.1	228.1	0.02	0.20	2.3	43.8	17.5	3.9	65.2
2006	90.6	63.4	154.0	0.01	0.10	26.9	18.5	41.6	1.3	61.4
2007	376.8	33.6	410.5	0.06	0.21	1.7	113.7	16.8	1.6	132.1
2008	68.0	81.4	149.4	0.00	0.10	25.2	18.6	50.2	1.7	70.6
2009	248.2 ^{a/}	21.1	269.3	0.00 ^{a/}	0.00	11.9	78.6	16.4	5.6	100.6
2010	217.1 ^{b/}	66.5 ^{a/}	283.5	NA ^{c/}	0.04 ^{a/}	16.7	46.2	44.4	0.4	91.0

a/ Preliminary: incomplete cohort data (age-5 unavailable).

b/ Preliminary: incomplete cohort data (age-4 and age-5 unavailable).

c/ Not estimated: incomplete cohort data (age-4 and age-5 unavailable).

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 1 of 4)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
	Age-3		
1985	113,000	276,000	0.41
1986	426,000 ^{b/}	1,304,409	0.33
1987	511,800	781,198	0.66
1988	370,800	756,261	0.49
1989	450,600	369,828	1.22
1990	479,000	176,133	2.72
1991	176,200	69,424	2.54
1992	50,000	39,502	1.27
1993	294,400	168,473	1.75
1994	138,000	119,913	1.15
1995	269,000	784,260	0.34
1996	479,800	192,272	2.50
1997	224,600	140,153	1.60
1998	176,000	154,799	1.14
1999	84,800	129,066	0.66
2000	349,600	617,098	0.57
2001	187,200	356,128	0.53
2002	209,000	513,561	0.41
2003	171,300	400,242	0.43
2004	72,100	159,560	0.45
2005	185,700	189,976	0.98
2006	44,100	90,606	0.49
2007	515,400	376,841	1.37
2008	31,600	67,993	0.46
2009	474,900	248,170	1.91
2010 ^{c/}	223,400	217,062	1.03
2011	304,600	-	-

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 2 of 4)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
	Age-4		
1985	56,875	57,500	0.99
1986	66,250	140,823	0.47
1987	206,125	341,875	0.60
1988	186,375	234,772	0.79
1989	215,500	177,245	1.22
1990	50,125	103,951	0.48
1991	44,625	37,172	1.20
1992	44,750	28,169	1.59
1993	39,125	15,037	2.60
1994	86,125	41,736	2.06
1995	47,000	28,725	1.64
1996	268,500	225,521	1.19
1997	53,875	62,820	0.86
1998	46,000	44,733	1.03
1999	78,750	30,456	2.59
2000	38,875	44,176	0.88
2001	247,000	133,801	1.85
2002	143,800	98,928	1.45
2003	132,400	192,156	0.69
2004	134,500	105,051	1.28
2005	48,900	38,079	1.28
2006	63,700	63,383	1.00
2007	26,100	33,615	0.78
2008	157,200	81,366	1.93
2009	25,200	21,118	1.19
2010 ^{c/}	106,300	66,452	1.60
2011	61,600	-	-

TABLE II-3. Comparisons of preseason forecasts and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 3 of 4)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
	Age-5		
1985	NA	11,113	NA
1986	NA	6,376	NA
1987	5,250	19,414	0.27
1988	13,250	14,632	0.91
1989	10,125	9,612	1.05
1990	7,625	7,767	0.98
1991	1,500	2,774	0.54
1992	1,250	1,444	0.87
1993	1,125	1,759	0.64
1994	500	1,468	0.34
1995	2,000	3,805	0.53
1996	1,125	787	1.43
1997	7,875	8,859	0.89
1998	3,250	2,382	1.36
1999	2,000	2,106	0.95
2000	1,375	1,051	1.31
2001	1,250	258	4.84
2002	9,700	6,933	1.40
2003	6,500	1,915	3.39
2004	9,700	17,170	0.56
2005	5,200	6,857	0.76
2006	2,200	5,236	0.42
2007	4,700	2,911	1.61
2008	1,900	2,900	0.66
2009	5,600	7,059	0.79
2010 ^{c/}	1,800	513	3.51
2011	5,000	-	-

TABLE II-3. Comparisons of preseason forecast and postseason estimates for ocean abundance of adult Klamath River fall Chinook. (Page 4 of 4)

Year (t)	Preseason Forecast ^{a/}	Postseason Estimate	Pre/Postseason
	Sept. 1 (t-1)	Sept. 1 (t-1)	
Total Adults			
1985	169,875 ^{d/}	344,613	0.49
1986	492,250 ^{d/}	1,451,608	0.34
1987	723,175	1,142,487	0.63
1988	570,425	1,005,665	0.57
1989	676,225	556,685	1.21
1990	536,750	287,851	1.86
1991	222,325	109,370	2.03
1992	96,000	69,115	1.39
1993	334,650	185,269	1.81
1994	224,625	163,117	1.38
1995	318,000	816,790	0.39
1996	749,425	418,580	1.79
1997	286,350	211,832	1.35
1998	225,250	201,914	1.12
1999	165,550	161,628	1.02
2000	389,850	662,325	0.59
2001	435,450	490,187	0.89
2002	362,500	619,422	0.59
2003	310,200	594,313	0.52
2004	216,300	281,781	0.77
2005	239,800	234,912	1.02
2006	110,000	159,225	0.69
2007	546,200	413,367	1.32
2008	190,700	152,259	1.25
2009	505,700	276,347	1.83
2010 ^{c/}	331,500	284,027	1.17
2011	371,100	-	-

a/ Original preseason forecasts for years 1985-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the assumed May 1 (t) number by the Sept. 1 (t-1) through May 1 (t) survival rate in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ A scalar of 0.75 was applied to the jack count to produce the forecast because, (1) most jacks returned to the Trinity River, and (2) the jack count was outside the database range.

c/ Postseason estimates are preliminary.

d/ Does not include age-5 adults.

TABLE II-4. Summary of management objectives and predictor performance for Klamath River fall Chinook.

Year(t)	Preseason Ocean Abundance Forecast ^{a/}		Postseason Ocean Abundance Estimate		Preseason Age-4 Harvest Rate Forecast ^{b/}		Postseason Age-4 Harvest Rate Estimate ^{c/}		Preseason Adult Harvest Forecast		Postseason Adult Harvest Estimate	
	Sept. 1 (t-1)		Sept. 1 (t-1)		Ocean	River	Ocean	River	Ocean	River	Ocean	River
1986	426,000	66,250	1,304,409	140,823	0.28	0.50	0.46	0.67	72,000	37,700	301,999	46,154
1987	511,800	206,125	781,198	341,875	0.28	0.53	0.43	0.44	121,200	78,200	277,224	73,265
1988	370,800	186,375	756,261	234,772	0.31	0.53	0.39	0.52	114,100	65,400	253,905	73,854
1989	450,600	215,500	369,828	177,245	0.30	0.49	0.36	0.70	128,100	67,600	125,117	54,340
1990	479,000	50,125	176,133	103,951	0.30	0.49	0.55	0.36	85,100	31,200	114,786	11,459
1991	176,200	44,625	69,424	37,172	0.13	0.28	0.18	0.45	16,700	12,800	9,872	13,581
1992	50,000	44,750	39,502	28,169	0.06	0.15	0.07	0.27	4,200	4,200	3,142	6,787
1993	294,400	39,125	168,473	15,037	0.12	0.43	0.16	0.49	20,100	22,500	11,355	12,808
1994	138,000	86,125	119,913	41,736	0.07	0.20	0.09	0.29	10,400	14,300	7,961	13,524
1995	269,000	47,000	784,260	28,725	0.07	0.32	0.14	0.19	13,500	18,500	32,233	21,637
1996	479,800	268,500	192,272	225,521	0.17	0.66	0.16	0.39	88,400	129,100	45,155	69,241
1997	224,600	53,875	140,153	62,820	0.10	0.43	0.06	0.26	17,600	26,500	8,656	17,764
1998	176,000	46,000	154,799	44,733	0.07	0.29	0.09	0.30	10,200	14,800	4,891	17,897
1999	84,800	78,750	129,066	30,456	0.10	0.28	0.09	0.45	12,300	18,100	5,116	16,942
2000	349,600	38,875	617,098	44,176	0.11	0.53	0.10	0.25	24,000	32,400	42,050	35,066
2001	187,200	247,000	356,128	133,801	0.14	0.61	0.09	0.29	45,600	105,300	21,747	50,780
2002	209,000	143,800	513,561	98,928	0.13	0.57	0.15	0.26	30,000	70,900	28,895	35,069
2003	171,300	132,400	400,242	192,156	0.16	0.50	0.21	0.28	30,600	52,200	70,684	39,715
2004	72,100	134,500	159,560	105,051	0.15	0.38	0.34	0.48	26,500	35,800	63,885	29,807
2005	185,700	48,900	189,976	38,079	0.08	0.16	0.20	0.19	7,100	9,600	12,826	10,001
2006	44,100	63,700	90,606	63,383	0.11	0.23	0.10	0.18	10,000	10,000	10,401	10,345
2007	515,400	26,100	376,841	33,615	0.16	0.63	0.21	0.56	30,200	51,400	30,244	33,884
2008	31,600	157,200	67,993	81,366	0.02	0.43	0.10	0.38	4,500	49,500	8,679	24,180
2009	474,900	25,200	248,170	21,118	0.00	0.57	0.00	0.40	100	61,700	52	34,040
2010 ^{d/}	223,400	106,300	217,062	66,452	0.12	0.49	0.04	0.40	22,600	46,600	4,235	33,031
2011	304,600	61,600	-	-	-	-	-	-	-	-	-	-

a/ Original preseason forecasts for years 1986-2001 were for May 1 (t); converted to Sept. 1 (t-1) forecasts by dividing the May 1 (t) number by the assumed Sept. 1 (t-1) through May 1 (t) survival rate assumed in those years: 0.5 age-3, 0.8 age-4, 0.8 age-5.

b/ Ocean harvest rate forecast is the fraction of the predicted ocean abundance expected to be harvested Sept. 1 (t-1) through August 31(t). River harvest rate forecast is the fraction of the predicted river run expected to be harvested in river fisheries. Original ocean harvest rate forecasts for year (t), 1986-2001, were based on a May 1 (t) ocean abundance denominator; converted to Sept. 1 (t-1) abundance denominator by multiplying former values by 0.8 (the assumed age-4 survival rate between Sept. 1 (t-1) and May 1 (t) in those years).

c/ Ocean harvest rate is the fraction of the postseason ocean abundance harvested Sept. 1 (t-1) through August 31 (t). River harvest rate is the fraction of the river run harvested by river fisheries.

d/ Postseason estimates are preliminary.

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 1 of 4)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))						River Fisheries (t)			
	KMZ			North of	South of	Subtotal	Ocean Total	Net	Sport	Total
	Troll	Sport	Subtotal	KMZ	KMZ					
HARVEST (numbers of fish)										
Age-3										
1986	35,632	4,876	40,508	73,777	122,913	196,690	237,198	8,100	18,100	26,200
1987	17,240	5,083	22,323	43,439	56,378	99,817	122,140	11,400	11,400	22,800
1988	15,999	5,165	21,164	24,317	107,971	132,288	153,452	12,500	15,600	28,100
1989	6,456	11,783	18,239	15,315	23,729	39,044	57,283	2,700	900	3,600
1990	81	4,357	4,438	36,579	11,006	47,585	52,023	1,300	1,400	2,700
1991	0	1,022	1,022	344	810	1,154	2,176	2,123	1,277	3,400
1992	0	0	0	972	0	972	972	970	251	1,221
1993	0	822	822	833	6,424	7,257	8,079	5,426	2,917	8,343
1994	42	604	646	0	3,387	3,387	4,033	4,543	965	5,508
1995	0	999	999	12,213	14,810	27,023	28,022	11,840	5,536	17,376
1996	0	0	0	0	9,314	9,314	9,314	12,363	3,661	16,024
1997	0	232	232	620	1,215	1,835	2,067	2,166	2,736	4,902
1998	0	6	6	298	466	764	770	2,231	5,781	8,012
1999	63	180	243	1,262	433	1,695	1,938	4,981	1,748	6,729
2000	404	3,282	3,686	8,604	25,203	33,807	37,493	22,458	4,893	27,351
2001	113	105	218	2,749	6,082	8,831	9,049	17,885	7,294	25,179
2002	220	784	1,004	1,501	9,915	11,416	12,420	11,734	6,258	17,992
2003	173	679	852	1,885	27,309	29,194	30,046	6,996	5,061	12,057
2004	402	971	1,373	9,719	7,331	17,050	18,423	4,679	2,051	6,730
2005	0	568	568	619	2,381	3,000	3,568	4,394	1,641	6,035
2006	0	477	477	32	341	373	850	2,388	13	2,401
2007	770	8,099	8,869	4,193	9,365	13,558	22,427	17,543	5,734	23,277
2008	0	0	0	0	0	0	0	3,225	608	3,833
2009 ^{al}	0	52	52	0	0	0	52	19,820	4,715	24,535
2010 ^{al}	84	23	107	0	1,335	1,335	1,442	13,190	1,884	15,074

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 2 of 4)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))						River Fisheries (t)			
	KMZ			North of	South of	Ocean Total	Net	Sport	Total	
	Troll	Sport	Subtotal	KMZ	KMZ					
HARVEST (numbers of fish)										
Age-4										
1986	7,745	1,113	8,858	23,486	31,913	55,399	64,257	17,000	2,900	19,900
1987	21,736	4,427	26,163	70,645	48,832	119,477	145,640	41,000	8,500	49,500
1988	11,870	3,596	15,466	26,381	50,296	76,677	92,143	38,600	6,200	44,800
1989	6,064	9,735	15,799	32,116	16,608	48,724	64,523	41,000	7,700	48,700
1990	3,997	2,919	6,916	39,627	10,624	50,251	57,167	6,000	2,200	8,200
1991	0	1,001	1,001	1,513	4,135	5,648	6,649	7,593	2,016	9,609
1992	171	55	226	1,783	12	1,795	2,021	4,360	723	5,083
1993	0	0	0	849	1,616	2,465	2,465	3,786	243	4,029
1994	0	1,124	1,124	1,168	1,499	2,667	3,791	6,666	818	7,484
1995	0	242	242	1,879	1,771	3,650	3,892	2,957	480	3,437
1996	773	3,464	4,237	10,337	20,741	31,078	35,315	43,959	9,080	53,039
1997	3	172	175	463	2,994	3,457	3,632	8,734	2,586	11,320
1998	0	105	105	3,942	0	3,942	4,047	7,164	1,822	8,986
1999	15	381	396	1,657	696	2,353	2,749	8,789	494	9,283
2000	117	895	1,012	2,327	1,076	3,403	4,415	6,733	756	7,489
2001	1,312	1,604	2,916	5,819	3,926	9,745	12,661	20,759	4,819	25,578
2002	1,938	827	2,765	2,811	9,416	12,227	14,992	11,929	4,063	15,992
2003	834	918	1,752	7,855	30,007	37,862	39,614	22,754	4,592	27,346
2004	1,421	1,215	2,636	11,504	21,949	33,453	36,089	17,623	1,751	19,374
2005	247	317	564	5,243	1,909	7,152	7,716	3,048	304	3,352
2006	196	725	921	4,192	985	5,177	6,098	7,569	42	7,611
2007	270	2,336	2,606	1,991	2,472	4,463	7,069	8,987	502	9,489
2008	6,376	1,105	7,481	546	113	659	8,140	17,891	1,260	19,151
2009	0	0	0	0	0	0	0	5,831	706	6,537
2010 ^{av}	37	114	151	924	1,547	2,471	2,622	16,682	1,134	17,816

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 3 of 4)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))						River Fisheries (t)			
	KMZ			North of	South of	Subtotal	Ocean Total	Net	Sport	Total
	Troll	Sport	Subtotal	KMZ	KMZ					
HARVEST RATE^{b/}										
Age-3										
1986	0.03	0.00	0.03	0.06	0.09	0.15	0.18	0.05	0.11	0.16
1987	0.02	0.01	0.03	0.06	0.07	0.13	0.16	0.13	0.13	0.25
1988	0.02	0.01	0.03	0.03	0.14	0.17	0.20	0.12	0.15	0.28
1989	0.02	0.03	0.05	0.04	0.06	0.11	0.15	0.05	0.02	0.07
1990	0.00	0.02	0.03	0.21	0.06	0.27	0.30	0.11	0.12	0.23
1991	0.00	0.01	0.01	0.00	0.01	0.02	0.03	0.21	0.13	0.34
1992	0.00	0.00	0.00	0.02	0.00	0.02	0.02	0.14	0.04	0.18
1993	0.00	0.00	0.00	0.00	0.04	0.04	0.05	0.11	0.06	0.17
1994	0.00	0.01	0.01	0.00	0.03	0.03	0.03	0.12	0.03	0.15
1995	0.00	0.00	0.00	0.02	0.02	0.03	0.04	0.06	0.03	0.09
1996	0.00	0.00	0.00	0.00	0.05	0.05	0.05	0.32	0.09	0.41
1997	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.06	0.08	0.14
1998	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.14
1999	0.00	0.00	0.00	0.01	0.00	0.01	0.02	0.17	0.06	0.23
2000	0.00	0.01	0.01	0.01	0.04	0.05	0.06	0.12	0.03	0.15
2001	0.00	0.00	0.00	0.01	0.02	0.02	0.03	0.18	0.07	0.25
2002	0.00	0.00	0.00	0.00	0.02	0.02	0.02	0.12	0.07	0.19
2003	0.00	0.00	0.00	0.00	0.07	0.07	0.08	0.07	0.05	0.13
2004	0.00	0.01	0.01	0.06	0.05	0.11	0.12	0.14	0.06	0.20
2005	0.00	0.00	0.00	0.00	0.01	0.02	0.02	0.10	0.04	0.14
2006	0.00	0.01	0.01	0.00	0.00	0.00	0.01	0.13	0.00	0.13
2007	0.00	0.02	0.02	0.01	0.02	0.04	0.06	0.15	0.05	0.20
2008	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.03	0.21
2009 ^{al}	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.06	0.31
2010 ^{al}	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.29	0.04	0.33

TABLE II-5. Harvest levels and rates of age-3 and age-4 Klamath River fall Chinook. (Page 4 of 4)

Year (t)	Ocean Fisheries (Sept. 1 (t-1) - Aug. 31 (t))						River Fisheries (t)			
	KMZ			North of	South of	Ocean Total	Net	Sport	Total	
	Troll	Sport	Subtotal	KMZ	KMZ					Subtotal
	HARVEST RATE^{b/}									
Age-4										
1986	0.05	0.01	0.06	0.17	0.23	0.39	0.46	0.57	0.10	0.67
1987	0.06	0.01	0.08	0.21	0.14	0.35	0.43	0.36	0.08	0.44
1988	0.05	0.02	0.07	0.11	0.21	0.33	0.39	0.45	0.07	0.52
1989	0.03	0.05	0.09	0.18	0.09	0.27	0.36	0.59	0.11	0.70
1990	0.04	0.03	0.07	0.38	0.10	0.48	0.55	0.26	0.10	0.36
1991	0.00	0.03	0.03	0.04	0.11	0.15	0.18	0.35	0.09	0.45
1992	0.01	0.00	0.01	0.06	0.00	0.06	0.07	0.23	0.04	0.27
1993	0.00	0.00	0.00	0.06	0.11	0.16	0.16	0.46	0.03	0.49
1994	0.00	0.03	0.03	0.03	0.04	0.06	0.09	0.26	0.03	0.29
1995	0.00	0.01	0.01	0.07	0.06	0.13	0.14	0.16	0.03	0.19
1996	0.00	0.02	0.02	0.05	0.09	0.14	0.16	0.32	0.07	0.39
1997	0.00	0.00	0.00	0.01	0.05	0.06	0.06	0.20	0.06	0.26
1998	0.00	0.00	0.00	0.09	0.00	0.09	0.09	0.24	0.06	0.30
1999	0.00	0.01	0.01	0.05	0.02	0.08	0.09	0.43	0.02	0.45
2000	0.00	0.02	0.02	0.05	0.02	0.08	0.10	0.22	0.02	0.25
2001	0.01	0.01	0.02	0.04	0.03	0.07	0.09	0.24	0.05	0.29
2002	0.02	0.01	0.03	0.03	0.10	0.12	0.15	0.19	0.06	0.26
2003	0.00	0.00	0.01	0.04	0.16	0.20	0.21	0.24	0.05	0.28
2004	0.01	0.01	0.03	0.11	0.21	0.32	0.34	0.43	0.04	0.48
2005	0.01	0.01	0.01	0.14	0.05	0.19	0.20	0.17	0.02	0.19
2006	0.00	0.01	0.01	0.07	0.02	0.08	0.10	0.18	0.00	0.18
2007	0.01	0.07	0.08	0.06	0.07	0.13	0.21	0.53	0.03	0.56
2008	0.08	0.01	0.09	0.01	0.00	0.01	0.10	0.36	0.03	0.38
2009	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.36	0.04	0.40
2010 ^{a/}	0.00	0.00	0.00	0.01	0.02	0.04	0.04	0.38	0.03	0.40

a/ Preliminary (incomplete cohort).

b/ Ocean harvest rates are the fraction of Sept. 1 (t-1) ocean abundance harvested in these fisheries. River harvest rates are the fraction of the river run (t) harvested in these fisheries.

TABLE II-6. Rogue River fall Chinook inriver run and ocean population indices.

Return Year	Inriver Run Index in Thousands of Fish ^{a/}					Ocean Impact Rate by Age ^{b/}		Rogue Ocean Population Index (ROPI) in Thousands of Fish ^{c/}			
	Age-2	Age-3	Age-4	Age-5	Total ^{d/}	Age-3	Age-4-5	Age-3	Age-4	Age-5	Total
1980	0.4	0.2	0.9	0.6	2.1	0.23	0.55	5.2	4.0	1.4	10.6
1981	1.1	3.3	1.0	0.3	5.7	0.21	0.53	9.2	3.0	0.7	12.9
1982	0.7	1.3	1.3	0.1	3.4	0.30	0.52	9.8	2.9	0.3	13.0
1983	0.3	1.1	1.5	0.0	2.9	0.19	0.60	8.6	4.4	0.1	13.1
1984	0.4	1.2	1.8	0.1	3.5	0.08	0.38	9.9	4.7	0.2	14.8
1985	2.5	1.3	3.5	0.6	7.9	0.11	0.25	9.7	6.3	0.9	16.9
1986	3.1	12.5	2.3	0.5	18.4	0.18	0.46	71.3	5.9	1.0	78.2
1987	2.6	7.8	18.1	0.4	28.9	0.16	0.43	80.3	36.3	0.6	117.2
1988	1.4	4.8	25.2	1.5	32.9	0.20	0.39	17.3	47.9	2.5	67.7
1989	0.5	1.3	4.0	2.0	7.8	0.15	0.36	8.4	7.2	3.2	18.8
1990	0.0	0.3	1.4	0.2	1.9	0.30	0.55	6.0	4.7	0.5	11.2
1991	0.2	0.4	1.9	0.5	3.0	0.03	0.18	3.5	3.2	0.6	7.3
1992	0.5	0.3	1.5	0.5	2.8	0.02	0.07	4.4	2.4	0.6	7.4
1993	0.3	3.5	1.5	0.5	5.8	0.05	0.16	16.1	3.2	0.6	19.9
1994	0.5	0.8	5.8	0.9	8.0	0.03	0.09	3.0	9.5	0.9	13.4
1995	0.2	0.6	1.4	2.0	4.2	0.04	0.13	4.3	1.7	2.3	8.3
1996	0.1	0.4	1.8	0.1	2.4	0.05	0.16	2.4	2.8	0.1	5.3
1997	0.1	0.3	1.0	0.3	1.7	0.01	0.06	5.2	1.5	0.3	7.0
1998	0.0	0.5	2.8	0.3	3.6	0.00	0.09	3.8	3.9	0.3	8.0
1999	0.2	0.3	1.6	0.5	2.6	0.01	0.09	1.5	2.7	0.6	4.8
2000	0.2	2.0	0.8	0.6	3.6	0.06	0.10	9.9	0.9	0.6	11.4
2001	0.8	2.3	4.2	0.0	7.3	0.03	0.09	14.1	5.9	0.0	20.0
2002	0.9	4.0	7.1	0.8	12.7	0.02	0.15	32.2	9.1	0.9	42.2
2003	0.9	2.3	12.0	0.4	15.6	0.08	0.21	14.4	22.1	0.5	37.0
2004	0.4	0.6	4.9	2.9	8.8	0.12	0.34	3.9	9.7	4.4	18.0
2005 ^{f/}	NA	NA	NA	NA	NA	0.02	0.20	7.6	5.0	0.8	13.4
2006 ^{f/}	NA	NA	NA	NA	NA	0.01	0.10	4.9	3.2	0.5	8.6
2007 ^{f/}	NA	NA	NA	NA	NA	0.06	0.21	5.8	3.8	0.6	10.2
2008 ^{f/}	NA	NA	NA	NA	NA	0.00	0.10	6.9	4.6	0.7	12.2
2009 ^{f/}	NA	NA	NA	NA	NA	0.00	0.00	6.1 ^{e/}	4.0	0.7	10.7 ^{e/}
2010 ^{f/}	NA	NA	NA	NA	NA	NA	0.04	9.8 ^{e/}	6.5 ^{e/}	1.1	17.3 ^{e/}
2011 ^{f/}	NA	NA	NA	NA	NA	-	-	9.5 ^{g/}	6.3 ^{g/}	1.0 ^{g/}	16.8 ^{g/}

a/ Index based on carcass counts in spawning survey index areas. Carcass counts in 1978, 1979, and 1980 adjusted for prespawning mortality. Age composition developed from carcass scale sampling.

b/ Exploitation rates since 1981 are based on Klamath River fall Chinook cohort analysis, 1977-1980 based on 1981-1983 average.

c/ Based on cohort reconstruction methods. Index values for 2011 predicted from regression equations; postseason estimates are not available.

d/ Excludes age-6 fish.

e/ Preliminary, complete cohort not available, mean maturity rate used to derive estimate.

f/ Spawning surveys were discontinued 2005.

g/ Preseason forecast.

TABLE II-7. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 1 of 3)

Year	March Preseason Forecast ^{a/}	April STT Modeled Forecast ^{b/}	Postseason Return	March Pre/Postseason	April Pre/Postseason
URB					
1984	90.10	93.00	131.40	0.69	0.71
1985	159.10	159.10	196.40	0.81	0.81
1986	285.90	286.10	281.60	1.02	1.02
1987	436.40	436.40	420.70	1.04	1.04
1988	450.70	446.50	339.90	1.33	1.31
1989	234.00	231.80	261.30	0.90	0.89
1990	127.20	126.90	153.60	0.83	0.83
1991	88.80	88.90	103.30	0.86	0.86
1992	68.40	66.30	81.00	0.84	0.82
1993	84.50	82.70	102.90	0.82	0.80
1994	85.40	94.70	132.80	0.64	0.71
1995	103.70	125.00	106.50	0.97	1.17
1996	88.90	94.20	143.20	0.62	0.66
1997	166.40	158.00	161.70	1.03	0.98
1998	150.80	141.80	142.30	1.06	1.00
1999	147.50	102.10	166.10	0.89	0.61
2000	171.10	208.20	155.70	1.10	1.34
2001	127.20	132.70	232.60	0.55	0.57
2002	281.00	273.80	276.90	1.01	0.99
2003	280.40	253.20	373.20	0.75	0.68
2004	292.20	287.00	367.90	0.79	0.78
2005	352.20	354.60	268.70	1.31	1.32
2006	253.90	249.10	230.40	1.10	1.08
2007	182.40	185.20	112.60	1.62	1.64
2008	162.50	165.90	196.90	0.83	0.84
2009	259.90	269.80	212.00	1.23	1.27
2010 ^{c/}	310.80	319.10	324.90	0.96	0.98
2011	398.20	-	-	-	-
LRW					
1984	16.70	NA	13.30	1.26	NA
1985	12.90	NA	13.30	0.97	NA
1986	15.70	NA	24.50	0.64	NA
1987	29.20	NA	37.90	0.77	NA
1988	43.30	42.10	41.70	1.04	1.01
1989	27.30	26.90	38.60	0.71	0.70
1990	23.70	23.40	20.30	1.17	1.15
1991	12.70	12.70	19.80	0.64	0.64
1992	17.40	16.70	12.50	1.39	1.34
1993	12.50	11.90	13.30	0.94	0.89
1994	14.70	13.20	12.20	1.20	1.08
1995	12.40	11.50	16.00	0.78	0.72
1996	8.80	8.10	14.60	0.60	0.55
1997	7.50	7.20	12.30	0.61	0.59
1998	8.10	7.00	7.30	1.11	0.96
1999	2.60	2.50	3.30	0.79	0.76
2000	3.50	2.70	10.20	0.34	0.26
2001	16.70	18.50	15.70	1.06	1.18
2002	18.70	18.30	24.90	0.75	0.73
2003	24.60	23.40	26.00	0.95	0.90
2004	24.10	24.20	22.30	1.08	1.09
2005	20.20	21.40	16.80	1.20	1.27
2006	16.60	16.60	18.10	0.92	0.92
2007	10.10	10.00	4.30	2.35	2.33
2008	3.80	3.80	7.10	0.54	0.54
2009	8.50	8.60	7.50	1.13	1.15
2010 ^{c/}	9.70	10.00	10.90	0.89	0.92
2011	12.50	-	-	-	-

TABLE II-7. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 2 of 3)

Year	March Preseason Forecast ^{a/}	April STT Modeled Forecast ^{b/}	Postseason Return	March Pre/Postseason	April Pre/Postseason
LRH					
1984	70.40	89.00	102.40	0.69	0.87
1985	81.50	86.70	111.00	0.73	0.78
1986	171.60	173.90	154.80	1.11	1.12
1987	294.90	298.70	344.10	0.86	0.87
1988	267.70	246.50	309.90	0.86	0.80
1989	104.90	97.50	130.90	0.80	0.74
1990	68.50	65.50	60.00	1.14	1.09
1991	71.40	73.10	62.70	1.14	1.17
1992	113.20	121.50	62.60	1.81	1.94
1993	79.30	77.70	52.30	1.52	1.49
1994	36.10	46.50	53.60	0.67	0.87
1995	35.80	42.40	46.40	0.77	0.91
1996	37.70	48.30	75.50	0.50	0.64
1997	54.20	68.70	57.40	0.94	1.20
1998	19.20	22.50	45.30	0.42	0.50
1999	34.80	38.20	40.00	0.87	0.96
2000	23.70	26.40	27.00	0.88	0.98
2001	32.20	30.50	94.30	0.34	0.32
2002	137.60	133.00	156.40	0.88	0.85
2003	115.90	116.90	155.00	0.75	0.75
2004	77.10	79.00	108.90	0.71	0.73
2005	74.10	78.44	78.30	0.95	1.00
2006	55.80	57.50	58.30	0.96	0.99
2007	54.90	54.40	32.70	1.68	1.66
2008	59.00	55.90	60.30	0.98	0.93
2009	88.80	88.20	76.70	1.16	1.15
2010 ^{c/}	90.60	85.60	103.00	0.88	0.83
2011	133.50	-	-	-	-
SCH					
1984	21.30	27.00	47.50	0.45	0.57
1985	34.90	37.10	33.20	1.05	1.12
1986	16.00	16.20	16.60	0.96	0.98
1987	9.10	9.20	9.10	1.00	1.01
1988	6.50	5.90	12.00	0.54	0.49
1989	29.50	23.00	26.80	1.10	0.86
1990	27.30	23.70	18.90	1.44	1.25
1991	56.30	61.40	52.40	1.07	1.17
1992	40.90	41.30	29.50	1.39	1.40
1993	19.90	18.20	16.80	1.18	1.08
1994	20.20	28.90	18.50	1.09	1.56
1995	17.50	22.50	33.80	0.52	0.67
1996	27.60	35.40	33.10	0.83	1.07
1997	21.90	25.70	27.40	0.80	0.94
1998	14.20	14.20	20.20	0.70	0.70
1999	65.80	61.00	50.20	1.31	1.22
2000	21.90	26.90	20.50	1.07	1.31
2001	56.60	61.90	125.00	0.45	0.50
2002	144.40	136.00	160.80	0.90	0.85
2003	96.90	101.90	180.60	0.54	0.56
2004	138.00	150.00	175.30	0.79	0.86
2005	114.10	115.79	93.10	1.23	1.24
2006	50.00	51.80	27.90	1.79	1.86
2007	21.80	21.30	14.60	1.49	1.46
2008	87.20	86.20	91.90	0.95	0.94
2009	59.30	56.50	49.00	1.21	1.15
2010 ^{c/}	169.00	162.90	130.80	1.29	1.25
2011	116.40	-	-	-	-

TABLE II-7. Predicted and postseason returns of Columbia River adult fall Chinook in thousands of fish. (Page 3 of 3)

Year	March Preseason Forecast ^{a/}	April STT Modeled Forecast ^{b/}	Postseason Return	March Pre/Postseason	April Pre/Postseason
MCB					
1990	69.50	69.30	58.90	1.18	1.18
1991	48.40	48.50	35.40	1.37	1.37
1992	42.50	40.70	31.10	1.37	1.31
1993	33.00	32.30	27.50	1.20	1.17
1994	23.90	26.70	33.70	0.71	0.79
1995	25.00	30.00	34.20	0.73	0.88
1996	40.80	43.20	59.70	0.68	0.72
1997	72.10	61.90	59.00	1.22	1.05
1998	47.80	44.90	36.80	1.30	1.22
1999	38.30	27.70	50.70	0.76	0.55
2000	50.60	61.60	36.80	1.38	1.67
2001	43.50	45.30	76.40	0.57	0.59
2002	96.20	91.80	108.40	0.89	0.85
2003	104.80	94.60	150.20	0.70	0.63
2004	90.40	88.80	117.60	0.77	0.76
2005	89.40	89.73	98.00	0.91	0.92
2006	88.30	86.60	80.40	1.10	1.08
2007	68.00	69.10	46.90	1.45	1.47
2008	54.00	55.10	75.50	0.72	0.73
2009	94.40	97.90	73.10	1.29	1.34
2010 ^{c/}	79.00	74.60	79.00	1.00	0.94
2011	100.00	-	-	-	-

a/ March preseason forecasts are ocean escapements based on terminal run size and stock-specific cohort relationships affected by the historical "normal" ocean fisheries, generally between 1979 and the most recent adequately complete broods.

b/ STT modeled forecasts adjust March preseason forecasts for Council-adopted ocean regulations each year and should provide a more accurate estimate of expected ocean escapement.

c/ Postseason estimates are preliminary.

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 1 of 4)

Year	Nooksack-Samish			East Sound Bay			Skagit			Skagit		
	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason
	Hatchery and Natural			Hatchery			Hatchery			Natural		
1993	50.4	32.3	1.53	3.2	3.8	0.84	1.0	1.4	0.71	14.0	6.9	2.00
1994	46.6	28.1	1.66	3.2	0.7	4.00	1.3	5.5	0.30	8.4	5.9	1.27
1995	38.5	22.3	1.73	3.5	0.2	17.50	1.6	3.4	0.48	5.0	9.2	0.52
1996	27.0	29.2	0.92	1.7	0.5	2.43	1.0	1.2	0.83	7.1	10.9	0.58
1997	34.0	41.7	0.99	1.2	1.2	1.00	0.1	0.0	-	6.4	6.1	1.03
1998	28.0	31.5	0.95	0.5	0.3	1.67	0.0	0.0	-	6.6	15.0	0.44
1999	27.0	42.1	0.66	2.3	0.3	7.67	0.0	0.0	-	7.6	5.3	1.46
2000	19.0	32.6	0.57	5.0	0.1	50.00	0.0	0.0	-	7.3	17.3	0.42
2001	34.9	64.7	0.55	1.6	0.9	16.00	0.0	0.0	-	9.1	14.1	0.65
2002	52.8	54.3	0.99	1.6	0.9	2.29	0.0	0.1	-	13.8	20.0	0.69
2003	45.8	30.0	1.51	1.6	0.2	8.00	0.0	0.3	-	13.7	10.3	1.38
2004	34.2	17.9	1.83	0.8	0.0	200.00	0.5	0.0	-	20.3	24.3	0.83
2005	14.5	15.9	1.07	0.4	0.0	13.33	0.7	0.4	3.50	23.4	23.4	0.99
2006	16.9	30.7	0.55	0.4	0.0	25.00	0.6	0.4	1.51	24.1	22.5	1.07
2007	18.8	32.7	0.57	0.4	0.0	66.67	1.1	0.4	2.75	15.0	13.0	1.15
2008	35.3	34.2	1.03	0.8	0.0	0.00	0.7	0.2	3.50	23.8	15.0	1.59
2009	23.0	25.7	0.89	0.1	0.0	25.00	0.6	0.1	6.00	23.4	12.5	1.87
2010 ^{b/}	30.3	NA	NA	2.3	NA	NA	0.9	NA	NA	13.0	NA	NA
2011	37.5	-	-	0.4	-	-	1.5	-	-	14.3	-	-

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 2 of 4)

Year	Stillaguamish ^{c/}			Snohomish ^{c/}			Snohomish ^{c/}			Tulalip ^{c/}		
	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason
		Natural			Hatchery			Natural			Hatchery	
1993	NA	1.3	-	1.6	2.7	0.58	4.9	5.5	0.89	2.8	1.4	2.03
1994	NA	1.3	-	1.8	5.4	0.33	4.5	5.0	0.90	2.8	1.8	1.59
1995	1.8	0.9	1.92	2.2	4.0	0.54	4.3	4.0	1.08	2.3	8.5	0.27
1996	1.3	1.2	1.04	6.7	4.6	1.47	4.2	5.9	0.71	2.7	11.5	0.24
1997	1.6	1.2	1.36	7.7	12.0	0.64	5.2	4.4	1.19	4.0	8.7	0.46
1998	1.6	1.6	1.03	6.5	4.7	1.37	5.6	6.4	0.88	2.5	7.2	0.35
1999	1.5	1.1	1.36	7.8	4.7	1.65	5.6	4.8	1.16	4.5	15.2	0.30
2000	2.0	1.7	1.21	6.2	1.9	3.20	6.0	6.1	0.98	5.0	8.3	0.60
2001	1.7	1.4	1.22	4.1	0.9	4.57	5.8	8.4	0.69	5.5	5.1	1.08
2002	2.0	1.6	1.25	6.8	2.6	2.66	6.7	7.3	0.92	5.8	5.2	1.12
2003	2.0	1.0	1.98	9.4	5.8	1.63	5.5	5.6	0.99	6.0	8.7	0.69
2004	1.9	1.6	1.19	3.3	6.4	0.52	10.6	11.2	0.95	6.8	6.5	1.05
2005	1.7	1.2	1.42	4.4	4.0	1.10	14.1	5.0	2.82	6.4	7.4	0.86
2006	1.0	1.3	0.77	2.8	4.3	0.65	11.0	8.8	1.25	9.3	5.8	1.60
2007	1.0	0.8	1.27	3.5	6.6	0.53	12.7	4.0	3.18	8.4	6.1	1.38
2008	0.6	1.8	0.34	3.8	6.2	0.61	7.4	8.4	0.88	2.7	3.9	0.69
2009	1.7	1.3	1.29	4.9	2.0	2.45	8.4	2.3	3.65	4.0	0.8	5.00
2010 ^{b/}	1.4	NA	NA	5.6	NA	NA	9.9	NA	NA	3.4	NA	NA
2011	1.9	-	-	5.2	-	-	7.4	-	-	3.5	-	-

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 3 of 4)

Year	South Puget Sound Hatchery			South Puget Sound Natural			Strait of Juan de Fuca Hatchery			Strait of Juan de Fuca Natural		
	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason	Preseason Forecast	Postseason Return	Pre/Postseason
1993	61.8	43.1	1.68	26.5	9.6	1.34	0.7	1.0	3.50	3.1	1.6	1.29
1994	52.7	49.9	1.08	18.0	10.5	0.60	3.9	1.2	2.44	1.0	1.0	2.00
1995	49.6	75.4	0.67	21.7	24.9	0.63	3.0	0.7	30.00	0.9	2.3	0.33
1996	51.9	53.2	0.89	19.0	16.5	0.53	2.8	1.4	14.00	0.9	2.0	0.29
1997	65.1	38.3	1.40	18.2	15.9	0.88	2.2	1.0	7.33	0.8	2.9	0.23
1998	67.8	49.6	1.24	21.8	14.6	0.79	1.7	1.7	1.00	0.9	2.1	0.47
1999	59.4	67.3	0.71	19.6	33.5	1.15	1.9	0.7	2.71	0.9	2.7	0.33
2000	77.5	47.4	1.39	17.5	39.5	1.26	2.0	1.2	1.67	1.1	1.7	0.65
2001	73.7	76.6	0.76	16.2	44.6	0.80	0.0	1.7	0.00	3.5	2.0	1.75
2002	90.8	69.2	1.07	16.9	58.5	0.79	0.0	1.6	0.00	3.6	2.2	0.97
2003	86.6	56.6	1.14	19.6	31.0	1.28	0.0	1.3	0.00	3.4	2.8	0.72
2004	86.5	66.4	1.16	17.5	24.5	0.61	0.0	1.4	0.00	3.5	4.1	0.85
2005	83.1	73.9	0.95	17.7	19.1	0.46	0.0	1.4	0.00	4.2	2.0	2.00
2006	85.8	105.1	0.82	21.3	42.2	0.50	0.0	1.2	0.00	4.2	3.0	1.39
2007	83.0	147.1	0.56	17.0	29.7	0.57	0.0	0.8	0.00	4.4	1.3	3.38
2008	101.6	95.1	1.07	21.1	30.9	0.68	0.0	0.7	0.00	4.5	1.2	3.75
2009	93.0	77.3	1.20	17.2	13.0	1.32	0.0	1.5	0.00	3.4	1.3	2.62
2010 ^{b/}	97.4	NA	NA	12.7	NA	NA	0.0	NA	NA	3.7	NA	NA
2011	118.6	-	-	8.9	-	-	0.0	-	-	4.5	-	-

TABLE II-8. Comparison of preseason forecasts and postseason estimates of Puget Sound run size for summer/fall Chinook in thousands of fish.^{a/} (Page 4 of 4)

Year	Preseason	Postseason	Pre/Postseason
	Forecast	Return	
	Hood Canal Hatchery and Natural		
1993	NA	9.2	-
1994	11.7	8.1	1.44
1995	11.5	7.8	1.47
1996	3.9	16.2	0.24
1997	9.0	30.2	0.30
1998	2.7	20.9	0.13
1999	6.7	30.4	0.22
2000	14.0	34.4	0.41
2001	19.2	38.8	0.49
2002	25.3	36.9	0.69
2003	24.0	59.4	0.40
2004	29.6	47.0	0.63
2005	30.5	39.3	0.78
2006	30.2	40.9	0.74
2007	47.5	44.0	1.08
2008	36.8	41.0	0.90
2009	42.6	44.0	0.97
2010 ^{b/}	45.0	41.0	1.10
2011	40.6	-	-

a/ Puget Sound run size is defined as the run available to Puget Sound net fisheries. Does not include fish caught by troll and recreational fisheries inside Puget Sound.

b/ Postseason returns are preliminary.

c/ These numbers are in terms of terminal run of Chinook returning to area 8A. This includes all adult Chinook harvested in the net fisheries in Areas 8A, 8D, the Stillaguamish and Snohomish Rivers; harvest in sport fisheries in Area 8D and the Stillaguamish and Snohomish Rivers; and escapement.




FIGURE II-1. The Sacramento Index (SI) and relative levels of its components. The Sacramento River fall Chinook escapement goal range of 122,000-180,000 adult spawners is noted on the vertical axis.




FIGURE II-2. Regression estimator for the SI based on previous year's escapement of Sacramento River fall Chinook jacks, 1990-2010. Years shown are SI years.






FIGURE II-3. Regression estimators for Klamath River fall Chinook ocean abundance (September 1) based on that year's river return of same cohort. Numbers in plots denote brood years.