

**APPENDIX A  
SUMMARY OF COUNCIL STOCK MANAGEMENT GOALS**

**LIST OF TABLES**

	<u>Page</u>
TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries .....	63
TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13.....	75
TABLE A-3. Fishery impact rate criteria for OCN coho stock components based on the harvest matrix resulting from the OCN work group 2000 review of Amendment 13.....	76



TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 1 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p><b>CALIFORNIA CENTRAL VALLEY</b> - All fall, late-fall, winter, and spring stocks of the Sacramento and San Joaquin Rivers and their tributaries. Management of this stock complex is based primarily on Sacramento River fall Chinook, which includes a large hatchery component and natural Sacramento River winter Chinook, which are listed as endangered. The San Joaquin system has been severely degraded by water development projects and pollution. Natural populations of spring Chinook there have been extirpated, and remaining spawning areas are utilized primarily by fall Chinook, which have comprised &lt;10% of the total Central Valley fall run.</p>			
<p><b>Sacramento River Fall</b></p>	<p>122,000-180,000 natural and hatchery adult spawners (MSY proxy adopted 1984). This objective is intended to provide adequate escapement of natural and hatchery production for Sacramento and San Joaquin fall and late-fall stocks based on habitat conditions and average run-sizes as follows: Sacramento River 1953-1960; San Joaquin River 1972-1977 (ASETF 1979; PFMC 1984; SRFCRT 1994). The objective is less than the estimated basin capacity of 240,000 spawners (Hallock 1977), but greater than the 118,000 spawners for maximum production estimated on a basin by basin basis before Oroville and Nimbus Dams (Reisenbichler 1986).</p>	<p>Yes. A conservation alert or overfishing concern will be based on a failure to meet 122,000 adult spawners.</p>	<p>Contributes to ocean fisheries off California, southern and central Oregon, Washington, and British Columbia. Council management actions on this stock are directed at fisheries south of Pt. Arena; impacts on this stock between Pt. Arena and Horse Mt. are incidental to management measures directed at Klamath River fall Chinook.</p>
<p><b>Sacramento River Spring</b> Threatened (1999)</p>	<p>Listed as threatened under ESA. NMFS ESA consultation standard/recovery plan. Present level of ocean fishery impacts limited by measures constraining harvest on Sacramento River winter and Klamath River fall Chinook.</p>	<p>No. NMFS ESA consultation standard provides interim rebuilding program MSY criteria undefined.</p>	<p>Contributes to ocean fisheries off California, but also known to occur off Oregon. Ocean fishery impacts primarily incidental to harvest of Sacramento River fall Chinook and may be lower due to differences in run timing. Stock has been affected by man-caused loss and deterioration of freshwater habitat.</p>
<p><b>Sacramento River Winter</b> Endangered (1994)</p>	<p>Listed as endangered under ESA. NMFS ESA consultation standard specifies duration and timing of commercial and recreational fisheries south of Pt. Arena.</p>	<p>No. NMFS ESA consultation standard provides interim rebuilding program.</p>	<p>Believed to contribute predominantly to ocean fisheries south of Pt. Arena. Ocean fishery impacts incidental to harvest of Sacramento River fall Chinook.</p>
<p><b>NORTHERN CALIFORNIA COAST</b> - All fall and spring stocks of California streams north of the entrance to San Francisco Bay. Management of this stock complex is based primarily on meeting spawning escapements for natural fall Chinook. Limited data is available except for the Klamath River. An assessment and monitoring program is under consideration by CDFG for stocks originating from the Smith, Eel, Mattole, and Mad Rivers, which might provide a more thorough management basis for the future. There are significant water diversion problems in several drainages. In the Klamath River Basin, there is significant hatchery production of fall Chinook, and less so of spring Chinook, resulting primarily from mitigation programs for dams constructed in both Upper Klamath and Trinity Rivers.</p>			
<p><b>Eel, Mattole, Mad, and Smith Rivers</b> (Fall and Spring) Eel, Mattole, and Mad River stocks - Threatened (1999)</p>	<p>Eel, Mattole, and Mad River stocks listed as threatened under ESA. Data insufficient to define MSY criteria. Indices of spawning abundance limited to one tributary of the Mad River and two tributaries of the Eel River. NMFS ESA consultation standard/recovery plan for Eel, Mattole, and Mad River stocks requires that the projected ocean harvest rates on age-4 Klamath River fall Chinook not exceed 16.0%.</p>	<p>Eel, Mattole, and Mad - No. NMFS ESA consultation standard provides interim rebuilding program MSY criteria undefined. Smith - Indirectly. Data insufficient to define MSY criteria. CDFG developing an assessment and monitoring program.</p>	<p>Very limited management data available. Believed to occur in ocean fisheries off northern California and southern Oregon. Ocean fishery impacts incidental to fisheries for Sacramento and Klamath Rivers fall Chinook. No preseason or postseason abundance estimates available.</p>

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 2 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
<b>--- CHINOOK ---</b>			
<b>Klamath River Fall</b> (Klamath and Trinity Rivers)	33% to 34% of potential adult natural spawners, but no fewer than 35,000 naturally spawning adults in any one year. Brood escapement rate must average 33% to 34% over the long-term, but an individual brood may vary from this range to achieve the required tribal/nontribal annual allocation. Objective designed to allow a wide range of spawner escapements from which to develop an MSY objective or proxy while protecting the stock during prolonged periods of reduced productivity. Adopted 1988 based on Hubbell and Boydstun (1985); KRTT (1986); PFMC (1988); minor technical modifications in 1989 and 1996 (Table I-1). Natural spawners to maximize recruitment are estimated at 41,000 to 106,000 adults (Hubbell and Boydstun 1985).	Yes. A conservation alert or overfishing concern will be based on a failure to meet the 35,000 floor.	Contributes primarily to ocean fisheries from Humbug Mt., Oregon to Horse Mt., California (the KMZ) and to Klamath River tribal and recreational fisheries. Coastwide impacts are considered in meeting allocation requirements for Indian tribes with federally recognized fishing rights and the inland fishery. Specific management measures for this stock generally are implemented from Pt. Sur, California to Cape Falcon, Oregon.
<b>Klamath River Spring</b> (Klamath and Trinity Rivers)	Undefined. Productive potential believed to be protected by fishery management objective for Klamath River fall Chinook, which includes an inside allocation to tribal and sport fisheries.	Indirectly. MSY criteria undefined.	Little information available on ocean distribution. Believed to occur in ocean fisheries off northern California and southern Oregon (based on Trinity River Hatchery fish).
<b>OREGON COAST</b> - All fall and spring stocks from Oregon streams south of the Columbia River. No preseason abundance estimates available. Management based primarily on an aggregate objective of 150,000 to 200,000 natural adult spawners (attainment of objective based on a postseason estimate of 60 to 90 natural adult spawners per mile in nine standard index streams). This objective is based on optimal escapement estimates for individual coastal rivers at habitat capacity (Thompson 1977). Lower end of the objective range is nearly twice the estimated MSY spawning escapement of 79,000 fall Chinook adults based on stock recruit analysis (McGie 1982). Significant hatchery production also exists within the coastal streams. Far-north migrating, naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.			
<b>Southern Oregon</b> (Aggregate of fall and spring stocks in all streams south of Elk River; Rogue River fall stock is used to indicate relative abundance and ocean contribution rates)	Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982). ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council.	Yes, based on postseason estimates of <60 natural adult spawners per mile. Conservation promoted by the objective for Klamath River fall Chinook, which includes a large inside allocation component that reduces ocean fishery exploitation rate in areas inhabited by these fish, and by ESA consultation standard for California coastal Chinook, which limits projected ocean harvest rates on age-4 Klamath River fall Chinook to $\leq 16.0\%$ .	Medium abundance. Data limited except for Rogue River fall stock. Stocks migrate southerly or remain local, and fall Chinook contribute to ocean fisheries off northern California and Oregon, less so for spring stocks.
<b>Central and Northern Oregon</b> (Aggregate of fall and spring stocks in all streams from the Elk River to just south of the Columbia River)	Unspecified portion of an aggregate 150,000 to 200,000 natural adult spawners for Oregon coast (Thompson 1977 and McGie 1982). ODFW developing specific conservation objectives for spring and fall stocks that may be implemented without plan amendment upon approval by the Council.	Yes, based on postseason estimates of <60 natural adult spawners per mile.	Variable between high and medium abundance. Stocks migrate northward and contribute to ocean fisheries off British Columbia and southeast Alaska, and to a lesser degree, off Washington and Oregon. Nehalem, Siletz, and Siuslaw stocks are subject to the PSC ISBM harvest limitations.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 3 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p><b>COLUMBIA RIVER BASIN</b> - All pertinent fall, summer, and spring stocks of the Columbia River and its tributaries. Stocks within this complex are noted by area of origin: lower river (below Bonneville Dam), mid-river (Bonneville to McNary Dams), and upper river (above McNary Dam). Spawner escapement goals for these stocks are set through procedures of the U.S. District Court in <i>U.S. v. Oregon</i> and subsequent court orders. These goals are set forth in the Columbia River Fishery Management Plan and are recognized in the Council's conservation objectives. Annual inside fishery management planning activities are conducted within the Columbia River Compact and other state and tribal management forums. The Columbia River Compact, initially established by Oregon and Washington to jointly administer commercial fisheries within the Columbia River, takes into account the impacts from other state and tribal fisheries (e.g., recreational, ceremonial, subsistence, etc.) authorized under <i>U.S. v. Oregon</i>. The majority of ocean Chinook harvest north of Cape Falcon is provided by Columbia River salmon stocks, primarily hatchery production of tule fall Chinook from the Bonneville Pool (Spring Creek) and lower river hatcheries, smaller numbers of upper river bright hatchery and natural fall Chinook, and some lower river hatchery spring Chinook (Cowlitz). Hatchery objectives are based on long-range production programs and/or mitigation requirements associated with displaced natural stocks. Threatened Snake River fall Chinook, which suffer from severe dam passage mortalities and extreme loss of freshwater habitat, are of prime concern in limiting ocean exploitation rates in all ocean fisheries north of Pigeon Pt., California. These limits act to provide considerable protection to other weak natural stocks subject to ocean fishery impacts. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.</p>			
<p><b>North Lewis River Fall</b> Threatened (1999)</p>	<p>NMFS ESA consultation standard/recovery plan (not established at time of printing). Mclsaac (1990) stock-recruit analysis supports MSY objective of 5,700 natural adult spawners.</p>	<p>No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program. Base period Council-area ocean fishery impacts around 7%.</p>	<p>Below average abundance in 2008. Present in ocean fisheries north of Cape Falcon to SE Alaska. Subject to the PSC ISBM harvest limitations.</p>
<p><b>Lower River Hatchery Fall</b></p>	<p>15,400 adults to meet egg-take goal or as determined by management entities. 41.0% total RER in 2008 for ESA listed lower Columbia River natural tule fall Chinook estimated from Cowlitz, Washougal, Kalama and Big Creek hatchery fall Chinook.</p>	<p>No (hatchery exception or listed stock). NMFS ESA consultation standard provides interim rebuilding program.</p>	<p>Below average abundance in 2008. Major contributor to ocean fisheries north of Cape Falcon to central British Columbia.</p>
<p><b>Lower River Hatchery (Spring)</b></p>	<p>2,700 adults to meet Cowlitz, Kalama, and Lewis Rivers broodstock needs.</p>	<p>No (hatchery exception).</p>	<p>Below average abundance in 2008. Present in ocean fisheries north of Cape Falcon to southeast Alaska.</p>
<p><b>Upper Willamette (Spring)</b> Threatened (1999)</p>	<p>NMFS ESA consultation standard/recovery plan (ODFW FMFP). Willamette River Management Plan provides an MSY proxy of 30,000 to 45,000 hatchery and natural adults over Willamette River falls, depending on run size.</p>	<p>No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program. Base period Council-area ocean fishery exploitation rate of &lt;5% prevents effective Council fishery management and rebuilding.</p>	<p>Below average abundance in 2008. Present in fisheries north of Cape Falcon to southeast Alaska.</p>
<p><b>Mid-Columbia Bright Hatchery (Fall)</b></p>	<p>None for ocean fishery management.</p>	<p>No (hatchery exception).</p>	<p>Below average abundance in 2008. Contributor to ocean fisheries off Washington, British Columbia, and southeast Alaska.</p>
<p><b>Spring Creek Hatchery (Fall)</b></p>	<p>7,000 adults to meet hatchery egg-take goal.</p>	<p>No (hatchery exception).</p>	<p>Average abundance in 2008. Major contributor to ocean fisheries north of Cape Falcon to southern British Columbia.</p>

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 4 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<b>COLUMBIA RIVER BASIN (continued)</b>			
<b>Klickitat, Deschutes, John Day, and Yakima Rivers (Spring)</b>	Hold ocean fishery impacts at or below base period (<1%) and recognize CRFMP objective - MSY proxy of 115,000 adults above Bonneville Dam, including upper and mid-Columbia and Snake River stocks (state and tribal management entities considering separate conservation objectives for these stocks).	Limited. Base period Council-area ocean fishery exploitation rate of <1% prevents effective Council fishery management and rebuilding. Major habitat restoration addressing water withdrawals and dam passage and blockages is necessary for rebuilding.	Below average abundance in 2008. No significance to ocean fisheries, infrequent occurrence in fisheries north of Cape Falcon to Alaska.
<b>Snake River Fall Threatened (1992)</b>	NMFS ESA consultation/recovery standard. Since 1995, Council has met a standard of limiting its fisheries so that the total exploitation rate on age-3 and age-4 Lyons Ferry Hatchery fall Chinook (representing Snake River fall Chinook) for all ocean fisheries (including Canada) has been ≤70.0% of the 1988-1993 average adult equivalent exploitation rate. Prior to listing, managed within objectives for upper Columbia River bright fall Chinook.	No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim rebuilding program. Recovering historic abundance unlikely, as dams block former primary spawning area.	Depressed. Present in ocean fisheries from central California to southeast Alaska with greatest contribution to Canadian fisheries. Primary impacts in Council fisheries north of Cape Falcon, but also extending to Pigeon Pt., California.
<b>Snake River Spring/Summer Threatened (1992)</b>	Not applicable for ocean fisheries.	No. Listed stock. Base period Council-area ocean fishery impacts rare (unmeasurable). Dam passage mortality must be reduced to allow stock recovery.	Depressed, recent upward trend. Rare occurrence in ocean fisheries from Washington to southeast Alaska.
<b>Upper River Bright (Fall)</b>	40,000 natural bright adults above McNary Dam (MSY proxy) adopted in 1984 based on CRFMP. The management goal was increased to 45,000 by Columbia River managers between 1986 and 1993. Since 1994, inriver fisheries management based on a NMFS ESA consultation standard exploitation rate to protect Snake River wild fall Chinook	Limited. Base period Council-area ocean fishery exploitation rate <4% prevents effective Council fishery management and rebuilding.	Below average abundance in 2008. Major contributor to ocean fisheries off Canada, and to a lesser extent, Washington and Oregon. Primary impact area north of Cape Falcon. Subject to the PSC ISBM harvest limitations.
<b>Upper River Summer</b>	Hold ocean fishery impacts at or below base period (<2%); recognize <i>U.S. v. Oregon</i> objective - MSY proxy of 29,300 adults to river mouth destined to for areas above Priest Rapids Dam (excludes Snake River stocks).	Limited. Base period Council-area ocean fishery exploitation rate <2% prevents effective Council fishery management and rebuilding. Dam passage mortalities must be reduced to allow rebuilding.	Long-term depressed abundance, significant upward trend in the last few years. Present in ocean fisheries north of Cape Falcon to southeast Alaska. Subject to the PSC ISBM harvest limitations.
<b>Upper Columbia River Spring Endangered (1999)</b>	None applicable to ocean fisheries. Ensure ocean fishery impacts remain rare and recognize CRFMP objective - MSY proxy of 115,000 adults above Bonneville Dam, including upper and mid-Columbia and Snake River stocks (state/tribal management entities considering separate objectives for these stocks).	No. Listed stock. Base period Council-area ocean fishery impacts rare (not measurable), making Council management and rebuilding ineffective. Reduce dam passage mortalities to allow rebuilding.	Long-term depressed abundance, recent upward trend. Captive broodstock programs started in 1997. No significance to ocean fisheries. Rare occurrence in ocean fisheries north of Cape Falcon to Canada.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 5 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<b>WASHINGTON COAST</b> - All pertinent fall, summer and spring stocks from coastal streams north of the Columbia River through the western Strait of Juan de Fuca (west of the Elwha River). This stock complex consists of several natural stocks, generally of small to medium sized populations, and some hatchery production (Willapa Bay and the Quinault River). Stocks in this complex tend to range further north than most Columbia River stocks and, while present in fisheries from Cape Falcon to southeast Alaska, are not significantly impacted by Council-area ocean fisheries. Preseason abundance estimates are generally not available for Council management. These stocks qualify as exceptions to the Council's overfishing criteria, due to very low fishery impacts. Spawning escapement goals for stocks managed within this complex, established in U.S. District Court by WDFW and the treaty tribes, are recognized in the Council's conservation objectives below. Objectives for Grays Harbor and the north coast river systems have been established pursuant to the U.S. District Court order in <u>Hoh v. Baldrige</u> . However, annual natural spawning escapement targets may vary from the conservation objectives below if agreed to by WDFW and the treaty tribes under the provisions of <u>Hoh v. Baldrige</u> and subsequent U.S. District Court orders. After agreement is reached on the annual targets, ocean fishery escapement objectives are established for each river, or region of origin, which include provisions for treaty allocation and inside, non-Indian fishery needs. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border.			
<b>Willapa Bay Fall</b> (Natural)	No FMP objective. WDFW goal of 4,400 natural spawners.	Limited (exploitation rate exception).	
<b>Willapa Bay Fall</b> (Hatchery)	9,800 adult return to hatchery.	No (hatchery exception).	
<b>Grays Harbor Fall</b>	14,600 natural adult spawners--MSP based on full seeding of spawning and rearing habitat (WDF 1979). 1,400 natural adult spawners.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Grays Harbor Spring</b>		Limited (exploitation rate exception).	
<b>Quinault Fall</b>	Hatchery production.	No (hatchery exception).	
<b>Queets Fall</b>	Manage terminal fisheries for 40% harvest rate, but no less than 2,500 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Queets Spring/Summer</b>	Manage terminal fisheries for 30% harvest rate, but no less than 700 natural adult spawners.	Limited (exploitation rate exception).	
<b>Hoh Fall</b>	Manage terminal fisheries for 40% harvest rate, but no less than 1,200 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Hoh Spring/Summer</b>	Manage terminal fisheries for 31% harvest rate, but no less than 900 natural adult spawners.	Limited (exploitation rate exception).	
<b>Quillayute Fall</b>	Manage terminal fisheries for 40% harvest rate, but no less than 3,000 natural adult spawners, the MSY level estimated by Cooney (1984).	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Quillayute Spring/Summer</b>	1,200 natural adult spawners for summer component (MSY).	Limited (exploitation rate exception).	
<b>Hoko Summer/Fall</b> (Western Strait of Juan de Fuca)	850 natural adult spawners, the MSP level estimated by Ames and Phinney (1977). May include adults used for supplementation program.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 6 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- CHINOOK ---			
<p><b>PUGET SOUND</b> - All fall, summer, and spring stocks originating from U.S. tributaries to Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek). This stock complex consists of numerous natural Chinook stocks of small to medium sized populations and significant hatchery production. Puget Sound stocks contribute to fisheries off British Columbia and are present into southeast Alaska, but are impacted to a minor degree by Council-area ocean fisheries. Base period, Council-area ocean fishery exploitation rates (adult equivalent) of 2% or less are below a management threshold which allows effective Council management of these stocks and they qualify as <b>exceptions</b> to the Council's overfishing criteria. The naturally spawning stocks within this complex are listed as threatened under the ESA. Naturally spawning stocks are also subject to the 1999 Chinook agreement of the Pacific Salmon Commission and may be subject to exploitation rate constraints in U.S. fisheries south of the Canada/Washington border. Management objectives for hatchery stocks are based on hatchery escapement needs. Fisheries in Puget Sound conducted under a Resource Management Plan (RMP) are exempted from ESA Section 9 take prohibitions under Limit 6 of the 4(d) rule. This RMP will expire on May 1 of this year. A new RMP is currently under review by NOAA Fisheries but this review will not be completed prior to the March Council meeting.</p>			
<b>Eastern Strait of Juan de Fuca Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>Skokomish Summer/Fall</b> (Hood Canal) Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>Nooksack Spring (early)</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Skagit Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Skagit Spring</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Stillaguamish Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Snohomish Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Cedar River Summer/Fall</b> (Lake Washington) Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). The preliminary 2004 consultation standard is an RER constraint total mortality in all fisheries not to exceed 31%.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.



TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 7 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
<b>PUGET SOUND</b> (continued)			
<b>White River Spring</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>Puyallup Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>Green River Summer/Fall</b> Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	Subject to the PSC ISBM harvest limitations.
<b>Nisqually River Summer/Fall</b> (South Puget Sound) Threatened (1999)	NMFS ESA consultation standard. Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>Mid Hood Canal Fall</b> Threatened (1999)	NMFS ESA consultation standard is expressed in terms of Recovery Exploitation Rate (RER). Guidance will be provided prior to the March Council meeting.	Limited (exploitation rate exception).	
<b>SOUTHERN BRITISH COLUMBIA</b> - Fall and spring stocks of British Columbia coastal streams and the Fraser River. Management based primarily on natural and hatchery fall Chinook. Base period, Council-area ocean fishery exploitation rates (adult equivalent) on the coastal stocks of 1% or less are below a management threshold which allows effective Council management of these stocks, and they qualify as <b>exceptions</b> to the Council's overfishing criteria.			
<b>Coastal Stocks</b>	Undefined for Council fisheries. Manage consistent with the Pacific Salmon Treaty.	No. Under Canadian authority and would also be an exploitation rate exception.	Medium abundance. Major contributors to ocean fisheries off British Columbia; significant contributors north into southeast Alaska and present off northern Washington.
<b>Fraser River</b>	Undefined for Council fisheries. Manage consistent with the Pacific Salmon Treaty.	No. Under Canadian authority.	Medium abundance. Major contributors to ocean fisheries off British Columbia; contributors off northern Washington; and present north into southeast Alaska. Harrison River stock subject to the PSC ISBM harvest limitations.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 8 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
<p><b>OREGON PRODUCTION INDEX AREA</b> - All Washington, Oregon, and California natural and hatchery coho stocks from streams south of Leadbetter Pt. Significant production from Columbia River and Oregon coastal hatcheries provide harvest in ocean fisheries throughout the Council management area. Ocean fisheries are usually limited primarily to meet natural escapement objectives. Treaty Indian obligations, non-Indian harvest opportunity, and hatchery requirements must also be factored in for the Columbia River stocks. Natural components have been severely depressed for several years due to a combination of previously high fishery impacts, major losses or degradation of freshwater habitat, and long-term marine conditions unfavorable to coho survival.</p>			
<p><b>Central California Coast</b> Threatened (1996)</p>	<p>NMFS ESA consultation standard/recovery plan. Since 1998, no retention of coho in commercial and recreational fisheries off California in conjunction with total marine fishery impacts of no more than 13% on Rogue/Klamath hatchery coho (surrogate stock). Objective undefined prior to listing.</p>	<p>No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim protection of productive capacity. Recovery limited by deterioration of significant portions of freshwater habitat, distribution at southern edge of coho range, and ongoing unfavorable marine conditions.</p>	<p>Very minor component of OPI area fisheries, limited potential for significant contribution to ocean and inland fisheries. Current impacts incidental in ocean fisheries off California. Development of monitoring and assessment program considered for Ten Mile River, Noyo River, Gualala River, Lagunitas Creek, and Scott Creek. Rogue/Klamath coho are believed to have a similar, but more northerly distribution.</p>
<p><b>Northern California</b> Threatened (1997)</p>	<p>NMFS ESA consultation standard/recovery plan. Since 1998, total marine fishery impacts limited to no more than 13.0% on Rogue/Klamath hatchery coho (surrogate stock) and no retention of coho in California ocean fisheries. Objective undefined prior to listing.</p>	<p>No. Listed stock, MSY criteria undefined. NMFS ESA consultation standard provides interim protection of productive capacity. Recovery may last more than 10 years even with no fishery impacts, due to loss or deterioration of significant portions of freshwater habitat and ongoing unfavorable marine conditions.</p>	<p>Depressed and listed. Very minor natural component of OPI area fisheries, potential for minor contribution to ocean fisheries off California and southern Oregon, and inland California fisheries. Current impacts incidental in ocean and inland fisheries (total non-retention south of Cape Falcon since 1994). CDFG considering monitoring to provide data for the Smith, Trinity, Eel, Mattole, and Klamath Rivers.</p>
<p><b>Oregon Coastal Natural</b> Comprised of Southern, South-Central, North-Central, and Northern Oregon stocks.</p>	<p>An allowable marine and freshwater exploitation rate of no more than 13% to 35%, depending on parent escapement and ocean survival trends, based on Amendment 13 of the Salmon FMP, or no more than 8% to 45% based on the OCN workgroup review of Amendment 13. Standard is ≤8.0% in 2008</p>	<p>No. Listed stock, rebuilding program initiated in 1998. The annual conservation objective should allow component stocks to rebuild when environmental conditions are favorable. Recovery for some components may last more than 10 years even with no fishery impacts, due to loss or deterioration of significant portions of freshwater habitat and ongoing unfavorable marine conditions.</p>	<p>Decline in 2007 after recent increases in abundance. Major natural component of OPI area and freshwater fisheries in Oregon coastal streams. Current impacts are primarily incidental in ocean fisheries under a total nonretention regulation south of Cape Falcon since 1994 (except 2007).</p>

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 9 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
<b>OREGON PRODUCTION INDEX (continued)</b>			
<b>Columbia River Late</b> (Hatchery)	Hatchery rack return goal of 17,200 adults.	No (hatchery exception).	Major component of ocean fisheries north of Cape Falcon. When abundant, significant contributors to ocean fisheries off Oregon north into Canada and Columbia River fisheries.
<b>Columbia River Early</b> (Hatchery)	Hatchery rack return goal of 18,800 adults.	No (hatchery exception).	Major component of OPI area fisheries. When abundant, significant contributors to ocean fisheries off California and north to Leadbetter Pt., Washington and to Columbia River fisheries. Current ocean fishery impacts from very limited retention fisheries north of Cape Falcon and incidental hook-and-release mortality in fisheries south of Cape Falcon.
<b>Columbia River</b> (Natural) Threatened, 2005	NMFS ESA consultation standard for 2008 is $\leq 8.0\%$ in total exploitation rate in marine and mainstem Columbia River fisheries.	No. Listed stock. NMFS ESA consultation standard provides interim rebuilding program.	Extinct above the Dalles Dam, small populations in Clackamas, and Sandy rivers in Oregon, and Cedar Creek (Lewis River) Washington. Lower river coho are also listed under the Oregon State ESA.
<p><b>WASHINGTON COASTAL</b> - All pertinent natural and hatchery stocks originating in Washington coastal streams north of the Columbia River through the western Strait of Juan de Fuca (West of the Elwha River). Management goals for Grays Harbor and Olympic Peninsula coho stocks include achieving natural spawning escapement objectives and treaty allocation requirements, although Grays Harbor also contains a significant amount of hatchery production. The conservation objectives for these stocks are based on MSY spawner escapements established pursuant to the U.S. District Court order in <u>Hoh v. Baldrige</u>. Annual natural spawning escapement targets and total escapement objectives are established by the WDFW and treaty tribes under the provisions of <u>U.S. v. Washington</u> and subsequent U.S. District Court orders. After agreement to annual targets is reached by the parties in this litigation, ocean fishery escapement objectives are established for each river, or region of origin, which include provisions for providing treaty allocation requirements and inside, non-Indian fishery needs. The conservation objectives for the Queets, Hoh, and Quillayute Rivers were developed as ranges intended to bracket the current best estimates of MSY escapement. The range of each objective reflects the degree of uncertainty inherent by using the high estimate of recruits-per-spawner and low estimate of carrying capacity for the lower bound and the low estimate of recruits-per-spawner with the high estimate of smolt carrying capacity for the upper end of the range. The ranges were subsequently adjusted upward for risk aversion and again for habitat considerations by 26% to 184% (Lestelle <i>et al.</i> 1984). These stocks are also subject to provisions of the 2002 PSC Coho Management Plan, which requires the United States and Canada to constrain total fishery exploitation rates to levels associated with the categorical status (low, moderate, and abundant) and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries are established through formulas specified in the PSC Management Plan. However, the salmon FMP management objectives determine the criteria for triggering a conservation alert or an overfishing concern; annual management objectives established pursuant to U.S. District Court orders and the PSC Coho Management Plan do not.</p>			
<b>Willapa Bay</b> (Hatchery)	Meet WDFW program objectives.	No (hatchery exception).	Contributes to ocean fisheries off northern Oregon north into Canada. Significant contributor to inside non-Indian commercial net and recreational fisheries. WDFW critically reviewing current management to determine if objectives for natural stocks are warranted.
<b>Grays Harbor</b>	35,400 natural adult spawners (MSP based on WDF [1979]) or annual target agreed to by WDFW and the Quinalt Indian Nation.	Yes. Conservation alert or overfishing concern based on fewer than 35,400 natural spawners.	Ocean distribution from Oregon to northern British Columbia. Harvested by treaty Indian, non-Indian commercial, and recreational fisheries in Grays Harbor and tributary rivers.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 10 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
<b>WASHINGTON COAST</b> (continued)			
<b>Queets</b>	MSY range of 5,800 to 14,500 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and the Quinault Indian Nation.	Yes. Conservation alert or overfishing concern based on fewer than 5,800 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver. Coho supplementation project conducted since the late 1970s.
<b>Hoh</b>	MSY range of 2,000 to 5,000 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and Hoh Tribe.	Yes. Conservation alert or overfishing concern based on fewer than 2,000 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver.
<b>Quillayute Fall</b>	MSY range of 6,300 to 15,800 natural adult spawners (Lestelle <i>et al.</i> 1984) or annual target agreed to by WDFW and the Quillayute Tribe.	Yes. Conservation alert or overfishing concern based on fewer than 6,300 natural spawners.	Ocean distribution from south-central Oregon to northwest Vancouver Island off British Columbia. Harvested by treaty Indian gillnet and non-treaty recreational fisheries inriver.
<b>Quillayute Summer</b> (Hatchery)	Meet hatchery program objectives.	No (hatchery exception).	Early river entry timing. Contributor to ocean fisheries off Washington north into British Columbia; present south to central Oregon.
<b>Western Strait of Juan de Fuca</b> (Sekiu, Hoko, Clallam, Pysht, East and West, and Lyre Rivers and miscellaneous streams west of the Elwha River)	11,900 natural adult spawners  PSC 2007 annual management objective: 40% (low status) exploitation rate.	Yes. Overfishing concern based on fewer than 11,900 natural spawners.	Little information on ocean distribution.

TABLE A-1. Conservation objectives and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 11 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
---COHO---			
<p><b>PUGET SOUND</b> - All pertinent natural and hatchery stocks originating from U.S. tributaries to Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek). The Puget Sound Salmon Management Plan defines management objectives and long-term goals for these stocks as developed by representatives from federal, state, and tribal agencies. Conservation objectives for specific stocks are currently based on either MSP principles for stocks managed primarily for natural production or upon hatchery escapement needs for stocks managed for artificial production. Puget Sound management procedures are outlined in a "Memorandum Adopting Salmon Management Plan" (<u>U.S. v. Washington</u>, 626 F. Supp. 1405 [1985]). The original conservation objectives were developed by a State/Tribal Management Plan Development Team following the Boldt Decision with the goal for natural spawning stocks defined as "the adult spawning population that will, on the average, maximize biomass of juvenile outmigrants subsequent to incubation and freshwater rearing under average environmental conditions." The methodology used to develop the objectives was based on assessment of the quantity and quality of rearing habitat and the number of adult spawners required to fully seed the habitat (Zillges 1977). Some objectives have subsequently been modified under fixed procedures set by the U.S. District Court and its Fisheries Advisory Board (Clark 1983 and PSSSRG 1997) and later determinations of the WDFW/Tribal Technical Committee. These natural stocks are also subject to provisions of the 2002 PSC Coho Management Plan, which requires the United States and Canada to constrain total fishery exploitation rates to levels associated with the categorical status (low, moderate, and abundant) and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries are established through formulas specified in the PSC Management Plan. However, the salmon FMP management objectives determine criteria for triggering a conservation alert or an overfishing concern; annual management objectives established pursuant to U.S. District Court orders and the PSC Coho Management Plan do not.</p>			
<b>Eastern Strait of Juan de Fuca</b> (Streams east of Salt Creek through Chimacum Creek))	FMP: MSP objective of 950 natural adult spawners (Clark 1983 modified by habitat apportionment of WDFW/Tribal Technical Committee in 1998) or annual target agreed to in U.S. District Court procedures. The Elwha and Dungeness rivers are not included in this objective but are managed on a harvest rate basis. PSC 2008: 40% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 950 natural spawners.	Little information on ocean distribution.
<b>Hood Canal</b>	FMP: MSP objective of 21,500 natural adult spawners (Clark 1983 modified since 1994 by WDFW/Tribal Technical Committee) or annual target agreed to in U.S. District Court procedures. PSC 2008: 45% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 21,500 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
<b>Skagit</b>	FMP: MSP objective of 30,000 natural adult spawners (Zillges 1977 and Clark 1983) or annual target agreed to in U.S. District Court procedures. (The spawner assessment methodology is currently being revised and may result in an objective significantly different from 30,000.) PSC 2008: 35% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 30,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
<b>Stillaguamish</b>	FMP: MSP objective of 17,000 natural adult spawners (Zillges 1977) or annual target agreed to in U.S. District Court procedures. PSC 2008: 50% (normal status) total exploitation rate.	Yes. Overfishing concern based on fewer than 17,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
<b>Snomish</b>	FMP: MSP objective of 70,000 natural adult spawners (Zillges 1977 as modified by WDFW/Tribal Technical Committee) or annual target agreed to in U.S. District Court procedures. PSC 2008: 40% (low status) total exploitation rate.	Yes. Overfishing concern based on fewer than 70,000 natural spawners.	Ocean distribution from Cape Falcon, Oregon to British Columbia.
<b>South Puget Sound</b> (Hatchery)	Hatchery rack return goal of 52,000 adults. Natural production goals under development.	No (hatchery exception).	Ocean distribution from Cape Falcon, Oregon to British Columbia.

TABLE A-1. **Conservation objectives** and management information for salmon stocks of significance to ocean salmon fisheries. Abundance information is based on recent year information. (Page 12 of 12).

Stock	Conservation Objective (to be met annually, unless noted otherwise)	Subject to Council Actions to Prevent Overfishing	Other Management Information
--- COHO ---			
<b>SOUTHERN BRITISH COLUMBIA COAST</b> - Stocks of southern British Columbia coastal streams (including Vancouver Island) and the Fraser River.			
<b>Coastal Stocks</b>	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty.	No. Not under Council management authority.	Contributes to ocean fisheries off British Columbia, north into southeast Alaska and present off northern Washington.
<b>Fraser River</b>	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty. For 2008, southern U.S. fisheries total exploitation rate of $\leq 10.0\%$ .	No. Not under Council management authority.	Contributes to ocean fisheries off British Columbia and Washington, and to Strait of Juan de Fuca and Puget Sound fisheries.
--- PINK (odd-numbered years) ---			
The Fraser River Panel of the PSC manages fisheries for pink salmon in the Fraser River Panel Area (U.S.) north of 48° N latitude to meet Fraser River natural spawning escapement and U.S./Canada allocation requirements. The Council manages pink salmon harvests in that portion of the EEZ, which is not in the Fraser River Panel Area (U.S.) waters consistent with Fraser River Panel management intent. Pink salmon management objectives must address meeting natural spawning escapement objectives, allowing ocean pink harvest within fixed constraints of coho and Chinook harvest ceilings and providing for treaty allocation requirements.			
<b>Puget Sound</b>	900,000 natural spawners or consistent with provisions of the Pacific Salmon Treaty (Fraser River Panel).	No. Minor impacts in Council fisheries and not under Council management authority.	Contributes to ocean fisheries off British Columbia and in Puget Sound. Present south into Oregon. Rare off California.
<b>Fraser River</b>	Manage Council fisheries that impact Canadian stocks consistent with provisions of the Pacific Salmon Treaty (Fraser River Panel).	No. Minor impacts in Council fisheries and not under Council management authority.	Contributes to ocean fisheries off British Columbia; present into southeast Alaska and off Washington and northern Oregon. Rare off California.

TABLE A-2. Allowable fishery impact rate criteria for OCN coho stock components under the Salmon Fishery Management Plan Amendment 13. (Page 1 of 1)

PARENT SPAWNER STATUS		MARINE SURVIVAL INDEX (based on return of jacks per hatchery smolt)			
		Low (<0.0009)	Medium (0.0009 to 0.0034)	High (>0.0034)	
		Allowable Total Fishery Impact Rate			
<b>High:</b>	Parent spawners achieved Level #2 rebuilding criteria, grandparent spawners achieved Level #1	≤15%	≤30% <sup>a/</sup>	≤35% <sup>a/</sup>	
<b>Medium:</b>	Parent spawners achieved Level #1 or greater rebuilding criteria	≤15%	≤20% <sup>a/</sup>	≤25% <sup>a/</sup>	
<b>Low:</b>	Parent spawners less than Level #1 rebuilding criteria	≤15% ≤10-13% <sup>b/</sup>	≤15%	≤15%	
OCN Coho Spawners by Stock Component					
Rebuilding Criteria	Northern	North-Central	South-Central	Southern	Total
Full Seeding at Low Marine Survival:	21,700	55,000	50,000	5,400	132,100
Level #2 (75% of full seeding):	16,400	41,300	37,500	4,100	99,300
Level #1 (50% of full seeding):	10,900	27,500	25,000	2,700	66,100
38% of Level #1 (19% of full seeding):	4,100	10,500	9,500	1,000	25,100
Stock Component (Boundaries)	Full Seeding of Major Basins at Low Marine Survival (Number of Adult Spawners)				
<b>Northern:</b> (Necanicum River to Neskowin Creek)	Nehalem	Tillamook	Nestucca	Ocean Tribs.	
	17,500	2,000	1,800	400	
<b>North-Central:</b> (Salmon River to Siuslaw River)	Siletz	Yaquina	Alsea	Siuslaw	Ocean Tribs.
	4,300	7,100	15,100	22,800	5,700
<b>South-Central:</b> (Siltcoos River to Sixes River)	Umpqua	Coos	Coquille	Coastal Lakes	
	29,400	7,200	5,400	8,000	
<b>Southern:</b> (Elk River to Winchuck River)	Rogue				
	5,400				

a/ When a stock component achieves a medium or high parent spawner status under a medium or high marine survival index, but a major basin within the stock component is less than 10% of full seeding, (1) the parent spawner status will be downgraded one level to establish the allowable fishery impact rate for that component, and (2) no coho-directed harvest impacts will be allowed within that particular basin.

b/ This exploitation rate criteria applies when (1) parent spawners are less than 38% of the Level #1 rebuilding criteria, or (2) marine survival conditions are projected to be at an extreme low as in 1994-1996 (<0.0006 jack per hatchery smolt). If parent spawners decline to lower levels than observed through 1998, rates of less than 10% would be considered, recognizing that there is a limit to further bycatch reduction opportunities.

TABLE A-3. Fishery **impact** rate criteria for **OCN coho** stock components based on the harvest matrix resulting from the **OCN work group** 2000 review of Amendment 13. (Page 1 of 1).

Parent Spawner Status <sup>a/</sup>	Marine Survival Index (based on return of jacks per hatchery smolt)						
	Extremely Low (<0.0008 )	Low (0.0008 to 0.0014 )	Medium (>0.0014 to 0.0040)	High (>0.0040 )			
<b>High</b> Parent Spawners > 75% of full seeding	<b>E</b> ≤ 8%	<b>J</b> ≤ 15%	<b>O</b> ≤ 30%	<b>T</b> ≤ 45%			
<b>Medium</b> Parent Spawners > 50% & ≤ 75% of full seeding	<b>D</b> ≤ 8%	<b>I</b> ≤ 15%	<b>N</b> ≤ 20%	<b>S</b> ≤ 38%			
<b>Low</b> Parent Spawners > 19% & ≤ 50% of full seeding	<b>C</b> ≤ 8%	<b>H</b> ≤ 15%	<b>M</b> ≤ 15%	<b>R</b> ≤ 25%			
<b>Very Low</b> Parent Spawners > 4 fish per mile & ≤ 19% of full seeding	<b>B</b> ≤ 8%	<b>G</b> ≤ 11%	<b>L</b> ≤ 11%	<b>Q</b> ≤ 11%			
<b>Critical</b> <sup>b/</sup> Parental Spawners ≤ 4 fish per mile	<b>A</b> 0 - 8%	<b>F</b> 0 - 8%	<b>K</b> 0 - 8%	<b>P</b> 0 - 8%			
Sub-aggregate and Basin Specific Spawner Criteria Data							
Sub-aggregate	Miles of Available Spawning Habitat	100% of Full Seeding	"Critical"		Very Low, Low, Medium & High		
			4 Fish per Mile	12% of Full Seeding	19% of Full Seeding	50% of Full Seeding	75% of full Seeding
Northern	899	21,700	3,596	NA	4,123	10,850	16,275
North - Central	1,163	55,000	4,652	NA	10,450	27,500	41,250
South - Central	1,685	50,000	6,740	NA	9,500	25,000	37,500
Southern	450	5,400	NA	648	1,026	2,700	4,050
Coastwide Total	4,197	132,100	15,636		25,099	66,050	99,075

a/ Parental spawner abundance status for the OCN aggregate assumes the status of the weakest sub-aggregate.

b/ "Critical" parental spawner status is defined as 4 fish per mile for the Northern, North-Central, and South-Central subaggregates. Because the ratio of high quality spawning habitat to total spawning habitat in the Rogue River Basin differs significantly from the rest of the basins on the coast, the spawner density of 4 fish per mile does not represent "Critical" status for that basin. Instead, "Critical" status for the Rogue Basin (Southern Sub-aggregate) is estimated as 12% of full seeding of high quality