

**APPENDIX B
HISTORICAL RECORD OF ESCAPEMENTS TO
INLAND FISHERIES AND SPAWNING AREAS**

LIST OF TABLES

	<u>Page</u>
TABLE B-1. California Central Valley natural fall Chinook salmon spawning escapements in numbers of fish	195
TABLE B-2. California Central Valley hatchery fall Chinook salmon spawning escapements in numbers of fish	196
TABLE B-3. Sacramento River late-fall, winter, and spring Chinook salmon spawning escapement estimates in numbers of fish.....	197
TABLE B-4. Summary of Klamath River fall Chinook salmon estimates in numbers of adults and jacks.....	198
TABLE B-5. Estimates of Yurok and Hoopa Valley reservation Indian gillnet Chinook harvest in numbers of fish	199
TABLE B-6. Shasta River fall Chinook salmon weir counts or spawning escapement estimates in numbers of fish	200
TABLE B-7. Summary of California North Coast salmon spawning stock surveys in numbers of fish	201
TABLE B-8. Peak spawning counts in index areas for selected south/local migrating Oregon coastal fall Chinook stocks.....	202
TABLE B-9. Counts of natural and hatchery spring Chinook salmon at Gold Ray Dam on the Rogue River and at Winchester Dam on the North Umpqua River in thousands of fish.....	203
TABLE B-10. Rogue River fall Chinook carcass counts in numbers of fish.....	204
TABLE B-11. Peak counts for north migrating Oregon coastal Chinook stocks on selected fall Chinook spawning index stream surveys.....	205
TABLE B-12. Estimates of minimum inriver run size, catch, and escapement in numbers of Columbia River adult spring Chinook destined for areas below Bonneville Dam	206
TABLE B-13. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult spring Chinook destined for areas above Bonneville Dam	207
TABLE B-14. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult summer Chinook destined for areas above Bonneville Dam	208
TABLE B-15. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult Spring Creek Hatchery (SCH) stock fall Chinook.....	209
TABLE B-16. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult lower river hatchery (LRH) stock fall Chinook	210
TABLE B-17. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult lower river wild (LRW) stock fall Chinook	211
TABLE B-18. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult upriver bright (URB) stock fall Chinook destined for areas above McNary Dam and the Deschutes River	212
TABLE B-19. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult mid-Columbia bright (MCB) stock fall Chinook destined for areas below McNary Dam, not including the Deschutes River	213
TABLE B-20. Estimates of minimum inriver run size and catch in numbers of adult spring, summer, and fall Chinook from the Columbia River	214
TABLE B-21. Estimates of minimum inriver run size, catch, and escapement in thousands of adult coho entering the Columbia River	217
TABLE B-22. Estimated catch and effort in the Buoy 10 fishery	218
TABLE B-23. Willapa Bay fall Chinook terminal run size, catch, and spawning escapement in numbers of fish	219
TABLE B-24. Willapa Bay coho terminal run size, catch, and spawning escapement in numbers of fish	220

LIST OF TABLES (continued)

		<u>Page</u>
TABLE B-25.	Grays Harbor Chinook terminal catch, spawning escapement, and run size in numbers of fish	221
TABLE B-26.	Grays Harbor coho terminal catch, spawning escapement, and run size estimates in numbers of fish	223
TABLE B-27.	Treaty Indian gillnet catch of Chinook, chum, and sockeye salmon in the Quinault River in numbers of fish	224
TABLE B-28.	Estimated inriver run size, catch and escapement for Quinault River coho in numbers of fish	225
TABLE B-29.	Estimated inriver run size, catch, and escapement of Queets River spring/summer Chinook in numbers of fish	226
TABLE B-30.	Estimated inriver run size, catch, and escapement of Queets River fall Chinook in numbers of fish	227
TABLE B-31.	Estimated terminal run size, catch, and escapement for Queets River coho in numbers of fish	228
TABLE B-32.	Estimated inriver run size, catch, and escapement for Hoh River spring/summer Chinook in numbers of fish	229
TABLE B-33.	Estimated inriver run size, catch, and escapement for Hoh River fall Chinook in numbers of fish	230
TABLE B-34.	Estimated inriver run size, catch, and escapement for Hoh River coho in numbers of fish	231
TABLE B-35.	Estimated inriver run size, catch, and escapement for Quillayute River spring/summer Chinook in numbers of fish	232
TABLE B-36.	Estimated inriver run size, catch, and escapement for Quillayute River fall Chinook in numbers of fish	233
TABLE B-37.	Estimated inriver run size, catch, and escapement for Quillayute River coho stocks in numbers of fish	234
TABLE B-38.	Puget Sound commercial net and troll fishery salmon catches in numbers of fish	236
TABLE B-39.	Summary of Puget Sound marine recreational salmon catch estimates in numbers of fish from catch record cards	238
TABLE B-40.	Puget Sound reported commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound Chinook stocks	239
TABLE B-41.	Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound coho stocks	243
TABLE B-42.	Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound pink stocks. ^{iv} (Page 1 of 4)	246
TABLE B-42.	Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound pink stocks. ^{iv} (Page 1 of 4)	247
TABLE B-43.	Puget Sound spring Chinook spawning escapement estimates in numbers of adult fish	251

TABLE B-1. California Central Valley natural fall Chinook salmon spawning escapements in numbers of fish.^{a/}

Year or Average	Lower Sacramento River										Sacramento River		San Joaquin River		Central Valley	
	Upper Sacramento River		Feather River		Yuba River		American River ^{b/}		Total		Totals		Totals		Central Valley	
	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks
1971-1975	58,462	18,289	40,221	9,745	10,877	1,615	41,726	3,695	92,824	15,055	151,286	33,344	13,462	1,345	164,748	34,690
1976-1980	67,011	17,905	33,954	3,544	7,387	1,563	28,509	1,344	69,850	6,452	136,861	24,357	2,886	763	139,747	25,120
1981-1985	57,793	22,432	36,252	5,243	12,825	5,146	32,332	4,954	81,409	15,343	139,202	37,775	34,930	10,721	174,132	48,496
1986-1990	87,397	17,244	38,709	6,426	9,261	2,444	24,420	3,323	72,390	12,193	159,787	29,437	10,853	4,377	170,640	33,814
1991-1995	60,151	11,497	32,578	4,355	8,309	2,131	28,549	4,151	69,436	10,637	129,586	22,134	2,626	904	132,212	23,038
1996	131,267 ^{c/}	11,650 ^{c/}	44,593	12,577	23,492	4,408	67,719	7,026	135,803	24,012	267,071	35,661	5,766	5,979	272,837	41,640
1997	167,354	13,736	47,009	3,538	19,202	6,746	46,036	6,159	112,246	16,444	279,600	30,180	17,983	1,146	297,583	31,326
1998	60,713 ^{c/}	5,137 ^{c/}	39,600 ^{d/}	3,400	26,737	4,353	41,094	13,698	107,431	21,451	168,144	26,588	13,119	6,292	181,263	32,880
1999	256,629	7,495	30,000 ^{d/}	7,500	18,778	5,452	48,311	8,688	97,089	21,640	353,718	29,135	10,708	7,185	364,426	36,320
2000	152,923	3,900	109,924	7,017	12,954	2,041	93,413	5,646	216,291	14,704	369,214	18,604	36,896	2,578	406,110	21,182
2001	130,440	5,132	169,588	9,114	21,567	1,825	167,062	13,553	358,217	24,492	488,657	29,624	23,899	3,705	512,555	33,330
2002	481,924 ^{e/}	9,009	93,766	11,397	18,406	4,796	95,711	10,635	207,883	26,828	689,806	35,837	21,852	3,788	711,658	39,626
2003	164,802	4,402	85,578	4,369	26,820	1,489	136,238	9,627	248,636	15,485	413,438	19,887	14,519	2,164	427,957	22,051
2004	70,557 ^{f/}	7,221 ^{f/}	48,580	5,591	9,260	5,208	75,090	13,774	132,930	24,573	203,487	31,794	7,249	3,311	210,736	35,105
2005	96,716 ^{f/}	3,267 ^{f/}	43,738	4,848	16,251	987	54,001	2,842	113,990	8,677	210,706	11,944	15,905	1,517	226,611	13,461
2006	85,882 ^{f/g/}	2,729 ^{f/}	73,585	1,845	7,998	233	21,755	1,145	103,338	3,223	189,220	5,952	5,691	692	194,911	6,644
2007 ^{i/}	32,854 ^{f/h/}	950 ^{f/g/}	21,541	321	2,482	77	9,850	150	33,873	548	66,727	1,498	1,374	76	68,101	1,574

a/ Upper Sacramento River jack estimates based on Red Bluff Diversion Dam samples. All other estimates are generally based on carcass surveys. (Adult and jack numbers are generally based on a 24-inch fork length cut-off [unpublished CDFG data.]) Upper Sacramento River estimates also include Tehama-Colusa Spawning Channel for 1971 to 1980. For years prior to 2004, all numbers in this table were reviewed and updated by CDFG in 2003 to reflect CDFG final project reports.

b/ Does not include Nimbus Weir counts.

c/ Total includes Butte Creek fall spawner survey data.

d/ Survey methodology was variable, may not be comparable to other surveys.

e/ Change in estimation methodology (due to extremely high Battle Creek escapement in 2002).

f/ Total includes fall spawner estimates in Butte, Clear, Deer, Mill, and lower Battle creeks.

g/ Total does not include Cow Creek (estimated 2006 fall escapement of 3,987 adults and 143 jacks).

h/ Total does not include Cow, Bear, or Cottonwood creeks (estimated 2007 total fall escapement of 2,930 Chinook).

i/ Preliminary.

TABLE B-2. California Central Valley hatchery fall Chinook salmon spawning escapements in numbers of fish.^{a/}

Year or Average	Sacramento Hatcheries								San Joaquin Hatcheries						Central Valley Hatchery Totals	
	Coleman ^{b/}		Feather River		Nimbus		Totals		Mokelumne River		Merced River		Totals		Adults	Jacks
	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults ^{c/}	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks		
1971-1975	1,373	1,167	3,882	1,387	7,791	1,311	13,661	4,065	305	156	460	19	765	175	14,427	4,240
1976-1980	4,239	1,292	4,261	1,043	7,238	1,990	17,198	4,760	271	59	346	23	617	82	17,814	4,842
1981-1985	11,557	3,734	6,845	884	10,072	2,257	29,832	7,689	759	734	797	449	1,556	1,183	31,388	8,872
1986-1990	11,507	2,288	5,837	1,947	5,685	1,349	23,028	5,584	278	286	299	140	577	426	23,605	6,010
1991-1995	11,948	2,295	10,537	2,762	6,414	1,447	28,899	6,505	1,077	554	239	233	1,316	788	30,215	7,292
1996	18,848	2,330	6,494	1,613	7,177	474	32,519	4,417	1,828	2,055	395	746	2,223	2,801	34,742	7,218
1997	44,590	6,080	13,358	1,770	5,328	322	63,276	8,172	6,305	189	838	108	7,143	297	70,419	8,469
1998	42,400	1,951	17,567	1,322	9,949	1,839	69,915	5,113	2,686	585	347	452	3,033	1,037	72,948	6,150
1999	23,194	3,776	12,822	1,104	6,207	3,553	42,224	8,432	1,611	1,542	650	987	2,261	2,529	44,484	10,962
2000	20,793	866	16,470	1,676	10,312	848	47,575	3,390	4,637	887	1,615	331	6,252	1,218	53,827	4,608
2001	23,710	988	24,001	871	9,688	1,956	57,399	3,815	4,467	1,427	1,137	523	5,604	1,950	63,003	5,765
2002	61,946	4,112	17,516	2,991	6,231	3,586	85,693	10,689	5,800	2,119	1,250	588	7,050	2,707	92,743	13,396
2003	82,708	5,555	13,615	1,352	11,875	3,012	108,198	9,919	5,108	3,009	392	157	5,500	3,166	113,698	13,085
2004	51,557	16,672	15,769	5,535	12,741	13,659	80,067	35,866	5,477	4,879	456	594	5,933	5,473	86,000	41,339
2005	142,135	2,604	20,597	1,787	20,569	1,780	183,301	6,171	5,035	528	346	75	5,381	603	188,682	6,774
2006	56,966	1,056	13,400	634	8,322	406	78,688	2,096	2,801	1,338	136	15	2,937	1,353	81,625	3,449
2007 ^{d/}	11,558	220	5,091	172	4,590	7	21,239	399	1,004	39	70	9	1,074	48	22,313	447
GOALS ^{e/}	9,000	-	5,000	-	6,000	-	20,000	-	5,000	-	1,000	-	6,000	-	26,000	-

a/ For years prior to 2004, all numbers in this table were reviewed and updated by CDFG in 2003 to reflect CDFG final project reports.

b/ Fall spawning fish. Some spring run are included.

c/ Total adults in Sacramento Hatcheries include Tehama-Colusa Fish Facility for 1971 to 1985.

d/ Preliminary.

e/ Hatchery-specific goals, not PFMC goals.

TABLE B-3. Sacramento River late-fall, winter, and spring Chinook salmon spawning escapement estimates in numbers of fish.

Year or Average	Upper Sacramento River									Grand Totals	
	Late Fall ^{a/b/}		Winter ^{a/b/}		Spring						
	Adults	Jacks	Adults	Jacks	Tributary ^{c/} Adults and Jacks ^{g/}	Sacramento River ^{d/} Adults Jacks		Feather River ^{e/f/} Adults Jacks		Adults	Jacks
1971-1975	18,193	1,087	22,863	9,063	5,194	5,098	1,718	366	-	51,714	11,650
1976-1980	9,662	1,798	13,499	2,640	1,201	8,335	2,571	375	-	33,073	7,009
1981-1985	8,102	1,746	5,027	921	1,061	9,798	4,241	1,446	133	25,434	7,040
1986-1990	10,047	1,761	1,369	390	1,658	8,795	1,930	2,884	406	24,753	4,487
1991-1995	3,844 ^{h/}	383 ^{h/}	586	78	2,813	410	165	3,441	465	11,094	1,090
1996	1,003 ^{h/}	382 ^{h/}	708	629	2,322	314	64	5,571	810	9,918	1,886
1997	4,166 ^{h/}	412 ^{h/}	528	352	1,303	36	90	2,970	683	9,003	1,537
1998	40,185 ^{i/}	5,055 ^{i/}	2,079	923	23,609	624	491	6,240	506	72,738	6,974
1999	24,475 ^{i/}	3,986 ^{i/}	822	2,466	6,104	142	117	3,530	201	35,073	6,770
2000	11,060 ^{i/}	3,507 ^{i/}	563	789	5,504	94	38	3,657	315	20,878	4,649
2001	23,956 ^{i/}	998 ^{i/}	1,696	3,827	21,623 ^{j/}	981	0 ^{k/}	4,052	83	52,308	4,908
2002	39,700 ^{i/}	401 ^{i/}	7,614	1,555	20,198 ^{j/}	430	53	3,982	207	71,924	2,216
2003	9,295 ^{i/}	191 ^{i/}	6,172	3,585	21,798 ^{j/}	0	0	8,373	389	45,638	4,165
2004	13,552 ^{i/}	370 ^{i/}	2,588	4,604	12,556 ^{j/}	763	326	3,630	572	33,089	5,872
2005	14,437 ^{i/}	2,598 ^{i/}	3,521	1,778	21,319 ^{j/}	21	9	1,811 ^{l/}	24 ^{l/}	41,109	4,409
2006	14,950 ^{i/}	330 ^{i/}	4,792	2,623	10,669 ^{j/}	0	0	2,052 ^{l/}	9 ^{l/}	32,463	2,962
2007 ^{m/}	17,842 ^{i/}	751 ^{i/}	3,004	3,140	8,951 ^{j/}	225	22	2,747 ^{l/}	5 ^{l/}	32,769	3,918

a/ Estimated number of jacks and adults based on sampling at Red Bluff Diversion Dam (unpublished CDFG data). Beginning in 1987 for late-fall and winter and 1994 for fall, estimates have been based on historical run patterns and partial counts at Red Bluff Diversion Dam, due to the raising of the dam gates during the last part of fall, late-fall runs, and the first part of the winter run.

b/ Variable numbers of late-fall and winter run are trapped at Keswick Dam and spawned at Coleman or Livingston Stone Hatcheries.

c/ Natural spawning spring run which are isolated from fall run. Primarily Mill, Deer, and Butte Creeks.

d/ Sacramento River spring run estimates are the total Red Bluff Diversion Dam counts minus the spring run numbers in the upper Sacramento tributaries. If this number is less than or equal to zero, then upper Sacramento River spring run estimates are zero.

e/ Primarily fish returning to Feather River Hatchery.

f/ Spring run Chinook are not distinguished from fall run in the natural spawning surveys. They are reported in the fall run natural escapement numbers.

g/ No data available for age composition of tributary spring run.

h/ Primarily number of fish spawned at Coleman hatchery 1995-1997. No data are available for natural spawners, as gates were raised during the time coinciding with the late-fall run.

i/ Data from carcass counts of natural spawners and fish spawned at Coleman hatchery.

j/ Methodology change from using snorkel survey to carcass survey for Butte Creek spring run estimates.

k/ Jack proportion could not be determined.

l/ Methodology change for distinguishing spring run Chinook at Feather River Hatchery was implemented in 2005. Fish arriving at the hatchery prior to the spring Chinook spawning period were tagged and returned to the river. The spring Chinook escapement estimate was the number of these tagged fish that subsequently returned to the hatchery during the spring Chinook spawning period.

m/ Preliminary.

TABLE B-4. Summary of Klamath River fall Chinook salmon estimates in numbers of adults and jacks.

Year or Average	Category	Total Inriver Run	Inriver Harvest			Nonlanded Fishery Mortality	Spawning Escapement								
			Indian	Sport	Total		Klamath River			Trinity River			Total		
							Hatchery	Natural	Total	Hatchery	Natural	Total	Hatchery	Natural	Total
1978-1980	Adults	63,306	14,621	2,777	17,398	1,329	3,886	21,277	25,163	3,823	15,593	19,416	7,709	36,871	44,579
	Jacks	23,731	1,379	3,385	4,764	189	544	8,224	8,768	1,515	8,495	10,010	2,059	16,719	18,778
1981-1985	Adults	63,230	17,128	5,096	22,224	1,593	8,812	16,313	25,125	2,934	11,354	14,288	11,746	27,667	39,413
	Jacks	29,811	1,287	6,447	7,734	243	1,162	6,227	7,389	4,888	9,556	14,444	6,050	15,783	21,833
1986-1990	Adults	151,203	36,669	15,145	51,814	3,498	13,194	21,543	34,737	11,912	49,242	61,154	25,106	70,785	95,891
	Jacks	20,227	446	4,924	5,370	139	1,009	3,460	4,469	2,285	7,964	10,248	3,294	11,423	14,718
1991-1995	Adults	80,666	10,574	3,094	13,668	983	12,980	26,594	39,574	5,104	21,339	26,442	18,084	47,932	66,016
	Jacks	12,038	291	2,741	3,032	81	1,140	3,216	4,356	1,134	3,435	4,569	2,274	6,651	8,925
1996	Adults	175,773	56,476	12,766	69,242	5,172	13,622	38,680	52,302	6,411	42,646	49,057	20,033	81,326	101,359
	Jacks	9,532	190	2,312	2,502	64	543	1,696	2,239	249	4,478	4,727	792	6,174	6,966
1997	Adults	83,736	12,087	5,676	17,763	1,167	13,275	34,637	47,912	5,387	11,507	16,894	18,662	46,144	64,806
	Jacks	7,993	35	2,409	2,444	52	452	1,380	1,832	820	2,845	3,665	1,272	4,225	5,497
1998	Adults	90,647	10,187	7,710	17,897	1,043	14,923	18,028	32,951	14,296	24,460	38,756	29,219	42,488	71,707
	Jacks	4,639	53	1,108	1,161	28	403	881	1,284	192	1,974	2,166	595	2,855	3,450
1999	Adults	51,048	14,660	2,282	16,942	1,322	9,290	11,660	20,950	5,037	6,797	11,834	14,327	18,457	32,784
	Jacks	19,248	271	1,616	1,887	57	4,830	6,293	11,123	2,027	4,154	6,181	6,857	10,447	17,304
2000	Adults	218,077	29,415	5,650	35,065	2,673	71,635	58,388	130,023	25,976	24,340	50,316	97,611	82,728	180,339
	Jacks	10,246	303	1,582	1,885	58	839	2,891	3,730	1,070	3,503	4,573	1,909	6,394	8,303
2001	Adults	187,333	38,645	12,134	50,779	3,608	37,204	40,944	78,148	17,908	36,890	54,798	55,112	77,834	132,946
	Jacks	11,343	399	1,500	1,899	66	1,364	6,378	7,742	267	1,369	1,636	1,631	7,747	9,378
2002	Adults	160,788 ^{a/}	24,574	10,495	35,069	2,351	23,667	54,225	77,892	3,516	11,410	14,926	27,183	65,635	92,818
	Jacks	9,226	126	870	996	29	1,294	1,529	2,823	1,037	2,338	3,375	2,331	3,867	6,198
2003	Adults	191,949	30,034	9,680	39,714	2,810	31,970	55,423	87,393	29,812	32,219	62,031	61,782	87,642	149,424
	Jacks	3,845	44	814	858	21	290	848	1,138	574	1,254	1,828	864	2,102	2,966
2004	Adults	79,191	25,803	4,003	29,806	2,325	10,582	10,959	21,541	12,399	13,120	25,519	22,981	24,079	47,060
	Jacks	9,691	168	2,741	2,909	71	937	891	1,828	1,044	3,839	4,883	1,981	4,730	6,711
2005	Adults	65,227	8,016	1,985	10,001	738	13,955	13,554	27,509	13,744	13,235	26,979	27,699	26,789	54,488
	Jacks	2,296	70	1,030	1,100	27	42	398	440	59	670	729	101	1,068	1,169
2006	Adults	61,374	10,283	62	10,345	1,344	11,604	14,264	25,868	7,918	15,899	23,817	19,522	30,163	49,685
	Jacks	26,935	415	5,527	5,942	149	2,386	6,516	8,902	4,076	7,866	11,942	6,462	14,382	20,844
2007 ^{b/}	Adults	130,506	27,381	5,901	33,282	2,501	16,969	21,280	38,249	18,023	38,451	56,474	34,992	59,731	94,723
	Jacks	1,661	21	356	377	9	180	232	412	34	829	863	214	1,061	1,275
GOAL	Adults														≥35,000

a/ Total inriver run includes an estimated 30,550 fish that died prior to spawning in September 2002.

b/ Preliminary.

TABLE B-5. Estimates of Yurok and Hoopa Valley reservation Indian gillnet Chinook harvest in numbers of fish.

Year	Area	Spring Run			Fall Run		
		Jack	Adult	Total	Jack	Adult	Total
2001	Commercial:Estuary	79	4,637	4,716	63	7,011	7,074
	Upper Klamath	1	58	59	1	51	52
	Subsistence:Estuary	152	8,846	8,998	198	21,956	22,154
	Middle Klamath	0	134	134	28	1,697	1,725
	Upper Klamath	19	1,504	1,523	49	2,976	3,025
	Trinity River	46	4,164	4,210	60	4,954	5,014
	Total	297	19,343	19,640	399	38,645	39,044
2002	Commercial:Estuary	7	1,852	1,859	7	8,952	8,959
	Upper Klamath	-	-	-	-	-	-
	Subsistence:Estuary	25	6,551	6,576	10	11,197	11,207
	Middle Klamath	70	1,310	1,380	10	729	739
	Upper Klamath	24	2,205	2,229	31	2,528	2,559
	Trinity River	40	3,052	3,062	68	1,168	1,236
	Total	166	14,970	15,136	126	24,574	24,700
2003	Commercial:Estuary	4	779	783	11	17,084	17,095
	Upper Klamath	0	0	0	0	0	0
	Subsistence:Estuary	10	1,800	1,810	4	5,604	5,608
	Middle Klamath	0	2,355	2,355	5	1,376	1,381
	Upper Klamath	0	1,730	1,730	12	3,199	3,211
	Trinity River	7	2,380	2,387	12	2,771	2,783
	Total	21	9,044	9,065	44	30,034	30,078
2004	Commercial:Estuary	2	408	410	13	14,251	14,264
	Upper Klamath	0	0	0	13	540	554
	Subsistence:Estuary	10	2,178	2,188	62	6,787	6,848
	Middle Klamath	6	2,346	2,352	14	577	591
	Upper Klamath	11	1,715	1,726	46	1,959	2,005
	Trinity River	62	1,944	2,006	20	1,689	1,709
	Total	91	8,591	8,682	168	25,083	25,971
2005	Commercial:Estuary	0	0	0	0	0	0
	Upper Klamath	0	0	0	0	0	0
	Subsistence:Estuary	0	477	477	21	2,293	2,314
	Middle Klamath	0	518	518	5	464	469
	Upper Klamath	0	1,320	1,320	33	2,851	2,884
	Trinity River	17	1,858	1,858	11	2,409	2,420
	Total	33	7,286	7,302	70	8,017	8,087
2006	Commercial:Estuary	0	0	0	0	0	0
	Upper Klamath	0	0	0	0	0	0
	Subsistence:Estuary	8	302	310	30	2,726	2,756
	Middle Klamath	3	1,113	1,116	93	1,310	1,403
	Upper Klamath	36	1,257	1,293	147	2,086	2,233
	Trinity River	58	1,632	1,690	145	4,161	4,306
	Total	105	4,304	4,409	415	10,283	10,698
2007 ^{a/}	Commercial:Estuary	0	2,300	2,300	15	21,087	21,102
	Upper Klamath	0	0	0	0	0	0
	Subsistence:Estuary	0	1,332	1,332	1	2,235	2,236
	Middle Klamath	0	200	200	0	411	411
	Upper Klamath	0	631	631	5	1,350	1,355
	Trinity River	6	1,349	1,355	0	2,298	2,298
	Total	6	5,812	5,818	21	27,381	27,402

a/ Preliminary.

TABLE B-6. Shasta River fall Chinook salmon weir counts or spawning escapement estimates in numbers of fish.^{a/}

Year or Average	Adults	Jacks	Total
1931-1935 ^{b/}	37,474	12,690	50,164
1936-1940	26,165	8,223	34,389
1941-1945	9,654	3,129	12,783
1946-1950	1,862	178	2,040
1951-1955	1,577	370	1,947
1956-1960	6,146	1,074	7,220
1961-1965	15,167	4,388	19,555
1966-1970	10,472	1,410	11,882
1971-1975	6,297	2,866	9,163
1976-1980 ^{c/}	6,506	3,194	9,700
1981-1985 ^{d/}	4,560 ^{d/}	1,942	6,503
1986-1990 ^{e/}	2,403	318	2,721
1991-1995	1,891	184	2,075
1991	716	10	726
1992	520	66	586
1993	1,341	85	1,426
1994	3,363	1,840	5,203
1995	12,816	695	13,511
1996	1,404	46	1,450
1997	1,667	334	2,001
1998	2,466	76	2,542
1999	1,296	1,901	3,197
2000	11,025	1,271	12,296
2001	8,452	2,641	11,093
2002	6,432	386	6,818
2003	4,134	155	4,289
2004	833	129	962
2005 ^{f/}	2,018	37	2,055
2006 ^{f/}	789	1,395	2,184
2007 ^{f/}	2,009	27	2,036

a/ From 1930-1937, 1957-1987 and 1991-1995, the counts were made near the river mouth. From 1938-1955, they were made 6.5 miles upstream from the mouth; considerable spawning occurred downstream from the racks in these years. From 1988-1990, escapements were estimated from mark-recapture data (spawning surveys).

b/ Commercial fishing in lower Klamath River closed by the state after the 1933 season.

c/ Gillnetting resumed in lower 20 miles of Klamath River by Hoopa Valley Indian Reservation fishers in 1976.

d/ Includes 276 females taken to Iron Gate Hatchery in 1981.

e/ Low water conditions appeared to hinder entry into the river in 1988.

f/ Preliminary.

TABLE B-7. Summary of California North Coast salmon spawning stock surveys in numbers of fish.

Year	Cañon Creek (Mad River) ^{a/b/}			Sprowl Creek (Eel River) ^{a/c/}			Tomki Creek (Eel River) ^{d/}
	Surveys	Chinook	Coho	Surveys	Chinook	Coho	Chinook
1978-1979	-	-	-	2	534	23	-
1979-1980	-	-	-	2	572	0	2,410
1980-1981	-	-	-	1	164	4	317
1981-1982	3	23	0	2	121	0	1,153
1982-1983	3	68	0	6	169	1	1,807
1983-1984	2	137	0	2	82	0	-
1984-1985 ^{e/}	1	16	0	6	67	13	1,292
1985-1986	10	514	14	6	320	0	3,558
1986-1987 ^{e/}	4	90	3	5	307	13	2,173
1987-1988	4	117	29	3	2,187	4	3,666
1988-1989	2	69	7	3	339	12	556
1989-1990 ^{e/}	4	9	9	5	89	14	-
1990-1991	1	0	3	2	0	0	-
1991-1992 ^{e/}	2	8	0	2	159	0	3
1992-1993 ^{e/}	3	57	1	2	142	2	15
1993-1994	3	20	0	4	171	36	5
1994-1995	3	33	3	7	52	0	21
1995-1996 ^{e/}	1	93	4	3	136	8	69
1996-1997	1	129	4	3	106	8	84
1997-1998	2	55	1	4	97	0	39
1998-1999	2	66	0	4	79	11	45
1999-2000 ^{e/}	8	162	1	7	34	1	24
2000-2001 ^{e/}	3	79	3	4	12	0	50
2001-2002	2	45	6	5	136	25	162 ^{f/}
2002-2003	3	402	1	6	267	17	5 ^{f/}
2003-2004 ^{e/}	2	79	1	5	106	8	137 ^{f/}
2004-2005 ^{e/}	4	86	0	5	199	36	115 ^{f/}
2005-2006	1	270	0	5	201	13	77 ^{f/}
2006-2007 ^{g/}	4	152	2	8	37	9	20 ^{f/}
2007-2008 ^{h/}	2	88	1	4	65	6	55 ^{f/}

a/ Numbers reflect maximum annual counts of live fish and carcasses with adults and jacks combined. Counts are not shown in years where visibility is too poor to conduct surveys.

b/ Survey area was from mouth to falls (2 miles).

c/ Survey area was the mainstem and West Fork (4.5 miles).

d/ Total run size estimate including jacks and adults.

e/ Low flows this season appeared to increase mainstem spawning and decrease tributary spawning.

f/ Survey methodology changed to using index sites and is not comparable to previous estimates.

g/ Cañon totals do not include 20 additional salmonids (species unknown) observed during poor visibility.

h/ Preliminary data. Cañon and Sprowl creek totals do not include 24 and 61 unknown salmonids, respectively.

TABLE B-8. Peak spawning counts in index areas for selected south/local migrating Oregon coastal fall Chinook stocks.

Year or Avg.	Deep Creek (Pistol River) (0.4 mile)		Big Emily Creek (Chetco River) (1.0 mile)		Bear Creek (Winchuck River) (0.8 mile)		Index (fish per mile)	
	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks
1961-1965	6	1	-	-	22	1	-	-
1966-1970	31	3	-	-	36	2	-	-
1971-1975	5	0	211	12	25	2	130	7
1976-1980	2	1	124	32	18	1	65	14
1981-1985	24	2	62	10	13	1	45	6
1986-1990	9 ^{a/}	1 ^{a/}	58	12	10	2	35	7
1991-1995	12	9	74	10	16	2	46	10
1996	81	9	79	7	27	5	85	10
1997	17	1	60	5	41	1	41	3
1998	46	11	52	3	19	2	53	7
1999	58	3	12	0	10	0	36	1
2000	26	3	63	6	11	1	45	5
2001	25	2	49	2	9	3	38	3
2002	62	7	70	3	15	0	67	5
2003	20	7	28	5	12	1	27	6
2004	97	19	29	4	11	1	62	11
2005	15	2	16	3	1	0	15	2
2006	22	3	24	2	5	1	23	3
2007 ^{b/}	44	0	14	4	6	1	29	3

a/ Pistol River was subject to several "slope failures" in 1986 resulting in severe short-term alterations in gravel bars and spawning index areas. Considerable debris and siltation severely limited Chinook surveys resulting in "0" counts in Deep Creek index areas through December.

b/ Preliminary.

TABLE B-9. Counts of natural and hatchery spring Chinook salmon at Gold Ray Dam on the Rogue River and at Winchester Dam on the North Umpqua River in thousands of fish.

Year or Avg.	Gold Ray Dam, Rogue River ^{a/}				Winchester Dam, Umpqua River ^{a/}			
	Natural	Hatchery	Total	Jacks ^{b/}	Natural	Hatchery	Total	Jacks ^{b/}
1942-1945	35.1	-	35.1	4.9	-	-	-	-
1946-1950	24.7	-	24.7	3.0	2.7	-	2.7	0.5
1951-1955	21.4	-	21.4	4.2	4.2	0.9	4.9	1.0
1956-1960	19.8	-	19.8	3.4	4.4	0.9	5.4	0.7
1961-1965	37.7	-	37.7	6.4	6.4	1.8	8.2	1.8
1966-1970	33.9	-	33.9	5.5	7.2	4.5	11.8	3.2
1971-1975	26.0	0.8	26.8	5.0	7.3	6.2	13.5	3.8
1976-1980	25.8	6.3	32.1	7.0	5.8	3.9	9.7	3.2
1981-1985	16.4	6.2	22.6	7.3	5.2	3.5	8.7	2.5
1986-1990	28.5	39.2	67.7	14.9	7.5	4.1	11.6	2.5
1991-1995	9.7	18.4	28.0	3.9	3.5	2.5	6.0	1.1
1996	10.3	26.3	36.6	3.4	4.3	2.2	6.5	1.0
1997	9.6	32.2	41.8	2.8	3.3	2.5	5.8	16.0
1998	3.7	12.3	16.0	2.8	4.0	2.9	6.9	1.5
1999	6.0	15.0	21.0	1.9	2.8	4.6	7.4	3.1
2000	3.4	26.8	30.2	3.1	3.4	9.2	12.6	4.6
2001	9.3	23.9	33.2	2.3	6.1	14.6	20.7	4.7
2002	7.0	40.8	47.8	3.2	6.8	17.4	24.2	3.1
2003	19.3	22.6	41.9	3.0	7.9	12.3	20.2	4.1
2004	13.3	26.0	39.3	3.8	5.4	10.1	15.4	2.5
2005	5.8	12.3	18.1	1.3	3.6	5.5	9.0	1.3
2006	4.8	7.0	11.7	2.2	2.6	3.5	6.1	1.7
2007 ^{c/}	3.5	7.7	11.2	1.6	2.4	4.2	6.6	1.7

a/ Jacks included in natural, hatchery, and total counts.

b/ Jacks include all Chinook less than 20 inches prior to 1978 and all Chinook less than 24 inches beginning in 1978.

c/ Preliminary.

TABLE B-10. Rogue River fall Chinook carcass counts in numbers of fish.

Year or Avg.	Carcass Counts		
	Adults	Jacks	Total
1977-1980	5,256	1,004	6,259
1981-1985	3,906	1,009	4,915
1986-1990	16,797	1,527	18,324
1990-1995	4,387	316	4,703
1996	2,448	121	2,569
1997	1,643	68	1,711
1998	3,601	40	3,641
1999	2,493	157	2,650
2000	3,366	226	3,592
2001	6,380	772	7,152
2002	11,836	905	12,741
2003	14,620	983	15,603
2004	5,326 ^{a/}	250	5,576
2005 ^{b/}	-	-	-
2006 ^{b/}	-	-	-
2007 ^{b/}	-	-	-

a/ In 2004, one of the standard survey sections was not sampled. In the previous two years, this section accounted for 33% of the total adult carcass counts.

b/ Surveys were not conducted.

TABLE B-11. Peak counts for north migrating Oregon coastal Chinook stocks on selected fall Chinook spawning index stream surveys.

Year or Average	Tributaries (River)																		Index Fish Per Mile	
	Humbug (Nehalem) (1.0 mile)		Tillamook (1.8 mile)		Niagara (Nestucca) (0.4 mile)		Sunshine (Siletz) (1.2 mile)		Grant (Yaquina) (1.7 mile)		Buck (Alsea) (1.0 mile)		Siuslaw Lake (0.8 mile)		W.F. Millicoma (Coos) (0.5 mile)		Salmon (Coquille) (0.8 mile)			
	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks	Adults	Jacks		
1961-1965	95	22	116	25	72	5	59	13	43	13	28	9	61	15	2	1	23	13	54	13
1966-1970	57	3	93	27	47	6	30	5	61	13	26	16	134	40	6	1	26	9	52	13
1971-1975	101	26	55	5	55	4	40	5	64	8	17	3	94	49	18	13	15	5	50	14
1976-1980	143	12	61	6	32	2	47	5	127	23	22	3	166	37	31	28	31	10	73	14
1981-1985	163	18	95	9	78	6	55	2	178	24	47	6	149	31	21	2	45	7	89	11
1986-1990	136	4	154	8	118	3	54	2	240	25	100	6	427	44	13	5	49	6	140	11
1991-1995	65	2	92	6	103	3	60	2	153	10	44	4	395	18	49	7	84	5	115	6
1996	86	2	60	0	40	0	122	0	a/	a/	62	2	614	29	92	3	29	3	147	5
1997	162	1	47	1	24	1	60	0	a/	a/	49	3	325	9	12	0	108	3	105	2
1998	93	2	42	1	42	0	83	3	a/	a/	78	0	176	2	29	11	191	7	98	3
1999	116	3	38	1	60	2	36	3	a/	a/	55	5	478	14	14	3	136	8	124	5
2000	175	3	40	3	32	2	63	1	a/	a/	38	3	205	18	5	0	83	9	85	5
2001	220	4	62	6	53	7	195	3	a/	a/	95	6	711	49	30	5	153	22	203	14
2002	311	1	137	3	124	1	221	1	a/	a/	114	6	834	22	51	12	218	9	268	7
2003	215	6	135	5	27	1	120	3	a/	a/	145	1	1,230	37	209	31	147	2	297	11
2004	196	3	71	1	76	1	19	0	a/	a/	91	5	988	16	40	4	101	5	211	5
2005	124	3	b/	b/	74	2	54	1	a/	a/	40	1	302	5	17	2	61	2	118	3
2006	31	0	65	0	67	0	118	0	a/	a/	22	0	165	0	7	1	129	8	106	1
2007 ^{c/}	91	1	34	2	20	0	6	0	a/	a/	17	1	132	2	14	3	2	0	42	1

a/ Survey discontinued in 1994; landowner would not allow access.

b/ Surveys were not conducted in 2005.

c/ Preliminary.

TABLE B-12. Estimates of minimum inriver run size, catch, and escapement in numbers of Columbia River adult spring Chinook destined for areas below Bonneville Dam. (Page 1 of 1).

Year or Average	Minimum Inriver Run Size	Tributary Runs									
		Lower River Catch ^{a/}		Willamette			Sandy	Cowlitz ^{c/}	Lewis ^{c/}	Kalama	Hatchery Escapement ^{d/}
		Commercial	Sport	Run Size	L. Willamette Sport Catch	Will. Falls Escapement ^{b/}					
1971-1975	84,000	13,800	3,700	53,300	17,000	34,300	-	11,900	200	1,100	20,000
1976-1980	92,160	6,160	2,720	51,240	14,380	31,420	975	19,680	2,980	2,020	26,580
1981-1985	130,000	6,680	1,840	67,700	15,620	35,580	1,940	19,960	4,220	3,740	28,840
1986-1990	176,103	11,980	4,330	103,640	21,140	58,760	2,425	10,691	11,340	1,877	32,460
1991-1995	122,059	3,680	2,875	68,600	18,180	32,580	5,088	6,801	5,870	1,976	23,700
1996	55,879	149	0	34,800	6,100	20,400	3,997	1,787	1,730	627	15,900
1997	54,324	300	0	35,300	1,900	26,200	4,625	1,877	2,196	505	18,100
1998	54,254	100	49	45,100	2,800	33,100	3,768	1,055	1,611	407	22,900
1999	64,964	349	0	54,200	5,500	38,900	3,985	2,069	1,753	977	25,900
2000	73,904	1,149	249	57,500	9,000	37,594	3,641	2,199	2,515	1,418	24,100
2001	102,558	3,700	4,300	80,300	7,600	52,700	5,329	1,649	3,777	1,784	29,000
2002	151,310	7,900	5,800	121,700	10,800	83,100	5,903	5,019	3,554	2,883	58,300
2003	166,505	1,900	8,200	126,600	13,500	87,600	5,600	15,890	5,104	4,528	45,725
2004	201,093	8,500	7,500	144,400	12,000	95,200	12,675	16,712	11,090	4,573	67,910
2005	86,728	3,400	4,400	61,000	5,800	35,453	7,475	9,200	3,400	3,100	32,891
2006	92,193	3,000	2,900	59,700	7,200	36,851	4,812	7,000	7,500	5,600	35,565
2007 ^{e/}	64,800	1,900	2,600	40,000	5,700	23,098	3,400	3,700	6,700	7,300	29,377

a/ Includes some upriver origin spring Chinook through 1980. Beginning in 1981, the lower river catch of lower river spring Chinook is based on mark recoveries rather than the timing of the catch, as in previous years. Since 1986, GSI and VSI techniques have been used for stock composition analysis. Commercial catch includes Select Area fisheries. Sport catch is mainstem Columbia River, does not include tributaries. Catch may include small numbers of jacks. Sport fishery closed in 1995 to 1997.

b/ Prior to 1988, the escapement goal at Willamette Falls was 30,000 to 35,000. Beginning in 1988, the goal was dependent on run size under the Willamette Basin Fish Management Plan. Since 2001, hatchery escapement targets are set in the Fisheries Management and Evaluation Plan developed by ODFW. Lower Willamette sport catch may include small numbers of jacks.

c/ Includes hatchery escapement, tributary recreational catch, and natural spawning escapement for 1975 to present. The years 1971-1973 are based on using the 1975-1976 Cowlitz River recreational fishery adult harvest rates.

d/ Includes hatcheries operated by all agencies. Values are included in the totals for the tributary runs.

e/ Preliminary.

TABLE B-13. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult spring Chinook destined for areas above Bonneville Dam.^{a/} This table includes Snake River summer Chinook. (Page 1 of 1)

Year or Avg.	Inriver Run Size	Lower River Catch ^{b/}		Mainstem Treaty Indian Catch				Snake River Escapement ^{e/}		U. Columbia	Hatchery Escapement
		Commercial	Sport	Bonneville Dam Count	Commercial ^{d/}	Ceremonial/ Subsistence	Zone 6 Escapement ^{d/}	Total	Wild	River Escapement ^{f/}	
1976-1980	55,897	185	0	55,712	259	1,714	53,740	9,317	6,413	8,138	5,703
1981-1985	70,057	1,706	393	67,959	1,024	2,545	64,390	18,295	10,679	13,943	12,887
1986-1990	107,535	2,378	1,356	103,800	186	6,771	96,843	29,893	9,755	15,359	27,883
1991-1995	63,040	511	710	61,819	15	3,730	58,074	14,275	7,013	12,173	13,883
1996	55,299	46	17	55,236	0	2,911	52,325	7,037	3,788	2,430	5,211
1997	123,824	53	13	123,758	14	8,309	115,435	44,849	4,406	6,780	46,089
1998	43,512	27	14	43,471	1	2,224	41,246	14,337	7,391	4,124	9,872
1999	42,582	28	21	42,533	1	1,983	40,549	6,741	2,856	4,150	7,303
2000	186,141	265	102	185,774	1,354	9,973	174,447	38,064	8,255	19,143	37,039
2001	437,910	2,543	22,714	412,653	43,715	10,985	357,953	188,145	45,335	50,379	167,281
2002	331,303	10,150	16,213	304,940	24,254	9,208	271,478	99,070	30,248	34,083	89,725
2003	242,638	3,524	9,615	229,499	9,205	9,090	211,204	87,999	32,366	18,136	66,540
2004	221,606	6,240	17,041	198,325	8,370	9,114	180,841	81,423	21,401	13,521	67,162
2005	106,920	2,288	7,235	97,397	1	6,163	91,233	33,258	10,156	14,148	32,838
2006	132,140	1,822	4,161	126,158	0	8,401	117,757	30,042	9,490	13,535	28,151
2007 ^{g/}	86,230	1,483	3,918	80,829	3	6,141	74,685	30,804	9,733	6,627	27,466
GOAL				115,000							

a/ Spring Chinook accounting ends on June 15. Chinook formerly managed separately as Snake River summer Chinook are now grouped with all upriver spring Chinook because of overlap in run timing. Snake River summer Chinook have been moved from Table B-14 to this table.

b/ Includes some lower river origin spring Chinook through 1980. Beginning in 1981, the lower river catch of upriver spring Chinook is based on mark recoveries rather than timing of the catch as in previous years. Since 1986, GSI techniques have been used for stock composition analysis. Catch includes estimated miscellaneous fishery-related impacts from test fisheries, commercial shad fisheries, and Select Area commercial gillnet fisheries beginning in 1979 and catch and release mortalities from selective fisheries beginning in 2001.

c/ Spring season fishery closed in 1975, 1976, and from 1978 to 2000. Spring Chinook landed during those years were from the winter season fishery.

d/ Bonneville Dam count minus Zone 6 mainstem commercial and ceremonial/subsistence treaty Indian harvest.

e/ Count at uppermost Snake River Dam (Little Goose in 1971-1974 and Lower Granite plus Tucannon wild escapement after 1974) plus harvest below Lower Granite Dam.

f/ Priest Rapids Dam count.

g/ Preliminary.

TABLE B-14. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult summer Chinook destined for areas above Bonneville Dam.^{a/} This table does not include Snake River summer Chinook. (Page 1 of 1)

Year or Avg.	Inriver Run Size	Lower River Catch ^{b/}		Bonneville Dam Count	Mainstem Treaty Indian Catch		Zone 6 Escapement ^{e/}	U. Columbia River Escapement ^{f/}
		Commercial ^{c/}	Sport		Commercial ^{d/}	Ceremonial/ Subsistence		
1976-1980	22,566	81	0	22,485	38	1,047	21,401	18,161
1981-1985	17,092	55	0	17,037	304	654	16,079	12,202
1986-1990	21,668	71	7	21,590	708	194	20,689	15,785
1991-1995	13,348	30	15	13,303	0	227	13,077	13,347
1996	12,333	15	27	12,291	0	374	11,917	10,995
1997	18,277	6	19	18,252	0	270	17,982	13,107
1998	16,332	1	27	16,304	0	335	15,969	13,387
1999	22,347	1	41	22,305	0	411	21,894	20,898
2000	23,169	0	25	23,144	0	209	22,935	22,306
2001	54,935	1	64	54,870	150	542	54,286	53,170
2002	92,820	8	1,503	91,309	42	2,019	89,225	96,326
2003	83,120	36	2,007	81,077	3,587	710	80,098	83,004
2004	65,446	236	1,240	63,970	8,004	390	63,542	67,060
2005	60,060	2,574	1,622	55,684	6,415	1,227	54,563	61,227
2006	76,196	5,017	3,360	67,819	15,771	548	67,231	57,236
2007 ^{g/}	37,190	1,122	2,369	33,699	4,564	811	32,828	30,644
GOAL	29,300 ^{h/}							

a/ Summer Chinook accounting begins on June 16. Chinook managed as Snake River summer Chinook prior to 2004 are now grouped with all upriver spring Chinook because of overlap in run timing. As of 2004, they have been moved from this table to Table B-13.

b/ Includes estimated miscellaneous fishery-related impacts from test fisheries, commercial shad fisheries, and terminal area commercial gillnet fisheries beginning in 1979. Includes catch and release mortality in selective fisheries beginning in 2002.

c/ No directed commercial summer Chinook fishery from 1964 to 2003. Landings during those years are bycatch from commercial shad and sockeye fisheries.

d/ No directed commercial summer Chinook fishery from 1965 to 2003. Landings during those years are bycatch from commercial sockeye fishery.

e/ Bonneville Dam count minus Zone 6 mainstem commercial and ceremonial/subsistence treaty Indian harvest.

f/ Priest Rapids Dam count.

g/ Preliminary.

h/ Comanager goal established in 2004 associated with regrouping Snake River summer Chinook with Snake River spring Chinook.

TABLE B-15. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult Spring Creek Hatchery (SCH) stock fall Chinook.^{a/} Page 1 of 1

Year or Average	Inriver Run Size	Harvest				Escapement	
		Bonneville Dam Count	Treaty Indian Commercial and Subsistence	Non-Indian		Natural	Hatchery ^{d/}
				Commercial ^{b/}	Sport		
1971-1975	105,700	67,600	29,000	37,900	300	2,900	17,000
1976-1980	116,522	83,000	32,533	31,794	131	3,884	21,972
1981-1985	63,342	49,780	24,637	9,747	580	2,711	15,955
1986-1990	16,673	10,200	6,080	2,920	820	1,500	4,600
1991-1995	30,192	25,564	11,360	2,067	1,280	1,460	9,700
1996	33,137	30,300	21,100	1,700	900	1,300	7,700
1997	27,377	23,300	10,329	0	2,981	4,612	8,688
1998	20,158	17,100	6,592	197	2,556	2,731	3,224
1999	50,189	46,800	28,197	258	2,617	3,338	14,488
2000	20,527	18,400	7,903	1,141	897	4,085	6,257
2001	124,951	115,800	52,124	3,693	3,302	5,063	36,663
2002	158,299	145,200	48,350	11,485	6,654	8,069	67,436
2003	180,592	161,735	48,204	9,850	7,659	27,894	56,935
2004	175,245	164,482	59,941	3,690	5,614	14,084	68,932
2005	103,526	98,322	49,471	3,981	3,049	4,667	31,977
2006	27,917	21,197	13,400	1,774	654	1,931	9,889
2007 ^{d/}	17,100	16,560	8,170	190	100	1,080	7,310
GOAL							7,000 ^{e/}

a/ Based on Columbia River fall Chinook database, WDFW, unpublished.

b/ Includes Select Area fisheries.

c/ Does not include strays to hatcheries below Bonneville Dam. Includes fall Chinook tules trapped at Bonneville Dam, 1986-1994 and 1998.

d/ Preliminary estimates based on inseason run updates.

e/ Escapement goal was changed from 8,200 fish to 7,000 fish, or 4,000 females, in 1994.

TABLE B-16. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult lower river hatchery (LRH) stock fall Chinook.^{a/} (Page 1 of 1)

Year or Average	Inriver Run Size	Harvest			Escapement	
		Treaty Indian	Non-Indian		Natural	Hatchery ^{d/}
		Commercial	Commercial ^{b/}	Sport ^{c/}		
1971-1975	175,900	0	78,100	5,400	49,200	43,200
1976-1980	145,377	20	59,400	4,380	36,940	44,620
1981-1985	107,163	851	25,604	4,486	37,755	36,846
1986-1990	199,938	655	93,794	17,420	38,774	48,821
1991-1995	55,519	238	2,871	4,998	19,915	27,419
1996	75,495	360	3,899	4,641	23,909	42,662
1997	57,393	0	2,369	7,704	22,663	24,657
1998	45,265	0	844	4,519	16,713	23,035
1999	39,933	0	2,234	6,118	12,551	19,030
2000	26,997	0	860	3,212	10,714	12,211
2001	94,331	0	4,428	7,443	39,434	42,996
2002	156,444	279	9,928	15,353	80,670	50,138
2003	154,983	0	9,216	14,213	97,089	34,465
2004	109,055	475	13,122	11,870	53,399	30,103
2005	78,293	186	9,219	10,140	33,598	25,042
2006	58,319	237	5,919	9,449	26,633	15,957
2007 ^{e/}	35,000	0	1,100	5,260	9,920	18,220
GOAL						Hatchery Production

a/ Based on Columbia River fall Chinook database, WDFW, unpublished.

b/ Includes select area fisheries.

c/ Includes tributary catches.

d/ Does not include strays to hatcheries above Bonneville Dam or fish trapped at Bonneville Dam.

e/ Preliminary estimates based on inseason run updates.

TABLE B-17. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult lower river wild (LRW) stock fall Chinook.^{a/} (Page 1 of 1)^f

Year or Average	Inriver Run Size	Harvest			Escapement	
		Treaty Indian	Non-Indian		Natural	Hatchery
		Commercial	Commercial	Sport ^{b/}		
1971-1975	59,700	0	27,900	2,100	29,400	100
1976-1980	26,963	20	11,720	1,220	13,720	240
1981-1985	16,287	0	1,940	1,320	12,480	480
1986-1990	32,600	60	10,689	3,251	18,383	181
1991-1995	14,761	0	2,159	2,433	10,101	68
1996	14,566	0	325	234	13,914	93
1997	12,323	0	0	1,082	11,241	0
1998	7,253	0	0	667	6,493	93
1999	3,349	0	18	0	3,257	74
2000	10,234	0	604	0	9,422	208
2001	15,721	0	1,382	729	13,610	0
2002	25,171	161	1,801	3,245	19,654	50
2003	26,021	0	3,391	4,962	17,668	0
2004	22,327	0	2,343	3,638	16,346	0
2005	16,767	0	2,240	2,632	11,725	170
2006	18,105	0	2,546	2,801	12,758	0
2007 ^{c/}	10,040	0	300	1,340	8,410	0
GOAL					5,700 ^{d/}	

a/ Based on Columbia River fall Chinook database, WDFW, unpublished.

b/ Includes tributary catches.

c/ Preliminary estimates based on inseason run updates.

d/ Escapement objective is for North Lewis River, but escapement numbers include other fish. The escapement objective for the North Lewis River was met for all years except 1998-1999.

TABLE B-18. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult upriver bright (URB) stock fall Chinook destined for areas above McNary Dam and the Deschutes River.^{a/} (Page 1 of 1)

Year or Average	Harvest						Escapement					
	Inriver Run Size	Bonneville Dam Count	Treaty Indian Commercial and Subsistence	Non-Indian		Natural ^{c/}	Hatchery	Deschutes above Sheares Falls	McNary Dam Count	Ice Harbor Dam Count	Total Lower Granite Count	SRW L. Granite Dam Count ^{d/}
				Commercial	Sport ^{b/}							
1971-1975	110,500	80,400	35,100	29,300	3,100	36,800	2,600	NA	39,500	5,600	-	-
1976-1980	92,301	72,360	32,160	19,180	980	29,480	1,980	NA	31,080	1,160	532	532
1981-1985	111,873	94,120	26,700	13,880	3,020	46,060	8,100	NA	51,042	1,583	586	450
1986-1990	291,407	222,337	100,379	61,499	13,613	90,709	13,231	5,023	107,252	4,369	691	289
1991-1995	105,302	99,028	20,813	5,000	5,095	51,424	9,419	5,545	61,362	3,352	903	473
1996	143,155	135,499	29,868	3,717	8,918	59,598	15,905	8,759	73,929	3,810	1,308	639
1997	161,735	152,941	42,637	1,429	11,506	68,889	13,114	20,678	67,192	2,752	1,451	797
1998	141,575	137,509	33,760	770	8,137	54,297	18,798	10,923	63,791	4,220	1,909	306
1999	165,889	155,756	38,822	2,133	15,173	48,372	30,272	3,997	78,356	6,586	3,381	905
2000	156,595	145,104	36,501	5,551	10,545	66,512	10,841	3,230	66,378	6,509	3,602	1,148
2001	232,366	219,801	35,422	8,151	12,648	92,194	21,143	11,161	110,517	13,635	8,915	5,163
2002	279,548	257,711	57,405	6,881	25,651	123,446	17,299	12,252	141,682	15,319	12,351	2,116
2003	374,154	341,208	49,060	15,930	25,918	176,865	12,356	12,590	179,970	20,903	11,732	3,856
2004	362,804	336,585	46,566	19,760	22,276	148,028	23,137	11,879	170,648	21,100	14,960	4,756
2005	278,539	256,119	45,776	8,464	23,980	115,612	23,299	14,092	131,550	14,677	11,170	2,954
2006	230,390	132,632	44,565	8,757	14,515	79,852	15,197	13,374	90,973	10,272	8,048	2,483
2007 ^{e/}	122,000	113,120	9,870	4,360	7,870	59,464	4,596	NA	51,990	13,408	10,195	NA
GOAL									40,000 ^{f/}			

a/ Based on Columbia River fall Chinook database, WDFW, unpublished. Does not include hatchery URB Chinook reared and released below McNary Dam.

b/ Includes tributary and mainstem catches.

c/ Includes Deschutes, Yakima, Upper Columbia, and Snake River escapements.

d/ Snake River wild; adjusted for stray hatchery fish. Includes wild fish hauled to Lyons Ferry Hatchery.

e/ Preliminary based on inseason run update.

f/ FMP goal. The U.S. v Oregon parties managed for an escapement of 45,000 between 1990 and 1993 at McNary Dam to account for increased hatchery brood stock needs and concern for the Snake River wild fall Chinook stock. Starting in 1994, inriver fisheries were based on ESA consultation standards, rather than a McNary Dam escapement goal.

TABLE B-19. Estimates of inriver run size, catch, and escapement in numbers of Columbia River adult mid-Columbia bright (MCB) stock fall Chinook destined for areas below McNary Dam, not including the Deschutes River.^{a/} (Page 1 of 1)

Year or Average	Inriver Run Size	Harvest				Escapement		
		Bonneville Dam Count	Treaty Indian Commercial and Subsistence	Non-Indian		Natural	Hatchery ^{c/}	
				Commercial	Sport ^{b/}			
1982-1985	10,275	4,925	1,875	1,675	100	0	3,450	
1986-1990	60,894	24,780	16,288	26,547	2,277	4,253	9,194	
1991-1995	32,352	19,360	6,014	4,151	1,622	7,327	10,631	
1996	59,698	38,100	12,443	5,306	3,387	15,618	15,868	
1997	58,932	36,600	11,596	3,320	6,517	15,916	15,808	
1998	37,328	29,900	5,570	1,063	5,697	11,380	8,401	
1999	50,788	40,400	10,581	1,543	5,927	17,213	7,334	
2000	37,191	25,600	7,186	3,739	3,507	10,135	7,884	
2001	76,504	48,100	16,821	7,023	9,580	14,600	13,701	
2002	108,198	57,600	24,358	9,437	12,737	27,005	21,946	
2003	150,042	97,179	27,830	20,432	12,804	38,204	24,175	
2004	122,496	79,866	23,392	9,178	11,167	27,779	26,210	
2005	100,333	60,464	23,158	6,590	10,727	14,271	30,991	
2006	80,470	31,402	22,705	4,577	3,567	12,501	19,745	
2007 ^{d/}	30,200	13,340	10,800	6,410	1,410	3,220	7,570	
GOAL							Hatchery Production	

a/ Based on Columbia River fall Chinook database, WDFW, unpublished. Does not include URB Chinook destined for areas above McNary Dam or the Deschutes River.

b/ Includes tributary and mainstem catches.

c/ Little White Salmon and Bonneville Hatcheries.

d/ Preliminary based on inseason run updates.

TABLE B-20. Estimates of minimum inriver run size and catch in numbers of adult spring, summer, and fall Chinook from the Columbia River. (Page 1 of 3)

Year	Minimum Inriver Run Size	Below Bonneville Dam					Bonneville Dam Counts	Above Bonneville Dam					Total Treaty Indian & Non-Indian		
		Non-Indian Sport		Non-Indian Commercial				Non-Indian Sport		Treaty Indian					
		Tributary ^{a/}	Buoy 10	Mainstem	Select Area ^{b/}	Mainstem		Mainstem	Tributary ^{c/}	Ticketed Commercial ^{d/}	Non-Ticketed Public Sales	Ceremonial & Subsistence ^{e/}		Non-Indian Total Sport	Commercial
Spring Chinook^{f/}															
'79-'80	146,497	11,427	g/	1,150	-	2,900	55,712	-	-	259	0	1,714	12,577	2,900	17,450
'81-'85	200,057	19,568	g/	2,233	-	8,197	67,959	-	777	1,024	0	2,545	22,422	8,197	34,188
'86-'90	283,638	39,688	g/	5,686	-	14,138	103,800	-	3,156	186	0	6,771	48,530	14,138	69,625
'91-'95	185,100	33,232	g/	3,010	376	4,042	61,819	-	652	15	0	3,730	36,894	4,343	44,982
1996	111,178	13,206	g/	17	789	154	55,236	-	264	0	0	2,911	13,487	943	17,341
1997	178,148	8,270	g/	13	1,821	309	123,758	-	7,326	14	0	8,309	15,609	2,130	26,062
1998	97,766	10,350	g/	63	2,313	100	43,471	-	1,717	1	0	2,224	12,130	2,413	16,768
1999	107,546	14,940	g/	21	1,980	351	42,533	-	220	1	0	1,983	15,181	2,331	19,496
2000	260,045	17,582	g/	351	6,631	1,237	185,774	0	11,502	1,354	0	9,973	29,435	7,868	48,630
2001	540,468	18,925	g/	27,014	9,719	5,279	412,653	93	56,685	22,019	21,696	10,985	102,717	14,998	172,415
2002	482,613	31,800	g/	22,013	12,251	17,383	304,940	1,237	25,859	17,930	6,324	9,208	80,908	29,634	144,004
2003	409,143	33,877	g/	17,815	8,783	4,659	229,499	1,302	21,179	6,363	2,842	9,090	74,173	13,442	105,910
2004	422,699	41,679	g/	24,541	11,643	14,489	198,325	1,419	22,508	5,256	3,114	9,114	90,147	26,132	133,763
2005	193,648	15,493	g/	11,635	2,553	5,646	97,397	448	6,480	1	0	6,163	34,056	8,199	48,419
2006	224,333	13,553	g/	7,061	7,581	4,689	126,158	648	2,396	0	0	8,401	23,658	12,270	44,329
2007 ^{h/}	151,030	11,422	g/	6,518	6,968	3,330	80,829	611	2,152	3	0	6,141	20,703	10,298	37,145
Summer Chinook^{f/h/}															
'79-'80	22,566	-	-	-	-	81	22,485	-	-	38	--	1,047	0	81	1,165
'81-'85	17,092	-	-	-	-	55	17,037	-	-	304	--	654	0	55	1,013
'86-'90	21,668	-	-	7	-	71	21,590	-	-	1,180	--	194	7	71	980
'91-'95	13,348	-	-	15	-	30	13,303	-	-	-	--	227	15	30	271
1996	12,333	-	-	27	-	15	12,291	-	-	-	-	374	27	15	416
1997	18,277	-	-	19	-	6	18,252	-	-	-	-	270	19	6	295
1998	16,332	-	-	27	-	1	16,304	-	-	-	-	335	27	1	363
1999	22,347	-	-	41	-	1	22,305	-	-	-	-	411	41	1	453
2000	23,169	-	-	25	-	0	23,144	-	-	-	-	209	25	0	234
2001	54,935	-	-	64	-	1	54,870	42	0	150	--	542	64	1	799
2002	92,820	-	-	1,503	-	8	91,309	65	0	42	--	2,019	1,568	8	3,637
2003	83,120	-	-	2,007	36	-	81,077	269	0	3,587	--	710	2,276	36	6,609
2004	65,446	-	-	1,240	3	233	63,970	38	157	8,004	--	390	1,435	236	10,065
2005	60,060	-	-	1,622	0	2,553	55,684	74	338	6,415	--	1,227	2,034	2,574	12,250
2006	76,196	-	-	3,360	9	5,008	67,819	40	216	15,771	--	548	3,616	5,017	24,952
2007 ^{h/}	37,190	0	0	2,369	0	1,122	33,699	60	0	4,564	0	811	2,429	1,122	8,926

TABLE B-20. Estimates of minimum inriver run size and catch in numbers of adult spring, summer, and fall Chinook from the Columbia River. (Page 2 of 3)

Year	Minimum Inriver Run Size	Below Bonneville Dam					Bonneville Dam Counts	Above Bonneville Dam					Total Treaty Indian & Non-Indian		
		Non-Indian Sport			Non-Indian Commercial			Non-Indian Sport		Treaty Indian					
		Tributary ^{a/}	Buoy 10	Mainstem	Select Area ^{b/}	Mainstem		Mainstem	Tributary ^{c/}	Ticketed Commercial ^{d/}	Non-Ticketed Public Sales	Ceremonial & Subsistence ^{e/}		Non-Indian Total Sport	Commercial
Fall Chinook^{f/}															
'79-'80	337,608	3,651	-	1,155	20,793	73,253	135,878	500	NA	32,568	--	--	5,306	113,253	151,127
'81-'85	307,206	4,158	7,176	1,528	8,560	45,490	150,768	2,795	NA	48,888	--	5,025	10,234	54,050	118,196
'86-'90	603,713	6,383	20,641	4,119	16,059	181,817	258,807	5,825	442	118,864	4,765	5,692	37,056	197,876	360,441
'91-'95	240,267	3,541	6,224	2,633	1,230	14,693	145,489	4,150	584	33,408	4,732	658	15,887	15,923	70,476
1996	332,051	1,474	1,322	9,178	4,879	11,937	205,358	5,125	1,479	41,718	21,533	504	18,578	16,816	99,149
1997	322,460	2,615	13,048	8,447	4,002	5,130	214,779	4,300	2,100	40,878	23,757	341	30,510	9,132	104,618
1998	255,379	120	5,441	10,232	1,700	2,350	189,085	4,297	2,324	28,096	16,923	--	22,414	4,050	71,483
1999	313,648	2,462	10,170	8,551	2,009	5,774	242,143	7,375	1,720	43,780	32,883	1,310	30,278	7,783	116,034
2000	257,917	320	4,549	7,422	2,007	11,539	192,793	4,324	1,988	37,514	13,635	269	18,603	13,546	83,567
2001	553,598	2,971	12,287	8,683	4,200	22,938	400,205	7,922	2,800	79,959	31,397	365	34,663	27,138	173,522
2002	739,022	7,789	18,273	21,235	7,899	34,428	473,692	11,171	5,940	96,277	33,918	457	64,408	42,327	237,387
2003	902,060	11,999	14,873	25,931	9,360	54,620	610,075	9,267	4,490	94,822	31,107	683	66,560	63,980	257,152
2004	806,121	8,379	15,201	16,968	12,400	40,373	583,499	10,297	4,215	111,833	15,379	416	55,060	52,773	235,461
2005	590,559	7,810	9,983	20,111	8,677	26,231	415,777	9,110	4,307	92,463	22,058	570	51,321	34,908	201,320
2006	422,433	7,052	1,620	13,447	4,822	23,144	299,726	5,136	3,969	58,842	18,849	391	31,224	27,966	137,272
2007 ^{h/}	221,800	3,520	3,820	6,090	4,800	11,950	157,946	3,420	160	34,001	11,085	270	17,010	16,750	79,116
Total Chinook															
'79-'80	506,672	13,253	-	1,728	20,793	39,608	214,075	250	0	16,581	0	2,760	15,230	59,608	94,178
'81-'85	524,355	23,726	7,176	3,761	8,560	53,742	235,764	1,677	621	50,216	0	8,224	32,655	62,302	153,397
'86-'90	909,020	46,071	20,641	9,812	16,059	196,025	384,197	5,825	3,245	119,758	953	12,656	85,593	212,085	431,046
'91-'95	438,715	36,773	6,224	5,658	1,531	18,765	220,612	4,150	1,236	33,424	4,732	4,482	52,796	20,295	115,729
1996	455,563	14,680	1,322	9,222	5,668	12,106	272,885	5,125	1,743	41,718	21,533	3,789	32,092	17,774	116,906
1997	518,884	10,885	13,048	8,479	5,823	5,445	356,789	4,300	9,426	40,892	23,757	8,920	46,138	11,268	130,975
1998	369,477	10,470	5,441	10,322	4,013	2,451	248,860	4,297	4,041	28,097	16,923	2,559	34,571	6,464	88,614
1999	443,541	17,402	10,170	8,613	3,989	6,126	306,981	7,375	1,940	43,781	32,883	3,704	45,499	10,115	135,982
2000	541,131	17,902	4,549	7,798	8,638	12,776	401,711	4,324	13,490	38,868	13,635	10,451	48,063	21,414	132,431
2001	1,149,001	21,896	12,287	35,761	13,919	28,218	867,728	8,057	59,485	102,128	53,093	11,892	137,444	42,137	346,736
2002	1,314,455	39,589	18,273	44,751	20,150	51,819	869,941	12,473	31,799	114,249	40,242	11,684	146,885	71,969	385,029
2003	1,394,323	45,876	14,873	45,753	18,179	59,279	920,651	10,838	25,669	104,772	33,949	10,483	143,009	77,458	369,671
2004	1,294,266	50,058	15,201	42,749	24,046	55,095	845,794	11,754	26,880	125,093	18,493	9,920	146,642	79,141	379,289
2005	844,267	23,303	9,983	33,368	11,230	34,430	568,858	9,632	11,125	98,879	22,058	7,960	87,411	45,681	261,989
2006	722,962	20,605	1,620	23,868	12,412	32,841	493,703	5,824	6,581	74,613	18,849	9,340	58,498	45,253	206,553
2007 ^{h/}	410,020	14,942	3,820	14,977	11,768	16,402	272,474	4,091	2,312	38,568	11,085	7,222	40,142	28,170	125,187

TABLE B-20. Estimates of minimum inriver run size and catch in numbers of adult spring, summer, and fall Chinook from the Columbia River. (Page 3 of 3)

- a/ For spring Chinook: includes lower and upper Willamette, Clackamas, Cowlitz, Kalama, Lewis, and Sandy Rivers. Sandy River harvest not available before 1990. Catch estimates may include small numbers of jacks. For summer Chinook: all tributaries are closed. For fall Chinook: all tributaries downstream from Bonneville Dam.
- b/ Youngs Bay Select Area began in 1992. Tongue Point and Blind Slough began in 1998. Select Area test fisheries began in 1991. Other Select Areas include Knappa in Oregon and Deep River in Washington.
- c/ Includes tributaries between Bonneville and McNary Dams, the Snake and Yakima rivers, Icicle and Ringold creeks.
- d/ Primarily mainstem fisheries between Bonneville and McNary dams, but also includes fish caught in miscellaneous commercial Indian fisheries such as Klickitat dip net and mainstem fisheries upstream from McNary Dam.
- e/ Primarily mainstem fisheries between Bonneville and McNary dams. Significant subsistence fisheries also occur in tributaries throughout the Columbia and Snake River basin, especially for spring Chinook, which are not included in these estimates.
- f/ Upriver spring Chinook accounting ends on June 15 and summer Chinook accounting begins on June 16.
- g/ Spring Chinook Buoy 10 area catch is included in mainstem sport.
- h/ Preliminary. Fall Chinook estimates are from inseason run updates.
- i/ Summer Chinook retention was prohibited for all mainstem non-Indian and treaty Indian fisheries until 2003. Small non-Indian incidental mortalities prior to 2003 are associated with recreational steelhead fisheries and commercial shad and sockeye fisheries. A few stray summer Chinook are caught in Select Area (terminal) fisheries that are open for late returning spring Chinook and early returning fall Chinook. Prior to 2003, Treaty Indians could retain summer Chinook for subsistence purposes.
- j/ Fall Chinook minimum run size includes LRH, LRW, SCH, URB, MCB, and SAB.

TABLE B-21. Estimates of minimum inriver run size, catch, and escapement in thousands of adult coho entering the Columbia River.^{a/} (Page 1 of 1)

Year or Average	Minimum Inriver Run Size	Below Bonneville Dam					Above Bonneville Dam			
		Lower River Catch ^{b/}			Lower River Escapement		Bonneville Dam Counts ^{e/}	Mainstem Commercial Treaty Catch	Zone 6 Escapement ^{f/}	Hatchery Escapement
		Commercial	Recreational		Hatchery ^{c/}	Tributary Dam Counts ^{d/}				
			Buoy 10	Mainstem						
1971-1975	373.4	199.4	-	11.8	117.1	9.5	35.6	9.1	26.6	11.6
1976-1980	263.3	123.6	-	10.1	102.2	3.6	23.8	2.6	21.2	7.0
1981-1985	305.5	132.0	30.6	11.4	101.3	4.6	31.9	2.6	29.2	12.5
1986-1990	689.2	392.2	82.3	14.5	148.8	5.8	46.3	5.5	40.7	11.5
1991-1995	301.4	115.9	55.9	10.7	91.3	3.7	23.6	2.0	21.6	6.1
1996	113.0	26.2	4.5	3.8	62.2	0.6	15.7	0.1	15.6	1.4
1997	148.1	19.4	20.4	11.6	69.7	2.8	24.2	0.6	23.6	4.4
1998	168.7	23.0	3.2	6.7	87.9	1.3	46.6	0.2	46.4	11.3
1999	274.1	79.0	9.0	19.9	124.5	1.0	40.7	1.7	39.0	10.0
2000	547.6	168.4	21.5	37.7	228.6	5.6	85.8	6.3	79.3	26.6
2001	1,108.3	253.1	132.0	78.0	377.3	8.2	259.8	5.5	254.0	80.6
2002	499.9	163.0	6.2	27.4	211.1	3.6	88.6	1.6	86.5	2.9
2003	677.3	257.3	54.2	23.3	205.4	11.2	125.7	3.1	122.6	3.9
2004	442.5	119.6	15.1	13.5	173.5	5.6	115.0	6.0	109.0	6.2
2005	342.0	94.8	6.9	10.5	143.3	3.3	83.3	4.7	78.6	2.3
2006	383.0	63.4	3.7	16.3	188.0	9.5	102.1	5.4	96.7	0.7
2007 ^{g/}	318.6	39.1	8.4	13.4	160.5	4.7	92.5	7.9	84.6	1.0
GOAL					Hatchery Production				Hatchery Production	

a/ These numbers match OPI databases. Adjustments were made to the escapement figures and catches.

b/ Includes some upriver origin coho. Mainstem recreational catches listed in this table include tributary catches and catches in the Chinook/Hammond area of 3,195 in 1989, 28 in 1990, and 1,151 in 1991.

c/ Includes hatcheries operated by all agencies.

d/ Willamette Falls, Clackamas River (North Fork Dam) and Sandy River (Marmot Dam).

e/ Includes additional small adults counted as jacks for 1983-1984 and 1986-1989.

f/ Bonneville Dam count minus Zone 6 mainstem commercial treaty Indian harvest.

g/ Preliminary.

TABLE B-22. Estimated catch and effort in the Buoy 10 fishery.^{a/} (Page 1 of 1)

Year	Angler Trips	Catch		Catch Per Trip
		Chinook	Coho	
1982-1985	30,996	4,040	30,547	0.97
1986-1990 ^{b/c/}	130,633	22,107	82,910	0.78
1991-1995 ^{d/}	79,475	5,689	55,895	0.50
1996	18,034	1,409	4,537	0.33
1997	55,725	13,153	20,357	0.60
1998	29,998	5,784	3,175	0.30
1999	49,581	9,850	8,861	0.38
2000 ^{e/}	72,518	6,085	21,478	0.38
2001 ^{e/}	125,884	12,709	132,038	1.15
2002 ^{e/}	84,457	19,441	6,233	0.30
2003 ^{e/}	88,827	16,316	54,440	0.80
2004 ^{e/}	68,818	16,016	15,169	0.45
2005 ^{e/}	55,182	9,286	6,878	0.29
2006 ^{e/}	40,688	1,706	3,687	0.13
2007 ^{e/f/}	36,064	3,776	8,356	0.34

a/ Prior to 1982, Buoy 10 area catches were not estimated separately and are included in the Columbia River marine area (Cape Falcon to Leadbetter Pt.) recreational catches. Estimates include bank anglers fishing from Clatsop Spit in Oregon and from the North Jetty in Washington. Effort and catch for the North Jetty fishery applied to the ocean quota for the Columbia River area until the ocean fishery closed.

b/ 1989 includes catch and effort data for the Chinook/Hammond fishery occurring during weeks 32 and 33. A total of 7,922 angler trips produced catches of 492 Chinook, 3,195 coho, and a catch rate of 0.47 fish per trip. Catches in this fishery were counted against the Buoy 10 quota.

c/ 1990 includes catch and effort data for the Chinook/Hammond fishery occurring during weeks 31 and 32. A total of 3,225 angler trips produced catches of 54 Chinook, 28 coho, and a catch rate of 0.03 fish per trip.

d/ 1991 includes catch and effort data for the Chinook/Hammond fishery occurring during weeks 31 and 32. A total of 2,759 angler trips produced catches of 39 Chinook, 1,151 coho, and a catch rate of 0.43 fish per trip.

e/ Includes catch and effort from the Astoria-Megler Bridge upstream to the new boundary from Tongue Point, Oregon to Rocky Point, Washington.

f/ Preliminary.

TABLE B-23. Willapa Bay fall Chinook terminal run size, catch, and spawning escapement in numbers of fish. (Page 1 of 1)

Year or Average	Non-local Stocks Catch ^{a/}	Terminal Catch		Spawning Escapement		Terminal Run Size ^{d/}
		Gillnet	Sport ^{b/}	Natural ^{c/}	Hatchery	
1976-1980	8,660	14,496	419	1,995	4,529	21,439
1981-1985	1,011	7,331	589	1,588	5,398	14,906
1986-1990	2,521	18,173	1,578	5,596	22,458	47,805
1991-1995	697	28,082	2,823	2,818	17,086	50,809
1996	0	37,065	3,024	2,153	12,079	54,321
1997	0	12,311	2,404	3,852	13,729	32,296
1998	0	6,765	2,178	3,114	8,658	20,715
1999	0	265	1,906	1,360	6,966	10,497
2000	0	5,902	1,399	2,303	10,455	20,059
2001	0	5,444	2,121	2,161	10,099	19,825
2002	36	9,452	2,532	1,729	13,680	27,429
2003	220	7,488	3,241	2,728	14,631	28,230
2004 ^{e/}	0	4,349	3,851	2,532	21,444	32,018
2005 ^{e/}	0	6,523	6,630	1,804	18,514	33,382
2006 ^{e/}	0	12,334	6,442	3,598	24,569	42,078
2007 ^{e/}	0	4,112	NA	NA	NA	NA
GOAL				4,400 ^{f/}	9,800 ^{f/}	

a/ Non-local gillnet is catch in Area 2G prior to Aug. 16.

b/ Adults. Sport catch since 1991 includes marine areas within Willapa Bay (e.g., Washaway Beach).

c/ Escapement estimates after 1984 are based on revised spawning habitat estimates. Natural = adult returns assumed to be from natural origin parents.

d/ Does not include catch of non-local stocks.

e/ Preliminary.

f/ WDFW goal; not an FMP goal.

TABLE B-24. Willapa Bay coho terminal run size, catch, and spawning escapement in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch		Spawning Escapement		Terminal Run Size ^{d/}
	Gillnet	Sport ^{a/}	Natural ^{b/}	Hatchery ^{c/}	
1976-1980	15,011	2,842	5,800	14,328	37,981
1981-1985	39,007	2,181	3,567 ^{e/}	26,640	69,968
1986-1990	69,199	2,591	e/	35,811	107,601
1991-1995	34,287	2,802	4,582 ^{e/}	27,205	65,211
1996	38,316	4,052	15,711	48,854	106,933
1997	1,550	806	4,934	6,691	13,981
1998	13,140	852	13,807	6,902	34,701
1999	5,467	2,836	12,355	22,823	43,481
2000	10,193	1,780	23,031	29,387	64,391
2001	31,837	5,707	48,414	54,359	140,317
2002	59,435	5,683	58,703	48,871	172,692
2003	66,460	5,881	49,386	66,115	187,842
2004 ^{f/}	16,533	2,325	44,463	16,443	55,903
2005 ^{f/}	50,031	3,867	30,080	40,945	124,923
2006 ^{f/}	19,914	811	14,413	7,565	42,640
2007 ^{f/}	8,218	NA	NA	NA	NA
GOAL			13,090 ^{g/}	6,100 ^{g/}	

a/ Adults. Sport catch since 1991 includes marine areas within Willapa Bay (e.g., Washaway Beach).

b/ Natural spawning escapement estimates in 1996, 1997, and 1998 do not include adult fish released upstream of hatchery racks.

c/ Hatchery rack number includes fish released upstream.

d/ Does not include natural spawning escapement between 1984 and 1994.

e/ Estimates of natural spawning escapement were not made between 1984 and 1994.

f/ Preliminary

g/ WDFW goal; not an FMP goal.

TABLE B-25. Grays Harbor Chinook terminal catch, spawning escapement, and run size in numbers of fish. (Page 1 of 2)

Year or Average	Terminal Catch					Spawning Escapement		Terminal Run Size ^{d/}
	Early Non-local Catch	Non-Indian Gillnet	Treaty Indian Gillnet	Chehalis Tribal Gillnet	Sport ^{a/}	Natural ^{b/}	Hatchery ^{c/}	
SPRING Chinook								
1976-1980	-	-	-	587	e/	600	-	1,187
1981-1985	-	-	-	57	5	924	-	963
1986-1990	-	-	e/	143	6	1,875	-	2,024
1991-1995	-	-	0	94	15	1,566	-	1,675
1996	-	-	104	127	52	4,462 ^{f/}	-	4,745
1997	-	-	52	172	160	4,460 ^{f/}	-	4,844
1998	-	-	6	164	121	2,288	-	2,579
1999	-	-	3	187	76	1,285	-	1,551
2000	-	-	17	174	91	3,135	-	3,417
2001	-	-	4	210	252	2,860	-	3,326
2002	-	-	76	419	124	2,598	-	3,217
2003	-	-	68	0	131	1,904	-	2,103
2004 ^{g/}	-	-	54	177	65	5,034	-	5,330
2005 ^{g/}	-	-	26	439	88	2,129	-	2,682
2006 ^{g/}	-	-	5	249	128	2,481	-	2,863
2007 ^{g/}	-	-	10	NA	NA	NA	-	NA
GOAL						1,400		

TABLE B-25. Grays Harbor Chinook terminal catch, spawning escapement, and run size in numbers of fish. (Page 2 of 2)

Year or Average	Terminal Catch					Spawning Escapement		Terminal Run Size ^{d/}
	Early Non-local Catch	Non-Indian Gillnet	Treaty Indian Gillnet	Chehalis Tribal Gillnet	Sport ^{a/}	Natural ^{b/}	Hatchery ^{c/}	
FALL Chinook								
1976-1980	4,433	3,642	3,108	1,006	1,128	7	413	13,736
1981-1985	602	964	3,524	465	268	10	742	6,575
1986-1990	694	4,122	10,414	597	1,340	20,692	1,319	39,178 ^{h/}
1991-1995	206	5,000	7,750	901	3,794	14,276	3,006	34,934 ^{h/}
1996	148	1,441	4,068	49	7,456	20,227	4,307	37,696 ^{h/}
1997	24	2,796	6,630	311	2,687	18,168	2,416	33,032 ^{h/}
1998	5	267	4,135	0	2,912	12,539	1,921	21,779 ^{h/}
1999	0	87	1,926	1	114	10,363	1,990	14,481 ^{h/}
2000	671	647	3,289	0	1,714	9,250	1,450	17,021
2001	0	2,523	3,885	0	3,210	9,491	1,121	20,230
2002	40	26	963	0	2,955	11,343	2,006	17,333
2003	0	99	851	0	1,031	19,417	2,858	24,256
2004 ^{g/}	0	108	3,497	0	6,158	31,770	2,695	44,228
2005 ^{g/}	0	218	2,260	21	465	19,499	3,285	25,748
2006 ^{g/}	0	0	3,738	NA	1,635	17,113	NA	NA
2007 ^{g/}	0	514	2,470	NA	NA	NA	NA	NA
GOAL						14,600		

a/ Age-3 and older.

b/ Age-3 and older, including hatchery fish spawning naturally.

c/ Includes naturally spawning fish taken for broodstock.

d/ Minimum estimate due to incomplete estimates of river recreational catch. Does not include non-local catch.

e/ Fewer than 50 fish.

f/ WDFW is not able to differentiate spawning time and believes this includes fall Chinook.

g/ Preliminary.

h/ Recreational catch estimates by WDFW reflect application of catch record card bias correction factor of 0.833. Quinault Indian Nation does not believe this factor is appropriate for this fishery. Unadjusted catch estimates are 1,000 for 1987; 2,400 for 1988; 2,500 for 1989; 2,400 for 1990; 4,500 for 1991; 2,600 for 1992; 4,200 for 1993; 4,300 for 1994; 6,500 for 1995; 6,800 for 1996; 3,400 for 1997; 3,500 for 1998; and 0.1 for 1999; terminal run sizes would be adjusted accordingly.

TABLE B-26. Grays Harbor coho terminal catch, spawning escapement, and run size estimates in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch				Spawning Escapement ^{b/}		Terminal Run Size		
	Non-Indian Gillnet	Treaty Indian Gillnet	Chehalis Tribal Gillnet	Sport ^{a/}	Natural	Hatchery	Natural	Hatchery	Total ^{c/}
	1976-1980	5,231	9,675	3,500	2,021	29,510	9,310	44,430	17,933
1981-1985	5,299	15,614	2,863	5,012	36,847	16,753	42,974	32,432	82,388
1986-1990	7,715	30,109	1,817	5,355	44,836	30,767	53,030	65,193	120,600
1991-1995	12,502	29,745	2,716	10,503	36,516	31,654	51,888	69,650	123,635
1996	10,096	51,812	2,915	20,846	63,572	49,378	83,263	100,764	198,619
1997	115	5,548	125	1,547	22,469	12,710	19,299	22,352	42,514
1998	795	13,586	361	2,123	35,551	16,903	36,076	32,435	69,319
1999	1,674	12,212	797	4,507	33,346	31,488	34,325	48,561	84,024
2000	4,995	10,947	331	5,122	38,054	28,253	39,997	47,070	87,702
2001	3,152	15,671	533	20,868	79,112	80,865	74,489	137,858	200,201
2002	6,853	14,518	666	13,083	108,695	45,365	104,738	81,615	189,180
2003	6,623	12,041	1,000	12,026	83,874	66,922	89,503	94,067	182,486
2004 ^{d/}	5,231	17,431	977	9,847	60,690	49,147	65,860	79,199	143,323
2005 ^{d/}	3,073	22,649	4,400	11,043	44,090	50,892	49,767	84,410	136,147
2006 ^{d/}	649	8,708	NA	2,151	14,401	18,617	NA	NA	NA
2007 ^{d/}	1,687	8,839	NA	NA	23,662	NA	NA	NA	NA
GOAL					35,400				

a/ Beginning in 1987, estimates provided by WDFW for recreational catch reflect punch card bias correction factor.

b/ "Natural" includes hatchery fish spawning in wild. "Hatchery" includes wild fish taken for broodstock.

c/ The combined natural and hatchery run size total may not add to the sum of the catch and escapements due to hatchery total run size including on-station and off-station escapements.

d/ Preliminary.

TABLE B-27. Treaty Indian gillnet catch of Chinook, chum, and sockeye salmon in the Quinault River in numbers of fish. (Page 1 of 1)

Year or Average	Spring/Summer Chinook ^{a/}	Fall Chinook ^{a/}	Chum	Sockeye
1976-1980	149	4,320	7,960	17,560
1981-1985	114	5,100	4,720	12,600
1986-1990	338	8,822	4,686	11,218
1991-1995	98	6,293	2,505	9,523
1996	41	5,221	594	1,244
1997	19	2,625	1,033	2,532
1998	75	6,124	4,699	3,440
1999	10	4,840	599	73
2000	0	3,421	755	0
2001	5	4,047	2,009	0
2002	36	4,542	1,151	16,939
2003	92	7,343	3,742	37,130
2004	142	10,662	2,916	6,990
2005 ^{b/}	24	7,648	1,283	116
2006 ^{b/}	16	7,044	862	1
2007 ^{b/}	<20	2,125	1,173	1

a/ Preliminary. Stock separation under review.

b/ Preliminary.

TABLE B-28. Estimated inriver run size, catch and escapement for Quinault River coho in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch ^{a/}			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport	Natural	Hatchery	Natural	Hatchery	Total
	1977-1980	9,750	-	-	3,425	3,107	8,465	7,750
1981-1985	10,700	-	-	3,237	6,239	7,809	12,657	20,466
1986-1990	13,777	-	-	3,185	4,239	8,024	13,200	21,224
1991-1995	7,963	-	-	4,319	8,046	6,205	13,472	19,678
1996	10,087	-	-	13,327	9,521	18,849	13,865	32,714
1997	365	-	-	3,150	1,054	3,339	1,118	4,457
1998	5,946	-	-	3,770	3,158	7,156	5,581	12,737
1999	15,491	-	-	12,666	14,617	19,138	23,101	42,239
2000	16,194	-	-	7,421	9,481	14,559	18,099	32,658
2001	25,348	-	-	21,565	30,689	30,016	47,115	77,131
2002	19,197	-	-	12,213	16,841	16,847	30,196	47,043
2003	22,546	-	-	4,710	16,841	9,546	34,132	43,678
2004	17,055	-	-	1,404	10,321	3,377	24,821	28,198
2005 ^{b/}	23,852	-	-	6,418	10,034	15,951	25,574	41,525
2006 ^{b/}	9,785	336	325	1,107	3,198	3,429	11,023	14,751
2007 ^{b/}	11,770	-	-	NA	NA	NA	NA	NA
GOAL	Hatchery Production							

a/ Ceremonial, subsistence, and recreational catch negligible. Includes dip-in fish destined for other river systems.

b/ Preliminary.

TABLE B-29. Estimated inriver run size, catch, and escapement of Queets River spring/summer Chinook in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{a/}	Natural ^{b/}	Hatchery	Natural	Hatchery	Total
1976-1980	267	18	53	851	24	1,176	37	1,078
1981-1985	243	20	27	890	31	956	44	1,209
1986-1990	646	46	67	1,527	0	2,287	0	2,287
1991-1995	64	5	10	610	0	689	0	688
1996	43	3	69	776	0	891	0	891
1997	72	10	71	540	0	693	0	693
1998	18	27	0	492	0	537	0	537
1999	12	41	0	373	0	426	0	426
2000	0	2	0	248	0	250	0	250
2001	0	17	0	548	0	565	0	565
2002	0	17	0	738	0	755	0	755
2003	0	6	0	189	0	195	0	195
2004 ^{c/}	0	15	0	604	0	619	0	619
2005 ^{c/}	0	8	0	298	0	306	0	306
2006 ^{c/}	0	6	0	330	0	336	0	336
2007 ^{c/}	0	6	0	352	0	358	0	358
GOAL				700 ^{d/}				

a/ River catch of adults.

b/ Natural escapement includes hatchery strays.

c/ Preliminary.

d/ Minimum. Terminal run managed at 30% exploitation rate of inriver run size.

TABLE B-30. Estimated inriver run size, catch, and escapement of Queets River fall Chinook in numbers of fish. (Page 1 of 1).

Average	Terminal Catch			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{a/}	Natural ^{b/}	Hatchery ^{c/}	Natural	Hatchery	Total
1976-1980	1,540	100	36	2,820	-	4,320	-	4,320
1981-1985	2,104	20	135	3,720	360	5,691	591	6,282
1986-1990	2,430	20	214	8,298	619	10,677	861	11,538
1991-1995	1,860	20	109	3,849	407	5,511	708	6,219
1996	1,307	20	238	3,441	927	4,693	1,234	5,927
1997	1,708	20	210	2,477	545	4,122	823	4,945
1998	804	20	347	3,951	58	5,009	164	5,173
1999	947	20	93	1,933	135	2,885	220	3,105
2000	262	20	50	3,572	333	3,752	395	4,147
2001	1,366	64	306	2,321	778	3,604	1,204	4,808
2002	2,887	69	20	2,097	492	4,377	1,184	5,562
2003	1,322	93	278	4,120	859	5,203	1,415	6,618
2004	1,228	93	376	3,576	1,529	4,782	2,020	6,802
2005	1,648	90	441	3,076	1,481	4,521	2,213	6,736
2006	1,079	57	71	2,338	713	3,253	1,005	4,259
2007 ^{d/}	634	NA	NA	1,924	454	NA	NA	NA
GOAL				2,500 ^{e/}				

a/ River sport catch of 3-year-olds and older. The 2000 sport fishery was closed to retention of unmarked Chinook. The 2002 sport fishery was closed to Chinook retention on October 18 due to unusually low water conditions.

b/ Includes fish taken for hatchery broodstock.

c/ This is an integrated wild/hatchery program. All broodstock are unmarked wild fish collected from spawning grounds.

d/ Preliminary. 2007 Escapements based on the CPUE model & actual versus expected catches. Spawning escapement estimates not complete.

e/ Minimum. Terminal run managed at 40% exploitation rate of inriver run size.

TABLE B-31. Estimated terminal run size, catch, and escapement for Queets River coho in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch ^{a/}			Escapement			Terminal Run Size			
	Gillnet	Ceremonial & Subsistence	River Sport ^{b/}	Natural ^{c/}	Supplemental ^{d/}	Hatchery	Natural ^{c/}	Supplemental	Hatchery	Total
1976-1980	2,440	60	140	3,460	-	1,000	5,100	-	1,640	6,740
1981-1985	2,385	20	104	5,457	-	2,654	6,414	-	3,794	10,208
1986-1990	8,455	18	241	4,824	2,128	3,366	6,357	2,988	9,357	17,507
1991-1995	4,423	285	273	5,019	954	3,351	5,888	1,113	6,912	13,690
1996	16,035	920	279	8,993	3,574	5,191	10,715	5,319	17,646	33,680
1997	3,087	222	106	1,851	d/	2,137	1,970	d/	5,086	7,056
1998	7,411	452	135	4,102	1,413	3,504	4,576	1,562	10,364	16,502
1999	3,974	381	119	4,791	521	3,551	5,029	557	7,061	12,647
2000	5,066	479	223	7,939	682	3,849	8,498	702	5,619	14,819
2001	13,722	1,280	1,554	23,793	1,084	6,594	28,303	2,081	13,589	43,974
2002	23,712	1,003	399	13,773	1,048	2,240	15,904	1,292	21,290	38,486
2003	12,692	918	743	9,846	704	7,394	13,223	1,344	15,707	30,273
2004 ^{e/}	8,189	655	1,287	8,709	0	3,260	11,470	243	10,494	22,207
2005 ^{e/}	20,810	980	873	6,539	432	7,810	9,801	534	25,935	36,270
2006 ^{e/f/}	6,190	291	46	5,400	0	1,447	NA	0	NA	NA
2007 ^{f/}	2,261	NA	NA	5,272	0	1,714	NA	0	NA	NA
GOAL				5,800-14,500						

a/ Includes dip-in fish from other river systems.

b/ Recreational catch of adults (coho over 20 inches).

c/ Natural escapement and run sizes estimates include fish taken for hatchery brood stock.

d/ 1991 and 1997 supplemental was included in natural escapement and run size.

e/ Poor conditions during the coho spawner survey season precluded conduct of an independent spawner escapement estimate.

f/ Preliminary. In-season effort model used to scale run size to observed catch and effort, natural escapement, and actual hatchery rack escapement.

TABLE B-32. Estimated inriver run size, catch, and escapement for Hoh River spring/summer Chinook in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch ^{a/}			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{b/}	Natural	Hatchery	Natural	Hatchery	Total
1976-1980	640	52	84	1,040	0	1,835	0	1,835
1981-1985	448	30	124	1,431	50	1,944	128	2,073
1986-1990	1,072	33	315	2,829	34	4,043	257	4,300
1991-1995	432	22	273	1,268	0	1,852	156	2,008
1996	471	30	267	1,371	16	2,083	114	2,197
1997	416	57	331	1,826	0	2,582	53	2,635
1998	294	20	288	1,287	0	1,880	28	1,908
1999 ^{c/}	155	20	52	928	99	1,081	171	1,252
2000 ^{d/}	87	38	21	492	0	529	116	645
2001 ^{d/}	134	39	43	1,159	0	1,231	101	1,332
2002 ^{e/}	587	37	372	2,464	0	3,375	85	3,460
2003 ^{e/}	296	20	206	1,228	0	1,646	104	1,750
2004 ^{e/}	401	20	102	1,786	0	2,239	70	2,309
2005 ^{e/}	323	36	73	1,193	0	1,389	217	1,606
2006 ^{e/f/}	576	37	109	904	0	1,061	571	1,632
2007 ^{e/f/}	692	68	93	817	0	1,019	651	1,670
GOAL				900 ^{g/}				

a/ Beginning in 1981, catch breakouts recalculated to account for Solduc hatchery yearling release dip-in fish.

b/ Recreational catch of adults (at least 24 inches total length).

c/ Sport fishery closed until July 14.

d/ Sport fishery closed through August 31 to retention of wild adult spring/summer Chinook. Sport catch reflects retention of hatchery fish only.

e/ Sport fishery open May 16 through August 31 from mouth to Willoughby Creek.

f/ Preliminary.

g/ Minimum. Terminal run managed at 31% harvest rate of inriver run size.

TABLE B-33. Estimated inriver run size, catch, and escapement for Hoh River fall Chinook in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{a/}	Natural ^{b/}	Hatchery	Natural	Hatchery	Total
1976-1980	760	36	37	2,080	-	2,960	-	2,960
1981-1985	849	36	59	2,745	20	3,684	100	3,764
1986-1990	2,000	32	213	4,500	33	6,819	88	6,907
1991-1995	871	27	233	2,774	0	3,590	65	3,655
1996	836	30	192	3,022	0	4,061	19	4,080
1997	1,114	35	164	1,773	0	3,034	52	3,086
1998	846	30	268	4,257	0	5,388	13	5,401
1999	596	30	413	1,924	0	2,941	22	2,963
2000	404	20	479	1,749	0	2,632	20	2,652
2001	946	40	600	2,560	0	4,116	120	4,236
2002 ^{c/}	1,461	30	134	4,415	82	5,716	406	6,122
2003 ^{d/}	517	30	216	1,649	32	2,319	99	2,418
2004 ^{d/}	815	30	400	3,211	26	4,410	72	4,482
2005 ^{d/}	970	21	229	4,180	14	5,316	77	5,393
2006 ^{d/}	579	30	204	1,532	NA	2,088	19	2,107
2007 ^{d/}	660	30	NA	1,655	NA	2,734	11	2,745
GOAL				1,200 ^{e/}				

a/ Recreational catch of age-3 and older fish.

b/ Includes fish taken for hatchery brood stock.

c/ Low water in October and early November delayed upstream migration, prompting closure of the sport fishery to Chinook retention on October 19 for the remainder of season. Tribal gillnet fishery closed weeks 44 and 45.

d/ Preliminary.

e/ Minimum. Terminal run managed at 40% harvest rate of inriver run size through 1996; for 1997 and 1998, fishing regimes were designed to target a range near 40%.

TABLE B-34. Estimated inriver run size, catch, and escapement for Hoh River coho in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch ^{a/}			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{b/}	Natural ^{c/}	Hatchery	Natural	Hatchery	Total
1976-1980	1,960	74	28	2,700	39	4,683	259	4,942
1981-1985	1,604	48	22	3,371	92	4,655	452	5,107
1986-1990	2,507	30	165	3,145	238	5,221	760	5,981
1991-1995	801	26	168	3,078	122	3,816	379	4,195
1996	972	50	101	4,858	0	5,835	146	5,981
1997 ^{d/}	85	25	4	1,386	0	1,449	51	1,500
1998	650	20	213	4,418	0	5,184	118	5,302
1999	1,706	25	256	4,594	0	6,293	308	6,601
2000	1,932	20	280	6,772	0	8,831	173	9,004
2001	3,909	40	786	10,773	840	14,801	1,547	16,348
2002 ^{e/}	3,114	30	401	9,009	1,922	11,254	3,222	14,476
2003	1,872	20	350	6,273	645	8,118	1,021	9,139
2004	1,255	20	437	4,702	14	6,291	137	6,428
2005	3,830	30	280	4,711	732	8,294	1,259	9,553
2006 ^{f/}	1,313	30	108	1,282	NA	2,159	436	2,595
2007 ^{f/}	1,764	30	NA	3,072	NA	5,186	51	5,237
GOAL				2,000 to 5,000				

a/ Includes dip-in fish from other river systems.

b/ Recreational catch of adults (coho over 20 inches).

c/ Natural escapement and run sizes estimates include fish taken for hatchery brood stock.

d/ Recreational fishermen were limited to Chinook only. Release of adult coho required. Tribal net fishery used large mesh to minimize coho impacts.

e/ Sport and tribal gillnet seasons reduced inseason in response to delayed upriver movement of coho caused by extreme low water conditions in October and early November. Closures were for two weeks.

f/ Preliminary.

TABLE B-35. Estimated inriver run size, catch, and escapement for Quillayute River spring/summer Chinook in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{d/}	Natural ^{b/}	Hatchery	Natural	Hatchery ^{c/}	Total
1976-1980	2,520	20	380	2,093	800	NA	NA	3,698
1981-1985	700	20	48	731	260	NA	NA	1,164
1986-1990	1,631	22	258	1,602	1,003	3,085	2,503	4,341
1991-1995	893	25	293	1,159	832	1,444	1,758	3,202
1996	136	50	257	1,170	226	1,388	426	1,814
1997	106	50	263	890	198	1,177	305	1,482
1998	199	50	128	1,599	247	1,829	369	2,198
1999	368	50	238	713	596	818	1,147	1,965
2000	254	50	307	989	227	1,149	678	1,827
2001	330	50	353	1,225	973	1,399	1,515	2,914
2002	419	50	367	1,002	836	1,100	1,573	2,673
2003 ^{d/}	184	50	343	1,219	1,250	1,308	1,738	3,046
2004 ^{d/}	217	50	341	1,093	763	1,259	1,195	2,454
2005 ^{d/e/f}	332	3	479	876	801	1,033	1,467	2,500
2006 ^{d/e/f}	688	0	340	553	1,032	604	2,002	2,606
2007 ^{d/e/f/g}	795	0	NA	498	1,007	563	1,737	2,300
GOAL				1,200 ^{h/}				

a/ Recreational catch of adults.

b/ Natural escapement includes hatchery strays and broodstock fish.

c/ Hatchery escapement and terminal run size exclude hatchery strays.

d/ Hatchery, no wild retention in sport fishery after 2002.

e/ Beginning in 2005, C&S catch taken during scheduled gillnet fishery is included in gillnet harvest numbers.

f/ Preliminary.

g/ Terminal run size estimates incomplete because inriver sport catch estimates are unavailable.

h/ FMP goal is adults; WDFW goal of 1,200 includes age-3 males (jacks).

TABLE B-36. Estimated inriver run size, catch, and escapement for Quillayute River fall Chinook in numbers of fish. (Page 1 of 1)

Year or Average	Terminal Catch			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{d/}	Natural ^{b/}	Hatchery ^{c/}	Natural	Hatchery ^{c/}	Total
1976-1980	2,640	20	220	4,220	144	6,540	640	7,180
1981-1985	2,075	50	131	6,282	77	8,219	305	8,525
1986-1990	5,475	50	564	12,238	112	18,004	379	18,383
1991-1995	713	50	289	5,670	11	6,705	29	6,733
1996	1,377	100	500	7,316	0	9,293	0	9,293
1997	282	50	310	5,405	0	6,047	0	6,047
1998	762	100	326	6,752	0	7,940	0	7,940
1999	1,129	100	195	3,334	0	4,758	0	4,758
2000	604	100	360	3,730	0	4,794	0	4,794
2001	1,650	100	659	5,136	0	7,545	0	7,545
2002	3,074	100	271	6,067	0	9,512	0	9,512
2003	1,345	100	626	7,398	0	9,469	23	9,492
2004	527	100	681	3,831	0	6,133	12	6,145
2005 ^{d/e/f}	1,414	0	502	6,406	0	8,319	32	8,351
2006 ^{d/e/f}	1,969	0	216	5,642	0	7,656	15	7,671
2007 ^{d/e/f}	905	0	NA	2,934	0	3,839	0	3,839
GOAL				3,000 ^{g/}				

a/ River recreational catch of age-3 and older fish.

b/ Includes fish taken for hatchery brood stock and hatchery strays.

c/ Hatchery escapement and terminal run size exclude hatchery strays.

d/ Preliminary.

e/ Terminal run size estimates incomplete since inriver sport catch estimates are unavailable.

f/ Beginning in 2005, C&S catch taken during scheduled gillnet fishery is included in gillnet harvest numbers.

g/ Minimum. Terminal run managed at 40% harvest rate.

TABLE B-37. Estimated inriver run size, catch, and escapement for Quillayute River coho stocks in numbers of fish. (Page 1 of 2)

Year or Average	Terminal Catch ^{a/}			Escapement		Terminal Run Size		
	Gillnet	Ceremonial & Subsistence	River Sport ^{b/}	Natural ^{c/}	Hatchery ^{d/}	Natural ^{c/}	Hatchery ^{d/}	Total
SUMMER COHO								
1976-1980	5,038	56	266	1,192	4,565	1,962	9,154	11,116
1981-1985	4,062	50	105	946	2,744	2,106	5,802	7,908
1986-1990	3,204	50	94	723	4,001	1,643	6,430	8,072
1991-1995	1,286	50	191	784	6,501	989	7,823	8,812
1996	2,552	50	189	465	3,400	801	5,855	6,656
1997	70	50	14	753	1,509	798	1,598	2,396
1998	1,310	50	93	346	1,688	593	2,894	3,487
1999	945	50	292	624	7,527	723	8,715	9,438
2000	1,188	50	278	1,001	3,745	1,237	5,025	6,262
2001	2,196	50	590	961	12,993	1,841	14,949	16,790
2002 ^{e/}	3,982	50	150	1,012	3,939	2,099	7,034	9,133
2003 ^{e/}	2,412	50	326	505	6,539	1,472	8,360	9,832
2004 ^{e/}	1,337	50	343	1,269	6,527	1,874	7,652	9,526
2005 ^{e/f/}	10,273	0	487	1,218	7,182	2,197	16,963	19,160
2006 ^{e/f/}	2,248	0	141	604	1,832	1,620	3,165	4,785
2007 ^{e/f/g/}	1,430	0	NA	792	4,778	1,383	5,630	7,013
GOAL	Hatchery Production							

TABLE B-37. Estimated inriver run size, catch, and escapement for Quillayute River coho stocks in numbers of fish. (Page 2 of 2)

Year or Average	Terminal Catch ^{a/}			Escapement		Terminal Run Size		Total
	Gillnet	Ceremonial & Subsistence	River Sport ^{b/}	Natural ^{c/}	Hatchery ^{d/}	Natural ^{c/}	Hatchery ^{d/}	
FALL COHO								
1976-1980	5,985	53	70	9,002	2,435	13,959	3,587	17,546
1981-1985	3,789	49	164	7,464	2,102	10,988	2,580	13,568
1986-1990	5,794	100	385	8,766	1,771	14,119	2,695	16,815
1991-1995	3,598	100	565	7,357	4,736	9,930	6,426	16,356
1996	8,419	100	1,336	11,009	11,515	14,596	17,783	32,379
1997	456	50	38 ^{g/}	4,623	2,645	5,021	2,791	7,812
1998	4,606	50	1,340	13,866	12,834	16,980	15,716	32,696
1999	22,946	50	1,054	9,365	13,528	19,524	27,515	47,039
2000	5,606	50	1,059	13,343	13,118	17,706	15,470	33,176
2001	23,991	50	2,620	18,876	23,892	36,714	32,715	69,429
2002	22,214	50	2,002	23,016	30,656	34,695	43,243	77,938
2003	13,949	50	2,533	14,756	13,799	25,188	19,899	45,087
2004	19,321	50	2,831	13,354	21,248	25,118	31,687	56,805
2005 ^{e/f}	29,530	0	3,420	11,501	24,137	22,125	46,463	68,588
2006 ^{e/f}	9,677	0	291	5,642	4,450	12,627	7,433	20,060
2007 ^{e/f/g}	9,367	10	NA	5,609	5,423	9,758	10,671	20,429
GOAL				6,300-15,800				

a/ Includes dip-in fish from other systems.

b/ Recreational catch of adults (coho over 20 inches).

c/ Natural escapement and run size estimates include fish taken for hatchery brood stock.

d/ Hatchery escapement and terminal run size exclude hatchery strays.

e/ Preliminary.

f/ Beginning in 2005, C&S catch taken during scheduled gillnet fishery is included in gillnet harvest numbers. Catch during designated C&S fisheries is listed separately.

g/ Terminal run size estimates incomplete since inriver sport catch estimates are unavailable.

h/ Regulations required nonretention of coho.

TABLE B-38. Puget Sound commercial net and troll fishery salmon catches in numbers of fish.^{a/} (Page 1 of 2)

Year or Average	Fishery	Chinook	Coho	Pink ^{b/}	Chum	Sockeye
1971-1975	Non-Indian	105,332	525,867	1,172,614	331,029	2,158,784
	Treaty Indian	57,672	224,743	61,818	78,266	38,225
	Total	163,005	750,610	1,234,433	409,295	2,197,009
1976-1980	Non-Indian	103,546	413,583	1,050,560	407,859	1,095,603
	Treaty Indian	135,592	492,549	185,831	296,057	277,771
	Total	239,138	906,132	1,236,391	703,916	1,373,374
1981-1985	Non-Indian	72,934	346,125	1,154,851	368,762	928,477
	Treaty Indian	155,966	608,241	829,340	387,951	912,408
	Total	228,899	954,366	1,984,191	756,713	1,840,885
1986-1990	Non-Indian	57,550	470,494	509,445	540,843	964,690
	Treaty Indian	176,966	812,712	590,138	662,215	1,028,361
	Total	234,516	1,283,206	1,099,583	1,203,058	1,993,051
1991-1995	Non-Indian	17,519	74,371	784,067	523,396	735,834
	Treaty Indian	82,513	316,784	832,948	607,028	741,058
	Total	100,033	391,155	1,617,015	1,130,424	1,476,892
1996	Non-Indian	9,046	19,218	2	530,372	50,474
	Treaty Indian	67,061	153,748	58	264,486	286,187
	Total	76,107	172,966	60	794,858	336,661
1997	Non-Indian	21,894	10,454	869,345	229,261	690,236
	Treaty Indian	56,638	133,150	1,007,380	188,850	678,489
	Total	78,532	143,604	1,876,725	418,111	1,368,725
1998	Non-Indian	12,428	12,538	352	505,349	229,313
	Treaty Indian	43,273	148,441	512	320,122	308,446
	Total	55,701	160,979	864	825,471	537,759
1999	Non-Indian	9,512	11,902	1,109	133,404	37
	Treaty Indian	83,686	102,278	51,432	117,763	20,495
	Total	93,198	114,180	52,541	251,167	20,532

TABLE B-38. Puget Sound commercial net and troll fishery salmon catches in numbers of fish.^{a/} (Page 2 of 2)

Year or Average	Fishery	Chinook	Coho	Pink ^{b/}	Chum	Sockeye
2000	Non-Indian	11,468	21,910	9	140,611	230,379
	Treaty Indian	71,551	386,714	346	159,477	315,628
	Total	83,019	408,624	355	300,088	546,007
2001	Non-Indian	18,029	28,299	463,083	824,328	85,112
	Treaty Indian	109,865	366,011	319,553	777,019	170,309
	Total	127,894	394,310	782,636	1,601,347	255,421
2002 ^{c/}	Non-Indian	17,628	24,459	7	1,117,666	141,456
	Treaty Indian	98,251	286,500	327	833,497	339,773
	Total	115,879	310,959	334	1,951,163	481,229
2003 ^{c/}	Non-Indian	8,567	18,105	683,393	764,132	90,618
	Treaty Indian	84,680	244,091	556,943	814,212	183,670
	Total	93,247	262,196	1,240,336	1,578,344	274,288
2004 ^{c/}	Non-Indian	5,042	39,481	4	1,174,295	81,031
	Treaty Indian	99,741	529,668	712	934,155	137,699
	Total	104,783	569,149	716	2,108,450	218,730
2005 ^{c/}	Non-Indian	6,236	19,694	144,567	383,127	65,931
	Treaty Indian	86,071	296,637	243,012	351,416	141,747
	Total	92,307	316,331	387,579	734,543	207,678
2006 ^{c/}	Non-Indian	13,298	9,863	6	877,758	223,980
	Treaty Indian	134,604	292,448	195	546,812	543,546
	Total	147,902	302,311	201	1,424,570	767,526
2007 ^{c/}	Non-Indian	6,785	13,435	200,687	680,250	6,266
	Treaty Indian	115,263	212,925	313,383	782,907	7,334
	Total	122,048	226,360	514,070	1,463,157	13,600

a/ Data does not reflect treaty Indian allocations. Includes U.S. and Canadian-origin salmon and fish caught in test fisheries.

b/ Odd-year averages for pink salmon.

c/ Preliminary.

TABLE B-39. Summary of Puget Sound marine recreational salmon catch estimates in numbers of fish from catch record cards.^{a/} (Page 1 of 1)

Year or Average	Chinook	Coho	Pink ^{b/}
1971-1975	225,650	119,301	14,855
1976-1980	253,763	202,983	47,029
1981-1985 ^{c/}	156,183	196,632	14,910
1986-1990 ^{c/d/e/}	127,860	251,087	40,884
1991-1995 ^{e/f/}	77,310	137,637	71,030
1996 ^{e/}	72,069	85,139	50
1997 ^{e/}	60,425	137,571	35,197
1998 ^{e/}	26,114	89,520	201
1999 ^{e/}	28,739	22,055	23,780
2000 ^{e/}	23,879	74,934	17
2001 ^{e/}	44,422	193,454	117,367
2002 ^{e/g/}	30,900	67,261	31
2003 ^{e/g/}	30,936	101,485	148,965
2004 ^{e/g/}	27,121	88,036	213
2005 ^{e/g/}	22,758	62,110	68,166
2006 ^{e/g/}	32,794	29,780	19
2007 ^{e/g/}	NA	NA	NA

a/ WDFW Statistical Areas 5 through 13, which include the Strait of Juan de Fuca, San Juan Islands, and inner Puget Sound.

b/ Odd-year averages for pink salmon.

c/ 1981-1987: Adjusted all Puget Sound and freshwater estimates by 0.833, due to previous estimates being 20% too high.

d/ 1988: Area 5, no adjustment. Areas 6-13 adjusted by 0.633, due to estimates being 58% too high.

e/ 1989-Present: Area 5, no adjustment. Areas 6-13 adjusted by 0.685, due to estimates being 46% too high.

f/ 1991, 1992, and 1993 catch record card estimates adjusted for results of 1987-1990 WDFW/tribal sports emphasis study.

g/ Preliminary.

TABLE B-40. Puget Sound reported commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound Chinook stocks.^{df} (Page 1 of 4)

Year or Average	Commercial Net Catches			Spawning Escapement			Puget Sound Run Size ^{df}		
	Hatchery ^{df}	Natural	Total	Hatchery ^{df}	Natural	Total	Hatchery ^{df}	Natural	Total
Strait of Juan de Fuca									
1981-1985	57	126	183	811	1,450	2,261	868	1,576	2,444
1986-1990	136	456	591	1,276	4,755	6,031	1,411	5,211	6,622
1991-1995	69	108	177	979	2,390	3,369	1,048	2,498	3,546
1996	6	8	14	1,380	1,944	3,324	1,386	1,952	3,338
1997	14	42	56	938	2,823	3,761	952	2,865	3,817
1998	6	6	12	1,676	2,056	3,732	1,682	2,062	3,744
1999	10	15	25	726	2,675	3,401	736	2,690	3,426
2000	3	5	8	1,244	1,683	2,927	1,247	1,688	2,935
2001 ^{df}	4	4	8	1,660	1,947	3,607	1,664	1,951	3,615
2002 ^{df}	5	6	11	1,558	2,182	3,740	1,563	2,188	3,751
2003 ^{df}	6	15	21	1,258	2,787	4,045	1,264	2,802	4,066
2004 ^{df}	5	16	21	1,364	4,044	5,408	1,369	4,060	5,429
2005 ^{df}	7	8	15	1,401	1,961	3,362	1,408	1,969	3,377
2006 ^{df}	NA	NA	4	1,220	3,116	4,336	NA	NA	4,340
2007 ^{df}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL						5,300			
Nooksack-Samish									
1981-1985	54,054	33,567	87,621	16,083	6,541	22,623	70,137	40,107	110,244
1986-1990	38,058	26,273	64,330	10,729	4,127	14,856	48,786	30,400	79,186
1991-1995	18,245	2,294	20,539	8,646	731	9,376	26,891	3,025	29,916
1996	18,028	1,327	19,355	9,026	866	9,892	27,054	2,193	29,247
1997	18,200	3,743	21,943	15,775	3,985	19,760	33,975	7,728	41,703
1998	16,239	5,006	21,245	7,706	2,539	10,245	23,945	7,545	31,490
1999	25,724	6,804	32,528	6,962	2,598	9,560	32,686	9,402	42,088
2000	26,207	2,274	28,481	3,732	432	4,164	29,939	2,706	32,645
2001 ^{df}	22,213	27,164	49,377	6,300	9,017	15,317	28,513	36,181	64,694
2002 ^{df}	11,636	27,079	38,715	4,280	11,307	15,587	15,916	38,386	54,302
2003 ^{df}	6,582	12,251	18,833	3,302	7,864	11,166	9,884	20,115	29,999
2004 ^{df}	4,700	5,913	10,613	2,966	4,325	7,291	7,666	10,238	17,904
2005 ^{df}	8,432	3,584	12,016	2,438	1,411	3,849	10,870	4,995	15,865
2006 ^{df}	16,222	8,682	24,904	3,501	2,277	5,778	19,723	10,959	30,682
2007 ^{df}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL				1,800					

TABLE B-40. Puget Sound reported commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound Chinook stocks.^{a/} (Page 2 of 4)

Year or Average	Commercial Net Catches			Spawning Escapement			Puget Sound Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Skagit									
1981-1985	595	9,162	9,757	787	11,109	11,896	1,382	20,271	21,653
1986-1990	251	4,043	4,294	815	12,452	13,267	1,066	16,495	17,561
1991-1995	463	1,583	2,046	2,402	6,284	8,686	2,865	7,868	10,732
1996	21	209	230	1,133	10,706	11,839	1,154	10,915	12,069
1997	0	1,145	1,145	0	4,951	4,951	0	6,096	6,096
1998	0	321	321	0	14,700	14,700	0	15,021	15,021
1999	0	262	262	0	5,002	5,002	0	5,264	5,264
2000	0	321	321	0	17,024	17,024	0	17,345	17,345
2001 ^{ar}	0	251	251	0	13,868	13,868	0	14,119	14,119
2002 ^{dr}	1	323	324	101	19,676	19,777	102	19,999	20,101
2003 ^{dr}	10	322	332	298	9,964	10,262	308	10,286	10,594
2004 ^{dr}	0	559	559	0	23,750	23,750	0	24,309	24,309
2005 ^{dr}	41	2,573	2,614	331	20,803	21,134	372	23,376	23,748
2006 ^{dr}	30	1,693	1,723	368	20,819	21,187	398	22,512	22,910
2007 ^{dr}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					14,900				
Hood Canal									
1981-1985	4,918	3,649	8,567	3,787	2,038	5,824	8,705	5,686	14,391
1986-1990	10,493	4,938	15,432	6,189	2,006	8,195	16,682	6,944	23,626
1991-1995	1,830	1,020	2,850	3,946	1,408	5,355	5,776	2,429	8,205
1996	31	4	35	7,103	1,028	8,131	7,134	1,032	8,166
1997	118	7	125	7,295	492	7,787	7,413	499	7,912
1998	111	5	116	13,436	1,640	15,076	13,547	1,645	15,192
1999	1,302	202	1,504	18,361	2,895	21,256	19,663	3,097	22,760
2000	1,446	151	1,597	8,816	1,792	10,608	10,262	1,943	12,205
2001 ^{ar}	3,650	625	4,275	13,042	3,002	16,044	16,692	3,627	20,319
2002 ^{dr}	3,285	433	3,718	12,881	1,725	14,606	16,166	2,158	18,324
2003 ^{dr}	4,521	391	4,912	13,643	1,512	15,155	18,164	1,903	20,067
2004 ^{dr}	9,121	1,676	10,797	15,666	3,663	19,329	24,787	5,339	30,126
2005 ^{dr}	16,558	1,719	18,277	19,782	2,775	22,557	36,340	4,494	40,834
2006 ^{dr}	NA	NA	30,405	15,714	1,549	17,263	NA	NA	47,668
2007 ^{dr}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL				3,400					

TABLE B-40. Puget Sound reported commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound Chinook stocks.^{a/} (Page 3 of 4)

Year or Average	Commercial Net Catches			Spawning Escapement			Puget Sound Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Stillaguamish-Snohomish^{e/}									
1981-1985	3,894	6,917	10,811	1,990	4,901	6,891	5,884	11,818	17,702
1986-1990	3,370	4,241	7,612	1,148	5,210	6,358	4,519	9,451	13,970
1991-1995	3,688	1,965	5,653	2,253	4,371	6,624	5,941	6,337	12,278
1996	11,429	27	11,456	4,618	7,138	11,756	16,047	7,165	23,212
1997	8,862	112	8,974	11,777	5,448	17,225	20,639	5,560	26,199
1998	7,207	68	7,275	4,691	7,844	12,535	11,898	7,912	19,810
1999	15,192	33	15,225	4,700	5,897	10,597	19,892	5,930	25,822
2000	8,275	31	8,306	1,931	7,739	9,670	10,206	7,770	17,976
2001 ^{d/}	5,115	293	5,408	872	9,513	10,385	5,987	9,806	15,793
2002 ^{d/}	5,192	57	5,249	2,542	8,808	11,350	7,734	8,865	16,599
2003 ^{d/}	8,793	143	8,936	5,655	6,435	12,090	14,448	6,578	21,026
2004 ^{d/}	5,746	102	5,848	6,124	12,112	18,236	11,870	12,214	24,084
2005 ^{d/}	7,459	151	7,610	3,592	5,447	9,039	11,051	5,598	16,649
2006 ^{d/}	4,063	148	4,211	4,802	9,562	14,364	8,865	9,710	18,575
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					7,300				
South Puget Sound									
1981-1985	25,101	9,101	34,201	23,341	6,371	29,712	48,442	15,472	63,913
1986-1990	25,697	20,036	45,733	36,998	18,108	55,106	62,695	38,144	100,839
1991-1995	19,393	13,066	32,459	30,556	14,488	45,044	49,950	27,554	77,503
1996	17,338	13,118	30,456	35,858	24,769	60,627	53,196	37,887	91,083
1997	10,542	5,124	15,666	27,717	26,393	54,110	38,259	31,517	69,776
1998	11,105	8,385	19,490	38,491	25,370	63,861	49,596	33,755	83,351
1999	16,722	9,612	26,334	50,600	31,812	82,412	67,322	41,424	108,746
2000	14,861	13,893	28,754	32,496	27,998	60,494	47,357	41,891	89,248
2001 ^{d/}	21,598	18,555	40,153	55,026	42,069	97,095	76,624	60,624	137,248
2002 ^{d/}	22,503	15,835	38,338	46,744	41,135	87,879	69,247	56,970	126,217
2003 ^{d/}	23,319	13,704	37,023	33,234	29,998	63,232	56,553	43,702	100,255
2004 ^{d/}	20,731	15,521	36,252	45,706	26,864	72,570	66,437	42,385	108,822
2005 ^{d/}	21,644	6,661	28,305	52,049	12,540	64,589	73,693	19,201	92,894
2006 ^{d/}	39,467	15,253	54,720	65,562	27,433	92,995	105,029	42,686	147,715
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL						34,900			

TABLE B-40. Puget Sound reported commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound Chinook stocks.^{a/} (Page 4 of 4)

a/ Includes treaty Indian and non-Indian net commercial catches during the adult accounting period. Source: Puget Sound run reconstruction model.

b/ Includes estimated off-station returns.

c/ Puget Sound run size is defined as the run available to Puget Sound net fisheries; spawning escapement plus Puget Sound net fishery catch. Does not include fish caught by troll and recreational fisheries inside Puget Sound.

d/ Preliminary.

e/ Since 1999, numbers include Tulalip hatchery returns, which are not added into escapement since no broodstock is taken at the hatchery.

TABLE B-41. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound coho stocks.^{a/} (Page 1 of 3)

Year or Average	Commercial Net Catches ^{c/}			Spawning Escapement			Terminal Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Strait of Juan de Fuca									
1981-1985	15,822	2,907	18,729	9,300	5,960	15,260	25,122	8,867	33,989
1986-1990	5,956	2,301	8,258	2,913	6,920	9,833	8,869	9,221	18,091
1991-1995	1,872	286	2,158	4,316	4,810	9,126	6,188	5,096	11,284
1996	4,176	81	4,257	7,563	2,440	10,003	11,739	2,521	14,260
1997 ^{d/}	224	61	285	13,166	13,065	26,231	13,390	13,126	26,516
1998 ^{d/}	5,079	1,249	6,328	5,166	16,822	21,988	10,245	18,071	28,316
1999 ^{d/}	3,424	719	4,143	6,253	8,484	14,737	9,677	9,203	18,880
2000 ^{d/}	7,727	1,988	9,715	19,233	22,654	41,887	26,960	24,642	51,602
2001 ^{d/}	10,686	2,640	13,326	24,768	35,274	60,042	35,454	37,914	73,368
2002 ^{d/}	7,744	1,876	9,620	10,398	22,375	32,773	18,142	24,251	42,393
2003 ^{d/}	2,910	745	3,655	15,004	20,991	35,995	17,914	21,736	39,650
2004 ^{d/}	3,601	873	4,474	5,461	20,987	26,448	9,062	21,860	30,922
2005 ^{d/}	3,296	762	4,058	4,123	11,102	15,225	7,419	11,864	19,283
2006 ^{d/}	NA	NA	1,065	605	NA	NA	NA	NA	NA
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL						14,800			
Nooksack-Samish									
1981-1985	122,433	17,539	139,972	27,720	7,700	35,420	150,153	25,239	175,392
1986-1990	140,733	21,839	162,572	23,087	8,020	31,107	163,821	29,859	193,680
1991-1995	48,056	13,878	61,934	19,793	10,835	30,629	67,849	24,713	92,563
1996	50,711	1,607	52,318	38,293	2,018	40,311	89,004	3,625	92,629
1997 ^{d/}	14,464	546	15,010	34,305	6,700	41,005	48,769	7,246	56,015
1998 ^{d/}	15,752	7,134	22,886	21,089	10,300	31,389	36,841	17,434	54,275
1999 ^{d/}	41,926	7,457	49,383	41,876	8,039	49,915	83,802	15,496	99,298
2000 ^{d/}	58,008	9,597	67,605	49,035	11,000	60,035	107,043	20,597	127,640
2001 ^{d/}	49,044	26,098	75,142	49,788	27,500	77,288	98,832	53,598	152,430
2002 ^{d/}	34,626	16,824	51,450	45,161	20,300	65,461	79,787	37,124	116,911
2003 ^{d/}	33,914	9,451	43,365	35,482	14,200	49,682	69,396	23,651	93,047
2004 ^{d/}	70,761	18,861	89,622	27,625	11,591	39,216	98,386	30,452	128,838
2005 ^{d/}	20,079	15,497	35,576	25,211	2,187	27,398	45,290	17,684	62,974
2006 ^{d/}	16,348	5,429	21,777	8,533	1,233	9,766	24,881	6,662	31,543
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL				17,900					

TABLE B-41. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound coho stocks.^{a/}
(Page 2 of 4)

Year or Average	Commercial Net Catches ^{c/}			Spawning Escapement			Terminal Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Skagit									
1981-1985	6,619	8,858	15,477	21,740	19,800	41,540	28,359	28,658	57,017
1986-1990	5,309	11,448	16,757	13,861	25,800	39,661	19,170	37,248	56,418
1991-1995	1,338	1,739	3,077	11,082	14,240	25,322	12,420	15,979	28,399
1996	719	332	1,051	17,983	8,300	26,283	18,702	8,632	27,334
1997 ^{d/}	119	1,412	1,531	4,443	38,999	43,442	4,562	40,411	44,973
1998 ^{d/}	750	9,564	10,314	11,046	73,678	84,724	11,796	83,242	95,038
1999 ^{d/}	494	6,778	7,272	3,024	27,341	30,365	3,518	34,119	37,637
2000 ^{d/}	1,526	11,778	13,304	13,935	62,898	76,833	15,461	74,676	90,137
2001 ^{d/}	1,658	17,933	19,591	16,852	87,017	103,869	18,510	104,950	123,460
2002 ^{d/}	2,204	11,742	13,946	19,096	55,968	75,064	21,300	67,710	89,010
2003 ^{d/}	4,193	18,644	22,837	9,118	69,221	78,339	13,311	87,865	101,176
2004 ^{d/}	7,434	27,943	35,377	11,822	138,804	150,626	19,256	166,747	186,003
2005 ^{d/}	3,251	16,052	19,303	12,139	34,658	46,797	15,390	50,710	66,100
2006 ^{d/}	1,006	4,429	5,435	1,266	14,451	15,717	2,272	18,880	21,152
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					30,000				
Hood Canal									
1981-1985	36,470	21,180	57,650	19,020	23,589	42,609	55,490	44,769	100,259
1986-1990	42,838	21,862	64,699	14,711	18,328	33,039	57,549	40,190	97,738
1991-1995	13,334	673	14,007	14,792	30,048	44,840	28,126	30,721	58,847
1996	4,066	137	4,203	27,337	37,051	64,388	31,403	37,188	68,591
1997 ^{d/}	3,171	2,966	6,137	37,208	95,756	132,964	40,379	98,722	139,101
1998 ^{d/}	6,385	15,587	21,972	13,761	100,818	114,579	20,146	116,405	136,551
1999 ^{d/}	3,641	1,246	4,887	14,113	16,563	30,676	17,754	17,809	35,563
2000 ^{d/}	11,169	11,888	23,057	24,940	27,239	52,179	36,109	39,127	75,236
2001 ^{d/}	10,384	10,278	20,662	39,243	94,773	134,016	49,627	105,051	154,678
2002 ^{d/}	9,768	8,373	18,141	39,330	69,300	108,630	49,098	77,673	126,771
2003 ^{d/}	9,647	23,764	33,411	33,221	170,255	203,476	42,868	194,019	236,887
2004 ^{d/}	19,366	67,318	86,684	26,692	146,873	173,565	46,058	214,191	260,249
2005 ^{d/}	34,789	26,788	61,577	31,064	38,066	69,130	65,853	64,854	130,707
2006 ^{d/}	NA	NA	48,924	NA	13,800	NA	NA	NA	NA
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					21,500				

TABLE B-41. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound coho stocks.^{a/}
(Page 3 of 4)

Year or Average	Commercial Net Catches ^{c/}			Spawning Escapement			Terminal Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Stillaguamish-Snohomish									
1981-1985	19,973	47,552	67,524	12,940	88,000	100,940	32,913	135,552	168,464
1986-1990	58,543	86,887	145,431	26,134	110,400	136,534	84,677	197,287	281,965
1991-1995	40,705	21,375	62,080	23,570	97,720	121,290	64,275	119,095	183,370
1996	23,406	7,159	30,565	23,583	59,200	82,783	46,989	66,359	113,348
1997 ^{d/}	19,311	5,881	25,192	22,667	69,110	91,777	41,978	74,991	116,969
1998 ^{d/}	14,522	10,207	24,729	18,715	177,300	196,015	33,237	187,507	220,744
1999 ^{d/}	16,635	1,634	18,269	11,578	68,300	79,878	28,213	69,934	98,147
2000 ^{d/}	84,221	5,682	89,903	31,338	122,510	153,848	115,559	128,192	243,751
2001 ^{d/}	58,375	17,138	75,513	41,516	334,630	376,146	99,891	351,768	451,659
2002 ^{d/}	49,488	18,372	67,860	12,732	187,305	200,037	62,220	205,677	267,897
2003 ^{d/}	1,997	7,290	9,287	14,925	228,290	243,215	16,922	235,580	252,502
2004 ^{d/}	52,047	36,558	88,605	13,984	310,904	324,888	66,031	347,462	413,493
2005 ^{d/}	21,872	14,558	36,430	13,627	134,804	148,431	35,499	149,362	184,861
2006 ^{d/}	4,722	27,101	31,823	6,147	93,517	99,664	10,869	120,618	131,487
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL - Snohomish					70,000				
GOAL - Stillaguamish					17,000				
South Puget Sound									
1981-1985	328,516	141,229	469,745	76,560	38,510	115,070	405,076	179,738	584,815
1986-1990	509,525	211,476	721,001	69,198	28,882	98,080	578,723	240,358	819,081
1991-1995	137,961	56,462	194,423	97,002	23,945	120,947	234,963	80,407	315,370
1996	56,117	13,503	69,620	107,463	21,991	129,454	163,580	35,494	199,074
1997 ^{d/}	26,963	51,491	78,454	61,274	40,500	101,774	88,237	91,991	180,228
1998 ^{d/}	49,006	14,264	63,270	33,290	18,052	51,342	82,296	32,316	114,612
1999 ^{d/}	15,917	5,429	21,346	26,559	10,008	36,567	42,476	15,437	57,913
2000 ^{d/}	136,840	59,254	196,094	139,838	51,192	191,030	276,678	110,446	387,124
2001 ^{d/}	109,772	60,354	170,126	127,179	37,688	164,867	236,951	98,042	334,993
2002 ^{d/}	96,462	34,212	130,674	115,145	18,296	133,441	211,607	52,508	264,115
2003 ^{d/}	95,300	32,508	127,808	94,890	51,654	146,544	190,190	84,162	274,352
2004 ^{d/}	171,316	46,632	217,948	133,614	43,147	176,761	304,930	89,779	394,709
2005 ^{d/}	108,472	31,740	140,212	83,761	33,620	117,381	192,233	65,360	257,593
2006 ^{d/}	114,533	29,395	143,928	49,610	21,449	71,059	164,143	50,844	214,987
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL				52,000					

TABLE B-41. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound coho stocks.^{a/}
(Page 4 of 4)

a/ Includes treaty Indian and non-Indian net commercial catches during the adult accounting period. Source: Puget Sound run reconstruction model.

b/ Includes estimated off-station returns.

c/ Terminal run size is defined as the run to terminal marine areas; spawning escapement plus sport and commercial net catch (inriver and terminal fishery catch). Prior to 1996, estimates are Puget Sound run size, which is defined as the run available to Puget Sound net fisheries; spawning escapement plus commercial net catch (inriver, terminal, and pre-terminal Puget Sound net fishery catch), but not including fish caught in Puget Sound troll and recreational fisheries.

d/ Preliminary.

TABLE B-42. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound pink stocks.^{a/}
(Page 2 of 4)

Year or Average	Commercial Net Catches			Spawning Escapement			Puget Sound Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Skagit									
1981	403	150,626	151,029	268	100,268	100,536	671	250,894	251,565
1983	4	19,023	19,027	128	470,128	470,256	132	489,151	489,283
1985	9	229,993	230,002	30	710,030	710,060	39	940,023	940,062
1987	1,090	421,176	422,266	1,535	593,535	595,070	2,625	1,014,711	1,017,336
1989	8	661,061	661,069	5	401,300	401,305	13	1,062,361	1,062,374
1991	0	188,927	188,927	0	351,000	351,000	0	539,927	539,927
1993	0	180,088	180,088	0	530,000	530,000	0	710,088	710,088
1995	0	568,561	568,561	0	857,000	857,000	0	1,425,561	1,425,561
1997	0	57,710	57,710	0	60,000	60,000	0	117,710	117,710
1999	0	32,626	32,626	0	320,000	320,000	0	352,626	352,626
2001	0	206,533	206,533	0	894,061	894,061	0	1,100,594	1,100,594
2003	0	232,732	232,732	0	567,080	567,080	0	799,812	799,812
2005 ^{d/}	0	20,147	20,147	0	60,000	60,000	0	80,147	80,147
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					330,000				
Hood Canal									
1981	380	1,241	1,621	1,557	6,551	8,108	1,937	7,792	9,729
1983	50	831	881	503	25,201	25,704	553	26,032	26,585
1985	138	2,854	2,992	1,456	64,101	65,557	1,594	66,955	68,549
1987	1,855	6,942	8,797	8,056	62,220	70,276	9,911	69,162	79,073
1989	7,799	26,946	34,745	2,500	60,970	63,470	10,299	87,916	98,215
1991	409	13,518	13,927	3,300	118,450	121,750	3,709	131,968	135,677
1993	623	1,917	2,540	11,497	35,647	47,144	12,120	37,564	49,684
1995	1,565	994	2,559	24,665	31,306	55,971	26,230	32,300	58,530
1997	2,436	910	3,346	21,493	8,363	29,856	23,929	9,273	33,202
1999	18	10	28	7,617	12,667	20,284	7,635	12,677	20,312
2001	713	703	1,416	71,539	98,338	169,877	72,252	99,041	171,293
2003	464	691	1,155	25,217	37,531	62,748	25,681	38,222	63,903
2005 ^{d/}	98	121	219	14,116	17,481	31,597	14,214	17,602	31,816
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					Not Agreed Upon				

TABLE B-42. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound pink stocks.^{a/}
(Page 3 of 4)

Year or Average	Commercial Net Catches			Spawning Escapement			Puget Sound Run Size ^{c/}		
	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total	Hatchery ^{b/}	Natural	Total
Stillaguamish-Snohomish									
1981	40	49,480	49,520	96	108,096	108,192	136	157,576	157,712
1983	51	57,452	57,503	283	324,383	324,666	334	381,835	382,169
1985	63	175,095	175,158	192	502,192	502,384	255	677,287	677,542
1987	173	111,881	112,054	418	271,418	271,836	591	383,299	383,890
1989	33	354,805	354,838	16	150,549	150,565	49	505,354	505,403
1991	139	82,150	82,289	447	260,000	260,447	586	342,150	342,736
1993	13	21,444	21,457	135	210,000	210,135	148	231,444	231,592
1995	5	33,871	33,876	26	309,600	309,626	31	343,471	343,502
1997	0	59,173	59,173	0	192,109	192,109	0	251,282	251,282
1999	0	13,443	13,443	0	461,543	461,543	0	474,986	474,986
2001	0	100,015	100,015	0	1,847,648	1,847,648	0	1,947,663	1,947,663
2003	0	187,286	187,286	0	1,577,001	1,577,001	0	1,764,287	1,764,287
2005 ^{d/}	0	19,193	19,193	0	600,124	600,124	0	619,317	619,317
2007 ^{d/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL - Stillaguamish					155,000				
GOAL - Snohomish					120,000				
South Puget Sound									
1981	1,569	9,818	11,387	791	12,715	13,506	2,360	22,533	24,893
1983	492	11,265	11,757	149	12,200	12,349	641	23,465	24,106
1985	119	5,335	5,454	13	34,700	34,713	132	40,035	40,167
1987	15	9,386	9,401	3	42,200	42,203	18	51,586	51,604
1989	361	36,999	37,360	452	62,220	62,672	813	99,219	100,032
1991	357	5,037	5,394	346	15,950	16,296	703	20,987	21,690
1993 ^{e/}	3	2,330	2,333	21	10,619	10,640	24	12,949	12,973
1995 ^{e/}	13	5,163	5,176	84	18,278	18,362	97	23,441	23,538
1997 ^{e/}	0	449	449	0	2,965	2,965	0	3,414	3,414
1999 ^{e/}	0	80	80	12	4,670	4,682	12	4,750	4,762
2001 ^{e/f/}	5	735	740	48	16,173	16,221	53	16,908	16,961
2003 ^{e/f/}	1	5,393	5,394	68	185,277	185,345	69	190,670	190,739
2005 ^{d/e/f/}	0	3,964	3,964	0	466,435	466,435	0	470,399	470,399
2007 ^{d/e/f/}	NA	NA	NA	NA	NA	NA	NA	NA	NA
GOAL					25,000				

TABLE B-42. Puget Sound commercial net fishery catches and spawning escapements in numbers of fish for hatchery and natural Puget Sound pink stocks.^{a/}
(Page 4 of 4)

a/ Includes treaty Indian and non-Indian net commercial catches during the adult accounting period. Source: Puget Sound run reconstruction model.

b/ Includes estimated off-station returns.

c/ Puget Sound run size is defined as the run available to Puget Sound net fisheries; spawning escapement plus Puget Sound net fishery catch. Does not include fish caught by troll and recreational fisheries inside Puget Sound.

d/ Preliminary.

e/ Nisqually escapement estimate incomplete.

f/ Large runs of pinks have returned to Green River in 2001, 2003, and 2005; however, no formal escapement methodology exists, and Green River pinks are not included in the run reconstruction model. When the model is revised, pre-terminal catch estimates for all stocks will be affected.

TABLE B-43. Puget Sound spring Chinook spawning escapement estimates in numbers of adult fish. (Page 1 of 1)

Year or Average	Stock						
	Skagit		NF Nooksack		SF Nooksack	White River	Quilcene
	Hatchery	Natural	Hatchery	Natural ^{a/}	Hatchery/ Natural	Hatchery ^{b/}	Hatchery ^{c/}
1981-1985	15	1,408	0	152	317	70	149
1986-1990	155	1,826	0	235	280	408	125
1991-1995	815	907	770	266	222	1,065	19
1996	856	1,051	1,070	534	203	1,625	12
1997	1,220	1,041	1,663	520	180	1,609	16
1998	1,054	1,086	1,370	368	157	2,710	5
1999	3,171	471	2,873	823	166	1,550	4
2000	1,102	1,021	1,204	1,245	284	2,363	0
2001	1,566	1,856	1,006	2,209	267	5,690	0
2002 ^{d/}	1,606	1,065	5,649	3,741	289	1,780	0
2003 ^{d/}	1,537	844	6,250	2,857	204	2,760	0
2004 ^{d/}	3,107	1,622	3,533	1,746	130	1,115	0
2005 ^{d/}	2,254	1,305	1,569	2,167	120	2,061	0
2006 ^{d/}	1,487	1,896	732	1,184	355	4,321	0
2007 ^{d/}	1,931	613	505	1,438	182	8,417	0
GOAL		3,000					

a/ Natural escapement estimates based on carcass counts expanded by a 3.48 multiplier developed from 5 years of redd count based estimates. Most natural spawners are hatchery fish spawning in the wild.

b/ This estimate includes adult Chinook returns to Hupp Springs, White River Hatchery and to the Buckley Trap.

c/ Program has been discontinued.

d/ Preliminary.

Page Left Intentionally Blank