

Table 2-3. Estimated time to rebuild and SPR harvest rate relative to alternative 2011-2012 ACLs for depleted west coast groundfish species.

Species	Ttarget in the FMP	ACL Alt.	Median Time to Rebuild	ACLs (mt)		SPR HR	T @ F=0	Current Tmax	Re-est. Tmax
				2011	2012				
Bocaccio (S of 40°10' N lat.)	2026	1	2019	0	0	F100%	2019	2033	2031
		2	2019	53	56	F95%			
		3	2020	109	115	F90%			
		4	2022	263	274	F77.7%			
		5	2024	373	384	F70%			
		6	2028	539	545	F60%			
Canary	2021	1	2024	0	0	F100%	2024	2035	2046
		2	2025	49	51	F94.4%			
		3	2026	69	72	F92.2%			
		4	2027	102	107	F88.7%			
		5	2028	155	162	F83.4%			
		6	2031	253	263	F74.4%			
			2035	308	318	F70%			
			2043	396	408	F63.4%			
	2046	415	426	F62.1%					
Cowcod	2072	1	2060	0	0	F100%	2060	2098	2097
		2	2064	2	2	F90%			
		3	2072	4	4	F77.9%			
		4	2097	9	9	F59.7%			
Darkblotched	2011	1	2016	0	0	F100%	2016	2040	2037
		2	2018	130	131	F81.8%			
		3	2021	222	222	F71.9%			
		4	2024	298	296	F64.9%			
		5	2026	332	329	F62.1%			
		6	2037	461	453	F52.8%			
POP	2017	1	2018	0	0	F100%	2018	2037	2045
		2	2020	180	183	F86.4%			
		3	2021	204	208	F84.8%			
		4	2021	265	269	F81.1%			
			2024	404	408	F73.6%			
			2031	635	635	F63.6%			
			2038	751	747	F59.5%			
		5	2045	836	829	F56.8%			
Widow	2015	1	2010	0	0		2010	2023	NA
		2	2010	200	200				
		3	2010	400	400				
		4	2010	600	600				
		5	2010	1,000	1,000				
		6	2010	3,000	3,000				
Yelloweye	2084	1	2047	0	0	F100%	2047	2090	2092
		2	2058	9	9	F86%			
		3	2065	13	13	F80.7%			
		4	2074	17	17	F76%			
		5	2087	20	21	F71.9%			
		6	2092	21	22	F70.9%			