

CHAPTER III

COHO SALMON MANAGEMENT

OREGON PRODUCTION INDEX AREA COHO STOCKS

Oregon Production Index (OPI) area coho stocks include all Washington, Oregon, and California natural and hatchery stocks from streams south of Leadbetter Point, Washington, although stocks produced north of Leadbetter Point are also intercepted in the OPI area. The largest naturally produced coho stock is OCN coho, which includes coho produced from Oregon river and lake systems south of the Columbia River. OCN coho are managed as a stock aggregate with four identified components. NMFS listed three coho ESUs within the OPI area as threatened: CCC coho listed October 1996, SONCC coho listed May 1997, and OCN coho listed August 1998. In 2002, NMFS began an update of all its listing determinations and in January of 2006 concluded that the OCN ESU did not warrant listing under the ESA. That determination was overruled by a U.S. Court decision in 2007, and subsequently relisted by NMFS as threatened in February 2008. Columbia River natural coho were listed as endangered under the Oregon State ESA in 2002, and as threatened under the Federal ESA on June 28, 2005. The primary hatchery stocks include a south migrating Columbia River (early) stock, a north migrating Columbia River (late) stock, public hatchery coho from the Oregon and northern California Coast, and a small cooperative program along the southern Oregon Coast known as the Salmon Trout Enhancement Program (STEP).

Management Objectives

In establishing ocean salmon fisheries that impact OPI area coho stocks, the Council was guided by the reasonable and prudent alternatives of NMFS 1999 Supplemental Biological Opinion and Incidental Take Statement for CCC and SONCC coho and the March 2007 NMFS ESA guidance letter for LCR natural coho, which required:

- 1.No directed coho fisheries or retention of coho in all commercial and recreational fisheries off California to protect threatened CCC coho.
- 2.Marine fishery impacts on threatened CCC and SONCC coho must be no more than 13.0% as indicated by projected impacts on RK hatchery coho.
- 3.Fishery impacts on threatened LCN coho must not exceed a coastwide marine and mainstem Columbia River exploitation rate of 20.0%.

Based on parent escapement levels and observed OPI smolt-to-jack survival for 2004 brood OPI smolts, the total allowable OCN coho exploitation rate for 2007 fisheries was no greater than 20.0% under both the Salmon FMP (Amendment 13) and the matrix developed by the OCN work group during their review of Amendment 13. The work group recommendation was accepted by the Council as expert biological advice in November 2000.

The Council was also guided by a treaty Indian/non-Indian sharing agreement for Columbia upriver coho stocks, which required passage of 50% of the run destined for areas above Bonneville Dam.

Regulations to Achieve Objectives

Historically, OPI area coho stocks contributed primarily to ocean fisheries off Oregon and northern California and, to a lesser degree, Washington and B.C. The Council prohibited retention of coho in all

fisheries south of the Oregon/California border, and adopted seasons the STT projected would result in exploitation rates of 5.8% for RK coho in marine fisheries, 11.3% for OCN coho in marine and freshwater fisheries combined, and 13.3% for LCN coho in marine fisheries.

Commercial Troll

Commercial troll fisheries had been closed to coho retention south of Cape Falcon since 1993. Chinook fishery closures and gear restrictions (four-spread requirement) were also used to reduce coho impacts. In 2007, there was a commercial coho fishery with a quota of 10,000 (non-mark selective). Non-retention mortality on coho resulting from commercial Chinook fisheries south of Cape Falcon and the coho retention fishery was projected to be equivalent to exploitation rates of 3.1% for OCN coho and 2.1% for LCN coho.

Non-Indian commercial troll fisheries from Cape Falcon to the U.S./Canada border in 2007 had an overall quota of 22,400 coho. The fisheries were restricted to mark-selective coho retention.

All species treaty Indian fisheries north of Cape Falcon were not restricted to mark-selective retention of coho, and operated on an overall quota of 38,000 coho.

Recreational

Retention of coho has been limited in the recreational fisheries south of Cape Falcon since 1993. Retention of coho has been prohibited off California since 1996 to protect ESA listed CCC coho. All coho directed recreational fisheries in the OPI area have been mark-selective since 1998. Adequate abundance of marked coho in the OPI area has resulted in allowable harvests of marked coho in Oregon and Washington within constraints for OCN and LCN coho. In 2007, recreational coho fisheries north of Cape Falcon operated with quotas of 12,230 in the Neah Bay area, 3,060 in the La Push area, 28,510 in the Westport area, and 71,450 in the Columbia River area. The recreational fishery between Cape Falcon and the OR/CA border operated with a quota of 50,000 (Table I-3).

Inside Harvest

Coho retention in all California fisheries was prohibited.

The 2007 inside recreational harvest of coho in Oregon coastal streams, as in recent years, was very restricted and generally limited to areas where surplus hatchery coho returns were expected. Mark-selective coho fisheries occurred in nine freshwater areas. Estimates of the 2007 inriver recreational coho harvest were not available. Historical estimates of the recreational harvest of adult coho in Oregon coastal estuaries and rivers, derived from ODFW salmon and steelhead angler catch record cards, are reported in Table III-1.

A limited fishery for naturally-produced coho was approved in Siltcoos and Tahkenitch Lakes. The recreational fishery opened October 1 and closed December 31 as scheduled. The final catch estimates were not available but are expected to be similar to the previous 3-year average of 330 adults in the Siltcoos Lake fishery and 65 adults in the Tahkenitch Lake fishery.

The 2007 Columbia River non-Indian commercial gillnet fishery harvested 39,100 adult coho, compared to 63,400 coho in 2006. Select Area fisheries in both Oregon and Washington accounted for 10,100 the total 2007 Columbia River commercial coho catch. The Columbia River treaty Indian mainstem commercial gillnet coho catch was approximately 7,900 fish, compared to the 2006 catch of 5,400 coho. All Columbia River coho commercial fisheries were non-mark-selective. Coho harvest statistics for Columbia River commercial and recreational fisheries are presented in Appendix B, Table B-21.

The Buoy 10 and mainstem recreational fisheries below Bonneville Dam harvested 21,800 adult coho compared to 20,000 adult coho in 2006. In 2007, Columbia River managers opened the Buoy 10 fishery August 1 for adipose fin-clipped coho. The fishery ran through December 31 with the upriver boundary at the Tongue Point, Oregon to Rocky Point, Washington line. The 2007 Buoy 10 harvest and effort totaled 8,400 coho and 36,100 angler trips (Table III-2). All Columbia River recreational fisheries were mark-selective for coho. Historical Buoy 10 catch and effort data are provided in Appendix B, Table B-22. Recreational coho harvest estimates for Columbia River tributaries were not available.

Escapement and Management Performance

The overall abundance estimate for OPI area stocks in 2007 was 536,600 down from 557,100 in 2006 and less than the ten-year average of 737,300 (Table III-3; Figure III-1).

Central California Coast and Northern California Coho

Spawner estimates were not available for CCC coho. Estimates were available for escapement to Klamath River Basin hatcheries, but not for coho spawning in natural areas. In 2007, a total of 2,605 coho returned to Trinity River Hatchery and 625 coho returned to Iron Gate Hatchery. These values compare to a combined goal of 2,000 adults.

Oregon Coast Natural Coho

The preliminary estimate of natural spawner escapement in 2007 to Oregon coastal river and lake systems from the Coquille River north (Oregon coast ESU) was 51,900 adult coho by SRS accounting. This compares to 128,800 adults in 2006. Historical spawner escapement estimates of naturally produced coho are reported in Table III-1 and have been adjusted to reflect SRS accounting.

Preliminary information based on SRS surveys indicate the lowest total natural spawning population on the Oregon coast since 1999. The estimate of the natural spawning population in 2007 was 57,100, including estimates from the Rogue River, which is part of the SONCC ESU (Table III-4, Figure III-2).

Preliminary postseason estimates of combined marine and freshwater exploitation on OCN coho was 10.6%, slightly less than the preseason projection of 11.3%, and well below the 20% maximum allowed under the FMP and the OCN workgroup matrix. Preliminary postseason estimates of marine exploitation on RK coho was 5.4%, slightly less than the preseason projection of 5.8%, and well below the 13.0% maximum ESA consultation standard.

Oregon Coastal Hatchery Coho

Preliminary estimates of total coho returns to Oregon coastal public hatcheries and STEP smolt production facilities were 3,600 adults (Table III-1). Hatchery egg-take goals were expected to be met at all public hatchery stations.

Columbia River Coho

The 2007 ocean escapement of adult early and late Columbia River coho stocks was 318,600 fish, compared to 383,000 adults in 2006 (Appendix B, Table B-21). The 2007 Columbia River coho abundance was sufficient to meet all hatchery brood stock escapement needs.

Preliminary postseason estimates of marine exploitation on LCN coho was 11.9%, less than the preseason projected 13.3%.

WASHINGTON COASTAL COHO STOCKS

Washington coastal coho stocks include all natural and hatchery stocks originating in Washington coastal streams north of the Columbia River through the western Strait of Juan de Fuca (west of the Elwha River, inclusive). The stocks in this group most pertinent to ocean salmon fishery management were Willapa Bay (hatchery), Grays Harbor, Quinalt (hatchery), Queets, Hoh, and Quillayute coho.

Management Objectives

Management goals for Grays Harbor and Olympic Peninsula coho stocks included achieving natural spawning escapement objectives and treaty Indian allocation requirements. The Council's conservation objectives for stocks managed for natural production were based on maximum sustainable yield (MSY) spawner escapements established pursuant to the U.S. District Court order in *Hoh versus Baldrige*. The conservation objectives for the Queets, Hoh, and Quillayute rivers were developed as ranges intended to bracket estimates of MSY escapement. The range reflects the degree of uncertainty inherent by using the high estimate of recruits-per-spawner, and the low estimate of carrying capacity for the lower bound, and the low estimate of recruits-per-spawner with the high estimate of smolt carrying capacity for the upper end of the range. Annual targets for natural spawning escapement and total escapement were established by WDFW and treaty Indian tribes under the provisions of *U.S. versus Washington* and subsequent U.S. District Court orders. After the annual agreement was reached, ocean fishery escapement objectives were established for each river, or region of origin. The agreement includes provisions for treaty Indian allocation requirements and inside non-Indian fishery needs.

Regulations to Achieve Objectives

Washington coastal coho stocks contribute primarily to ocean fisheries off Washington and B.C. Those stocks did not play a primary role in 2007 Council area ocean fishery management because of impact constraints on Interior Fraser (Thompson River, B.C.) and LCN coho stocks, and treaty Indian/non-Indian in-river sharing of Columbia upriver coho. Overall harvest quotas were limited to levels well below those of the late 1980s and early 1990s. All non-Indian ocean coho fisheries north of Cape Falcon were mark-selective. Treaty Indian fisheries were not mark-selective.

Willapa Bay Coho

Inside Harvest

Historical terminal run size, harvest and escapement data for Willapa Bay coho are presented in Appendix B, Table B-24. The gillnet catch of coho in Willapa Bay in 2007 totaled 8,218 fish. Based on the preseason forecast for a terminal run of 51,200 fish, the scheduled commercial fisheries were expected to harvest approximately 16,152 total coho.

From July 1, 2007 through July 31, 2007, Willapa Bay (Marine Area 2-1) was open for recreational fishing, concurrent with the Ocean Marine Area 2 (ocean rules applied). August 1, 2007 through August 15, 2007 Willapa Bay was open to recreational fishing with a daily-bag-limit of six salmon with no more than two adults, and barbed hooks were allowed. August 16, 2007 through January 31, 2008, Willapa Bay was open to recreational fishing with a daily-bag-limit of six salmon, no more than three adults, of which only two could be Chinook. Chum retention was prohibited. Barbed hooks were allowed when fishing for salmon. Marine and freshwater recreational harvest estimates were not yet available for 2007. Expected harvest in recreational fisheries based on preseason forecast abundance was 1,376. Marine Area 2-1 and freshwater recreational harvest estimates for 2006 harvest estimates totaled 811 fish.

Freshwater recreational fisheries in the Willapa Bay watershed were open for salmon fishing from August 1, 2007 through January 31, 2008 with a daily-bag-limit of six salmon, no more than two adults, one of which may be a wild adult coho. Chum retention was prohibited.

Escapement and Management Performance

Willapa Bay coho were managed primarily for natural production. Estimates of natural spawning escapement for 2007 were not available. The most recent but still preliminary natural escapement estimate available was 14,413 in 2006. Escapement to Willapa Bay hatcheries in 2007 was estimated at 7,565 coho, which met the WDFW escapement objective of 6,100 spawners.

Grays Harbor Coho

Inside Harvest

Historical terminal run size, harvest and escapement data for Grays Harbor coho are presented in Appendix B, Table B-26. The run size forecast for Grays Harbor coho, after accounting for ocean fishery impacts, was 118,161 fish (53,555 wild and 64,606 hatchery). Nearly 10,500 coho (wild, hatchery, and net-pen origin) were harvested in treaty Indian and non-Indian gillnet fisheries. This included 8,839 coho in the Quinault Indian Nation fisheries, 1,687 in the non-Indian gillnet fishery, and small numbers in the Chehalis tribal fishery.

Recreational harvest estimates for 2007 were not available. Marine Area 2.2 was open from October 1 to October 31 for two salmon daily, one of which may be a wild coho and one of which may be a Chinook; from November 1 to November 30, the Area was open for two salmon daily, one of which may be a wild coho with no Chinook or chum retention. The Chehalis River and its tributaries downstream of the bridge crossing at the town of Porter were open for retention of up to two adult coho (regardless of mark status) from April 16 through July 31 and October 1 through November 30. The Chehalis River and its tributaries upstream of the bridge crossing at the town of Porter were open to retention of up to two adult coho (regardless of mark status) April 16 through July 31 and October 1 through November 30. In December 2007 and January and February 2008, openings varied by system, but coho harvest was limited to one unmarked coho in a two-adult coho bag limit or release wild adult coho in a two-adult coho bag limit. The Humptulips recreational fishery required the release of all wild adult coho (December 1, 2007 through January 31, 2008).

The Quinault Indian Nation operated two separately scheduled gillnet fisheries in the area of the Lower Humptulips and in the area of the Lower Chehalis, as described in Chapter II under the section labeled Grays Harbor Chinook, for both Chinook and coho, as well as chum. The expected coho fishery impacts were limited by the expected abundance and harvest of Chinook in those fisheries. The Humptulips area fishery harvested 3,717 coho, while the Chehalis area fishery harvested 5,122 coho. Harvest levels were about 66% of pre-season expected levels in both fisheries.

Escapement and Management Performance

Grays Harbor coho were managed for natural production with a spawning escapement goal of 35,400. Natural spawning escapement estimates for 2006 and 2007 were 14,401 and 23,662, respectively.

Quinault River Coho

Inside Harvest

Historical terminal run size, harvest, and escapement for Quinault River coho are presented in Appendix B, Table B-28. The treaty Indian gillnet fishery targeted hatchery Chinook and coho from early September through mid-November. A total of 11,770 coho were harvested by the gillnet fishery in 2007.

Escapement and Management Performance

Quinault River coho were managed for hatchery production. Escapement estimates for Quinault River coho in 2007 were not available. The Quinault National Fish Hatchery egg-take objectives for 2007 were achieved.

Queets River Coho

Inside Harvest

Historical terminal run size, harvest, and escapement for Queets River coho are presented in Appendix B, Table B-31. Queets River fisheries were managed under preseason agreement with WDFW based on preseason abundance estimates and planned Council ocean fisheries. The treaty Indian gillnet fishery was structured to target returning hatchery and wild coho during September and early October. The total harvest of fall coho in the gillnet fishery was 2,261 commercially landed fish, substantially below an expected catch of above 11,600. The gillnet harvest was comprised of an unknown mix of early-timed hatchery fish with a larger proportion of later-timed wild fish. Recreational fisheries operated with standard bag limits (no restriction on coho based on mark status) and schedules in the Queets, Clearwater, and Salmon Rivers. A 2007 catch estimate was not available.

Escapement and Management Performance

The preliminary natural coho spawner survey escapement estimate for 2007 was 5,272, below the escapement objective of 5,800 to 14,500 established for this stock. The preliminary 2006 estimate of natural coho escapement was 5,400. The inseason effort model suggests substantially reduced hatchery return run size of 1,714. Releases of supplemental coho were discontinued after 2004 so there were no returns of those fish in 2007.

Hoh River Coho

Inside Harvest

Historical terminal run size, catch, and escapement data for Hoh River coho are presented in Appendix B, Table B-34. The terminal run size of Hoh River wild coho was projected to be 4,678. The treaty gillnet fishery occurred from the week of May 7 to the week of August 27, as described in Chapter II under the section labeled Hoh River Chinook. The tribal fishery took approximately 1,764 coho, with 1,713 estimated to be wild, including dip-in wild fish. The non-Indian recreational fishery extended from September 1 through November 30, with the area below Willoughby Creek open and a daily-bag-limit of six salmon, two of which could be adults and no mark selective coho restriction. The portion of the river between Willoughby Creek and Morgan's Crossing opened October 16 to reduce impacts on spawning spring/summer Chinook in that reach. The river above Morgan's Crossing did not open for recreational salmon fishing. A catch estimate for the recreational fishery is not yet available.

Escapement and Management Performance

The preliminary spawning escapement estimate for coho in the Hoh River was 3,072, which was within the escapement goal range (2,000-5,000) established for this stock.

Quillayute River Coho

Inside Harvest

Historical terminal run size, catch, and escapement data for Quillayute River summer and fall coho are presented in Appendix B, Table B-37.

The recreational and tribal fisheries for coho were established by preseason agreement between Washington Department of Fish and Wildlife (WDFW) and the Quileute Tribe. A total of 1,430 (578 wild) summer coho were harvested in the Quileute Tribe's commercial and ceremonial and subsistence fisheries. An estimate of the 2007 recreational catch is not yet available.

The Quileute Tribal harvest of fall coho for 2007 was 9,377 (ceremonial and subsistence included). Tribal net fisheries harvested approximately 4,129 wild coho. An estimate of the 2007 recreational catch is not yet available.

WDFW reduced the impacts of the recreational fishery on wild summer and fall coho by requiring mark-selective fisheries for coho through October. The Quileute Tribe did not have a closure in their fishery this year, but as in past years, reduced their fishery to 29 hours per week during July and August.

Escapement and Management Performance

The summer coho run in the Quillayute is managed primarily for its hatchery component, which returns in August and September. The summer coho rack return was 4,778. This was well above the goal of 300. An additional 13 wild summer coho were collected as broodstock. The preliminary estimate for natural summer coho escapement was 792.

The preliminary 2007 escapement estimate for natural fall coho was 5,609, below the escapement goal of 6,300 to 15,800 established for this stock. An additional 20 wild fall coho were collected as broodstock. The hatchery rack return of 5,423 exceeded the goal of 600 adults.

PUGET SOUND COHO STOCKS

Puget Sound coho salmon stocks include natural and hatchery stocks originating from U.S. tributaries in Puget Sound and the eastern Strait of Juan de Fuca (east of Salt Creek, inclusive). The primary stocks in this group that are most pertinent to ocean salmon fishery management were eastern Strait of Juan de Fuca, Hood Canal, Skagit, Stillaguamish, Snohomish, and South Puget Sound (hatchery) coho.

Management Objectives

The Council's conservation objectives were based on the Puget Sound Salmon Management Plan, which defined management objectives and long-term goals for these stocks as developed by representatives from Federal, state, and tribal agencies. Conservation objectives for specific stocks were based on either maximum sustainable production for stocks managed primarily for natural production or on hatchery escapement needs for stocks managed for artificial production. The original conservation objectives were developed by a State/Tribal Management Plan Development Team following the Boldt Decision with the goal for natural spawning stocks defined as "the adult spawning population that will, on the average, maximize biomass of juvenile outmigrants subsequent to incubation and freshwater rearing under average

environmental conditions." The methodology used to develop the objectives was based on assessment of the quantity and quality of rearing habitat and the number of adult spawners required to fully seed the habitat. Some objectives have subsequently been modified by the U.S. District Court Fisheries Advisory Board and later determinations of the WDFW/Tribal Technical Committee. Annual escapement targets for these coho stocks were developed through procedures established in U.S. District Court. Puget Sound management procedures are outlined in a "Memorandum Adopting Salmon Management Plan" (*U.S. versus Washington*, 626 F. Supp. 1405 [1985]). A transition to exploitation rate management is currently under consideration by the involved managers.

The PSC adopted a management plan for coho salmon originating in Washington and southern B.C. river systems in 2002. The plan was directed at the conservation of key management units, four from southern B.C. (Interior Fraser, Lower Fraser, Strait of Georgia Mainland, Strait of Georgia Vancouver Island) and nine from Washington (Skagit, Stillaguamish, Snohomish, Hood Canal, Strait of Juan de Fuca, Quillayute, Hoh, Queets, and Grays Harbor). Under the plan, the United States and Canada were required to constrain total fishery exploitation rates to levels associated with the categorical status and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries were established through formulas specified in the plan. Categorical status was employed by the PST under the 2002 Coho Agreement to indicate general ranges of allowable total exploitation rates for U.S. and Canadian coho management units in 2007. Three categories were employed: low (total exploitation rate <20%), moderate (total exploitation rate 20%-40%), and abundant (total exploitation rate >40%). In 2007, the southern U.S. exploitation rate objectives for Skagit and Snohomish coho stocks were limited to no more than 35% and 40%, respectively. The pre-season predicted southern U.S. exploitation rates on these stocks were 34% (3.6% in Council area fisheries) and 39% (4.9% in Council area fisheries) respectively. Council area fisheries were constrained by other stocks in 2007, including Interior Fraser and LCN coho. Inside fisheries however, primarily in Puget Sound, were constrained to meet PSC objectives for Skagit and Snohomish coho stocks.

Regulations to Achieve Objectives

Puget Sound coho stocks contribute primarily to ocean fisheries off Washington and B.C. Those stocks did not play a primary role in 2007 ocean fishery management considerations, since management of impacts to Interior Fraser (Thompson River, B.C. Canada) and Columbia River stocks were more constraining. The mark-selective regulations in ocean and Puget Sound recreational fisheries served to increase harvest of marked hatchery fish while minimizing impacts on wild Puget Sound coho, LCN coho, OCN coho, and Interior Fraser coho.

Inside Harvest

Commercial inside harvest of Puget Sound coho was managed on the basis of six regional management units: Strait of Juan de Fuca, Nooksack-Samish, Skagit, Stillaguamish-Snohomish, South Puget Sound, and Hood Canal. Harvest of coho for each management unit is regulated according to the natural spawning escapement or hatchery program escapement goal for that unit. Commercial net and troll harvest (treaty Indian and non-Indian) for all coho stocks combined is presented in Appendix B, Table B-38. The 2007 total Puget Sound commercial catch of coho was 226,360 fish, compared to a catch of 302,490 coho in 2006. Non-Indian harvest was 19,321 coho, compared to a catch of 10,042 coho in 2006. Treaty Indian net and troll fisheries harvested 212,925 coho, compared to a catch of 292,448 coho in 2006.

Historical coho catches in the Puget Sound recreational fishery for the years from 1971 through 2007 are listed in Appendix B, Table B-39. Catch estimates for the 2007 Puget Sound recreational fishery were not available.

Escapement and Management Performance

No post season estimates of southern U.S. inside harvest impacts on coho stocks subject to the PSC coho management plan were available. Preliminary escapement information indicates natural Puget Sound coho escapements were generally higher in 2007 than the very low levels observed in 2006, and escapement levels for the hatchery programs were generally adequate, with a few exceptions.

BRITISH COLUMBIA COHO STOCKS

Management Objectives

The PSC adopted a management plan for coho salmon originating in Washington and southern B.C. river systems in 2002. The plan is directed at the conservation of key management units, four from southern B.C. (Interior Fraser, Lower Fraser, Strait of Georgia Mainland, Strait of Georgia Vancouver Island) and nine from Washington (Skagit, Stillaguamish, Snohomish, Hood Canal, Strait of Juan de Fuca, Quillayute, Hoh, Queets, and Grays Harbor). Under the plan, the United States and Canada are required to constrain total fishery exploitation rates to levels associated with the categorical status and target exploitation rates of the key management units as determined by domestic managers. Ceilings on exploitation rates by intercepting fisheries were established through formulas specified in the plan. Categorical status is employed by the PST under the 2002 Coho Agreement to indicate general ranges of allowable total exploitation rates for U.S. and Canadian coho management units. Three categories are employed: low (total exploitation rate <20%), moderate (total exploitation rate 20%-40%), and abundant (total exploitation rate >40%).

Regulations to Achieve Objectives

In 2007, the "low" status of Interior Fraser coho required the total exploitation rate on this stock by southern U.S. fisheries not to exceed 10.0%. This requirement constrained both Council area and inside fisheries. The pre-season expectation was that the total southern U.S. fishery exploitation rate on Interior Fraser coho would be 10.0% (4.1% in Council area fisheries). The mark-selective regulations in ocean and Puget Sound recreational fisheries served to increase harvest of marked hatchery fish while minimizing impacts on wild Interior Fraser coho.

Inside Harvest

Harvest of coho in inside waters affecting B.C. coho stocks occurred in Puget Sound fisheries, which were described in the previous section of this chapter.

Escapement and Management Performance

No post season estimates of southern U.S. inside harvest impacts on coho stocks subject to the PSC coho management plan were available. Preseason expectations were for an inside exploitation rate of 5.9% on Interior Fraser coho.

COASTWIDE GOAL ASSESSMENT SUMMARY

Conservation objective achievement assessments were not available for many coho stocks; however, those that were available all met their objectives except for Queets River coho and Quillayute fall coho. Skagit River coho spawning escapement estimates were not available but the preseason expectation was for a return less than the objective. OPI and Washington coastal areas generally experienced coho returns below the numbers forecasted.

A summary of 2007 performance for coho salmon by stock in relation to the Council's conservation objectives is presented in Table III-5.

TABLE III-1. Estimated returns to Oregon coastal streams and lakes in thousands of adult coho (SRS spawner accounting).
(Page 1 of 1)

Year	Returns to Hatcheries			Count at North Fork Umpqua Winchester Dam	Number of OCN Spawners ^{a/}			Inside Harvest Impacts ^{c/}	Ocean Escapement to Oregon Coast ^{a/}
	Private	Public	STEP ^{b/}		Lakes	Rivers	Total		
1970	-	36.2	-	0.2	20.5	51.2	71.7	39.8	147.9
1971	-	29.1	-	0.6	29.2	65.6	94.8	24.1	148.6
1972	-	12.9	-	0.3	10.0	24.1	34.1	16.6	63.9
1973	-	18.4	-	0.4	17.6	37.8	55.4	15.4	89.6
1974	-	35.1	-	0.4	6.4	28.1	34.5	13.5	83.5
1975	-	4.9	-	0.5	5.6	34.8	40.4	13.5	59.3
1976	-	38.7	-	0.3	1.5	39.2	40.7	19.6	99.3
1977	4.2	6.5	-	0.4	5.8	13.7	19.5	13.5	44.1
1978	12.3	5.6	-	0.5	1.6	18.2	19.8	4.5	42.7
1979	49.2	22.2	-	0.4	6.6	38.4	45.0	1.5	118.3
1980	38.7	21.9	-	0.2	4.7	25.6	30.3	6.3	97.4
1981	117.8	21.2	-	0.1	2.5	30.1	32.6	9.9	181.6
1982	184.7	14.8	-	2.7	7.9	68.3	76.2	14.7	293.1
1983	133.9	9.5	-	1.2	3.3	19.4	22.7	6.8	174.1
1984	115.4	28.6	-	3.2	14.7	59.7	74.4	17.4	239.0
1985	332.0	15.8	-	4.0	7.6	66.3	73.9	15.7	441.4
1986	453.7	35.8	2.5	9.6	11.8	58.2	70.0	30.3	601.9
1987	119.3	12.3	0.2	2.2	4.2	25.9	30.1	7.7	171.8
1988	116.1	33.7	1.2	1.2	5.8	51.0	56.8	13.3	222.3
1989	46.9	37.3	1.2	3.0	4.8	41.6	46.4	15.1	149.9
1990	35.6	15.4	1.6	2.3	4.4	16.5	20.9	9.5	85.3
1991	35.1	39.6	4.9	3.9	7.3	30.4	37.6	31.5	117.5
1992	-	23.3	0.6	5.0	2.0	40.2	42.2	18.7	89.8
1993	-	20.2	2.0	2.3	10.1	45.2	55.3	13.3	93.2
1994	-	23.4	1.8	2.0	5.8	38.3	44.2	2.4	73.7
1995	-	25.2	0.4	3.3	11.2	42.8	54.0	3.6	86.5
1996	-	23.8	1.0	6.3	13.5	60.5	74.0	4.0	109.1
1997	-	17.6	0.2	1.8	8.6	14.8	23.4	4.3	47.3
1998	-	15.2	0.2	4.6	11.1	20.6	31.8	5.2	56.9
1999	-	13.3	0.4	1.3	12.7	36.3	49.0	2.8	66.8
2000	-	15.0	0.5	9.3	12.7	55.9	68.7	4.5	97.9
2001	-	38.1	1.2	21.9	19.7	151.0	170.7	10.1	242.0
2002	-	30.9	2.6	7.4	22.2	238.4	260.6	8.1	309.5
2003	-	15.9	3.6	10.7	16.7	211.6	228.3	6.7	265.2
2004	-	13.2	0.8	7.2	18.7	151.2	169.9	6.3	197.3
2005	-	10.0	0.3	8.9	14.7	139.4	154.1	5.9	179.2
2006	-	9.8	0.1	7.0	24.4	104.5	128.8	2.2	148.0
2007 ^{d/}	-	3.6	0.0	2.7	8.9	43.0	51.9	1.3	59.5

a/ Does not include estimates for the southern OCN component (Rogue River). Spawner escapements to rivers prior to 1990 were estimated by a nonrandom standard index of streams north of the Rogue River. A total coastwide spawner escapement methodology based on SRS was initiated in 1990 and implemented concurrently with the standard index methodology. The SRS methodology indicated that actual escapements were less than estimated by the standard rivers index. The spawner index data for years prior to 1990 have been recalibrated in this table to be comparable with the SRS estimates.

b/ Oregon coastal Salmon Trout Enhancement Program (STEP) production from hatchery smolt rearing sites only.

c/ Freshwater sport catch from ODFW salmon/steelhead angler tag information and represents only those fish greater than 24 inches. Includes estimated mortality from hook-and-release.

d/ Preliminary.

TABLE III-2. Estimated weekly effort (in angler trips) and catches of Chinook and coho in the 2007 Buoy 10 recreational fisheries (all data are preliminary).^{a/} (Page 1 of 1)

Week Number	Ending Date of Period	Angler Trips	Catch		Catch Per Trip
			Chinook	Coho	
31	Aug.-5	449	0	4	0.01
32	Aug.-12	400	0	11	0.03
33	Aug.-19	2,065	0	91	0.04
34	Aug.-26	11,139	2,211	1,780	0.36
35	Sept.-2	14,437	1,374	3,516	0.34
36	Sept.-9	5,343	190	2,466	0.50
37	Sept.-16	1,484	1	255	0.17
38	Sept.-23	600	0	228	0.38
39	Sept.-30	147	0	5	0.03
40-43	Oct.-28	0	0	0	NA
Total		36,064	3,776	8,356	0.34

a/ Includes boat-based and shore-based fisheries from the new (2000) upstream boundary at the Tongue Point/Rocky Point line downstream to the Buoy 10 line including Clatsop Spit, the South Jetty of the Columbia River, and the North Jetty of the Columbia River after the ocean closed. Fishery was open August 1-September 28 for Chinook and adipose fin-clipped coho, with the daily-bag-limit of two adult salmon, only one of which may be a Chinook, except Chinook retention was prohibited from August 1-21 and September 4-28. From September 29-December 31 the daily-bag-limit of two adult salmon, only one of which may be a Chinook, was modified to allow the retention of two additional fin-clipped coho.

TABLE III-3. Oregon production index (OPI) area coho harvest impacts, spawning, abundance, and exploitation rate estimates by SRS accounting in thousands of fish.^{a/} (Page 1 of 1)

Year or Avg.	Oregon and California Coastal Returns								Ocean Exploitation Rate Based on OPI Abundance ^{d/}	OCN Exploitation Rate Based on Postseason FRAM
	Ocean Fisheries ^{b/}		Hatcheries and Freshwater		Private Hatcheries	Columbia River Returns	Abundance	OCN Spawners		
	Troll	Sport	Harvest ^{c/}							
1970-1975	1,629.6	558.4	45.8	55.2	-	460.4	2,749.3	0.80	-	
1976	2,936.1	977.7	62.6	40.7	-	337.0	4,347.6	0.90	-	
1977	664.4	412.1	21.4	19.5	4.2	93.8	1,220.4	0.89	-	
1978	1,104.2	524.6	12.6	19.8	12.3	307.5	1,977.4	0.83	-	
1979	1,056.6	334.4	27.4	45.0	49.2	276.5	1,789.5	0.79	-	
1980	506.9	526.4	32.1	30.3	38.7	301.6	1,436.4	0.73	-	
1981	830.9	339.9	34.1	32.6	117.8	170.2	1,555.0	0.81	-	
1982	740.9	300.4	37.1	76.2	184.7	453.1	1,763.4	0.62	-	
1983	429.6	275.0	18.2	22.8	133.9	111.2	1,070.0	0.79	-	
1984	95.8	174.2	51.2	74.5	115.4	425.9	881.5	0.32	-	
1985	166.4	280.4	45.4	73.9	332.0	367.2	1,373.4	0.43	-	
1986	643.5	320.6	81.8	70.0	453.7	1,549.1	3,026.7	0.34	-	
1987	469.1	296.2	45.3	30.1	119.3	316.6	1,377.9	0.60	-	
1988	844.7	297.2	62.3	56.8	116.1	670.8	1,989.2	0.56	-	
1989	646.9	425.5	62.3	46.4	46.9	712.8	1,871.2	0.55	-	
1990	277.6	357.1	30.6	24.3	35.6	196.7	1,128.5	0.69	-	
1991	450.6	469.9	84.0	38.6	35.1	934.3	1,823.2	0.45	-	
1992	67.5	256.5	52.8	44.4	-	215.9	610.0	0.51	-	
1993	13.2	140.8	40.6	55.7	-	113.9	342.1	0.42	-	
1994	2.7	3.0	30.0	49.6	-	168.9	250.5	0.02	0.07	
1995	5.4	43.5	38.6	57.7	-	74.1	215.9	0.23	0.12	
1996	7.0	31.8	47.9	78.6	-	113.0	297.3	0.15	0.08	
1997	5.5	22.4	27.2	31.7	-	148.1	204.6	0.12	0.12	
1998	3.5	12.8	29.7	34.1	-	168.7	265.2	0.06	0.08	
1999	3.6	36.5	20.9	50.4	-	274.1	414.0	0.12	0.09	
2000	25.9	74.6	32.9	79.6	-	547.6	901.0	0.13	0.07	
2001	38.1	216.8	82.5	182.9	-	1,108.3	1,438.6	0.16	0.07	
2002	14.9	118.7	56.3	268.4	-	499.9	990.5	0.14	0.12	
2003	28.8	252.4	47.8	235.0	-	677.3	1,183.6	0.23	0.14	
2004	26.2	159.4	38.7	194.4	-	442.5	826.8	0.25	0.15	
2005	10.5	58.2	42.8	164.1	-	342.0	592.1	0.12	0.11	
2006	4.5	47.5	31.7	132.8	-	383.0	557.1	0.06	0.11	
2007 ^{e/}	26.7	128.5	11.7	57.1	-	318.6	536.6	0.31	0.11	

a/ The OPI area includes ocean and inside harvest impacts and escapement to streams and lakes south of Leadbetter Pt., Washington.

b/ Includes estimated nonretention mortality: troll fishery--hook-and-release mortality for 1982-2005 and drop-off mortality for all years; sport fishery--hook-and-release mortality for 1994-2005 and drop-off mortality for all years.

c/ Includes returns from Salmon-Trout Enhancement Program (STEP) smolt releases.

d/ Ocean fishery impacts on private hatchery stock and returns to private hatcheries are excluded in calculating the OPI area stock aggregate ocean exploitation rate index.

e/ Preliminary.

TABLE III-4. OCN adult coho salmon conservation objective, fishery impacts, and spawner escapement, based on stratified random survey (SRS) methodology. (Page 1 of 1)

Year	Fishery Impact (Total Marine and Freshwater Exploitation Rate)			Adjusted SRS Adult Coho Spawner Population Estimates in Thousands of Spawners by Stock Component ^{a/}					Adult Coho Spawners Per Spawner Habitat Mile				
	Conservation Objective ^{b/}	Preseason Projection	Postseason Estimate ^{c/}	North		South		Coastwide	North		South		Coastwide Average
				Northern ^{d/}	Central ^{e/}	Central ^{f/}	Southern ^{g/}		Northern ^{d/}	Central ^{e/}	Central ^{f/}	Southern ^{g/}	
1990	-	-	-	2.2	5.6	13.5	3.1	24.3	2	5	8	7	6
1991	-	0.460	0.454	9.3	6.7	21.6	1.0	38.6	10	6	13	3	9
1992	-	0.420	0.511	2.4	15.4	24.4	2.2	44.4	3	13	15	5	11
1993	-	0.260	0.423	4.5	7.8	43.1	0.4 ^{h/}	55.7	5	7	27	1 ^{h/}	14
1994	≤0.20	0.111	0.068	3.5	9.8	30.9	5.4	49.6	4	8	19	13	12
1995	≤0.20	0.118	0.124	3.9	13.6	36.5	3.8	57.7	4	12	22	9	14
1996	≤0.20	0.125	0.083	3.3	18.1	52.6	4.6	78.6	4	16	32	11	19
1997	≤0.20	0.110	0.124	2.1	2.8	18.4	8.3	31.7	2	2	11	20	8
1998	≤0.13	0.119	0.078	2.6	3.3	25.9	2.3	34.1	3	3	16	6	8
1999	≤0.15	0.087	0.087	8.9	11.8	28.3	1.4	50.4	10	10	17	4	12
2000	≤0.15	0.082	0.073	17.9	14.3	36.5	11.0	79.6	20	12	23	27	19
2001	≤0.08	0.074	0.070	33.5	25.2	112.0	12.2	182.9	37	22	69	30	45
2002	≤0.15	0.123	0.120	52.5	104.0	104.1	7.8	268.4	58	89	64	19	66
2003	≤0.15	0.144	0.140	59.6	68.9	99.8	6.8	235.0	66	59	62	16	57
2004	≤0.15	0.147	0.150	33.1	40.4	96.4	24.5	194.4	37	35	59	60	47
2005	≤0.15 ^{i/}	0.111	0.110	16.5	51.4	86.3	10.0	164.1	18	44	53	24	40
2006	≤0.15 ^{i/}	0.096	0.110	24.1	21.2	83.5	3.9	132.8	27	18	51	10	32
2007 ^{j/}	≤0.20	0.113	0.106	15.1	10.0	26.8	5.2	57.1	17	9	17	13	14

a/ A spawner escapement methodology study based on SRS has been in effect since 1990 in which coho salmon population estimates have been made for Oregon coastal river systems from the Coquille River and north. Spawner population estimates include an adjustment for observation error.

b/ Prior to 1994, the conservation objective was expressed in terms of the total escapement of OCN spawners in index numbers rather than as an exploitation rate. The index escapement objectives from 1981 through 1993 are provided in Table III-2 of the Review of 1998 Ocean Salmon Fisheries and Table 1 of Amendment 11. From 1994 through 1997, Amendment 11 specified that at low stock sizes, only incidental harvest of OCN coho could occur and that impacts could not exceed 20%. Beginning in 1998, the OCN conservation objective has been as specified in Amendment 13 which is also the basis for the NMFS jeopardy standards under the Endangered Species Act listing.

c/ From the coho FRAM, except the estimates prior to 1994 represent the OPI composite exploitation rate for hatchery and natural stocks.

d/ Estimate based on 899 miles of spawner habitat within Nehalem, Tillamook, and Nestucca Rivers and other direct ocean tributaries from Necanicum River through Neskowin Creek.

e/ Estimate based on 1,163 miles of spawner habitat within Siletz, Yaquina, Alsea, and Siuslaw Rivers and other direct ocean tributaries from the Salmon through Siuslaw Rivers.

f/ Estimate based on 1,622 miles of spawner habitat within Umpqua, Coos, and Coquille Rivers. Also includes spawners using tributaries to Siltcoos, Tahkenitch, and Tenmile Lakes.

g/ Estimate based on a mark-recapture methodology and 410 miles of spawner habitat within the Rogue River.

h/ Unreliable estimate.

i/ Preliminary.

j/ The Salmon FMP specified an allowable marine and freshwater exploitation rate of 20%, however, the OCN workgroup matrix specified 15% and the Council chose to manage at the more conservative level for 2005 and 2006.

TABLE III-5. Performance of coho salmon stocks in relation to 2007 conservation objectives (preliminary data). (Page 1 of 2)

System and Stock	2007 FMP Conservation Objective	Achievement
OPI Area Coho		
(Columbia River and coastal stocks south of Leadbetter Point)	Natural spawner escapement objectives as provided below; meet hatchery egg-take goals; meet treaty Indian obligations.	Hatchery egg-take goals achieved. No information available on catch allocation.
Northern California (Threatened) and CCC (Threatened)	No directed coho fisheries or retention of coho south of Humbug Mt. Marine exploitation rate $\leq 13.0\%$ as indicated by R/K hatchery stocks. Council adopted a projected exploitation rate on R/K hatchery coho of 5.8%.	No directed coho fisheries or retention of coho south of Humbug Mt. Postseason marine exploitation rate estimate of 5.4% was below 13.0% maximum.
OCN	Combined marine and freshwater exploitation rate $\leq 20.0\%$. Council adopted a projected exploitation rate on OCN of 11.3%	Postseason marine and freshwater exploitation rate estimate of 10.6% was below 20.0% maximum.
Columbia River Natural (Threatened)	Combined marine and mainstem Columbia River exploitation rate $\leq 20\%$. Council adopted management measures resulted in a projected marine exploitation rate of 13.3%.	Postseason marine and mainstem Columbia River exploitation rate estimate of 18.6% was below 20.0% maximum; the 11.9% marine exploitation rate was below preseason expectation.
Washington Coast Coho		
	Natural spawner escapement objectives as provided below and in state/tribal agreements; meet hatchery egg-take goals; meet treaty Indian obligations.	Hatchery egg-take goals achieved. No information available on catch allocation.
Grays Harbor	35,400 natural adult spawners.	Preliminary estimate of 23,662 fails to meet the escapement floor.
Queets	5,800 to 14,500 natural adult spawners.	Preliminary estimate of 5,272 fails to meet the escapement floor.
Hoh	2,000 to 5,000 natural adult spawners.	Preliminary estimate of 3,072 is within the goal range.
Quillayute Fall	6,300 to 15,800 natural adult spawners.	Preliminary estimate of 5,609 fails to meet the escapement floor.

TABLE III-5. Performance of coho salmon stocks in relation to 2007 conservation objectives (preliminary data).
 (Page 2 of 2)

System and Stock	2007 FMP Conservation Objective	Achievement
Puget Sound Coho	Natural spawner escapement objectives as provided below and in state/tribal agreements; meet hatchery egg-take goals; meet treaty Indian obligations and inside non-Indian fishery needs for six management units.	Data not available for 2007 natural spawner escapements. Hatchery egg-take goals likely will be met.
Strait of Juan de Fuca	≤40% total exploitation rate. 12,800 adult spawners.	Preseason expected ocean escapement of 26,600 adult fish for eastern and western Strait of Juan de Fuca combined and a 12% total exploitation rate.
Hood Canal	≤65% total exploitation rate. 21,500 natural adult spawners.	Preseason expected ocean escapement of 30,900 adult fish and a 46% total exploitation rate.
Skagit	≤35% total exploitation rate. 30,000 natural adult spawners.	Preseason expected ocean escapement of 21,700 adult fish and a 34% total exploitation rate.
Stillaguamish	≤50% total exploitation rate. 17,000 natural adult spawners.	Preseason expected ocean escapement of 52,000 adult fish. 39% total exploitation rate.
Snohomish	≤40% total exploitation rate. 70,000 natural adult spawners.	Preseason expected ocean escapement of 70,100 adult fish and a 39% total exploitation rate.




Figure III-1. Oregon Production Index (OPI) area coho abundance estimates by stratified random surveys (SRS) accounting methods (1970-2007).




Figure III-2. Oregon coastal natural (OCN) adult coho spawners per habitat mile by coastal region based on SRS accounting methods, 1990-2007.

Page Left Intentionally Blank